



Research article

Phytochemical screening and antibacterial activities of *Tectona grandis* L. f. (Teak) leaves on microorganisms isolated from decayed food samples

O. T. Ogunmefun*, E. A. Ekundayo, F. C. Akharaiyi and D. Ewthenodere

Department of Biological Sciences, Afe Babalola University, Ado-Ekiti, Nigeria

*Corresponding Author: yinkatayo_08@yahoo.com

[Accepted: 19 September 2017]

Abstract: Bacteria were isolated from decayed food samples (tomatoes, cooked beans and rice) collected from Afe Babalola University, Ado Ekiti (ABUAD) cafeteria and characterized. Some of these isolated microorganisms could pose serious harm to humans including animals and they are normally treated with commercial antibiotics. However, the majority of bacteria are resistant to many antibiotics therefore, the use of plant extracts with therapeutic potential against resistant bacteria is necessary. In this investigation, eight bacteria were isolated from decayed food samples. The bacterial isolates were identified as *Bacillus cereus* and *B. siamensis* from rice sample; *Klebsiella oxytoxa*, *Salimicrobium halophilium* and *Nocardia brasiliensis* from beans sample; *Bacillus subtilis*, *Enterobacter taylore*, and *Brevibacillus agri* from tomatoes. The leaf samples of *Tectona grandis* were screened qualitatively and quantitatively for the phytochemicals while the crude methanol and chloroform extracts were used as antimicrobial agents against the isolated microorganisms. Alkaloids, carotenoids and tannins were present in large amount. The bacterial isolates were more susceptible to commercial antibiotics than that of methanol extracts of *T. grandis*. The methanol extracts of *T. grandis* have a higher antimicrobial activity than the chloroform extracts.

Keywords: Chloroform extracts - Food samples - Methanol extracts - Microorganisms - *Tectona grandis*.

[Cite as: Ogunmefun OT, Ekundayo EA, Akharaiyi FC & Ewthenodere D (2017) Phytochemical screening and antibacterial activities of *Tectona grandis* L. f. (Teak) leaves on microorganisms isolated from decayed food samples. *Tropical Plant Research* 4(3): 376–382]

INTRODUCTION

Food decay is any undesirable change in food that makes the food to lose its aesthetic value and becomes unacceptable to humans for consumption. Food is also said to be decayed when the original nutritional value, texture and flavor are altered through microbial activities. Various factors could be responsible for the deterioration of various foods (Istifanus *et al.* 2014). Consumption of this decayed or contaminated food results in food borne disease such as diarrhoea, gastroenteritis, respiratory infections and meningitis, if ingested (Mead *et al.* 1999, Humphrey *et al.* 2007, Barth *et al.* 2009). Also, toxins from these microorganisms can be found in contaminated food products and are pathogenic leading to disease (Edwin *et al.* 2013). Although, bacteria are the main and important source of food spoilage, but other microbes such as fungi, viruses and protozoa have also been implicated (Kumar *et al.* 2011). The most common pathogens causing serious infection in food include *Staphylococcus aureus*, *Pseudomonas* sp., *Streptococcus* sp., *Proteus* sp., *Clostridium* sp. and Coliforms (Lawrence *et al.* 1999). High pH (4.9–6.5), water and nutrient contents enhance microbial growth such as bacteria and fungi, which degrade the nutrients through enzymes production (Trias *et al.* 2008, Matthew 2011, Ogunbanwo *et al.* 2014). These also heighten spoilage susceptibility thereby reducing the nutritional and market values (Bello *et al.* 2016). Most of the pathogenic microorganisms causing diseases have developed resistance to modern antibiotics (Sharma *et al.* 2012). Bacteria generally have the genetic ability to transmit diseases and acquire resistance to drugs used as therapeutic agents (Cohen *et al.* 1992). Therefore, there is needed to look for new antimicrobials of plant origin.

Medicinal plants are gaining popularity in usage due to a large number of people in search of health remedies with little or no side effect which is the problem of most chemically synthesized drugs (Prance 1991, Bajpai *et al.* 2016). Considerable attention is presently given to the use of eco-friendly and bio-friendly products from plant origin for the prevention and cure of human and animal health challenges (diseases) (Gijtenbeek *et al.* 1999, Johnson & William 2002). Plants based antimicrobials represent a vast untapped source of medicines and further exploration of plant antimicrobials needs to occur. Antimicrobials of plant origin have enormous therapeutic potentials (Evans & Turnbull 2004). The majority of bacteria are resistant to many antibiotics therefore, the use of plant extracts against resistant bacteria leads to new choice for the treatment of infectious diseases (Purushotham *et al.* 2010).

Tectona grandis L. f. (Teak) is a tropical tree species distributed naturally in countries including India, Myanmar, Thailand, Myanmar, Nigeria and other tropical countries including Indonesia (Mahesh *et al.* 2016). Teak is one of the world's premier hardwood tree species, highly famous for its quality, profile and durability of timber (Sumthong *et al.* 2008). The generic name of *T. grandis* comes from 'Tekka', which is the Malabar name while the specific name, 'grandis' is a Latin word for 'large' or 'great'. *Tectona grandis* is a large, deciduous tree reaching over 30 m in height in favourable conditions (Orwa *et al.* 2009). "Fruit is a drupe with 4 chambers; round, hard and woody, enclosed in an inflated, bladder-like covering; pale green at first, then brown at maturity" (Orwa *et al.* 2009). It occurs naturally in various types of tropical deciduous forests. In seasonal climates, *T. grandis* is deciduous while trees grown in non-seasonal climates are semi-deciduous. It is often a dominant member of a mixed deciduous forest, where its main associates are *Xylocarpus* spp., *Azadirachta indica* (Kurz) Craib, *Terminalia* spp. and *Lagerstroemia* spp. *Tectona grandis* generally occurs scattered but can form almost pure stands under favourable conditions. Young plants show a remarkable capability to recover after fire (Orwa *et al.* 2009).

Diverse anti-microbial and substances that can serve as insect control agents can be produced by higher tropical plants (Nakamura *et al.* 1991, Downum *et al.* 1993, Lis-Balchin *et al.* 1996). Substances such as flavonoids, alkaloids, terpenoids are the secondary metabolites produced by the plants as a chemical defense against pests and diseases attacks. It is estimated only 10% of these tropical plants have been investigated for their pesticidal activity. Teak is one of these plants which produce secondary metabolic products containing phenolic compounds. Astiti *et al.* (1998) found that the water extract of teak leaf obviously inhibited the growth of *Monilia* species, the causative agent of wood decay. The sporulation of *Alternaria cajani* and *Helminthosporium* sp. was inhibited by the *T. grandis* leaf methanol crude extract at concentration of 5 mg.ml⁻¹ (Shalini & Srivastava 2008). The frontal leaves of the plant are widely used in folklore for the treatment of various kinds of wounds, especially burn wounds. They are also useful to treat haemostatic, depurative, anti-inflammatory and vulnerary and also useful in inflammation, leprosy, skin disease, pruritus, stomatitis, indolent ulcer, haemorrhages and haemopstysis (Purushotham *et al.* 2010). The presence of high amounts of tannins in the *T. grandis* leaves may be responsible for the pro-healing action of the extract and its antioxidant property is attributed to it (Mrityunjoy *et al.* 2007). All these studies clearly showed the antibacterial potential of *T. grandis*, thus the present study was designed to find out the effect of its leaves on the microorganisms isolated from decayed food samples.

MATERIALS AND METHODS

Collection of food samples

Food samples (fresh tomatoes, cooked beans, cooked rice and fresh pepper) were collected from Afe Babalola University, Ado-Ekiti (ABUAD) cafeteria/kitchen environment. These samples were allowed to decay in ABUAD Microbiology Department laboratory.

Collection of plant sample

The leaves of *Tectona grandis* L. f. were obtained from ABUAD surroundings and were identified by Dr. (Mrs.) O. T. Ogunmefun, a botanist in the Department of Biological Sciences of Afe Babalola University Ado - Ekiti, Ekiti State, Nigeria.

Preparation of extracts

The collected plant materials were air dried at room temperature, ground with blender after which dry mass of 887.2 g of the leaf sample was obtained (Purushotham & Sankar 2013). A 350 g each of powdered leaf sample was soaked in 700 ml of chloroform and methanol separately using cold extraction method for 72 hours.

The extracts were separated from the solvents using a rotary evaporator. The extracts were concentrated on water bath at a low temperature of 40°C (Ajaiyeoba *et al.* 2001, Ogbale *et al.* 2013).

Phytochemical screening of *Tectona grandis*

The powdered leaves of *T. grandis* were subjected to both qualitative and quantitative phytochemical tests for plant secondary metabolites namely alkaloids, anthraquinones, tannins, saponins, cardiac glycosides, flavonoids, carotenoids and phenols according to the method described by Harborne (1998) and Trease & Evans (2004).

Isolation, characterization and identification of microbial species from the decayed food samples

Five-fold serial dilution was carried out on the food samples. An aliquot (1 ml) of the fifth dilution was poured on nutrient agar and then incubated at 37°C for 24 hours. Colonies of bacterial isolates were selected from each plate respectively and purified by sub-culturing on nutrient agar plates. Bacterial isolates were preserved in slants at 4°C for temporary storage and were then identified both morphologically and biochemically (Prescott *et al.* 2002).

Determination of the antibacterial activities of *Tectona grandis* using agar well diffusion technique

A sterile swab stick was placed into the broth bacterial culture corresponding to MacFarland standard of a specific organism and then spread on already prepared sterile nutrient agar plate. This was repeated for all the isolates in duplicates. The plates were allowed to dry for approximately 5 minutes. The wells were then filled with dilutions of 50 mg.ml⁻¹, 100 mg.ml⁻¹, 150 mg.ml⁻¹, 200 mg.ml⁻¹, 250 mg.ml⁻¹, 300 mg.ml⁻¹, 350 mg.ml⁻¹ and 400 mg.ml⁻¹ with *T. grandis* extract reconstituted with 10% Dimethyl sulphoxide (DMSO) and sterilized by 0.45 µm membrane filter. The plates were incubated 37°C for 24 hours. The diameters of the inhibitory zones were measured in millimeters. The DMSO was used as a negative control while streptomycin was used as the positive control.

RESULTS

In this study, seven bacterial species were obtained from all the decayed food samples based on their morphological and biochemical characteristics. Five of the isolates were Gram positive while the remaining two isolates were Gram negative. The isolates were identified as *Bacillus cereus*, *B. subtilis*, *B. siamensis*, *Brevibacillus agri*, *Salimicrobium halophilum*, *Klebsiella oxytoxa*, *Norcadia brasiliensis* and *Enterobacter taylore*. The frequency of occurrence of the isolate is presented in table 1.

Table 1. Frequency of occurrence of bacterial isolates from decayed food samples.

Bacterial isolates	Food samples		
	Beans	Rice	Tomatoes
<i>Bacillus cereus</i>	-	+	-
<i>Bacillus siamensis</i>	-	+	+
<i>Bacillus subtilis</i>	-	-	+
<i>Klebsiella oxytoxa</i>	+	-	+
<i>Salimicrobium halophilum</i>	+	-	-
<i>Enterobacter taylore</i>	-	-	+
<i>Norcadia brasiliensis</i>	+	-	-
<i>Brevibacillus agri</i>	-	-	+

Note: +, Present; -, Absent.

Saponins, alkaloids, tannins among others were present while cardiac glycoside was not observed in the qualitative screening. However, the quantitative analysis showed that 20 mg per 100 g of cardiac glycoside was present. Alkaloids were most abundant followed by tannins and carotenoids respectively. The detail results of the qualitative and quantitative phytochemicals of *T. grandis* are presented in tables 2 and 3.

The leaf extracts obtained with chloroform and methanol were used as antibacterial agents against the bacterial isolates. The results showed that the extracts exhibited weak antibacterial activities on the isolates (Table 4). The antibacterial activities of streptomycin used as positive control ranged from 11.00–30.00 mm respectively and were without a doubt effective against food pathogens than the extract. All the bacterial isolates were resistant to DMSO used as negative control (Fig. 1). At concentrations of 50, 100, 150 and 200 mg.ml⁻¹, the extracts were not effective but at concentrations of 250, 300, 350 and 400 mg.ml⁻¹ there were clear zones of inhibition (Table 4).

Table 2. Qualitative phytochemical analysis of *Tectona grandis* leaves.

Parameter	Observation
Alkaloids	+++
Flavonoids	++
Tannins	+++
Saponins	++
Cardiac Glycosides	-
Total Phenolics	++
Anthraquinones	+
Carotenoids	++

Note: +++, Most abundant; ++, Abundant; +, Present; -, Absent.

Table 3. Quantitative phytochemical analysis of *Tectona grandis* leaves.

Phytochemical constituents	Amount of Phytochemical constituents
Alkaloids (mg per 100 g)	1778.30
Flavonoids (mg per 100 g)	666.60
Tannins (mg per 100 g)	1583.00
Saponins (mg per 100 g)	406.60
Cardiac Glycosides (mg per 100 g)	20.00
Total Phenolics (GAE per g)	69.50
Carotenoids (μg per 100 g)	1268.33




Figure 1. Comparative antibacterial activities of methanol extract of *Tectona grandis* and Streptomycin at 37°C on nutrient agar after 24 hours of incubation. [R, Rice; B, Beans; T, Tomatoes]

DISCUSSIONS

Different spoilage bacteria including fungi grow well at room temperature. A large number of microorganisms and their waste products cause the objectionable changes in odor, taste and texture in food samples (Kumar *et al.* 2011). The presence of these bacterial isolates may not be unconnected with the high moisture contents of the food substances which aided the spoilage as opined by Akharaiyi & Adeyanju (2016). *Bacillus cereus* has been associated with the production of toxins, diarrheal and emetic in food, which causes food poisoning. It is found in dust, soil and raw food and can survive under normal cooking conditions as produces heat resistant spore (Rajkowski & Bennett 2003). Bello *et al.* (2016) observed the prominence of *B. subtilis* in spoilage of tomato fruits. Ghosh (2009) also suggested that the prevalence of microbial contamination could be aggravated by poor sanitation including cross-contamination. Biological contaminant of bacterial origin represents a major cause of foodborne illnesses (Edema *et al.* 2005). The presence of *Bacillus subtilis* has been correlated with food poisoning. *Klebsiella* spp. and *Enterobacter* spp. call for concern as these organisms

are frequently associated with poor sanitary practices and could indicate the danger of possible food borne infection (Oranusi *et al.* 2007, Eni *et al.* 2010). *Klebsiella* spp. has been isolated in food such as fruits and vegetables, meat, milk and dairy products, salads, and drinking water, coming from the general environments of soil, dust, air and water and from social environments.

Table 4. Antibacterial activity of methanol extract of *Tectona grandis* leaves.

Bacterial isolates	Concentration of crude extract (mg.ml ⁻¹)							
	50	100	150	200	250	300	350	400
P3B3	R	R	R	R	10.00	10.00	11.00	11.50
P2B1	R	R	13.00	13.00	15.00	15.00	18.00	18.00
P3B3	R	R	R	2.00	4.00	6.00	11.00	12.00
P2B2	R	R	R	R	R	R	14.00	14.00
P1B1	R	R	R	R	R	R	R	R
RIB1	R	R	R	16.00	11.00	15.00	15.00	15.50
RB1	R	R	R	R	R	R	R	R
R1B2	R	R	R	R	R	R	R	R
R1B1a	R	R	R	R	R	R	R	R
T1B2	R	R	10.50	13.00	14.50	1.50	14.00	15.00
T3B1	R	R	R	12.00	12.50	13.50	15.00	18.50
T2B1	1.00	2.00	2.00	4.00	4.00	5.00	6.00	7.00
T1B1	R	R	3.00	4.00	4.00	6.00	6.00	8.00
T3B2	R	R	2.00	3.00	5.00	5.00	6.00	10.00

Note: **PB**, Isolates obtained from beans; **RB**, Isolates obtained from rice; **TB**, Isolates obtained from tomatoes; **B**, Bacteria; **R**, Resistance.

Higher tropical plants can produce diverse antimicrobial and anti-insect substances (Nakamura *et al.* 1991, Downum *et al.* 1993, Lis-Balchin *et al.* 1996). Substances such as flavonoids, alkaloids and terpenoids are secondary metabolites produced by plants as a chemical defence against pests and diseases' attacks. Tannins are astringent in nature and are used for treating intestinal disorders such as diarrhea and dysentery (Dharmananda 2003); they are referred to as natural antibiotics from plants.

The water extract of *T. grandis* leaf conspicuously inhibited the growth of *Monilia* sp., which is a causal organism attributing in wood decay (Astiti 1998). The methanol crude extract of *T. grandis* leaves at 5 mg.ml⁻¹ concentration inhibited the sporulation of *Alternaria cajani* and *Helminthosporium* sp. by 86.6 and 90.0% respectively (Shalini & Srivastava 2008).

CONCLUSION

The methanol extract of *Tectona grandis* leaf significantly inhibited bacterial growth. Chloroform extract was least effective on bacterial isolates as compared to methanol extracts. Teak extracts can be considered as one of the alternatives to chemical food preservatives and for controlling food decay caused by these isolates. A further study is needed to isolate and identify the active compounds that are responsible for the antimicrobial potency of *T. grandis*.

ACKNOWLEDGEMENT

All the authors are thankful to the Department of Biological Sciences, Afe Babalola University, Ado-Ekiti, Nigeria for providing necessary support. There is no conflict of interest among authors.

REFERENCES

- Ajaiyeoba EO, Onocha PA & Olanrewaju OT (2001) *In vitro* anthelmintic properties of *Buchholzia coriacea* and *Gynandropsis gynandra* extract. *Pharmaceutical Biology* 39: 217–220.
- Akharaiyi FC & Adeyanju A (2016) Implication of human handling on packaged sausage rolls during sale. *Annals of University Dunarea de Jos of Galati Fascicle VI-Food Technology* 40: 90–103.
- Astiti NPA (1998) Effectiveness water extract of the leaf of *Tectona grandis* L. against the growth of *Monilia* sp. *Journal of Biology Udayana* 2: 1–12.
- Bajpai O, Pandey J & Chaudhary LB (2016) Ethnomedicinal Uses of Tree Species by Tharu Tribes in the Himalayan Terai Region of India. *Research Journal of Medicinal Plant* 10(1): 19–41.
- Barth M, Hankinson TR, Zhuang H & Breidt F (2009) Microbiological spoilage of fruits and vegetables. In: Sperber WH & Doyle MP (eds) *Compendium of the Microbiological Spoilage of Foods and Beverages*, www.tropicalplantresearch.com

- Food Microbiology and Food Safety*. Springer Science Business Media, LLC, pp. 135–183.
- Bello OB, Bello IS, Aminu D, Olawuyi OJ, Afolabi-Balogun NB, Lawal AO, Azeez AH & Habib U (2016) Antibiotic sensitivity of bacterial and fungal isolates from tomato (*Solanum lycopersicum* L.) fruit. *Tropical Plant Research* 3(1): 112–119.
- Cohen B, McGuffin ME, Pfeifle C, Segal, D. & Cohen, SM (1992) Apetorus, a gene required for Imaginary disc development in *Drosophila* encodes a member of the LIM family of developmental regulatory proteins. *Genes and Development* 6(5): 715–729.
- Dharmananda S (2003) *Gall nuts and the uses of tannins in Chinese medicine*. In: Proceedings of Institute for Traditional Medicine, Portland, Oregon.
- Downum KR, Romeo JT & Sataford HA (1993) *Phytochemistry, 28, Potential of tropical plants*. Academic Press, New York, 105 p.
- Edema MO, Atayese AO, Idowu AO (2005) *Microbiological Quality of microwave processed Foods*. In: the Book of Abstract of the 29th Annual Conference & General Meeting on Microbes As Agents of Sustainable Development, organized by Nigerian Society for Microbiology (NSM), UNAAB, Abeokuta from 6–10th Nov. p. 17.
- Edwin A, Bergin L, Ilsedore C, Uma G, Ke Z, Gaeoffrey A, Joel DG, Sean MA, Neal JE II, Thomas H, Karin O, Khus P, Chunhua Q, Colette S & Van Dishoeck EF (2013) An old disk that can still form a planetary system. *Nature* 493: 644.
- Eni AO, Oluwawemitan IA & Oranusi US (2010) Microbial quality of fruits and vegetables sold in Sango Ota, Nigeria. *African Journal of Food Science* 4(5): 291–296.
- Evans J & Turnbull JW (2004) *Plantation forestry in the Tropics*. Oxford University Press, New York, NY, USA, pp. 482.
- Ghosh A (2009) Identification of microorganisms responsible for spoilage of tomato (*Lycopersicum esculentum*) fruit. *Journal of Phytology* 1(6): 414–416.
- Gijtenbeek JMM, Vanden Bent MJ & Vent CJ (1999) Cyclosporine neurotoxicity. *Journal of Neurology* 246: 339–346.
- Harborne JB (1998) *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. Chapman and Hall Ltd., London, 605 p.
- Humphrey T, O'Brien S, Madsen M (2007) Campylobacters as zoonotic pathogens: a food production perspective. *International Journal of Food Microbiology* 117(3): 237–57.
- Istifanus MF, Ogbonna AI, Sila MD & Ogbonna CIC (2014) Species of microorganisms associated with decayed tubers of Irish potato in storage in Plateau State, Nigeria. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 8(12)1: 54–57.
- Johnson WC & Williams OW (2002). Warfarin toxicity. *Journal of Vascular Surgery* 35: 413–421.
- Kumar A, Bhushan V, Verma S, Srivastav G & Kumar S (2011) Isolation and Characterization of Microorganisms Responsible for Different Types of Food Spoilages. *International Journal of Research in Pure and Applied Microbiology* 1(2): 22–31.
- Lawrence PA, Casal J & Struhl G (1999) Hedgehog and engrailed: pattern formation and polarity in *Drosophila* abdomen. *Development* 126(11): 2431–2439.
- Lis-Balchin M & Hart S (1999) Studies on the mode of action of the essential oil of Lavender *Lavandula angustifolia* P. Miller. *Phytotherapy Research* 13(6): 540–542.
- Mahesh S, Vaishnav V, Kumar P, Mohammad N & Ansari SA (2016) Characterization and validation of teak plus trees ramets of national teak germplasm bank, Chandrapur, Maharashtra through microsatellites. *Tropical Plant Research* 3(1): 213–220.
- Matthew T (2011) Post harvest microbial deterioration of tomato (*Lycopersicon esculentum*) fruits. *Report and Opinion* 3(4): 52–57.
- Mead PS, Slutsker L, Dietz V, McCraig LF, Bresee JS, Shapiro C, Griffin PM & Tauxe RV (1999) Food-relationship illness and death in the United States. *Emerging Infectious Diseases* 51: 607–625.
- Mrityunjy M, Naira nayeem A, Jagadish VK & Mohammed A (2007) Evaluation of *Tectona grandis* leaves for wound healing activity. *Pakistan Journal of Pharmaceutical Sciences* 20(2): 120–124.
- Nakamura C, Skarascis GV & Romagossa I (1991) *Cytogenetics and breeding in sugarbeet*. In: Tsuchiya T & Gupta PK (eds) *Chromosome engineering in plants, genetics, breeding and Evolution*. Elsevir, New York, USA, pp. 295–314.

- Ogbole OO, Adeniji JA, Ajaiyeoba EO & Adu DF (2013) Anti-polio virus activity of Medicinal plants selected from the Nigerian ethno-medicine. *Academic Journals* 12(24): 3878–3883.
- Ogunbanwo ST, Fadahunsi IF & Molokwu AJ (2014) Thermal stability of lactic acid bacteria metabolites and its application in preservation of tomato pastes. *Malaysian Journal of Microbiology* 10(1): 15–23.
- Oranusi S, Galadima M, Umoh VJ & Nwanze PI (2007) Food safety evaluation in boarding schools in Zaria, Nigeria, using the HACCP system. *Scientific Research and Essay* 2(10): 426–433.
- Orwa CA, Mutua R, Kindt R & Anthony S (2009) Agroforestry Database: A tree reference and selection guide version 4.0. Available from: <http://www.worldagroforestry.org/sites/treebs/treedatabase.asp> (accessed: 12 Dec. 2016).
- Prance GT (1991) What is ethnobotany today? *Journal of Ethnopharmacology* 32: 209–216.
- Prescott LM, Harley JP & Klein DA (2002) *Microbiology*, 5th Edition. New York, McGraw-Hill, pp. 1026.
- Purushotham KG & Sankar L (2013) Screening of in vitro Antibacterial activity of *Tectona grandis* on burn pathogens. *International Journal of Pharmacy and Biological Sciences* 3(3): 488–492.
- Purushotham KG, Arun P, Jayarani JJ, Vasanthakuopari R, Sankar L & Reddy BR (2010) Synergistic in vitro antibacterial activity *Tectona grandis* leaves with tetracycline. *International Journal of PharmTech Research* 2: 519–523.
- Rajkowski KT & Bennett RW (2003) *Bacillus cereus*. In: Miliotis MD & Bier JW (ed) *International Handbook of foodborne Pathogens*. Marcel Dekker, Inc. New York, pp. 40–51.
- Shalini SR (2008) Antifungal Activity Screening and HPLC Analysis of Crude Extract from *Tectona grandis*, *Shilajit*, *Valeriana wallachi*. *The International Journal of Alternative Medicine* 5: 2.
- Sharma N, Gupta PC & Rao CV (2012) Nutrient content, mineral content and antioxidant activity of *Amaranthus viridis* and *Moringa oleifera* leaves. *Research Journal of Medicinal Plant* 6(3): 253–259.
- Sumthong P, Romero-Gonzalez RR & Verpoorte R (2008) Identification of anti-wood rot compounds in Teak (*Tectona grandis* L.f) Sawdust extract. *Journal of Wood Chemistry Technology* 28: 247–260.
- Trease GE & Evans WC (2004) *Pharmacognosy*. Baillevera Tindall, London, 707 p.
- Trias R, Baeras L, Montesinos E & Badosa E (2008) Lactic acid bacteria from fresh fruit and vegetables as biocontrol agents of phytopathogenic bacteria and fungi. *International Journal of Microbiology* 11(4): 231–236.