

KAIRI KIIK

Reproduction and behaviour  
of the endangered European mink  
(*Mustela lutreola*) in captivity





**KAIRI KIIK**

Reproduction and behaviour  
of the endangered European mink  
(*Mustela lutreola*) in captivity



Department of Zoology, Institute of Ecology and Earth Sciences, Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of Doctor of Philosophy in zoology at the University of Tartu on the 06<sup>th</sup> of November 2017, by the Scientific Council of the Institute of Ecology and Earth Sciences, University of Tartu.

Supervisors: Prof. Toomas Tammaru, University of Tartu, Estonia  
Dr. Tiit Maran, Tallinn Zoological Gardens, Estonia

Opponent: Dr. María Díez León, University of London, United Kingdom

Commencement: Room 301, 46 Vanemuise Street, Tartu, on the 07<sup>th</sup> of February 2018 at 10.15 a.m.

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu

ISSN 1024-6479  
ISBN 978-9949-77-649-8 (print)  
ISBN 978-9949-77-650-4 (pdf)

Copyright: Kairi Kiik, 2018

University of Tartu Press  
[www.tyk.ee](http://www.tyk.ee)

## TABLE OF CONTENTS

LIST OF ORIGINAL PUBLICATIONS .....	6
1. INTRODUCTION.....	7
2. MATERIAL AND METHODS .....	11
2.1. Study species: European mink.....	11
2.2. The captive population of European mink at Tallinn Zoological Gardens.....	13
2.2.1. Management of the captive population.....	13
2.2.2. Data collection .....	13
3. RESULTS .....	16
4. DISCUSSION .....	19
SUMMARY .....	24
KOKKUVÕTE.....	25
ACKNOWLEDGEMENTS .....	27
REFERENCES.....	29
PUBLICATIONS .....	35
CURRICULUM VITAE .....	92
ELULOOKIRJELDUS.....	94

## LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following papers referred to in the text by Roman numerals.

- I** Kiik, K., Maran, T., Nemvalts, K., Sandre, S.-L., Tammaru, T. 2017. Reproductive parameters of critically endangered European mink (*Mustela lutreola*) in captivity. *Animal Reproduction Science* 181: 86–92.
- II** Nagl, A., Kneidinger, N., Kiik, K., Lindeberg, H., Maran, T., Schwarzenberger, F. 2015. Noninvasive monitoring of female reproductive hormone metabolites in the endangered European mink (*Mustela lutreola*). *Theriogenology* 84: 1472–1481.
- III** Kiik, K., Maran, T., Nagl, A., Ashford, K., Tammaru, T. 2013. The causes of the low breeding success of European mink (*Mustela lutreola*) in captivity. *Zoo Biology* 32: 387–393.
- IV** Kiik, K., Maran, T., Kneidinger, N., Tammaru, T. 2016. Social behaviour of endangered European mink (*Mustela lutreola*) litters in captivity. *Applied Animal Behaviour Science* 182: 61–71.
- V** Haage, M., Bergvall, U. A., Maran, T., Kiik, K., Angerbjörn, A. 2013. Situation and context impacts the expression of personality: The influence of breeding season and test context. *Behavioural Processes* 100: 103–109.

All published papers are used for the purpose of academic procedure and are reprinted with the permission of the copyright owners.

The participation of the author in preparing the listed publications (\* denotes a moderate contribution, \*\* a high contribution, \*\*\* the leading role)

	I	II	III	IV	V
Original idea	**	*	**	***	*
Study design	***	*	***	***	**
Data collection	**	**	**	***	**
Data analysis	***	*	***	***	*
Manuscript preparation	***	*	***	***	*

# 1. INTRODUCTION

Loss of biodiversity is a major concern. Some conservation measures have been notable successes, but the current level of conservation action is outweighed by the magnitude of the threat (Hoffmann et al., 2010). According to the IUCN Red List 22.2% of those mammal species assessed, are globally threatened or have become during the last few decades (IUCN 2017). The best method to improve the situation is to protect ecosystems and wild populations of species, that is, to apply *in-situ* conservation. However, this is not always possible as some threatened species have already too small, fragmented populations. Eliminating all the factors causing the decline of a species is not always achievable, or may even be impossible. For example, the last black-footed ferrets (*Mustela nigripes*) were threatened by habitat loss, overhunting, a decline in prey species and an epidemic disease (Clark, 1987). In such cases *in-situ* conservation might not be appropriate, as the results of conservation actions may take too long to yield results. The last chance to save a species from extinction may be the *ex-situ* conservation, i.e. bringing the last extant individuals into an artificial environment (Canessa et al., 2016; Snyder et al., 1996).

Unfortunately, the *ex-situ* conservation is not an easy way to save a species from extinction. This method causes the animals much stress during every phase: catching the wild specimens; transporting them to the new, artificial environment; and the management in captivity with all the new interactions with surroundings and conspecifics (Morgan and Tromborg, 2007). The *ex-situ* conservation is much more expensive compared with many other conservation activities and therefore the high cost of keeping a population in an artificial environment plays also a role in the moderate success of this method (Primack and Sher, 2016). Animals should be kept in captivity as few generations as possible, otherwise they start to adapt to the captive environment in terms of morphological (O'Regan and Kitchener, 2005) and behavioural changes, which result in the development of a domestic phenotype (Price, 1999).

Saving species in captivity has generally been more complicated and less successful than improving the natural habitat of the animals. Hence, it has been considered important that the decision to remove animals from the wild is made after a thorough investigation of the situation (Dolman et al., 2015). Conde and co-workers (2011) claim that *ex-situ* conservation programmes have had a low success rate, mainly because they have been initiated too late when only a few individuals remained. Captive breeding should therefore not be considered the last chance of saving a species from extinction, but as an important tool that should be instigated before a species faces extinction (Conde et al., 2011).

Although maintaining captive populations is generally difficult and problematic, there are several successful examples (Soorae, 2011). The most prominent are perhaps those involving the golden lion tamarin (*Leontopithecus rosalia*; Kierulff et al., 2008), Przewalski's horse (*Equus ferus przewalskii*; Xia et al., 2014) and the black-footed ferret (Clark, 1987).

The case of black-footed ferret has been an important and encouraging example for all *ex-situ* projects and especially for that on the European mink. The black-footed ferret was once common throughout West Central North America, from southern Canada to northern Mexico (Clark, 1987). The species is strongly dependent on different species of prairie dogs (*Cynomys* spp.) as prey, and these ferrets also use the burrows of this rodent as shelter and denning sites (Miller et al., 1988). At the beginning of 20<sup>th</sup> century, the number of wild black-footed ferrets decreased rapidly due to overhunting and the extermination of prairie dog colonies by humans (Biggins and Schroeder, 1988). In 1967, the black-footed ferret was listed on the U.S. Endangered Species List and in 1982 on the IUCN Red List as endangered (IUCN 2017). In 1979, the last known wild individuals died and the species was thought to be extinct. However, in 1981 an additional population was found, which consisted of approximately 100 animals. To secure the survival of the species, a captive breeding programme was established. In 1985 the wild population was decimated by an epidemic of Sylvatic Plague (*Yersinia pestis*), and the remaining individuals were captured (Reading et al., 1966). A captive population of 26 ferrets was established, 18 of them could be bred. Owing to the high degree of interrelatedness among those 18 animals, the genetic diversity corresponded to no more than the equivalent of seven founders. To preserve the genetic information of these founders, semen and ovaries were taken from these animals and stored frozen for future use. These frozen semen samples are still used to fertilize females (Howard et al., 2016). In general, keeping the captive population has been successful and to date more than 9000 black-footed ferrets have been born in captivity (Bortner, 2017; Marinari et al., 2006), and more than 4000 ferrets have been reintroduced to more than 20 locations in the United States, Canada, and Mexico. There are four self-sustaining populations, unfortunately, the number of prairie dogs is often not high, which in turn affects the success of reintroduction (Salkeld, 2017). Therefore, the re-established wild populations of black-footed ferrets remain small, fragmented, and still require intensive management (Santymire et al., 2014).

The captive breeding of black-footed ferrets was not without problems. In captivity, more than 50% of males failed to sire offspring. This negatively affects the genetic diversity of the captive population and may lead to increased incidence of deleterious traits affecting the overall viability of the population. An improper mating posture, abnormally high aggression during the mating attempt and underdeveloped testes have been identified as the main factors that induce reproductive failure in males (Wolf et al., 2000). In 2000, it was reported that reproductive failure was not related to semen quality, but in 2006–2007 the percentage of normal sperm morphology in ejaculates had decreased from 50% (in 2000) to 16%. In addition, the proportion of females that became pregnant after successful copulation declined from 70% (1987–1998) to 46% (2013). The reduced breeding success may be ascribed to inbreeding depression but also changes in diet have been discussed (Santymire et al., 2015).



There is an ongoing discussion regarding the reasons why so many carnivorous species are difficult to breed in captivity. One frequently cited reason is the large size of individual territories in the wild, which implies that the animals are adapted to covering long distances daily (Clubb and Mason, 2007; Quirke et al., 2012; Zhang et al., 2004). It is difficult to recreate similar territory sizes in captivity. In many cases, captive males show more problematic behaviour than females (Paper IV; MacKinnon et al., 2008; Wolf et al., 2000), which may be explained by the formers' larger home ranges (Kroshko et al., 2016; Palomares et al., 2017). Maybe this forced reduction in home range is the reason males cannot cope with small enclosures, which might be a factor in the development of abnormal behaviour primarily in the male sex (MacKinnon et al., 2008; Zhang et al., 2004; Wolf et al., 2000).

One of the most important issues in terms of keeping captive carnivores, is to provide them with suitable conditions during the breeding season. There are several examples of difficulties encountered with managing animals during the mating season, but the exact problems frequently remain unknown. For example, the Asiatic golden cat (*Catopuma temminckii*) is difficult to breed in captivity. The possible reasons of their failure to mate successfully might be socialization problems, as most individuals of this captive population have not had proper contact with conspecifics. Social isolation and a lack of early socialization with conspecifics may have affected these cats' ability to interact appropriately with each other (Olsen, 2012).

\*\*\*

The present thesis is focused on a critically endangered carnivore, the European mink (*Mustela lutreola*). This mustelid was once common almost all over Europe but, it has disappeared from most parts of its former range. The main causes of this drastic decline are habitat loss, overhunting, and invasion of the dominant American mink (*Neovison vison*; Maran, 2007; Maran and Henttonen, 1995). As the number of European mink decreased rapidly in the wild, a decision was made to launch a captive breeding programme to prevent their extinction (Maran et al., 2017).

Among the other mustelids, the European mink is relatively well studied. The ecological information available can be used in the planning of conservation measures. There are studies about diet (Maran et al., 1998; Põdra et al., 2013b; Sidorovich et al., 1998; Sidorovich et al., 2010), habitat preference, activity, and home range in the wild (Danilov and Tumanov, 1976; Harrington et al., 2014; Novikov, 1939 cited in Youngman, 1990; Sidorovich and Macdonald, 2001; Palomares et al., 2017). Nevertheless, information about mating behaviour, the litter period, and litter dispersal, is scarce.

To ensure a successful captive breeding programme, the aim of the present research was to study the reproductive biology of the European mink in captivity. We studied hormonal profiles and the factors that influence the reproductive success of females, mating problems, social behaviour during the

litter period, and individual differences in reactions to different test stimuli. Breeding biology was described in detail, including the timing of the mating season, gestation period, litter size, and hormonal changes during the breeding season (Papers **I** and **II**). In addition, the reproductive output of females was analysed. This included the number and size of litters delivered during each female's lifetime, as well as the factors (e.g. age, weight of the mother) affecting these parameters (Paper **I**).

The main problem in successfully managing the captive population of European mink was low mating success. In our study, 67% of all studied captive born males over 2004–2010 failed to pass their genes on to succeeding generations. The problem persists, with only approximately 40% of males breeding successfully every season (**III**). This negatively affects the effective population size, which leads to a reduction in genetic diversity and inbreeding depression. First, it is most important to determine the reasons for breeding failures, as only then can one make suggestions on how to improve the situation. For this reason, we conducted a detailed study of all mating attempts. We analysed which male and female traits are the main determinants of mating success. The origin of the animal, its gender, age, the identity of its partner, and the number of mating attempts in a single year were recorded (**III**).

The environment of young can have an important effect on their future behaviour (Bautista et al., 2010; Crews et al., 2009; Rödel and von Holst, 2009). In a study of American mink, it was found that the amount of time spent on social play may affect the development of adulthood social behaviour, and thereby future mating behaviour (Ahloy Dallaire and Mason, 2017). This may also be the case with the European mink. Therefore, the litter period of the European mink was a special focus to find any indications of future normal/abnormal mating behaviour. The aim was to describe the social play behaviour of European mink cubs in detail and compare it with related species. We aimed to find clues to any behavioural distortion in the captive population. Also, a more general classification of the behavioural acts of the cubs (non-contact play, duration of bites, and solitary behaviour) during the litter period was recorded. In addition, mothers' play with their cubs and aggressive behaviour of the mother was monitored and its temporal dynamics analysed (**IV**).

As do humans, other species can show individual variation in behaviour (Gosling, 2001; Sih et al., 2004). Also individuals of an endangered species may vary in how fast and successfully they adapt to a novel environment, as when introduced to captive conditions. Individual temperament may be an important factor in the success of conservation actions (McDougall et al., 2006). Therefore, we studied the personality types of European mink, with the aim of linking them to any behavioural problems observed during the mating season. As the first step, we addressed the question of whether it is possible to distinguish different personality types among a captive population of European mink. We performed a set of five experiments in two situations (non-breeding and breeding season, **V**).

## 2. MATERIAL AND METHODS

### 2.1. Study species: European mink

The European mink is a small semiaquatic carnivore that primarily inhabits small forest streams that do not completely freeze during winter. Their diet is comprised mostly of fish, amphibians, and crustaceans, but also includes small mammals (Maran et al., 1998; Sidorovich et al., 1998). The European mink was once spread throughout Europe, extending from the Ural Mountains to eastern Spain, and Finland to the Black Sea (Maran, 2007). Currently, the European mink is a critically endangered species (IUCN 2017) and one of the most endangered mammals in Europe (Temple and Terry, 2007). It is estimated that fewer than 5000 European mink persist in the wild, in less than 3% of their former range. The only remaining populations are in Danube Delta and in northern Spain (Maran et al., 2017), but even these populations are threatened by invasion of American mink (Macdonald and Harrington, 2003).

The decline of European mink has multiple causes in different regions and times, of which the introduction of American mink is just one. European mink started to vanish in central Europe as early as the end of 19<sup>th</sup> century. This was probably due to large-scale land-improvement and the dredging of rivers, which reduced the amount of suitable habitat. Declines in the first half of the 20<sup>th</sup> century in Eastern European countries (Poland, Hungary, the former Czechoslovakia), might be connected to the extensive development of agriculture, whereas in Finland it may have been caused by intensive hunting and active muskrat (*Ondatra zibethica*) trapping during the 1930s. In Estonia, major land improvements (e.g. the draining of small rivers) caused habitat loss, and thereby fragmentation of mink populations. European mink was still extant during the late 1980s in northern and eastern parts of Estonia, whereas in the south and west invasion of American mink had led to their local extinction. The last confirmed record of European mink is from 1996, after which wild populations have been regarded as extinct in Estonia (Maran, 1991). European mink had already experienced major declines before American mink were introduced in during the 1950s–1970s. This invasion of an alien mink was probably the terminal factor in the European mink becoming virtually extinct in Europe (Maran and Henttonen, 1995).

American mink were brought to Europe in the early 20<sup>th</sup> century as numbers of European mink were already too low for the fur-trapping industry and because of the comparatively higher value of American mink pelts. Introduction of American mink was not considered a threat to our native species as in those times they were not regarded as two distinct species (Maran et al., 2016b). American and European mink share the semi-aquatic habitat, and are similar in appearance and ecology despite the fact that they are not sister species. The American mink is even classified as belonging to a different genus, *Neovison* (Abramov, 2000; IUCN 2017). With respect to the diet of these two species,

European mink are slightly more specialized than American mink, but their diet overlaps to a considerable extent (Sidorovich et al., 1998). When these two species coexist, then encounters are probably aggressive, dominated by the larger American mink, and could even result in the death of European mink (Põdra et al., 2013a). As a consequence, American mink force European mink to withdraw to suboptimal habitats, where the amount of prey might not be sufficient for females with a litter and reproduction may fail. This, in turn, induces further fragmentation and decreases population sizes (Sidorovich et al., 1999; Sidorovich and Macdonald, 2001).

In 1992, the European Endangered Species Program (EEP) for the European mink was launched by the European Association of Zoos and Aquaria (EAZA), to secure the survival of the species. The first goal was to establish a viable captive population that breeds regularly. The long-term aim of this programme was to maintain in European zoos and other breeding facilities 85% of the original heterozygosity of the founding individuals over 50 years. Fourteen different institutions participate in this programme, with an overall stock of 210 specimens (as of 31.12.2016). The coordinator of this program is Tallinn Zoo, wherein the largest captive population of European mink is located. The Tallinn population is based on 22 founders, which were obtained from Russia through collaboration with the Central Forest Biosphere Reserve (Tver, Vologda, Pskov, Novgorod and Leningrad Regions, Russia; Maran, 2003). The first breeding event occurred in Tallinn in 1986, but regular breeding was not achieved until the mid-1990s. Currently, the size of the population in Tallinn has stabilised at the near maximum level that the facility can sustain – 112 individuals, with genetic diversity (GD) retained at 93%, a generation time of 1.62 years, and an effective population size of 65.6 individuals (as of September 2017). Extension of the facility is limited mainly by a shortage of funds for new constructions, but also by the high costs (food, husbandry etc) of maintaining a high number of mink in captivity (Maran, 2003).

In addition to the preservation of the genetic diversity of the species in captivity, the EEP also serves as a source for mink reintroduction projects in Estonia. The aim is to ensure that at least some areas have extant European mink populations, and that they must stay inaccessible to American mink. This means reintroducing individuals into a sufficiently large area with suitable habitats to maintain a self-sustaining population of European mink. Preferably, this should be an island surrounded by distribution barriers that make the invasion of American mink impossible (Maran, 2003). In 1998, it was decided to establish a special island reserve for European mink on Estonia's second largest island: Hiiumaa. The island lies 22 km from the mainland and the habitats there should be able to accommodate about 100 individuals. After trapping out the American mink that had escaped from a fur farm that no longer exists on the island, the island was pronounced American mink free. The distance to the mainland is too far for American mink to spread there independently (Maran, 2003). In 2000, the first European mink were released onto the island and such introduction attempts have occurred annually ever

since. So far (2000–2016), 578 animals have been released (Maran et al., 2016a). Monitoring radio-tracked European mink has revealed that released mink adapt to the wild and establish stable home ranges 1–1.5 months after release. Survival has been highest when the animals were maintained in large natural enclosures before release. The main causes of death have been predation by other carnivores (Maran et al., 2009). After years of purposeful work, there is now a small breeding population on the island (Maran et al., 2016a).

## **2.2. The captive population of European mink at Tallinn Zoological Gardens**

### **2.2.1. Management of the captive population**

All the studies included in this thesis were made at Tallinn Zoo. In Tallinn the captive European mink are housed individually in 2m × 2m × 4m enclosures, which are located side by side. The enclosures are separated by wire mesh walls with the lower parts covered by plywood to reduce visual contact between the animals. Each enclosure contains a nest box, a water pool (64cm × 35cm × 30cm) and several other items, such as branches, tree stumps, and tubes for the purpose of environmental enrichment. The animals are fed with rodents (frozen), fish, birds (chicks), or minced beef meat.

The animals are usually visited once a day, when they are fed and their enclosures cleaned. All animals are weighted once a month, when they are transported to and from a weighting room using a transport box. When the breeding season approaches, those females selected to be bred are regularly monitored for the onset of oestrus (**II**). When oestrus is detected, the male selected as a mate is introduced into her cage. The couples are observed for approximately an hour or until the copulation occurs, and then left together a few days (Papers **I**, **II** and **III**). Pregnant females and their future litters are left undisturbed in their enclosures until the cubs are about two months old, when they are separated from their mother (**IV**). At the beginning of September the litters are separated and all the cubs get their own enclosure.

While performing each of the studies, care was taken that the normal routine of the animals was not disturbed, or the changes were minimal. The well-being of the animals was always our first priority.

### **2.2.2. Data collection**

The breeding data used in papers **I** and **III** were received from the European mink studbook database, maintained in the Single Population Analysis and Record Keeping System (SPARKS; [www.species360.org](http://www.species360.org)). The purpose of this software is to assist in assembling, editing, analysing, and producing reports on the status of a captive population held at multiple facilities. The database

contains individual data on each animal: time and place of birth, gender, parents, any movement between breeding facilities, breeding, offspring produced and time and place of death.

To analyse the reproductive success of females, data on all the litters females had produced over 20-year period was gathered from SPARKS. 165 different females had produced at least one litter during that time. To calculate juvenile mortality, data about deaths before an animal's first breeding season was also retrieved from that database. Information about the pedigree of each mink was also retrieved to calculate the heritability of litter size. The dataset covered 204 individuals (I). In paper III, the SPARKS database was used to find the number of successful and unsuccessful breeding events, to analyse the effect of mink origin on breeding success. The compiled data set contained information on 33 wild and 224 captive born mink.

In addition, data from the breeding database – updated annually by the research lab at Tallinn Zoo – were used in paper I to retrieve data about cubs born and cubs that survived until litter separation (at the beginning of September). In paper III, such breeding logs were used to gather information about all the mating attempts from 2004–2010 (N=579), to analyse the determinants of mating success. A breeding log is filled in per mating attempt, with the data recorded including the identity of the male and female, behavioural observations, and the outcome (successful or not).

### ***Hormonal study***

To determine seasonal hormonal profiles in pregnant and nonpregnant females (II), a non-invasive monitoring method was used. Faecal samples were collected over 16 months – to include two consecutive breeding seasons – from 15 females selected for breeding. Oestrogen and pregnane metabolite levels were measured using enzyme immunoassays following an established protocol for total oestrogen and 20-oxo-pregnanes. To study induced ovulation, three other females that were in oestrus were treated with human chorionic gonadotropin (hCG; Sigma C-1063; 90 IU, 0.36 mL, intramuscular) and faecal samples collected 15 days later. To correlate vaginal cytology for oestrus detection with faecal oestrogen metabolite measurements, faecal samples and vaginal smears were collected at one to three day intervals from March to May from 11 females that were not bred that year.

### ***Litter behaviour***

To study social behaviour during the juvenile period of European mink, 13 litters were observed 10 times over a two-month period. During each observation event, the litter was video recorded for 30 minutes from the top of the enclosure. The recordings were used to compile a detailed ethogram of social and solitary play behaviour of European mink cubs. Litters were disturbed as

little as possible to avoid influencing their behaviour. Therefore, the pregnant females were placed into enclosures over which observation platforms had been built earlier. The litters stayed in the same enclosure over the whole period (IV).

### ***Personality traits***

To determine personality trait domains and examine how they are affected by situation and context, 80 animals were tested during autumn, and second time in spring before the breeding season (n=68). The animals were experimentally tested five times during both these periods: three times in their own enclosure, and twice in a novel environment. The animals were transported to a novel environment by catching them in a transport box, which is a standard procedure used also in connection to monthly for weighting.

Types of behaviour were scored only the first time they were observed, for example, when the animal first approached the novel object. We gave higher scores if the focal behaviour occurred early during the experiment. Individuals who, for example, attacked a novel object immediately, differed in their behaviour from individuals that dared to attack the object just before the end of the trial. The test period was 4 minutes divided into three 80 second periods. The number of points given depended on the period in which certain behaviour first occurred: 3 points in the first, 2 points in second, and 1 point in last period. Latency before the animal left the nest or transport box was measured in seconds (maximum 240 seconds; Paper V).

### 3. RESULTS

#### *Reproduction*

Breeding of European mink occurs during spring. Females come to oestrus from middle March to middle May, with the peak in April (**I**). When the first oestrus period remained undetected or the mating attempt failed, females could have another oestrus later in spring (**II**).

The gestation period was on average 43.8 days ( $\pm 1.4$  SD), the mode being 43 days. During the 20-year period, 267 litters were delivered, of 1171 total cubs. Average litter size was 4.42 ( $\pm 1.48$  SD), and the mode was four cubs. Eighty-five (7.3%) cubs died before their first mating season. Most of the deaths ( $n=79$ ; 93% of all deaths) occurred before litter separation. The average number of juveniles per litter that survived to litter separation was 4.08 ( $\pm 1.48$  SD), with 4.07 ( $\pm 1.5$  SD) surviving to their first breeding season (**I**).

We analysed the determinants of reproductive success among females with the aim of providing recommendations for selecting which females to breed. We found that the age of the mother had a negative effect on litter size. However, the negative effect of age appeared only when the mother was older than four years: younger age classes did not differ in breeding success. The weight of the mother had a positive effect on the size of the litter. All the large litters ( $> 6$  cubs) were produced by females that were heavier than 650g. We also found that the weight of mothers increased over the 20 years of the programme. We did not find any signs of the cost of reproduction: the number of litters a female had delivered earlier in her life did not have an effect on her litter size in the focal year; the number of cubs born and number of cubs survived were not affected by the mothers reproductive history from the preceding year. When the effects of age and weight were accounted for, individual females did not vary in terms of litter size, also the heritability of litter size was estimated to be low (**I**).

#### *Hormonal study*

With the aim of improving the breeding success of European mink, a detailed study of oestrus, ovulation and pregnancy was conducted. The overall pattern of total oestrogen and 20-oxo-pregnane in pregnant and nonpregnant animals followed a seasonal profile. During the breeding season, the mean oestrogen metabolite level was elevated as compared to before the onset of the breeding season. After conception, both oestrogen and 20-oxo-pregnane concentrations increased. Both metabolites reached the highest levels during gestation. After treatment with hCG, 20-oxo-pregnane levels increased resembling the hormonal profile of pregnant females. During lactation, 20-oxo-pregnane metabolite levels were higher than in nonpregnant females. Litter size was correlated with 20-oxo-pregnane levels but not with oestrogen (**II**).



### ***Mating success***

By comparing the breeding success of wild and captive born males, we found wild born males were significantly more successful than captive born males (success rate 89% and 35%, respectively). Hence, the poor breeding success should primarily be attributed to males born in captivity (III). In the detailed study of the data on mating attempts, we determined that male aggressive and passive behaviour were both primary causes of mating failures. We revealed that mating success, males' aggressiveness and passivity depend more on the male than the female partner. Additionally, we did not find any evidence that the behaviour of an individual was dependent on the identity of its partner. We concluded that passivity and aggressiveness are both alternative expressions of males' behavioural inability to mate. Aggressively behaving males are passive in non-aggressive situations and *vice versa* (III).

### ***Behaviour of cubs during the litter period***

To study the social behaviour of the cubs, 13 litters were observed during active times that amounted to a total of 3900 minutes of surveillance (IV). Overall, the most frequent type of behaviour was social play. Among the different litters, an approximately equal share of time was spent on social play. The play itself was diverse, containing all the different elements described in related species. We did not reveal any significant association between the time spent on biting and the age of the litter. Only the social play of the cubs (excluding bites) decreased significantly over time and the pattern was similar across the litters. There was a significant change in the duration of mothers' behaviours with litter maturation: the duration of mothers' play with the litter decreased and aggressive behaviour of the mother increased. We also found a positive correlation between time cubs spent biting siblings and mothers' play with the litter (IV).

### ***Personality traits***

During the study in which we assessed personality types, the behavioural measurements obtained from the tests were grouped into three factors, which taken together explained 55.9% of the variance (V). The first domain was called "boldness" as it was related to behaviours such as "approach to object" and "latency to leave the nestbox in the presence of a novel object". The second domain was mostly related to responses to the mirror image of the animals in the novel arena, such as "looking at and sniffing the mirror image", or "marking the area". It seemed to be related to non-aggressive social behaviour outside the animal's own territory. Therefore, this domain was called "sociability". The third domain was associated with explorative behaviour in the novel environment such as "number of zones visited in a novel arena", "hissing", "marking" and "digging". This factor was called "exploration".

The scores of all these three personality trait domains were repeatable between the seasons. Between the seasons, gender had a significant relationship with boldness and there was a significant interaction between gender and season in terms of boldness. Whilst males got bolder females got shyer with the breeding season approaching (**V**).

## 4. DISCUSSION

Overall, management of the captive population of the critically endangered European mink at Tallinn Zoo has been successful. A considerable amount of offspring (1171 cubs) have been born during the 20-year period (1996–2015). Mortality of the cubs has been minimal (85 cubs), with most cubs surviving until their first breeding season. An overview of the date of the onset of the breeding season, gestation period, average litter size, and litter mortality was given in our study (I) that involved a much larger sample size compared with earlier studies (Amstislavsky et al., 2009; Ternovskii, 1977) but the results were similar. Gestation period did not differ much from that of related species: it was similar to the European polecat (*Mustela putorius*), the steppe polecat (*M. eversmanni*; Nowak, 2004), and the black-footed ferret (Williams et al., 1992).

The average size of captive European mink litter was 4.4 (I) which is close to the value reported from Russia (4.7) by Danilov and Tumanov (1976), although these values are higher than those of wild population in Belarus (3.8; McDonald and Harris, 2002), and France and Spain (3.4; Fournier-Chambrillon et al., 2010). Consistently, when comparing the data about the litter sizes of related black-footed ferret then in captivity it was larger (3.4 cubs; Vargas and Anderson, 1998) than in nature (3.17 cubs; Ayers et al., 2014).

In female European mink, the hormonal profiles during the different reproductive stages were described and the hormonal differences between pregnant and nonpregnant females analysed (II). The results were in accordance with the earlier conclusion that European mink is a seasonally polyestrous species (Amstislavsky et al., 2009). The European mink is regarded as an induced ovulator, as are many other mustelids such as the polecat, black-footed ferret, river otters and the sable (Chang and Yanagimachi, 1963; Mead et al., 1988; Rozhnov et al., 2008; Stenson, 1988). The results of this study support this assumption because 20-oxo-pregnane levels did not increase in females that were not mated. Furthermore, 20-oxo-pregnane levels increased in hCG-treated females, closely resembling hormone profiles of pregnant females during the first 2 weeks of gestation. This indicates that hCG can be applied to induce ovulation and perhaps using as a new tool to support population management of the European mink. This has been successfully used in black-footed ferret (Howard et al., 1997) and domestic ferret (Chang, 1968).

An essential criterion of the successful management of a captive population is a sufficient number of offspring produced annually. Thus, per female mammals, the litter size and offspring survival are the parameters of primary interest, and predicting these is thus of a high practical value. We revealed that the most important factor influencing female minks' reproductive output was age, but only when the female was older than four years (I). The fact that the litter size decreases in time when the animal gets older is shown also in similar species like captive and wild American mink (Koivula et al., 2010; Melero et al., 2015). Nevertheless, in our study the litter size was not age-dependent in younger than

five year old females, which is in contrast to the findings in American mink (Koivula et al., 2010).

We also found a positive association between litter size and the weight of females (I), which has also been shown among various other species. The relationship between body weight and reproduction is mainly explained by variation in the nutritional status of the breeding animals (Frynta et al., 2011; Gese et al., 2016; Persson, 2005). We noticed that the body weight of European mink increased by approximately 10% over 20-years of observation. We hypothesise that this was likely the result of an improvement in the quality of the food provided to the animals, but unfortunately, the older nutritional data are too incomplete to test this. Nonetheless, the phenomenon of changing body weight among long-term captive population certainly requires future study.

In our study population, we found no evidence of the cost of reproduction (I). In the literature, it has been suggested that the cost of reproduction should be more apparent when the life-history trait in focus has a high variance. This is typically the case in long-lived species, to which the European mink can be considered to belong. Therefore, we probably could find some evidence of the cost of reproduction in wild European mink populations. In other members of the Mustelidae, the existence of reproductive costs have been reported, e.g. among wild American mink (García-Díaz and Lizana, 2013) and wild wolverines (Persson, 2005). We suggest that the high quality and constant availability of food in captivity eliminates the physiological mechanisms behind the cost of reproduction, so that a failure to observe this phenomenon should not be surprising. As a summary of these results, we suggest that when selecting females for breeding, there is little need to consider aspects other than genetic relatedness (I).

Captive populations of endangered species are mostly rather small in number and have minimal or non-existent immigration. Due to genetic drift, such populations are expected to randomly lose alleles. This can lead to significantly reduced genetic diversity and leave most of the loci in the homozygotic state, resulting in inbreeding depression (Frankham, 2010). Therefore, it is essential that all designated animals, which have been selected for breeding, with the aim to maintain maximum genetic diversity, will actually reproduce. Unfortunately, problems with breeding are common among captive species (Abello et al., 2011; Carlstead et al., 1999; Martin and Shepherdson, 2012). In paper III, the records of mating behaviour were examined. It was revealed that breeding failures mostly occur due to the abnormal behaviour of captive born males: some behave passively or are aggressive. In less than one third of the mating attempt studied, the males behaved normally, i.e. mated with the female. In most those cases, the female delivered a litter (III).

Problems with mating behaviour in captivity are common among many carnivorous (Wielebnowski et al., 2002; Wolf et al., 2000; Zhang et al., 2004). The suggested reasons of limited success include physiological problems (Wolf et al., 2000; Zhang et al., 2004), improper keeping conditions and environments with too low complexity (Diez-Leon et al., 2013; Zhang et al., 2004). In addition

familiarity with the breeding partner can play a significant role (Peng et al., 2009). The distortion of reproductive behaviour is frequently expressed as abnormal mating behaviour and it is often difficult to determine the causes (Coelho et al., 2012; Wolf et al., 2000). We can only suppose that the causes are associated with the captive environment and the conditions which exist there.

The early environment of an animals' life can cause a long-lasting effect on the individual's ontogeny (Hudson et al., 2011). As the behaviour of the European mink litters have not been described earlier, we compiled a detail ethogram of social behaviour among European mink litters. Their social behaviour was found to be complex (**IV**) and the behavioural acts observed closely resembled those reported for related species (Clark et al., 1986; Poole, 1966, 1967, 1978). This could indicate that the environmental conditions in enclosures are sufficiently variable and "normal" that young animals will and can perform different kind of play elements (Nogueira et al., 2011; Talbot et al., 2014).

We found that the most frequent type of behaviour during the litter period of the European mink was social play. Time spent on biting (which could indicate to aggressive behaviour) did not increase over time. The behaviour of the mother did change over time: the share of caring and playful behaviour decreased when the cubs grew older, and she also became more aggressive. Her aggressive behaviour became so frequent that we had to separate the mother from the litter to prevent injuries she could cause to her cubs. Likely, such an increase in the aggressive behaviour of the mother was associated with the litter dispersal. When the cubs got old enough they should start to seek their own territories and the mother's aggressive repulsing behaviour might encourage them to do so. Therefore we think that the change in the behaviour of the mothers was normal and would probably also happen in nature. In general, we found that the behavioural repertoire of captive-bred European mink litters was complex and consisted of all expected elements. We did not find litters with significantly differing behaviours and no unexpectedly high aggression was found. Therefore, we can conclude that our study did not reveal any juvenile behavioural distortion in European mink captive population (**IV**).

An increasing number of studies report the existence of personality types among various nonhuman species (Cheng et al., 2017; Kanda et al., 2017; Lehmkuhl Noer et al., 2016). In the case of the European mink, there might well be some environmental or management factor that affects some animals, so that their will develop abnormal mating behaviour. We identified personality types among European mink for the first time. These types were designated as boldness, sociability and exploration. We found them to be individual-specific and repeatable both within and across seasons. These results gave us an important knowledge base to test for possible relationships between these types and male reproductive behaviour in captivity (**V**).

\*\*\*

Summarizing the results from the perspective of *ex-situ* species conservation, I would highlight the following:

- at Tallinn zoo, fecundity and juvenile survival of European mink has been high compared to other available data on this and related species. General conditions at the zoo can therefore be considered appropriate, with little need for further improvements;
- there is little variance in the reproductive success of individual females. Females used for breeding can therefore be selected solely using the criterion of maintaining maximal genetic diversity;
- the main bottleneck threatening the success of the *ex-situ* breeding programme is the abnormal mating behaviour of captive born males;
- even if the reason for behavioural problems has often been in the social environment experienced during the juvenile development, the present study did not find any evidence of that in the case of the European mink.

Based on our results, I wish to outline several issues which deserve to be studied further.

- Even if there was nothing „suspicious” in the behaviour of juvenile European mink (**IV**), it would greatly help to obtain data on the behaviour of the juveniles and their parents in nature as a comparison.
- Also, I suggest focusing on the phenomena related to litter dispersal. This stage of their life cycle has unfortunately not been studied in detail, and no species-specific information exists regarding wild populations. However, some attempts along this stage have been made with captive European mink in Germany (Festl et al., 2003). The results indicate that social contacts with littermates persist after litter separation which may play an important role in the development of normal social behaviour in adulthood. Therefore, different timing and methods of litter separation should be studied in relation to future mating success.
- In our third paper (**III**), we found that being captive born is the main correlate of not mating normally. It has been demonstrated that environmental enrichment reduces stressful behaviours among captive American mink and improves their mating performance (Diez-Leon et al., 2013). It has also been found that the environmental enrichment is most important for young male black-footed ferrets as their stress level was reduced at most compared with other age classes and sexes (Poessel et al., 2011). We suggest that young European mink males should be hosted straight after litter separation in large, environmentally rich enclosures, and the effect of such treatment on mating success analysed.
- Even though we did not find evidence that mating success depends on the female partner, the effect of mate preference should be studied. It has been shown that mate preference positively affects copulation success, litter size

and offspring survival (Martin and Shepherdson, 2012; Martin-Wintle et al. 2015). We should compare reproductive success between females mated with a male from a neighbouring enclosure to success with males from a non-neighbouring enclosure, and between females mated with preferred and non-preferred neighbours. Preference should be determined based on behavioural interactions between the animals.

- The results of the fifth paper (V) suggest the need for even more thorough studies of individual behaviour. The methods used to determine the personality types of mink can be applied in future studies, consistent with the suggestions that captive breeding programmes should, beside genetics, also consider individual temperament when deciding on the selection of breeding pairs. Actions that ignore personality can lead to the captive population developing towards domestication through unintentionally favouring some types of temperament. This may affect the success of captive breeding programmes and future reintroduced populations (McDougall et al., 2006). Using results of paper V, we could develop an express method to quickly determine personality types, with the potential of using this information as part of a management protocol.
- The revealed increase in the weight of *ex-situ* females and the correlated increase in litter size (I), should be examined in the context of the effect of these temporal trends on mating success, health, and future reintroduction strategies.

## SUMMARY

Saving critically endangered species from extinction is the main goal of *ex-situ* conservation methods. Such methods are frequently inevitable, even when removing animals from their natural habitat and bringing them into an artificial environment is a difficult way to achieve this goal. Experience obtained from successfully managed populations is extremely important to ensure the success of other such programmes. The important and delicate events of animals' lives such as mating, gestation, birth and the litter period are also the most problematic in captivity.

In this thesis I gave an overview of European mink behaviour and reproduction. The results suggest practical actions to improve the breeding success of this species, and form the basis for future studies on captive European mink population.

At present, the most critical aspect of the captive breeding programme of this species is ensuring high mating success. We revealed that the main reason why breeding fails is due to the number of captive born males with inadequate (i.e. aggressive or passive) behaviour during the mating attempt. We found that mating success depends more on the male than on female partner (**III**). The females' hormonal profiles were analysed and no abnormalities were found (**II**). Females' breeding success in terms of litter size depends mainly on the age and weight of the animal, which corresponds to a pattern which is perhaps universal among mammals. No signs of reproductive cost were found. This was probably due to the high quality and always available food in captivity which might eliminate the physiological mechanisms behind the cost of reproduction (**I**).

Abnormal behaviour could develop due to an unsuitable environment during early life stages. As the social behaviour of the European mink litters were completely undescribed, a detailed ethogram of play behaviour during the litter period was compiled. The observed play was diverse and contained all behavioural acts described in related species. None of the litters were characterised by deviating behaviour and we found no evidence of unexpected aggressive behaviour. Therefore, our results do not provide evidence of behavioural distortion among captive European mink cubs (**IV**).

As advised in the literature, keeping a population in captivity should rely more on individual-based approaches. We determined three personality trait domains among the European mink captive population: boldness, sociability and exploration. We demonstrated that the situation and context influences the expression of personality trait domains and plasticity can be large, but that the concept of personality type remains applicable due to the repeatability of the measurements. Establishing different personality traits among mink shows that animals do differ in their behaviour in a consistent manner. These behavioural differences may affect the way individuals interact with the environment which may also lead to an inability to display proper behaviour during mating attempts (**V**).

As management of species in captivity is a challenging task, any salient knowledge published about other species is of high value, and can become most useful throughout the wide area of *ex-situ* conservation.



# KOKKUVÕTE

## Ohustatud Euroopa naaritsa (*Mustela lutreola*) sigimine ja käitumine tehiskeskkonnas

Ohustatud loomaliigi kaitsmine väljaspool tema looduslikku elupaika (*ex-situ* liigikaitse) võetakse ette enamasti alles siis, kui ülejäänud võimalused tema väljasuremise vältimiseks on ammendatud. Kahjuks on taolise meetodi edukus väga madal, lisaks on ta keeruline ja kallis. Seda eelkõige seetõttu, et loomad on vaja välja püüda nende looduslikust keskkonnast ning tagada neile edukas toimetulek uudsetes tingimustes. Hoolimata raskustest on *ex-situ* liigikaitse pahatihti ainuke ja viimane sobiv lahendus. On mitmeid näiteid, kus taolise tegevusega on edu saavutatud: kadumas olnud liigi populatsioon on taas loodusesse asustatud. Selliste edukate ettevõtmiste käigus omandatud kogemus ja teadmised on äärmiselt olulised analoogiliste projektide õnnestumise tagamiseks.

Käesolevas doktoritöös keskendume ühele kriitiliselt ohustatud imetajaliigile, Euroopa naaritsale. Naarits oli varasemalt levinud peaaegu üle kogu Euroopa, kuid nüüdseks on säilinud vaid mõned üksikud, kahaneva arvukusega populatsioonid. Liigi kadumise peamisteks põhjusteks olid üleküttimine, sobivate elupaikade vähenemine ja Ameerika naaritsa invasioon. Kindlustamaks liigi edasikestmist loodi tehiskeskkonda naaritsa populatsioon. Eesmärgiks on loomi paljundada, alal hoida veel alles olevat geneetilist mitmekesisust ja taasasustada naaritsaid neile sobivatesse elupaikadesse looduses.

Tehistingimustes peetava populatsiooni käekäigu seisukohalt on eelõige olulised sigimishooaeg, tiinus, poegimine ja pesakonnaperiood. Paraku on eelnimetatutega seotud ka kõige suuremad probleemid. Nii on see ka Euroopa naaritsa puhul: kõige problemaatilisem on edukas sigimine. Paljud loomad ei saa järglasi, mistõttu populatsiooni geneetiline mitmekesisus langeb murettekitava kiirusega.

Euroopa naaritsa paljundusprogrammis on praegusel hetkel kõige kriitiliseks piisava paaritumisedukuse tagamine. Käesolevas töös leidsime, et eelkõige on võimalik sigimisprobleeme seletada isaste naaritsate ebaadekvaatse käitumisega paaritumiskatses. Nimelt käitub osa vangistuses sündinud isastest sigimishooajal ebanormaalselt, s.t. on emase vastu agressiivne või jääb passiivseks ega ilmuta huvi partneri vastu (**III**). Uurides emaslooma suguhormoone, leidsime, et kõik on normaalne ja hormonaaltsükkel järgib naaritsal ja teistel lähedastel liikidel eelnevalt kirjeldatud rada (**II**). Emaste sigimisedukus, mida kirjeldasime läbi pesakonna suuruse ja poegade ellujäämuse, sõltus eelkõige emase vanusest ja kaalust. Sellised seosed on imetajate hulgas üsnagi universaalsed. Lisaks otsisime tõendeid sigimise hinnast naaritsa tehiskeskkonnas peetaval populatsioonil, kuid neid me ei leidnud. Kuna mitmetel sarnastel looduses elavatel liikidel on sigimise hind tõendatud, siis eeldasime, et kõrge kvaliteediga toidu pidev kättesaadavus mõjutab neid füsioloogilisi mehhanisme, mis mõnes teises olukorras sigimise hinda põhjustaksid (**I**).

Ebanormaalne käitumine võib areneda varajases eluetapis kogetud ebasobiva keskkonna tõttu. Euroopa naaritsa poegade sotsiaalset käitumist ei ole varem kirjeldatud. Koostasime mängukäitumise detailse etogrammi, st nimekirja käitumisaktidest, mida noorloomade mäng sisaldab. Naaritsapoegade käitumine oli mitmekesine ja sisaldas kõiki käitumisakte, mis on kirjeldatud ka sugulasliikidel. Ühtegi pesakonda ei saanud iseloomustada kui käitumiselt hälbivat ning me ei leidnud ka tõendeid ebanormaalselt agressiivsest käitumisest. Seetõttu ei paku meie tulemused tõendeid käitumuslikust ebanormaalsusest vangistuses peetavatel Euroopa naaritsa poegadel (IV).

Üha rohkem hakatakse pöörama tähelepanu sellele, et tehiskeskkonnas peetavate populatsioonide majandamine peaks sisaldama rohkem isendipõhist lähenemist. Tegemaks algust taolise tegevusega, määrasime loomaia naaritsate iseloomutüübid. Eristasime kolm iseloomu mõõdet: julgus, sotsiaalsus ja uuri-vus. Näitasime, et situatsioon ja kontekst mõjutavad iseloomutüübi ilmumist ja plastilisus on seega suur. Siiski saab tänu mõõtmiste kõrgele korratavusele rääkida naaritsate iseloomutüüpidest (V).

## ACKNOWLEDGEMENTS

I am most grateful to Toomas Tammaru for providing the knowledgebase to design of scientific studies, statistical analysis and an overall insight into the world of science. I am also enormously grateful to Tiit Maran, who gave me the opportunity to study this magnificent species, the European mink. I am thankful that Tiit involved me in many different projects connected with the European mink, which extended my horizon outside of my own specific topic.

In addition, I wish to express my gratitude to all the people with whom I have worked at the breeding facility of Tallinn Zoo. The company was terrific and I learned so much from all my colleagues. Those blessed minutes with them behind a cup of tea, accompanied by an enjoyable chat often kept me going. Great thanks to Astrid Nagl, who gave me the deciding push to start with my scientific work. Thank you for your guidance and your patience with answering all my questions. Seeing your work and how the arisen ideas are being put into practice was really helpful to me. Thank you Nadja Kneidinger, for being a superb team worker even on the coldest and rainiest of days. Chats with you, discussing and speculating about future study questions and possible improvements to the situation, were always cheerful and full of many important ideas. Great thanks to Marianne Haage. Working with you was always very educational. You were extra precise when collecting data and it was great to see that there is a solution to every problem and obstacle.

And of course I thank all the staff at the Zoo: Merje Põlma, Kadri Ashford, Kertu Namsing, Kristel Nemvalts. Without you it would not have been possible. Most of all I thank the fantastic keepers, who are invaluable as they provide every-day care to these magnificent animals.

I also want to thank my great office companions that helped me endure the hard days full of work behind a computer: Leidi, Karmen, Ragne, and Egle. It was a great joy to have a nice scientific chat about a completely different topic now and then, and sometimes a less scientific one.

I am grateful to Andro, Anne, Katarina, Katrin, Mehis, Triin K. and Triin T. for just being there, giving me the chance to enjoy your wonderful companionship.

I am thankful to the Tartu Students Nature Protection circle and all its members for giving me extraordinary memories from wonderful hiking trips and who did not let me forget that studying the nature is one thing, but you must be in nature to be really part of it.

I thank my parents for letting me make my own choices, no matter what they were or how long they took, for the freedom to be me and become the person that I am, and for their constant support despite no guarantee of ever seeing an outcome. I thank Taivo, for being my first example of a scientist and for unwillingly pushing me to do my own PhD degree.

Last but certainly not least, I would very much like to thank Mikk and Pihla for being in my world, showing me which parts of life are the most important and giving me answers to a lot of questions that emerged during my time studying life sciences.

\*\*\*

This study was financially supported by institutional research funding IUT 20–33 of the Estonian Ministry of Education and Research, and the Lutreola foundation.

## REFERENCES

- Abello, M. T., Blasco, A., Colell, M. 2011. Could sexually experienced Gorillas *Gorilla gorilla gorilla* help hand-reared Gorillas to breed successfully? *International Zoo Yearbook* 45: 237–249.
- Abramov, A. V. 2000. A taxonomic review of the genus *Mustela* (Mammalia, Carnivora). *Zoosystematica Rossica* 8:357–364.
- Ahloy Dallaire, J., Mason, G. J. 2017. Juvenile rough-and-tumble play predicts adult sexual behaviour in American mink. *Animal Behaviour* 123: 81–89.
- Amstislavsky, S., Lindeberg, H., Ternovskaya, Y., et al. 2009. Reproduction in the European mink, *Mustela lutreola*: Oestrous cyclicity and early pregnancy. *Reproduction in Domestic Animals* 44: 489–498.
- Ayers, C. R., Belant, J. L., Eads, D. A., Jachowski, D. S., Millspaugh, J. J. 2014. Investigation of factors affecting black-footed ferret litter size. *Western North American Naturalist* 74: 108–115.
- Bautista, A., García-Torres, E., Prager, G., Hudson, R., Rödel, H. G. 2010. Development of behavior in the litter huddle in rat pups: Within- and between-litter differences. *Developmental Psychobiology* 52: 35–43.
- Biggins, D.E., Schroeder, M.H., 1988. Historical and present status of the black-footed ferret. In: Uresk, D. W., Schenbeck, G. L., Cefkin, R. (Eds.), Eighth Great Plains Wildlife Damage Control Workshop Proceedings. USDA For. Ser. Gen. Tech. Rep. RM-154. Fort Col-lins, Colorado, pp. 93–97.
- Bortner, R. 2017. US Fish and Wildlife Service, Black-footed Ferret Recovery Program. January 16, 2017. Black-footed ferret Managed Care Operations Manual, 288 pp.
- Canessa, S., Converse, S. J., West, M., et al. 2016. Planning for *ex situ* conservation in the face of uncertainty. *Conservation Biology* 30: 599–609.
- Carlstead, K., Fraser, J., Bennett, C., Kleiman, D. G. 1999. Black rhinoceros (*Diceros bicornis*) in US zoos: II. Behavior, breeding success, and mortality in relation to housing facilities. *Zoo Biology* 18: 35–52.
- Chang, M. C., Yanagimachi, R. 1963. Fertilization of ferret ova by deposition of epididymal sperm into the ovarian capsule with special reference to the fertilizable life of ova and the capacitation of sperm. *Journal of Experimental Zoology* 154: 175–187.
- Chang, M. C. 1968. Reciprocal insemination and egg transfer between ferrets and mink. *Journal of Experimental Zoology* 168: 49–59.
- Cheng, Q., Qu, J., Zhang, H., Zhang, Y. 2017. Personality variation and adaptation to the environment of the plateau pika (*Ochotona curzoniae*). *Acta Theriologica Sinica* 37: 115–123.
- Clark, T. W., Richardson, L., Forrest, S. C., Casey, D. E.III, T. M. C. 1986. Descriptive ethology and activity patterns of black-footed ferrets. *Great Basin Naturalist Memoir* 8: 115–134.
- Clark, T. W. 1987. Black-footed ferret recovery a progress report. *Conservation Biology* 1: 8–11.
- Clubb, R., Mason, G. J. 2007. Natural behavioural biology as a risk factor in carnivore welfare: How analysing species differences could help zoos improve enclosures. *Applied Animal Behaviour Science* 102: 303–328.
- Coelho, C. M., Schetini de Azevedo, C., Young, R. J. 2012. Behavioral responses of maned wolves (*Chrysocyon brachyurus*, *Canidae*) to different categories of

- environmental enrichment stimuli and their implications for successful reintroduction. *Zoo Biology* 31: 453–469.
- Conde, D. A., Flesness, N., Colchero, F., Jones, O. R., Scheuerlein, A. 2011. An emerging role of zoos to conserve biodiversity. *Science* 331: 1390–1391.
- Crews, D., Rushworth, D., Gonzalez-Lima, F., Ogawa, S. 2009. Litter environment affects behavior and brain metabolic activity of adult knockout mice. *Frontiers in Behavioral Neuroscience* 3: 12.
- Danilov, P. I., Tumanov, I. L. 1972. The male reproductive cycles of some Mustelidae. *Zoologicheskii Zhurnal*, 51: 781– 880 (in Russian).
- Diez-Leon, M., Bowman, J., Bursian, S., et al. 2013. Environmentally enriched male mink gain more copulations than stereotypic, barren-reared competitors. *PLoS ONE* 8(11), e80494.
- Dolman, P. M., Collar, N. J., Scotland, K. M., Burnside, R. J. 2015. Ark or park: The need to predict relative effectiveness of ex situ and in situ conservation before attempting captive breeding. *Journal of Applied Ecology* 52: 841–850.
- Festl, W., Bodenstern, C., Seebass, C. 2003. Captive breeding of european mink (*Mustela lutreola*; Linnè, 1761) Effects of keeping methods and stress factors on the reproduction success. In: International Conference on the Conservation of European Mink (*Mustela lutreola*). Logroño Spain, Proceedings Book. Gobierno de la Rioja, pp. 85–94.
- Fournier-Chambrillon, C., Bifulchi, A., Mazzola-Rossi, E., et al. 2010. Reliability of stained placental scar counts in farmed American mink and application to free-ranging mustelids. *Journal of Mammalogy* 91: 818–826.
- Frankham, R. 2010. Challenges and opportunities of genetic approaches to biological conservation. *Biological Conservation* 143: 1919–1927.
- Frynta, D., Fraňková, M., Čížková, B., et al. 2011. Social and life history correlates of litter size in captive colonies of precocial spiny mice (*Acomys*). *Acta Theriologica* 56: 289–295.
- García-Díaz, P., Lizana, M. 2013. Reproductive aspects of American minks (*Neovison vison*) in central Spain: Testing the effects of prey availability. *Mammalian Biology* 78: 111–117.
- Gese, E. M., Roberts, B. M., Knowlton, F. F. 2016. Nutritional effects on reproductive performance of captive adult female coyotes (*Canis latrans*). *Animal Reproduction Science* 165: 69–75.
- Gosling, S. D. 2001. From mice to men: What can we learn about personality from animal research? *Psychological Bulletin* 127: 45–86.
- Harrington, L. A., Pödra, M., Macdonald, D. W., Maran, T. 2014. Post-release movements of captive-born European mink *Mustela lutreola*. *Endangered Species Research* 24: 137–148.
- Hoffmann, M., Hilton-Taylor, C., Angulo, A., et al. 2010. The impact of conservation on the status of the world's vertebrates. *Science* 330: 1503–1509.
- Howard, J. G., Wolf, K., Vargas, A., et al. 1997. Enhanced reproductive efficiency and pregnancies after artificial insemination in black-footed ferrets. *Proceedings of the Annual Meeting of the American Association of Zoo Veterinarians*, pp. 351–352.
- Howard, J. G., Lynch, C., Santymire, R. M., Marinari, P. E., Wildt, D. E. 2016. Recovery of gene diversity using long-term cryopreserved spermatozoa and artificial insemination in the endangered black-footed ferret. *Animal Conservation* 19: 102–111.

- Hudson, R., Bautista, A., Reyes-Meza, V., Montor, J. M., Rödel, H. G. 2011. The effect of siblings on early development: A potential contributor to personality differences in mammals. *Developmental Psychobiology* 53: 564–574.
- IUCN 2017. The IUCN Red List of Threatened Species. Version 2017–2. <<http://www.iucnredlist.org>>. Downloaded on 14 September 2017.
- Kanda, L. L., Abdulhay, A., Erickson, C. 2017. Adult wheel access interaction with activity and boldness personality in Siberian dwarf hamsters (*Phodopus sungorus*). *Behavioural Processes* 138: 82–90.
- Kierulff, M. C. M., Rylands, A. B., de Oliveira, M. M. 2008. *Leontopithecus rosalia*. The IUCN Red List of Threatened Species 2008: e.T11506A3287321. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T11506A3287321.en>. Downloaded on 23 October 2017.
- Koivula, M., Strandén, I., Mäntysaari, E. A. 2010. Genetic and phenotypic parameters of age at first mating, litter size and animal size in Finnish mink. *Animal* 4: 183–188.
- Kroshko, J., Clubb, R., Harper, L., Mellor, E., Moehrensclager, A., Mason, G. 2016. Stereotypic route tracing in captive Carnivora is predicted by species-typical home range sizes and hunting styles. *Animal Behaviour* 117: 197–209.
- Lehmkuhl Noer, C., Kjær Needham, E., Wiese, A. S., Johannes Skovbjerg Balsby, T., Dabelsteen, T. 2016. Personality matters: Consistency of inter-individual variation in shyness-boldness across non-breeding and pre-breeding season despite a fall in general shyness levels in farmed American mink (*Neovison vison*). *Applied Animal Behaviour Science* 181: 191–199.
- Macdonald, D. W., Harrington, L. A. 2003. The american mink: The triumph and tragedy of adaptation out of context. *New Zealand Journal of Zoology* 30: 421–441.
- MacKinnon, K., Newberry, R., Wielebnowski, N., et al. 2008. Behavior and corticosteroids predict breeding success in the clouded leopard In: Herrick, J., Fletchall, N. & Swanson, W., editors. *Felid Taxon Advisory Group (TAG), Association of Zoos & Aquariums, 2008 Annual Report* 32–33.
- Maran, T. 1991. Distribution of the European mink, *Mustela lutreola*, in Estonia. *Folia Theriologica Estonica* 1: 1–17.
- Maran, T., Henttonen, H. 1995. Why is the European mink (*Mustela lutreola*) disappearing? – A review of the process and hypotheses. *Annales Zoologici Fennici* 32: 47–54.
- Maran, T., Kruuk, H., Macdonald, D. W., Polma, M. 1998. Diet of two species of mink in Estonia: Displacement of *Mustela lutreola* by *M. vison*. *Journal of Zoology* 245: 218–222.
- Maran, T. 2003. European mink: setting of goal for conservation and the Estonian case study. *Galemys* 15: 1–11.
- Maran, T. 2007. Conservation biology of the European mink, *Mustela lutreola* (Linnaeus 1761): Decline and causes of extinction. Tallinn University Dissertations on Natural Sciences 15, Tallinn University Press.
- Maran, T., Põdra, M., Põlma, M., Macdonald, D. W. 2009. The survival of captive-born animals in restoration programmes – Case study of the endangered European mink *Mustela lutreola*. *Biological Conservation* 142: 1685–1692.
- Maran, T., Pitsal, S., Kotsur, G., Põdra, M. 2016a. Euroopa naaritsa (*Mustela lutreola*) seirearuanne. Hiiumaa – 2016.

- Maran, T., Skumatov, D., Gomez, A., Põdra, M., Abramov, A.V., Dinets, V. 2016b. *Mustela lutreola*. The IUCN Red List of Threatened Species 2016: e.T14018A45199861. <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T14018A45199861.en>. Downloaded on 05 October 2017.
- Maran, T., Podra, M., Harrington, L., A., Macdonald, D., W. 2017. European mink – restoration attempts for a species on the brink of extinction. In: *Biology and Conservation of Musteloids*. Edited by Macdonald, D., W., Newman, C. & Harrington, L., A. Oxford University Press. 370–388.
- Marinari, P. E., Kreeger, J. S., Roelle, J., Miller, B., Godbey, J., Biggins, D. 2006. An adaptive management approach for black-footed ferrets in captivity. In *Recovery of the Black-footed Ferret: Progress and Continuing Challenges*. Proceedings of the Symposium on the Status of the Black-footed Ferret and Its Habitat. Edited by Roelle, J. E., Miller, B. J., Godbey, J. L. and Biggins, D. E. Fort Collins, Colo., 28–29 January 2004. 23–27.
- Martin, M. S., Shepherdson, D. J. 2012. Role of familiarity and preference in reproductive success in *ex situ* breeding programs. *Conservation Biology* 26: 649–656.
- Martin-Wintle, M. S., Shepherdson, D., Zhang, G., Zhang, H., Li, D., Zhou, X., Li, R., Swaisgood, R. R. 2015. Free mate choice enhances conservation breeding in the endangered giant panda. *Nature Communications*, 6. doi: 10.1038/ncomms10125.
- Macdonald, R. A., Harris, S. 2002. Population biology of stoats *Mustela erminea* and weasels *Mustela nivalis* on game estates in Great Britain. *Journal of Applied Ecology* 39: 793–805.
- McDougall, P. T., Réale, D., Sol, D., Reader, S. M. 2006. Wildlife conservation and animal temperament: Causes and consequences of evolutionary change for captive, reintroduced, and wild populations. *Animal Conservation* 9: 39–48.
- Mead, R. A., Joseph, M. M., Neirinckx, S. 1988. Optimal dose of human chorionic gonadotropin for inducing ovulation in the ferret. *Zoo Biology* 7: 263–267.
- Melero, Y., Robinson, E., Lambin, X. 2015. Density- and age-dependent reproduction partially compensates culling efforts of invasive non-native American mink. *Biological Invasions* 17: 2645–2657.
- Miller, B. J., Anderson, S. H., DonCarlos, M. W., Thorne, E. T. 1988. Biology of the endangered black-footed ferret and the role of captive propagation in its conservation. *Canadian Journal of Zoology* 66: 765–773.
- Morgan, K. N., Tromborg, C. T. 2007. Sources of stress in captivity. *Applied Animal Behaviour Science* 102: 262–302.
- Nogueira, S. S. C., Soledade, J. P., Pompéia, S., Nogueira-Filho, S. L. G. 2011. The effect of environmental enrichment on play behaviour in white-lipped peccaries (*Tayassu pecari*). *Animal Welfare* 20: 505–514.
- Novikov, G. A. 1939: The European mink. *Izd. Leningradskogo Gos. Univ., Leningrad* (in Russian).
- Nowak, R., M. 2004. Walker's Carnivores of the World. The Johns Hopkins University Press 141–150.
- Olsen, D. 2012. Introduction of Asiatic golden cat *Catopuma temminckii* at Taronga Conservation Society Australia. *International Zoo Yearbook* 46: 201–208.
- O'Regan, H. J., Kitchener, A. C. 2005. The effects of captivity on the morphology of captive, domesticated and feral mammals. *Mammal Review* 35, 215–230.
- Palomares, F., López-Bao, J. V., Telletxea, G., et al. 2017. Activity and home range in a recently widespread European mink population in Western Europe. *European Journal of Wildlife Research* 63 (5), 78.



- Peng, J., Jiang, Z., Lu, X., et al. 2009. Mate preference and sexual selection in giant panda, *Ailuropoda melanoleuca* in captivity. *Folia Zoologica* 58: 409–415.
- Persson, J. 2005. Female wolverine (*Gulo gulo*) reproduction: Reproductive costs and winter food availability. *Canadian Journal of Zoology* 83: 1453–1459.
- Poessel, S. A., Biggins, D. E., Santymire, R. M., Livieri, T. M., Crooks, K. R., Angeloni, L. 2011. Environmental enrichment affects adrenocortical stress responses in the endangered black-footed ferret. *General and Comparative Endocrinology* 172: 526–533.
- Poole, T. B. 1966. Aggressive play in polecats. *Symposia of the Zoological Society of London* 18: 23–44.
- Poole, T. B. 1967. Aspects of Aggressive Behaviour in Polecats. *Zeitschrift für Tierpsychologie* 24: 351–369.
- Poole, T. B. 1978. An analysis of social play in polecats (*Mustelidae*) with comments on the form and evolutionary history of the open mouth play face. *Animal Behaviour* 26: 36–39.
- Price, E. O. 1999. Behavioral development in animals undergoing domestication. *Applied Animal Behaviour Science*, 65, 245–271.
- Primack, B. R., Sher, A., A. 2016. Bringing species back from the brink. In: Meyers, R., editors. *An Introduction to Conservation Biology*. Sinauer Associates, Inc. 234–263.
- Põdra, M., Gómez, A., Palazón, S. 2013a. Do American mink kill European mink? Cautionary message for future recovery efforts. *European Journal of Wildlife Research* 59: 431–440.
- Põdra, M., Maran, T., Sidorovich, V. E., Johnson, P. J., Macdonald, D. W. 2013b. Restoration programmes and the development of a natural diet: A case study of captive-bred European mink. *European Journal of Wildlife Research* 59: 93–104.
- Quirke, T., O’Riordan, R. M., Zuur, A. 2012. Factors influencing the prevalence of stereotypical behaviour in captive cheetahs (*Acinonyx jubatus*). *Applied Animal Behaviour Science* 142: 189–197.
- Reading, R. P., Clark, T. W., Vargas, A., Hanebury, L. R., Miller B. J., Biggins, D. 1996. Recent directions in black-footed ferret recovery. *Endangered Species Update* 13: 1–6.
- Rozhnov, V. V., Chernova, I. E., Naidenko, S. V. 2008. Approaches to pregnancy diagnosis in the sable (*Martes zibellina*, Mustelidae, Carnivora) by noninvasive methods: Postimplantation period. *Biology Bulletin* 35: 615–618.
- Rödel, H. G., von Holst, D. 2009. Features of the early juvenile development predict competitive performance in male European rabbits. *Physiology and Behavior* 97: 495–502.
- Salkeld, D. J. 2017. Vaccines for Conservation: Plague, Prairie Dogs & Black-Footed Ferrets as a Case Study. *EcoHealth* 14:3, 432–437.
- Santymire, R. M., Lavin, S. R., Branvold-Faber, H., Kreeger, J., Marinari, P. 2015. Effect of dietary vitamin E and prey supplementation on semen quality in male black-footed ferrets (*Mustela nigripes*). *Theriogenology* 84: 217–225.
- Sidorovich, V., Kruuk, H., Macdonald, D. W., Maran, T. 1998. Diets of semi-aquatic carnivores in northern Belarus, with implications for population changes. Cambridge University Press
- Sidorovich, V., Kruuk, H., Macdonald, D. W. 1999. Body size, and interactions between European and american mink (*Mustela lutreola* and *M. vison*) in Eastern Europe. *Journal of Zoology* 248: 521–527.

- Sidorovich, V., Macdonald, D. W. 2001. Density dynamics and changes in habitat use by the European mink and other native mustelids in connection with the American mink expansion in Belarus. *Netherlands Journal of Zoology* 51: 107–126.
- Sidorovich, V. E., Polozov, A. G., Zalewski, A. 2010. Food niche variation of European and American mink during the American mink invasion in north-eastern Belarus. *Biological Invasions* 12: 2207–2217.
- Sih, A., Bell, A., Johnson, J. C. 2004. Behavioral syndromes: An ecological and evolutionary overview. *Trends in Ecology and Evolution* 19: 372–378.
- Snyder, N. F. R., Derrickson, S. R., Beissinger, S. R., et al. 1996. Limitations of captive breeding in endangered species recovery. *Conservation Biology* 10: 338–348.
- Soorae, P. S. (ed.) 2011. *Global Re-introduction Perspectives: More case studies from around the globe*. Gland, Switzerland: IUCN/SSC Re-introduction Specialist Group and Abu Dhabi, UAE: Environment Agency-Abu Dhabi xiv + 250 pp.
- Stenson, G. B. 1988. Oestrus and the vaginal smear cycle of the river otter, *Lutra canadensis*. *Journal of Reproduction and Fertility* 83: 605–610.
- Zhang, G., Swaisgood, R. R., Zhang, H. 2004. Evaluation of behavioral factors influencing reproductive success and failure in captive giant pandas. *Zoo Biology* 23: 15–31.
- Talbot, S., Freire, R., Wassens, S. 2014. Effect of captivity and management on behaviour of the domestic ferret (*Mustela putorius furo*). *Applied Animal Behaviour Science* 151: 94–101.
- Temple, H. J., Terry, A. 2007. *The Status and Distribution of European Mammals*. Office for Official Publications of the European Communities, Luxembourg.
- Ternovskii, D. V. 1977. *The Biology of the Mustelidae*. Akademii Nauk, Novosibirsk.
- Vargas, A., Anderson, S. H. 1998. Black-footed ferret (*Mustela nigripes*) behavioral development: Aboveground activity and juvenile play. *Journal of Ethology* 16: 29–41.
- Wielebnowski, N. C., Fletchall, N., Carlstead, K., Busso, J. M., Brown, J. L. 2002. Non-invasive assessment of adrenal activity associated with husbandry and behavioral factors in the North American clouded leopard population. *Zoo Biology* 21: 77–98.
- Williams, E. S., Thorne, E. T., Kwiatkowski, D. R., Lutz, K., Anderson, S. L. 1992. Comparative vaginal cytology of the estrous cycle of black-footed ferrets (*Mustela nigripes*), Siberian polecats (*M. eversmanni*), and domestic ferrets (*M. putorius furo*). *Journal of veterinary diagnostic investigation: official publication of the American Association of Veterinary Laboratory Diagnosticians, Inc* 4: 38–44.
- Wolf, K. N., Wildt, D. E., Vargas, A., Marinari, P. E., Ottinger, M. A., Howard, J. G. 2000. Reproductive inefficiency in male black-footed ferrets (*Mustela nigripes*). *Zoo Biology* 19: 517–528.
- Xia, C., Cao, J., Zhang, H., Gao, X., Yang, W., Blank, D. 2014. Reintroduction of Przewalski's horse (*Equus ferus przewalskii*) in Xinjiang, China: The status and experience. *Biological Conservation* 177: 142–147.
- Youngman, P. M. 1990. *Mustela lutreola*. *Mammalian Species* 362: 1–3.

## **PUBLICATIONS**

## CURRICULUM VITAE

**Name:** Kairi Kiik  
**Date of birth:** 18.07.1985  
**Citizenship:** Estonian  
**Address:** Department of Zoology, Institute of Ecology and Earth Sciences, Vanemuise 46, 51014 Tartu, Estonia  
**E-mail:** kairi.kiik@ut.ee

**Education:**  
2011–2018 University of Tartu,  
doctoral studies in Zoology and Hydrobiology  
2009–2011 University of Tartu,  
master's degree in Zoology and Hydrobiology  
2005–2009 University of Tartu, bachelor's degree in Biology  
1992–2004 Suure-Jaani Gymnasium

**Language skills:**  
Estonian (native), English (fluent), German (advanced), Russian (beginner)

**Work in non-governmental organisations:**  
2017 Estonian Theriological Society, organizing the 10th Baltic  
Theriological Conference  
2009–2011 Tartu Students Nature Protection circle, board member

**Research interests:**  
*Ex-situ* conservation, social behaviour of mammals

**Publications:**  
**Kiik, K.**, Maran, T., Nagl, A., Ashford, K., Tammaru, T. 2013. The causes of the low breeding success of European mink (*Mustela lutreola*) in captivity. *Zoo Biology* 32: 387–393.  
Haage, M., Bergvall, U. A., Maran, T., **Kiik, K.**, Angerbjörn, A. 2013. Situation and context impacts the expression of personality: The influence of breeding season and test context. *Behavioural Processes* 100: 103–109.  
Nagl, A., Kneidinger, N., **Kiik, K.**, Lindeberg, H., Maran, T., Schwarzenberger, F. 2015. Noninvasive monitoring of female reproductive hormone metabolites in the endangered European mink (*Mustela lutreola*). *Theriogenology* 84: 1472–1481.  
**Kiik, K.**, Maran, T., Kneidinger, N., Tammaru, T. 2016. Social behaviour of endangered European mink (*Mustela lutreola*) litters in captivity. *Applied Animal Behaviour Science* 182: 61–71.

**Kiik, K.,** Maran, T., Nemvalts, K., Sandre, S. L., Tammaru, T. 2017. Reproductive parameters of critically endangered European mink (*Mustela lutreola*) in captivity. *Animal Reproduction Science* 181: 86–92.

**Conference presentations:**

**Kiik, K.,** Maran, T., Nemvalts, K., Sandre, S.-L., Tammaru, T. “Breeding of European mink (*Mustela lutreola*) in captivity”, 10th Baltic Theriological Conference, Tartu, Eesti, 28–30.09.2017, oral presentation

**Kiik, K.,** Maran, T Nagl, A., Kneidinger, N., Tammaru, T. “Behavioural studies of critically endangered European mink (*Mustela lutreola*) in captivity”, Behaviour 2017 – 35th International Ethological Conference, Estoril, Portugal 30.7–04.08.2017, oral presentation

**Kiik, K.,** Tammaru, T., Kneidinger, N., Maran, T. “Behavioural interactions in European mink, *Mustela lutreola*, captive-bred litter – implications to conservation”, The Nordic Evolutionary Psychology Meeting 2013, 12–13.09.2013, Tartu, Estonia, poster presentation

**Kiik, K.,** Maran, T., “Reproductive parameters of captive European mink (*Mustela lutreola*)”, Conference on the Biology and Conservation of Wild Mustelids, Skunks, Procyonids and Red Pandas, 18 –21.03.2013, Oxford, UK, poster presentation

**Kiik, K.,** Maran, T., Nagl, A., Ashford, K., Tammaru, T. “Breeding success of European Mink (*Mustela lutreola*) in captivity”, The 3<sup>rd</sup> European Congress of Conservation Biology, 'Conservation on the Edge', 28.08.2012–01.09.2012, Glasgow, Scotland, UK, poster presentation

## ELULOOKIRJELDUS

**Nimi:** Kairi Kiik  
**Sünniaeg:** 18.07.1985  
**Kodakonsus:** Eesti  
**Aadress:** Zooloogia osakond, Maateaduste ja Ökoloogia Instituut,  
Vanemuise 46, 51014 Tartu, Eesti  
**E-post:** kairi.kiik@ut.ee

**Haridustee:**  
2011–2018 Tartu Ülikool, Zooloogia ja hüdrobioloogia doktoriõpe  
2009–2011 Tartu Ülikool, Loodusteaduste magister,  
Zooloogia ja hüdrobioloogia  
2005–2009 Tartu Ülikool, Loodusteaduste bakalaureus, Bioloogia  
1992–2004 Suure-Jaani Gümnaasium

**Keelteoskus:**  
eesti keel (emakeel), inglise keel (väga hea), saksa keel (kesktase), vene keel (algtase)

### Töö mittetulundusühingutes:

2017 Eesti Terioloogia Selts, X Balti Terioloogia konverentsi korraldamine  
2009–2011 Tartu Üliõpilaste Looduskaitsering, juhatuse liige

### Peamised uurimisvaldkonnad:

*Ex-situ* liigikaitse, imetajate sotsiaalne käitumine

### Publikatsioonid:

- Kiik, K.**, Maran, T., Nagl, A., Ashford, K., Tammaru, T. 2013. The causes of the low breeding success of european mink (*Mustela lutreola*) in captivity. Zoo Biology 32: 387–393.
- Haage, M., Bergvall, U. A., Maran, T., **Kiik, K.**, Angerbjörn, A. 2013. Situation and context impacts the expression of personality: The influence of breeding season and test context. Behavioural Processes 100: 103–109.
- Nagl, A., Kneidinger, N., **Kiik, K.**, Lindeberg, H., Maran, T., Schwarzenberger, F. 2015. Noninvasive monitoring of female reproductive hormone metabolites in the endangered European mink (*Mustela lutreola*). Theriogenology 84: 1472–1481.
- Kiik, K.**, Maran, T., Kneidinger, N., Tammaru, T. 2016. Social behaviour of endangered European mink (*Mustela lutreola*) litters in captivity. Applied Animal Behaviour Science 182: 61–71.

**Kiik, K.,** Maran, T., Nemvalts, K., Sandre, S. L., Tammaru, T. 2017. Reproductive parameters of critically endangered European mink (*Mustela lutreola*) in captivity. *Animal Reproduction Science* 181: 86–92.

**Konverentsi ettekanded:**

**Kiik, K.,** Maran, T., Nemvalts, K., Sandre, S.-L., Tammaru, T. “Breeding of European mink (*Mustela lutreola*) in captivity”, 10th Baltic Theriological Conference, Tartu, Eesti, 28–30.09.2017, suuline ettekanne

**Kiik, K.,** Maran, T Nagl, A., Kneidinger, N., Tammaru, T. “Behavioural studies of critically endangered European mink (*Mustela lutreola*) in captivity”, Behaviour 2017 – 35th International Ethological Conference, Estoril, Portugal 30.07–04.08.2017, suuline ettekanne

**Kiik, K.,** Tammaru, T., Kneidinger, N., Maran, T. “Behavioural interactions in European mink, *Mustela lutreola*, captive bred litter – implications to conservation”, The Nordic Evolutionary Psychology Meeting 2013, 12–13.09.2013, Tartu, Estonia, stendiettekanne

**Kiik, K.,** Maran, T., “Reproductive parameters of captive European mink (*Mustela lutreola*)”, Conference on the Biology and Conservation of Wild Mustelids, Skunks, Procyonids and Red Pandas, 18 –21.03.2013, Oxford, UK, stendiettekanne

**Kiik, K.,** Maran, T., Nagl, A., Ashford, K., Tammaru, T. “Breeding success of European Mink (*Mustela lutreola*) in captivity”, The 3<sup>rd</sup> European Congress of Conservation Biology, ‘Conservation on the Edge’, 28.08.2012–01.09.2012, Glasgow, Scotland, UK, stendiettekanne

## DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käärd.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.



20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand**. Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa**. Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa**. Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik**. Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo**. Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo**. Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots**. Health state indices of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero**. Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees**. Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks**. Cholecystokinin (CCK) – induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and serotonin. Tartu, 1999, 123 p.
52. **Ebe Sild**. Impact of increasing concentrations of O<sub>3</sub> and CO<sub>2</sub> on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva**. Electron microscopical analysis of the synaptone-mal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna**. Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro**. Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane**. Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm**. Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg**. Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild**. The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu**. Studies of the TOL plasmid transcription factor XylS. Tartu, 2000, 88 p.
61. **Dina Lepik**. Modulation of viral DNA replication by tumor suppressor protein p53. Tartu, 2000, 106 p.

62. **Kai Vellak**. Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu, 2000, 122 p.
63. **Jonne Kotta**. Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu, 2000, 160 p.
64. **Georg Martin**. Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000, 139 p.
65. **Silvia Sepp**. Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira**. On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000, 96 p.
67. **Priit Zingel**. The role of planktonic ciliates in lake ecosystems. Tartu, 2001, 111 p.
68. **Tiit Teder**. Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu, 2001, 122 p.
69. **Hannes Kollist**. Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu, 2001, 80 p.
70. **Reet Marits**. Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu, 2001, 112 p.
71. **Vallo Tilgar**. Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002, 126 p.
72. **Rita Hõrak**. Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002, 108 p.
73. **Liina Eek-Piirsoo**. The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002, 74 p.
74. **Krõõt Aasamaa**. Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002, 110 p.
75. **Nele Ingerpuu**. Bryophyte diversity and vascular plants. Tartu, 2002, 112 p.
76. **Neeme Tõnisson**. Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002, 124 p.
77. **Margus Pensa**. Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003, 110 p.
78. **Asko Lõhmus**. Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003, 168 p.
79. **Viljar Jaks**. p53 – a switch in cellular circuit. Tartu, 2003, 160 p.
80. **Jaana Männik**. Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003, 140 p.
81. **Marek Sammul**. Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003, 159 p.
82. **Ivar Ilves**. Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003, 89 p.

83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003, 109 p.
84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003, 158 p.
85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003, 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004, 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004, 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004, 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004, 117 p.
90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004, 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004, 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004, 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004, 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004, 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004, 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004, 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004, 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004, 103 p.
99. **Mikk Heidemaa.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004, 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N<sub>2</sub> fixation in some Estonian lakes. Tartu, 2004, 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004, 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004, 142 p.
103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.

104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005, 100 p.
106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005, 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005, 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005, 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005, 121 p.
110. **Juhan Javois.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005, 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005, 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005, 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005, 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006, 124 p.
116. **Priit Kupper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006, 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006, 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006, 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006, 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006, 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006, 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007, 123 p.

125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007, 143 p.
126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007, 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007, 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007, 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007, 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007, 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007, 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007, 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007, 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in European populations. Tartu, 2007, 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007, 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007, 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transporters. Tartu, 2008, 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008, 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008, 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008, 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008, 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008, 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.

146. **Lauri Saag**. Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
147. **Ulvi Karu**. Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm**. Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks**. Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu**. Acclimation of stomatal structure and function in tree canopy: effect of light and CO<sub>2</sub> concentration. Tartu, 2008, 108 p.
151. **Janne Pullat**. Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš**. Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko**. Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast**. Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats**. Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova**. The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik**. Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver**. Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja**. Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast**. Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak**. Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Remmel**. Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe**. Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe**. Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.

165. **Liisa Metsamaa**. Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
166. **Pille Säälük**. The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil**. Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik**. Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark**. Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap**. Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan**. Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe**. Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi**. Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson**. Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts**. Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis**. Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov**. Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster**. Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap**. Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar**. Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül**. Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk**. Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü**. Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla**. Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.



185. **Gyaneshwer Chaubey**. The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp**. Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
187. **Virve Sõber**. The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro**. The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold**. Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert**. Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu**. Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik**. ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber**. Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper**. Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak**. Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo**. Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel**. Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus**. Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius**. Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värvi**. Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Välik**. Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe**. Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht**. Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.

205. **Teele Jairus**. Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov**. PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.
207. **Marina Grigороva**. Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar**. The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild**. Oxidative defences in immunoecological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar**. The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag**. Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla**. Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve**. Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler**. The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova**. Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar**. Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp**. Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero**. Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram**. Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Annely Lorents**. Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik**. Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prouš**. Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.
223. **Triinu Visnapuu**. Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.

224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.
225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Placental gene expression in normal and complicated pregnancy. Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.
243. **Merli Pärnoja.** Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus.** Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.

245. **Pille Mänd**. Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants. Tartu, 2013, 128 p.
246. **Mario Plaas**. Animal model of Wolfram Syndrome in mice: behavioural, biochemical and psychopharmacological characterization. Tartu, 2013, 144 p.
247. **Georgi Hudjašov**. Maps of mitochondrial DNA, Y-chromosome and tyrosinase variation in Eurasian and Oceanian populations. Tartu, 2013, 115 p.
248. **Mari Lepik**. Plasticity to light in herbaceous plants and its importance for community structure and diversity. Tartu, 2013, 102 p.
249. **Ede Leppik**. Diversity of lichens in semi-natural habitats of Estonia. Tartu, 2013, 151 p.
250. **Ülle Saks**. Arbuscular mycorrhizal fungal diversity patterns in boreo-nemoral forest ecosystems. Tartu, 2013, 151 p.
251. **Eneli Oitmaa**. Development of arrayed primer extension microarray assays for molecular diagnostic applications. Tartu, 2013, 147 p.
252. **Jekaterina Jutkina**. The horizontal gene pool for aromatics degradation: bacterial catabolic plasmids of the Baltic Sea aquatic system. Tartu, 2013, 121 p.
253. **Helen Vellau**. Reaction norms for size and age at maturity in insects: rules and exceptions. Tartu, 2014, 132 p.
254. **Randel Kreitsberg**. Using biomarkers in assessment of environmental contamination in fish – new perspectives. Tartu, 2014, 107 p.
255. **Krista Takkis**. Changes in plant species richness and population performance in response to habitat loss and fragmentation. Tartu, 2014, 141 p.
256. **Liina Nagirnaja**. Global and fine-scale genetic determinants of recurrent pregnancy loss. Tartu, 2014, 211 p.
257. **Triin Triisberg**. Factors influencing the re-vegetation of abandoned extracted peatlands in Estonia. Tartu, 2014, 133 p.
258. **Villu Soon**. A phylogenetic revision of the *Chrysis ignita* species group (Hymenoptera: Chrysididae) with emphasis on the northern European fauna. Tartu, 2014, 211 p.
259. **Andrei Nikonov**. RNA-Dependent RNA Polymerase Activity as a Basis for the Detection of Positive-Strand RNA Viruses by Vertebrate Host Cells. Tartu, 2014, 207 p.
260. **Eele Õunapuu-Pikas**. Spatio-temporal variability of leaf hydraulic conductance in woody plants: ecophysiological consequences. Tartu, 2014, 135 p.
261. **Marju Männiste**. Physiological ecology of greenfinches: information content of feathers in relation to immune function and behavior. Tartu, 2014, 121 p.
262. **Katre Kets**. Effects of elevated concentrations of CO<sub>2</sub> and O<sub>3</sub> on leaf photosynthetic parameters in *Populus tremuloides*: diurnal, seasonal and interannual patterns. Tartu, 2014, 115 p.

263. **Küllil Lokko**. Seasonal and spatial variability of zoopsammon communities in relation to environmental parameters. Tartu, 2014, 129 p.
264. **Olga Žilina**. Chromosomal microarray analysis as diagnostic tool: Estonian experience. Tartu, 2014, 152 p.
265. **Kertu Lõhmus**. Colonisation ecology of forest-dwelling vascular plants and the conservation value of rural manor parks. Tartu, 2014, 111 p.
266. **Anu Aun**. Mitochondria as integral modulators of cellular signaling. Tartu, 2014, 167 p.
267. **Chandana Basu Mallick**. Genetics of adaptive traits and gender-specific demographic processes in South Asian populations. Tartu, 2014, 160 p.
268. **Riin Tamme**. The relationship between small-scale environmental heterogeneity and plant species diversity. Tartu, 2014, 130 p.
269. **Liina Remm**. Impacts of forest drainage on biodiversity and habitat quality: implications for sustainable management and conservation. Tartu, 2015, 126 p.
270. **Tiina Talve**. Genetic diversity and taxonomy within the genus *Rhinanthus*. Tartu, 2015, 106 p.
271. **Mehis Rohla**. Otolith sclerochronological studies on migrations, spawning habitat preferences and age of freshwater fishes inhabiting the Baltic Sea. Tartu, 2015, 137 p.
272. **Alexey Reshchikov**. The world fauna of the genus *Lathrolestes* (Hymenoptera, Ichneumonidae). Tartu, 2015, 247 p.
273. **Martin Pook**. Studies on artificial and extracellular matrix protein-rich surfaces as regulators of cell growth and differentiation. Tartu, 2015, 142 p.
274. **Mai Kukumägi**. Factors affecting soil respiration and its components in silver birch and Norway spruce stands. Tartu, 2015, 155 p.
275. **Helen Karu**. Development of ecosystems under human activity in the North-East Estonian industrial region: forests on post-mining sites and bogs. Tartu, 2015, 152 p.
276. **Hedi Peterson**. Exploiting high-throughput data for establishing relationships between genes. Tartu, 2015, 186 p.
277. **Priit Adler**. Analysis and visualisation of large scale microarray data, Tartu, 2015, 126 p.
278. **Aigar Niglas**. Effects of environmental factors on gas exchange in deciduous trees: focus on photosynthetic water-use efficiency. Tartu, 2015, 152 p.
279. **Silja Laht**. Classification and identification of conopeptides using profile hidden Markov models and position-specific scoring matrices. Tartu, 2015, 100 p.
280. **Martin Kesler**. Biological characteristics and restoration of Atlantic salmon *Salmo salar* populations in the Rivers of Northern Estonia. Tartu, 2015, 97 p.
281. **Pratyush Kumar Das**. Biochemical perspective on alphaviral nonstructural protein 2: a tale from multiple domains to enzymatic profiling. Tartu, 2015, 205 p.

282. **Priit Palta**. Computational methods for DNA copy number detection. Tartu, 2015, 130 p.
283. **Julia Sidorenko**. Combating DNA damage and maintenance of genome integrity in pseudomonads. Tartu, 2015, 174 p.
284. **Anastasiia Kovtun-Kante**. Charophytes of Estonian inland and coastal waters: distribution and environmental preferences. Tartu, 2015, 97 p.
285. **Ly Lindman**. The ecology of protected butterfly species in Estonia. Tartu, 2015, 171 p.
286. **Jaanis Lodjak**. Association of Insulin-like Growth Factor I and Corticosterone with Nestling Growth and Fledging Success in Wild Passerines. Tartu, 2016, 113 p.
287. **Ann Kraut**. Conservation of Wood-Inhabiting Biodiversity – Semi-Natural Forests as an Opportunity. Tartu, 2016, 141 p.
288. **Tiit Örd**. Functions and regulation of the mammalian pseudokinase TRIB3. Tartu, 2016, 182. p.
289. **Kairi Käiro**. Biological Quality According to Macroinvertebrates in Streams of Estonia (Baltic Ecoregion of Europe): Effects of Human-induced Hydromorphological Changes. Tartu, 2016, 126 p.
290. **Leidi Laurimaa**. *Echinococcus multilocularis* and other zoonotic parasites in Estonian canids. Tartu, 2016, 144 p.
291. **Helerin Margus**. Characterization of cell-penetrating peptide/nucleic acid nanocomplexes and their cell-entry mechanisms. Tartu, 2016, 173 p.
292. **Kadri Runnel**. Fungal targets and tools for forest conservation. Tartu, 2016, 157 p.
293. **Urmo Võsa**. MicroRNAs in disease and health: aberrant regulation in lung cancer and association with genomic variation. Tartu, 2016, 163 p.
294. **Kristina Mäemets-Allas**. Studies on cell growth promoting AKT signaling pathway – a promising anti-cancer drug target. Tartu, 2016, 146 p.
295. **Janeli Viil**. Studies on cellular and molecular mechanisms that drive normal and regenerative processes in the liver and pathological processes in Dupuytren's contracture. Tartu, 2016, 175 p.
296. **Ene Kook**. Genetic diversity and evolution of *Pulmonaria angustifolia* L. and *Myosotis laxa sensu lato* (Boraginaceae). Tartu, 2016, 106 p.
297. **Kadri Peil**. RNA polymerase II-dependent transcription elongation in *Saccharomyces cerevisiae*. Tartu, 2016, 113 p.
298. **Katrin Ruisu**. The role of RIC8A in mouse development and its function in cell-matrix adhesion and actin cytoskeletal organisation. Tartu, 2016, 129 p.
299. **Janely Pae**. Translocation of cell-penetrating peptides across biological membranes and interactions with plasma membrane constituents. Tartu, 2016, 126 p.
300. **Argo Ronk**. Plant diversity patterns across Europe: observed and dark diversity. Tartu, 2016, 153 p.

301. **Kristiina Mark.** Diversification and species delimitation of lichenized fungi in selected groups of the family Parmeliaceae (Ascomycota). Tartu, 2016, 181 p.
302. **Jaak-Albert Metsoja.** Vegetation dynamics in floodplain meadows: influence of mowing and sediment application. Tartu, 2016, 140 p.
303. **Hedvig Tamman.** The GraTA toxin-antitoxin system of *Pseudomonas putida*: regulation and role in stress tolerance. Tartu, 2016, 154 p.
304. **Kadri Pärtel.** Application of ultrastructural and molecular data in the taxonomy of helotialean fungi. Tartu, 2016, 183 p.
305. **Maris Hindrikson.** Grey wolf (*Canis lupus*) populations in Estonia and Europe: genetic diversity, population structure and -processes, and hybridization between wolves and dogs. Tartu, 2016, 121 p.
306. **Polina Degtjarenko.** Impacts of alkaline dust pollution on biodiversity of plants and lichens: from communities to genetic diversity. Tartu, 2016, 126 p.
307. **Liina Pajusalu.** The effect of CO<sub>2</sub> enrichment on net photosynthesis of macrophytes in a brackish water environment. Tartu, 2016, 126 p.
308. **Stoyan Tankov.** Random walks in the stringent response. Tartu, 2016, 94 p.
309. **Liis Leitsalu.** Communicating genomic research results to population-based biobank participants. Tartu, 2016, 158 p.
310. **Richard Meitern.** Redox physiology of wild birds: validation and application of techniques for detecting oxidative stress. Tartu, 2016, 134 p.
311. **Kaie Lokk.** Comparative genome-wide DNA methylation studies of healthy human tissues and non-small cell lung cancer tissue. Tartu, 2016, 127 p.
312. **Mihhail Kurašin.** Processivity of cellulases and chitinases. Tartu, 2017, 132 p.
313. **Carmen Tali.** Scavenger receptors as a target for nucleic acid delivery with peptide vectors. Tartu, 2017, 155 p.
314. **Katarina Oganjan.** Distribution, feeding and habitat of benthic suspension feeders in a shallow coastal sea. Tartu, 2017, 132 p.
315. **Taavi Paal.** Immigration limitation of forest plants into wooded landscape corridors. Tartu, 2017, 145 p.
316. **Kadri Õunap.** The Williams-Beuren syndrome chromosome region protein WBSCR22 is a ribosome biogenesis factor. Tartu, 2017, 135 p.
317. **Riin Tamm.** In-depth analysis of factors affecting variability in thiopurine methyltransferase activity. Tartu, 2017, 170 p.
318. **Keiu Kask.** The role of RIC8A in the development and regulation of mouse nervous system. Tartu, 2017, 184 p.
319. **Tiia Möller.** Mapping and modelling of the spatial distribution of benthic macrovegetation in the NE Baltic Sea with a special focus on the eelgrass *Zostera marina* Linnaeus, 1753. Tartu, 2017, 162 p.
320. **Silva Kasela.** Genetic regulation of gene expression: detection of tissue- and cell type-specific effects. Tartu, 2017, 150 p.

321. **Karmen Süld.** Food habits, parasites and space use of the raccoon dog *Nyctereutes procyonoides*: the role of an alien species as a predator and vector of zoonotic diseases in Estonia. Tartu, 2017, p.
322. **Ragne Oja.** Consequences of supplementary feeding of wild boar – concern for ground-nesting birds and endoparasite infection. Tartu, 2017, 141 p.
323. **Riin Kont.** The acquisition of cellulose chain by a processive cellobiohydrolase. Tartu, 2017, 117 p.
324. **Liis Kasari.** Plant diversity of semi-natural grasslands: drivers, current status and conservation challenges. Tartu, 2017, 141 p.
325. **Sirgi Saar.** Belowground interactions: the roles of plant genetic relatedness, root exudation and soil legacies. Tartu, 2017, 113 p.
326. **Sten Anslan.** Molecular identification of Collembola and their fungal associates. Tartu, 2017, 125 p.
327. **Imre Taal.** Causes of variation in littoral fish communities of the Eastern Baltic Sea: from community structure to individual life histories. Tartu, 2017, 118 p.
328. **Jürgen Jalak.** Dissecting the Mechanism of Enzymatic Degradation of Cellulose Using Low Molecular Weight Model Substrates. Tartu, 2017, 137 p.