

ON RARELY g -CONTINUOUS FUNCTIONS

MIGUEL CALDAS AND SAEID JAFARI
Universidade Federal Fluminense, Brazil and
College of Vestsjaelland Syd, Denmark

ABSTRACT. Popa introduced the notion of rare continuity. In this paper, we introduce a new class of functions called rarely g -continuous functions and investigate some of its fundamental properties. This type of continuity is a generalization of both rare continuity and weak continuity.

1. INTRODUCTION

Popa [11] introduced the notion of rare continuity as a generalization of weak continuity [8] which has been further investigated by Long and Herrington [10] and Jafari [6] and [7]. Levine [9] introduced the concept of generalized closed sets of a topological space and a class of topological spaces called $T_{1/2}$ -spaces. Dunham [4], Dunham and Levine [5] and Caldas [2] further studied some properties of generalized closed sets and $T_{1/2}$ -spaces.

The purpose of the present paper is to introduce the concept of rare g -continuity in topological spaces as a generalization of rare continuity and weak continuity. We investigate several properties of rarely g -continuous functions. The notion of $I.g$ -continuity is also introduced which is weaker than g -continuity and stronger than rare g -continuity. It is shown that when the codomain of a function is regular, then the notions of rare g -continuity and $I.g$ -continuity are equivalent.

2. PRELIMINARIES

Throughout this paper, X and Y are topological spaces. Recall that a rare set is a set R such that $\text{Int}(R) = \emptyset$. A nowhere dense set, is a set R which $\text{Int}(\text{Cl}(R)) = \emptyset$ if $\text{Cl}(R)$ is codense. Levine [9] introduces the notion of g -closed sets: A set A in X is called g -closed if $\text{Cl}(A) \subset G$ whenever $A \subset G$ and G is

2000 *Mathematics Subject Classification.* 54B05, 54C08, 54D05.

Key words and phrases. Rare set, g -open, rarely continuous, rarely almost compact.

open in X . The complement of a g -closed set is called g -open [9]. The family of all g -open (resp. open) sets will be denoted by $GO(X)$ (resp. $O(X)$). We set $GO(X, x) = \{U \mid x \in U \in GO(X)\}$ and $O(X, x) = \{U \mid x \in U \in O(X)\}$.

DEFINITION 2.1. A function $f : X \rightarrow Y$ is called:

- i) *Weakly continuous* [8] (resp. *weakly- g -continuous* [3]) if for each $x \in X$ and each open set G containing $f(x)$, there exists $U \in O(X, x)$ (resp. $U \in GO(X, x)$) such that $f(U) \subset \text{Cl}(G)$.
- ii) *Rarely continuous* [11] if for each $x \in X$ and each $G \in O(Y, f(x))$, there exist a rare set R_G with $G \cap \text{Cl}(R_G) = \emptyset$ and $U \in O(X, x)$ such that $f(U) \subset G \cup R_G$.
- iii) *g -continuous* [1] if the inverse image of every closed set in Y is g -closed in X .

3. RARE g -CONTINUITY

DEFINITION 3.1. A function $f : X \rightarrow Y$ is called *rarely g -continuous* if for each $x \in X$ and each $G \in O(Y, f(x))$, there exist a rare set R_G with $G \cap \text{Cl}(R_G) = \emptyset$ and $U \in GO(X, x)$ such that $f(U) \subset G \cup R_G$.

EXAMPLE 3.2. Let X and Y be the real line with indiscrete and discrete topologies respectively. The identity function is rarely g -continuous.

Note that, every weakly continuous function is rarely continuous and every rarely continuous function is rarely g -continuous.

Question 1. Is there any example showing that a function is rarely g -continuous but not rarely continuous?

THEOREM 3.3. The following statements are equivalent for a function $f : X \rightarrow Y$:

- (1) The function f is rarely g -continuous at $x \in X$.
- (2) For each set $G \in O(Y, f(x))$, there exists $U \in GO(X, x)$ such that $\text{Int}[f(U) \cap (Y \setminus G)] = \emptyset$.
- (3) For each set $G \in O(Y, f(x))$, there exists $U \in GO(X, x)$ such that $\text{Int}[f(U)] \subset \text{Cl}(G)$.

PROOF. (1) \Rightarrow (2): Let $G \in O(Y, f(x))$. By $f(x) \in G \subset \text{Int}(\text{Cl}(G))$ and the fact that $\text{Int}(\text{Cl}(G)) \in O(Y, f(x))$, there exist a rare set R_G with $\text{Int}(\text{Cl}(G)) \cap \text{Cl}(R_G) = \emptyset$ and a g -open set $U \subset X$ containing x such that $f(U) \subset \text{Int}(\text{Cl}(G)) \cup R_G$. We have $\text{Int}[f(U) \cap (Y - G)] = \text{Int}[f(U)] \cap \text{Int}(Y - G) \subset \text{Int}[\text{Cl}(G) \cup R_G] \cap (Y - \text{Cl}(G)) \subset (\text{Cl}(G) \cup \text{Int}(R_G)) \cap (Y - \text{Cl}(G)) = \emptyset$.

(2) \Rightarrow (3): It is straightforward.

(3) \Rightarrow (1): Let $G \in O(Y, f(x))$. Then by (3), there exists $U \in GO(X, x)$ such that $\text{Int}[f(U)] \subset \text{Cl}(G)$. We have $f(U) = [f(U) - \text{Int}(f(U))] \cup \text{Int}(f(U)) \subset [f(U) - \text{Int}(f(U))] \cup \text{Cl}(G) = [f(U) - \text{Int}(f(U))] \cup G \cup (\text{Cl}(G) - G) = [f(U) - \text{Int}(f(U))] \cap (Y - G) \cup G \cup (\text{Cl}(G) - G)$.

Set $R^* = [f(U) - \text{Int}(f(U))] \cap (Y - G)$ and $R^{**} = (\text{Cl}(G) - G)$. Then R^* and R^{**} are rare sets. More $R_G = R^* \cup R^{**}$ is a rare set such that $\text{Cl}(R_G) \cap G = \emptyset$ and $f(U) \subset G \cup R_G$. This shows that f is rarely g -continuous. \square

We define the following notion which is a new generalization of g -continuity.

DEFINITION 3.4. *A function $f : X \rightarrow Y$ is $I.g$ -continuous at $x \in X$ if for each set $G \in O(Y, f(x))$, there exists $U \in GO(X, x)$ such that $\text{Int}[f(U)] \subset G$.*

If f has this property at each point $x \in X$, then we say that f is $I.g$ -continuous on X .

EXAMPLE 3.5. Let $X = Y = \{a, b, c\}$ and $\tau = \sigma = \{X, \emptyset, \{a\}\}$. Then a function $f : X \rightarrow Y$ defined by $f(a) = f(b) = a$ and $f(c) = c$ is $I.g$ -continuous.

Question 2. Are there examples showing that a function is $I.g$ -continuous but not g -continuous and a function is rarely g -continuous but not $I.g$ -continuous?

REMARK 3.6. Since, if $f : X \rightarrow Y$ is g -continuous, then for each point $x \in X$ and each open set V containing $f(x)$, there exists $U \in GO(X, x)$ such that $f(U) \subset V$ ([1], Proposition 2). Then, it should be noted that $I.g$ -continuity is weaker than g -continuity and stronger than rare g -continuity.

THEOREM 3.7. *Let Y be a regular space. Then the function $f : X \rightarrow Y$ is $I.g$ -continuous on X if and only if f is rarely g -continuous on X .*

PROOF. We prove only the sufficient condition since the necessity condition is evident (Remark 3.6).

Let f be rarely g -continuous on X and $x \in X$. Suppose that $f(x) \in G$, where G is an open set in Y . By the regularity of Y , there exists an open set $G_1 \in O(Y, f(x))$ such that $\text{Cl}(G_1) \subset G$. Since f is rarely g -continuous, then there exists $U \in GO(X, x)$ such that $\text{Int}[f(U)] \subset \text{Cl}(G_1)$ (Theorem 3.3). This implies that $\text{Int}[f(U)] \subset G$ and therefore f is $I.g$ -continuous on X . \square

We say that a function $f : X \rightarrow Y$ is $r.g$ -open if the image of a g -open set is open.

THEOREM 3.8. *If $f : X \rightarrow Y$ is an $r.g$ -open rarely g -continuous function, then f is weakly g -continuous.*

PROOF. Suppose that $x \in X$ and $G \in O(Y, f(x))$. Since f is rarely g -continuous, there exist a rare set R_G with $\text{Cl}(R_G) \cap U = \emptyset$ and $U \in GO(X, x)$ such that $f(U) \subset G \cup R_G$. This means that $(f(U) \cap (Y \setminus \text{Cl}(G))) \subset R_G$. Since the function f is $r.g$ -open, then $f(U) \cap (Y \setminus \text{Cl}(G))$ is open. But the rare set R_G has no interior points. Then $f(U) \cap (Y \setminus \text{Cl}(G)) = \emptyset$. This implies that $f(U) \subset \text{Cl}(G)$ and thus f is weakly g -continuous. \square

THEOREM 3.9. *If $f : X \rightarrow Y$ is rarely g -continuous function, then the graph function $g : X \rightarrow X \times Y$, defined by $g(x) = (x, f(x))$ for every x in X is rarely g -continuous.*

PROOF. Suppose that $x \in X$ and W is any open set containing $g(x)$. It follows that there exist open sets U and V in X and Y , respectively, such that $(x, f(x)) \in U \times V \subset W$. Since f is rarely g -continuous, there exists $G \in GO(X, x)$ such that $\text{Int}[f(G)] \subset \text{Cl}(V)$. Let $E = U \cap G$. It follows that $E \in GO(X, x)$ and we have $\text{Int}[g(E)] \subset \text{Int}(U \times f(G)) \subset U \times \text{Cl}(V) \subset \text{Cl}(W)$. Therefore, g is rarely g -continuous. \square

DEFINITION 3.10. *Let $A = \{G_i\}$ be a class of subsets of X . By rarely union sets [6] of A we mean $\{G_i \cup R_{G_i}\}$, where each R_{G_i} is a rare set such that each of $\{G_i \cap \text{Cl}(R_{G_i})\}$ is empty.*

Recall that, a subset B of X is said to be rarely almost compact relative to X [6] if every open cover of B by open sets of X , there exists a finite subfamily whose rarely union sets cover B .

A topological space X is said to be rarely almost compact [6] if the set X is rarely almost compact relative to X .

A topological space X is called GO -compact [1] if every cover of X by g -open sets has a finite subcover.

Question 3. Characterize the notion of rarely almost compactness. Find example/s to show the relation of compactness and rarely almost compactness.

THEOREM 3.11. *Let $f : X \rightarrow Y$ be rarely g -continuous and K a GO -compact set relative to X . Then $f(K)$ is rarely almost compact subset relative to Y .*

PROOF. Suppose that Ω is an open cover of $f(K)$. Let B be the set of all V in Ω such that $V \cap f(K) \neq \emptyset$. Then B is an open cover of $f(K)$. Hence for each $k \in K$, there is some $V_k \in B$ such that $f(k) \in V_k$. Since f is rarely g -continuous there exist a rare set R_{V_k} with $V_k \cap \text{Cl}(R_{V_k}) = \emptyset$ and a g -open set U_k containing k such that $f(U_k) \subset V_k \cup R_{V_k}$. Hence there is a finite subfamily $\{U_k\}_{k \in \Delta}$ which covers K , where Δ is a finite subset of K . The subfamily $\{V_k \cup R_{V_k}\}_{k \in \Delta}$ also covers $f(K)$. \square

Recall that a space X is called $T_{1/2}$ -space [4] if every g -closed set in X is closed in X .

THEOREM 3.12. *Let $f : X \rightarrow Y$ be rarely g -continuous and X a $T_{1/2}$ -space. Then f is rarely continuous.*

A space X is called a door space if every subset of X is either open or closed.

W. Dunham [[4] Corollary 3.7] proved the following result:

LEMMA 3.13. *A door space is a $T_{1/2}$ -space.*

THEOREM 3.14. *Let $f : X \rightarrow Y$ be a rarely g -continuous and X be a door space. Then f is rarely continuous.*

PROOF. It is an immediate consequence of Lemma 3.13 and Theorem 3.12. \square

LEMMA 3.15 (Long and Herrington [10]). *If $g : Y \rightarrow Z$ is continuous and one-to-one, then g preserves rare sets.*

THEOREM 3.16. *If $f : X \rightarrow Y$ is rarely g -continuous and $g : Y \rightarrow Z$ is continuous and one-to-one, then $g \circ f : X \rightarrow Z$ is rarely g -continuous.*

PROOF. Suppose that $x \in X$ and $(g \circ f)(x) \in V$, where V is an open set in Z . By hypothesis, g is continuous, therefore there exists an open set $G \subset Y$ containing $f(x)$ such that $g(G) \subset V$. Since f is rarely g -continuous, there exist a rare set R_G with $G \cap \text{Cl}(R_G) = \emptyset$ and a g -open set U containing x such that $f(U) \subset G \cup R_G$. It follows from Lemma 3.15 that $g(R_G)$ is a rare set in Z . Since R_G is a subset of $Y \setminus G$ and g is injective, we have $\text{Cl}(g(R_G)) \cap V = \emptyset$. This implies that $(g \circ f)(U) \subset V \cup g(R_G)$. Hence the result. \square

Recall that a function $f : X \rightarrow Y$ is called pre- g -open if $f(U)$ is g -open in Y for every g -open set U of X .

THEOREM 3.17. *Let $f : X \rightarrow Y$ be a pre- g -open surjection and $g : Y \rightarrow Z$ a function such that $g \circ f : X \rightarrow Z$ is rarely g -continuous. Then g is rarely g -continuous.*

PROOF. Let $y \in Y$ and $x \in X$ such that $f(x) = y$. Let $G \in O(Z, (g \circ f)(x))$. Since $g \circ f$ is rarely g -continuous, there exist a rare set R_G with $G \cap \text{Cl}(R_G) = \emptyset$ and $U \in GO(X, x)$ such that $(g \circ f)(U) \subset G \cup R_G$. But $f(U)$ (say V) is a g -open set containing $f(x)$. Therefore, there exist a rare set R_G with $G \cap \text{Cl}(R_G) = \emptyset$ and $V \in GO(Y, y)$ such that $g(V) \subset G \cup R_G$, i.e., g is rarely g -continuous. \square

REFERENCES

- [1] K. Balachandran, P. Sundaram and H. Maki, *On generalized continuous maps in topological spaces*, Mem. Fac. Sci. Kochi Univ. (Series Math.) **12** (1991), 5-13.
- [2] M. Caldas, *On g -closed sets and g -continuous mappings*, Kyungpook Math. J. **33** (1993), 205-209.
- [3] M. Caldas, S. Jafari and T. Noiri, *Properties of weakly g -continuous functions* (under preparation).
- [4] W. Dunham, *$T_{1/2}$ -spaces*, Kyungpook Math. J. **17** (1977), 161-169.
- [5] W. Dunham and N. Levine, *Further results on generalized closed sets in topology*, Kyungpook Math. J. **20** (1980), 164-175.
- [6] S. Jafari, *A note on rarely continuous functions*, Univ. Bacáu. Stud. Cerc. St. Ser. Mat. **5** (1995), 29-34.

- [7] S. Jafari, *On some properties of rarely continuous functions*, Univ. Bacáu. Stud. Cerc. St. Ser. Mat. **7** (1997), 65-73.
- [8] N. Levine, *A decomposition of continuity in topological spaces*, Amer. Math. Monthly **60** (1961), 44-46.
- [9] N. Levine, *Generalized closed sets in topology*, Rend. Circ. Mat. Palermo (2) **19** (1970), 89-96.
- [10] P. E. Long and L. L. Herrington, *Properties of rarely continuous functions*, Glasnik Mat. Ser. III **17(37)** (1982), 147-153.
- [11] V. Popa, *Sur certain decomposition de la continuité dans les espaces topologiques*, Glasnik Mat. Ser. III **14(34)** (1979), 359-362.

M. Caldas

Departamento de Matemática Aplicada
Universidade Federal Fluminense
Rua Mário Santos Braga, s/n - CEP: 24020-140
Niteroi-RJ, Brazil
E-mail: gmamccs@vm.uff.br

S. Jafari

College of Vestsjaelland Syd
Herrestraede 11
4200 Slagelse, Denmark
E-mail: jafari@stofanet.dk

Received: 8.12.2004.

Revised: 12.1.2005.