

PUPPE EXACT SEQUENCE AND ITS APPLICATION IN THE FIBREWISE CATEGORY \mathbf{MAP}

YOSHIFUMI KONAMI AND TAKUO MIWA

Izumo-Nishi High School and Shimane University, Japan

ABSTRACT. In this paper, we study Puppe exact sequence and its application in the fibrewise category \mathbf{MAP} . The application shows that we can prove the generalized formula for the suspension of fibrewise product spaces. Further, introducing an intermediate fibrewise category \mathbf{TOP}_B^H , we give an another proof of the original formula in \mathbf{TOP}_B using the concepts of \mathbf{TOP}_B^H .

1. INTRODUCTION

For a base space B , the category \mathbf{TOP}_B is the fibrewise topology over B . For General Topology of Continuous Maps or Fibrewise General Topology, see B.A. Pasynkov [7],[8]. In [1], D. Buhagiar studied fibrewise topology in the category of all continuous maps, called \mathbf{MAP} by him (as a way of thinking of a category, \mathbf{MAP} can be seen in earlier works, see for example [10]). The study of fibrewise topology in \mathbf{MAP} is a generalization of it in the category \mathbf{TOP}_B . In the previous paper [4], we studied fibrewise (pointed) cofibrations and fibrations in the category \mathbf{MAP} . In this paper, we continue the previous work. In section 3, we prove that Puppe sequence is exact in \mathbf{MAP} . In section 4, we study an application of Puppe exact sequence. In this study of section 4, we need not to consider any generalized concept of fibrewise non-degenerate spaces [5; section 22], and we can prove the generalized formula for the suspension of fibrewise product spaces. In section 5, we introduce an intermediate fibrewise category \mathbf{TOP}_B^H which combines \mathbf{MAP} with \mathbf{TOP}_B , and we prove an extended theorem in \mathbf{TOP}_B^H of [5] Proposition 22.11 by using theorems in \mathbf{TOP}_B^H (see Theorem 5.4 and Proposition 5.5). As a corollary

2000 *Mathematics Subject Classification.* 55R70, 55P05, 55R65.

Key words and phrases. Fibrewise Topology, category \mathbf{MAP} , fibrewise pointed homotopy, fibrewise pointed cofibration, puppe exact sequence.

of the extended theorem and the fact that a fibrewise non-degenerate space is H -fibrewise well-pointed (see Proposition 5.5), we give another proof of [5] Proposition 22.11.

The objects of **MAP** are continuous maps from any topological space into any topological space. For two objects $p : X \rightarrow B$ and $p' : X' \rightarrow B'$, a morphism from p into p' is a pair (ϕ, α) of continuous maps $\phi : X \rightarrow X'$, $\alpha : B \rightarrow B'$ such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\phi} & X' \\ p \downarrow & & \downarrow p' \\ B & \xrightarrow{\alpha} & B' \end{array}$$

is commutative. We note that this situation is a generalization of the category \mathbf{TOP}_B since the category \mathbf{TOP}_B is isomorphic to the particular case of **MAP** in which the spaces $B' = B$ and $\alpha = id_B$. We call an object $p : X \rightarrow B$ an **M-fibrewise space** and denote (X, p, B) . Also, for two **M-fibrewise spaces** $(X, p, B), (X', p', B')$, we call the morphism (ϕ, α) from p into p' an **M-fibrewise map**, and denote $(\phi, \alpha) : (X, p, B) \rightarrow (X', p', B')$.

Furthermore, in this paper we often consider the case that an **M-fibrewise space** (X, p, B) has a section $s : B \rightarrow X$, we call it an **M-fibrewise pointed space** and denote (X, p, B, s) . For two **M-fibrewise pointed spaces** $(X, p, B, s), (X', p', B', s')$, if an **M-fibrewise map** $(\phi, \alpha) : (X, p, B) \rightarrow (X', p', B')$ satisfies $\phi s = s' \alpha$, we call it an **M-fibrewise pointed map** and denote $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$.

In this paper, we assume that all spaces are topological spaces, all maps are continuous and id is the identity map of $I = [0, 1]$ into itself. Moreover, we use the following notation : For any $t \in I$, the maps $\sigma_t : X \rightarrow I \times X$ and $\delta_t : B \rightarrow I \times B$ are defined by

$$\sigma_t(x) = (t, x), \quad \delta_t(b) = (t, b) \quad (x \in X, b \in B).$$

For other undefined terminology, see [3], [4] and [5].

2. **M-FIBREWISE POINTED HOMOTOPY**

In this section, first we shall define an **M-fibrewise pointed homotopy**, **M-fibrewise pointed cofibration** and **M-fibrewise pointed cofibred pair** which are introduced in [4]. Next, we shall introduce some concepts, for example, **M-fibrewise pointed mapping cylinder**, **M-fibrewise pointed collapse**, **M-fibrewise pointed cone** and **M-fibrewise pointed nulhomotopic**. Last, we shall prove some propositions. These concepts and propositions are used in latter section. We begin with the following definitions.

DEFINITION 2.1. (1) ([4; Definition 5.1]) Let
 $(\phi, \alpha), (\theta, \beta) : (X, p, B, s) \rightarrow (X', p', B', s')$

be \mathbf{M} -fibrewise pointed maps. If there exists an \mathbf{M} -fibrewise pointed map $(H, h) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B', s')$ such that (H, h) is an \mathbf{M} -fibrewise homotopy of (ϕ, α) into (θ, β) (that is: $H\sigma_0 = \phi$, $H\sigma_1 = \theta$, $h\delta_0 = \alpha$, $h\delta_1 = \beta$), we call it an \mathbf{M} -fibrewise pointed homotopy of (ϕ, α) into (θ, β) . If there exists an \mathbf{M} -fibrewise pointed homotopy of (ϕ, α) into (θ, β) , we say (ϕ, α) is \mathbf{M} -fibrewise pointed homotopic to (θ, β) and write $(\phi, \alpha) \simeq_{(\mathbf{P})}^{\mathbf{M}} (\theta, \beta)$.

- (2) ([4; Definition 5.2]) An \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ is called an \mathbf{M} -fibrewise pointed homotopy equivalence if there exists an \mathbf{M} -fibrewise pointed map $(\theta, \beta) : (X', p', B', s') \rightarrow (X, p, B, s)$ such that

$$(\theta\phi, \beta\alpha) \simeq_{(\mathbf{P})}^{\mathbf{M}} (id_X, id_B), \quad (\phi\theta, \alpha\beta) \simeq_{(\mathbf{P})}^{\mathbf{M}} (id_{X'}, id_{B'}).$$

Then we denote $(X, p, B, s) \cong_{(\mathbf{P})}^{\mathbf{M}} (X', p', B', s')$.

It is obvious that the relations $\simeq_{(\mathbf{P})}^{\mathbf{M}}$ and $\cong_{(\mathbf{P})}^{\mathbf{M}}$ are equivalence relations. Now, we define \mathbf{M} -fibrewise pointed cofibration and \mathbf{M} -fibrewise pointed cofibred pair as follows.

DEFINITION 2.2. ([4; Definition 5.6]) An \mathbf{M} -fibrewise pointed map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X, p, B, s)$ is an \mathbf{M} -fibrewise pointed cofibration if (u, γ) has the following \mathbf{M} -fibrewise homotopy extension property : Let $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ be an \mathbf{M} -fibrewise pointed map and $(H, h) : (I \times X_0, id \times p_0, I \times B_0, id \times s_0) \rightarrow (X', p', B', s')$ an \mathbf{M} -fibrewise pointed homotopy such that the following two diagrams

$$\begin{array}{ccc} X_0 & \xrightarrow{\sigma_0} & I \times X_0 & & B_0 & \xrightarrow{\delta_0} & I \times B_0 \\ u \downarrow & & \downarrow H & & \gamma \downarrow & & \downarrow h \\ X & \xrightarrow{\phi} & X' & & B & \xrightarrow{\alpha} & B' \end{array}$$

are commutative. Then there exists an \mathbf{M} -fibrewise pointed homotopy $(K, k) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B', s')$ such that $K\kappa_0 = \phi$, $K(id \times u) = H$, $k\rho_0 = \alpha$, $k(id \times \gamma) = h$, where $\kappa_0 : X \rightarrow I \times X$ and $\rho_0 : B \rightarrow I \times B$ are defined by $\kappa_0(x) = (0, x)$ and $\rho_0(b) = (0, b)$ for $x \in X$, $b \in B$.

DEFINITION 2.3 ([4; Definition 2.3 and p.210]). (1) Let (X, p, B) be an \mathbf{M} -fibrewise space. If $X_0 \subset X$, $B_0 \subset B$ and $p(X_0) \subset B_0$, we call $(X_0, p|_{X_0}, B_0)$ an \mathbf{M} -fibrewise subspace of (X, p, B) . We sometimes use the notation (X_0, p_0, B_0) instead of $(X_0, p|_{X_0}, B_0)$. By the same way, we define an \mathbf{M} -fibrewise pointed subspace.

- (2) For an \mathbf{M} -fibrewise pointed subspace (X_0, p_0, B_0, s_0) of (X, p, B, s) , the pair $((X, p, B, s), (X_0, p_0, B_0, s_0))$ is called by an \mathbf{M} -fibrewise pointed pair. If X_0 is closed in X and B_0 is closed in B , it is called a closed \mathbf{M} -fibrewise pointed pair. For an \mathbf{M} -fibrewise

pointed pair $((X, p, B, s), (X_0, p_0, B_0, s_0))$, if the inclusion map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X, p, B, s)$ is an \mathbf{M} -fibrewise pointed cofibration, we call the pair $((X, p, B, s), (X_0, p_0, B_0, s_0))$ an \mathbf{M} -fibrewise pointed cofibred pair.

The following proposition, which will be used in section 4, can be proved by the same method of the proof in [4; Theorem 3.9].

PROPOSITION 2.4. *Let $((X, p, B, s), (X_0, p_0, B_0, s_0))$ and $((X', p', B', s'), (X'_0, p'_0, B'_0, s'_0))$ be two closed \mathbf{M} -fibrewise pointed cofibred pairs. Then*

$((X \times X', p \times p', B \times B', s \times s'), (X_0 \times X'_0 \cup X \times X'_0, \bar{p}, B_0 \times B'_0 \cup B \times B'_0, \bar{s}))$

is also an \mathbf{M} -fibrewise pointed cofibred pair, where

$$\bar{p} = p \times p' | X_0 \times X'_0 \cup X \times X'_0, \bar{s} = s \times s' | B_0 \times B'_0 \cup B \times B'_0.$$

For cotriad, see [5]. We can also define the \mathbf{M} -fibrewise push-out of a cotriad as same as the fibrewise push-out in [5] as follows:

DEFINITION 2.5 (cf. [4; p.208–9]). *For an \mathbf{M} -fibrewise pointed map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X_1, p_1, B_1, s_1)$, we can construct the \mathbf{M} -fibrewise pointed push-out (M, p, B, s) of the cotraids*

$$(I \times X_0, \text{id} \times p_0, I \times B_0, \text{id} \times s_0) \xleftarrow{(\sigma_0, \delta_0)} (X_0, p_0, B_0, s_0) \xrightarrow{(u, \gamma)} (X_1, p_1, B_1, s_1)$$

where (σ_0, δ_0) is an \mathbf{M} -fibrewise embedding to 0-level, as follows : $M = (I \times X_0 + X_1) / \sim$ and $B = (I \times B_0 + B_1) / \approx$, where $(0, a) \sim u(a)$ for $a \in X_0$ and $(0, b) \approx \gamma(b)$ for $b \in B_0$, and $p : M \rightarrow B$ and $s : B \rightarrow M$ are defined, respectively, by

$$p(x) = \begin{cases} [\gamma p_0(a)] & \text{if } x = [u(a)], a \in X_0 \\ [t, p_0(a)] & \text{if } x = [t, a], t \neq 0 \\ [p_1(x)] & \text{if } x \in X_1 - u(X_0), \end{cases}$$

$$s(b) = \begin{cases} [s_1 \alpha(d)] & \text{if } b = [\alpha(d)], d \in B_0 \\ [t, s_0(d)] & \text{if } b = [t, d], t \neq 0 \\ [s_1(b)] & \text{if } b \in B_1 - \alpha(B_0), \end{cases}$$

where $[*]$ is the equivalence class. Then it is easily verified that p and s are well-defined and continuous. We call the \mathbf{M} -fibrewise pointed push-out of the cotriad the \mathbf{M} -fibrewise pointed mapping cylinder of (u, γ) , and denote by $M(u, \gamma)$.

Now we shall consider the case in which (X_0, p_0, B_0, s_0) is an \mathbf{M} -fibrewise pointed subspace of (X_1, p_1, B_1, s_1) and $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow$

(X_1, p_1, B_1, s_1) is the inclusion. We can define an \mathbf{M} -fibrewise pointed map $(e, \epsilon) : (M, p, B, s) \rightarrow (0 \times X_1 \cup I \times X_0, id \times p_1, 0 \times B_1 \cup I \times B_0, id \times s_1)$ by

$$e(x) = \begin{cases} (0, a) & \text{if } x = [u(a)], a \in X_0 \\ (t, a) & \text{if } x = [t, a], t \neq 0 \\ (0, x) & \text{if } x \in X_1 - u(X_0), \end{cases}$$

$\epsilon(b)$ is defined by a similar way. Moreover if X_0 is closed in X_1 and B_0 is closed in B_1 , the maps e and ϵ are homeomorphisms and we may identify (M, p, B, s) with $(0 \times X_1 \cup I \times X_0, id \times p_1, 0 \times B_1 \cup I \times B_0, id \times s)$.

For each \mathbf{M} -fibrewise pointed map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X_1, p_1, B_1, s_1)$, we can define an \mathbf{M} -fibrewise pointed map $(k, \xi) : (M, p, B, s) \rightarrow (I \times X_1, id \times p_1, I \times B_1, id \times s_1)$ by

$$k(x) = \begin{cases} (0, u(a)) & \text{if } x = [u(a)], a \in X_0 \\ (t, u(a)) & \text{if } x = [t, a], t \neq 0 \\ (0, x) & \text{if } x \in X_1 - u(X_0), \end{cases}$$

$$\xi(b) = \begin{cases} (0, \gamma(d)) & \text{if } b = [\gamma(d)], d \in B_0 \\ (t, \gamma(d)) & \text{if } b = [t, d], t \neq 0 \\ (0, b) & \text{if } b \in B_1 - \gamma(B_0). \end{cases}$$

Then we can obtain the following proposition by the same method of the proof in [4; Theorem 3.1].

PROPOSITION 2.6. *The \mathbf{M} -fibrewise pointed map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X_1, p_1, B_1, s_1)$ is an \mathbf{M} -fibrewise pointed cofibration if and only if there is an \mathbf{M} -fibrewise pointed map $(L, l) : (I \times X_1, id \times p_1, I \times B_1, id \times s_1) \rightarrow (M, p, B, s)$ such that $Lk = id_M, l\xi = id_B$, where (M, p, B, s) is the same one in Definition 2.5 and $(k, \xi) : (M, p, B, s) \rightarrow (I \times X_1, id \times p_1, I \times B_1, id \times s_1)$ is the same one in the above.*

The following lemma is used in the next section.

LEMMA 2.7. *For an \mathbf{M} -fibrewise pointed map $(u, \gamma) : (X_0, p_0, B_0, s_0) \rightarrow (X_1, p_1, B_1, s_1)$, let $M(u, \gamma) = (M, p, B, s)$ be the \mathbf{M} -fibrewise pointed mapping cylinder of (u, γ) constructed in Definition 2.5. Then the \mathbf{M} -fibrewise pointed map $(\sigma_1, \delta_1) : (X_0, p_0, B_0, s_0) \rightarrow M(u, \gamma)$ is an \mathbf{M} -fibrewise pointed cofibration, where (σ_1, δ_1) is defined by $\sigma_1(x) = (1, x), \delta_1(b) = (1, b)$.*

PROOF. Note that we can define an \mathbf{M} -fibrewise pointed map $(k, \xi) : M(\sigma_1, \delta_1) \rightarrow (I \times M, id \times p, I \times B, id \times s)$ by the same method in Proposition 2.6. We shall prove this lemma by using Proposition 2.6. For this purpose,

we now define an \mathbf{M} -fibrewise pointed function

$$\begin{aligned} (J, j) : (I \times (X_1 + I \times X_0), id \times (p_1 + id \times p_0), \\ I \times (B_1 + I \times B_0), id \times (s_1 + id \times s_0)) \\ \longrightarrow ((0 \times X_1 + \overline{I \times I} \times X_0, 0 \times p_1 + \overline{id \times id} \times p_0, \\ 0 \times B_1 + \overline{I \times I} \times B_0, 0 \times s_1 + \overline{id \times id} \times s_0) \end{aligned}$$

where $\overline{I \times I} = (0 \times I) \cup (I \times 1)$ and $\overline{id \times id} = id \times id|_{\overline{I \times I}}$.

Let $r : I \times I \rightarrow \overline{I \times I}$ be a retraction defined by the projection from the point $(2, 0)$. Then (J, j) is defined by

$$\begin{aligned} J(t, x) &= (0, x) & (x \in X_1) \\ J(t, (t', x_0)) &= (r(t, t'), x_0) & (t, t' \in I, x_0 \in X_0) \\ j(t, b) &= (0, b) & (b \in B_1) \\ j(t, (t', b_0)) &= (r(t, t'), b_0) & (t, t' \in I, b_0 \in B_0). \end{aligned}$$

Then it is easy to see that for $x_0 \in X_0$

$$J(t, (0, x_0)) = (r(t, 0), x_0) = (0, (0, x_0)) = (0, u(x_0)) = J(t, u(x_0)).$$

similarly $j(t, (0, b_0)) = j(t, \gamma(b_0))$ and (J, j) is an \mathbf{M} -fibrewise pointed map. Therefore it is easily verified that (J, j) induces the \mathbf{M} -fibrewise pointed retraction

$$(L, l) : (I \times M, id \times p, I \times B, id \times s) \rightarrow M(\sigma_1, \delta_1)$$

such that $Lk = id_{\overline{M}}, l\xi = id_{\overline{B}}$, where \overline{M} and \overline{B} are the total space and the base space of $M(\sigma_1, \delta_1)$, respectively. Thus by using Proposition 2.6, we complete the proof. \square

We now define \mathbf{M} -fibrewise pointed collapse, \mathbf{M} -fibrewise pointed cone and \mathbf{M} -fibrewise pointed nulhomotopic.

DEFINITION 2.8. (1) Let (X, p, B, s) be an \mathbf{M} -fibrewise pointed space and (X_0, p_0, B_0, s_0) a closed \mathbf{M} -fibrewise pointed subspace. Let \tilde{X} be a set $\cup_{b \in B} X_b / X_{0b}$, where $X_b = p^{-1}(b)$ and $X_{0b} = p_0^{-1}(b)$ for $b \in B$ (or $b \in B_0$). We introduce the set \tilde{X} the quotient topology of X and put $\tilde{B} = B$. If we define maps $\tilde{p} : \tilde{X} \rightarrow \tilde{B}$ and $\tilde{s} : \tilde{B} \rightarrow \tilde{X}$ indexed by p and s respectively, then $(\tilde{X}, \tilde{p}, \tilde{B}, \tilde{s})$ is an \mathbf{M} -fibrewise pointed space. We call $(\tilde{X}, \tilde{p}, \tilde{B}, \tilde{s})$ an \mathbf{M} -fibrewise pointed collapse of (X, p, B, s) with respect to (X_0, p_0, B_0, s_0) and denoted by

$$(X, p, B, s) /_{\mathbf{M}} (X_0, p_0, B_0, s_0).$$

(For fibrewise collapse, see [5; section 5].)

(2) For an \mathbf{M} -fibrewise pointed space (X, p, B, s) , we call the \mathbf{M} -fibrewise pointed collapse

$$(I \times X, id \times p, I \times B, id \times s) /_{\mathbf{M}} (1 \times X, id \times p | 1 \times X, 1 \times B, id \times s | 1 \times B)$$

the \mathbf{M} -fibrewise pointed cone of (X, p, B, s) and denote by $\Gamma(X, p, B, s)$.
(We denote the total space of $\Gamma(X, p, B, s)$ by CX .)

DEFINITION 2.9. Let $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ be an \mathbf{M} -fibrewise pointed map. Then we call (ϕ, α) to be \mathbf{M} -fibrewise pointed nulhomotopic if there is an \mathbf{M} -fibrewise pointed map $(c, \alpha_c) : (X, p, B, s) \rightarrow (X', p', B', s')$ such that $c = s'\alpha_c p$ and $(\phi, \alpha) \simeq_{(\mathbf{P})}^{\mathbf{M}} (c, \alpha_c)$.

We now prove the following proposition which is used in next section.

PROPOSITION 2.10. Let $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ be an \mathbf{M} -fibrewise pointed map. Then (ϕ, α) is \mathbf{M} -fibrewise pointed nulhomotopic if and only if $(\phi, \alpha) \circ (i, \epsilon)^{-1}$ can be extended to an \mathbf{M} -fibrewise pointed map of $\Gamma(X, p, B, s)$ to (X', p', B', s') , where $(i, \epsilon) : (X, p, B, s) \rightarrow \Gamma(X, p, B, s)$ is the natural embedding to 0-level of $\Gamma(X, p, B, s)$.

PROOF. “Only if” part: Let (ϕ, α) be \mathbf{M} -fibrewise pointed nulhomotopic. By Definition 2.9, there is an \mathbf{M} -fibrewise pointed map $(c, \alpha_c) : (X, p, B, s) \rightarrow (X', p', B', s')$ such that $c = s'\alpha_c p$, and there is an \mathbf{M} -fibrewise pointed homotopy

$$(H, h) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B', s')$$

such that $(H_0, h_0) = (\phi, \alpha)$ and $(H_1, h_1) = (c, \alpha_c)$. Now, we can define an \mathbf{M} -fibrewise pointed map

$$(\tilde{\phi}, \tilde{\alpha}) : \Gamma(X, p, B, s) \rightarrow (X', p', B', s')$$

by

$$\tilde{\phi}([t, x]) = H(t, x), \quad \tilde{\alpha}(t, b) = h(t, b).$$

Because it is obvious that $(\tilde{\phi}, \tilde{\alpha})$ is an \mathbf{M} -fibrewise pointed map by the facts

$$p'\tilde{\phi}([t, x]) = p'H(t, x) = h(t, p(x)) = \tilde{\alpha}(t, p(x)) = \tilde{p}([t, x])$$

where \tilde{p} is the projection from the total space of $\Gamma(X, p, B, s)$ to the base space $I \times B$, and

$$\tilde{\phi}(id \times s)(t, b) = \tilde{\phi}([t, s(b)]) = H(t, s(b)) = H(id \times s)(t, b) = s'h(t, b) = s'\tilde{\alpha}(t, b)$$

since (H, h) is \mathbf{M} -fibrewise pointed. Furthermore, it is easy to see that $(\tilde{\phi}, \tilde{\alpha}) \circ (i, \epsilon) = (\phi, \alpha)$, so $(\phi, \alpha) \circ (i, \epsilon)^{-1}$ can be extended to an \mathbf{M} -fibrewise pointed map of $\Gamma(X, p, B, s)$ to (X', p', B', s') .

“If” part: Let an \mathbf{M} -fibrewise pointed map

$$(\tilde{\phi}, \tilde{\alpha}) : \Gamma(X, p, B, s) \rightarrow (X', p', B', s')$$

be an extension of the \mathbf{M} -fibrewise pointed map $(\phi, \alpha) \circ (i, \epsilon)^{-1}$. Now, we can define \mathbf{M} -fibrewise pointed functions

$$(H, h) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B', s')$$

$$(c, \alpha_c) : (X, p, B, s) \rightarrow (X', p', B', s')$$

by

$$\begin{aligned} H(t, x) &= \tilde{\phi}([t, x]) \\ h(t, b) &= \tilde{\alpha}(t, b) \\ c &= H_1 \\ \alpha_c &= h_1. \end{aligned}$$

Then it is clear that $(H, h), (c, \alpha_c)$ are continuous and $H_0 = \phi, h_0 = \alpha$. Further, since $(\tilde{\phi}, \tilde{\alpha})$ is an \mathbf{M} -fibrewise pointed map, we have

$$H(id \times s)(t, b) = H(t, s(b)) = \tilde{\phi}([t, s(b)]) = \tilde{\phi}(id \times s)(t, b) = s' \tilde{\alpha}(t, b) = s' h(t, b).$$

Therefore (H, h) is an \mathbf{M} -fibrewise pointed map. Furthermore by the fact

$$\begin{aligned} s' \alpha_c p(x) &= s' \alpha_c(p(x)) = s' \tilde{\alpha}(1, p(x)) = s' \tilde{\alpha} \bar{p}([1, x]) \\ &= \tilde{\phi}([1, x]) = H_1(x) = c(x), \end{aligned}$$

it is easy to see that $s' \alpha_c p = c$. Thus, (ϕ, α) is \mathbf{M} -fibrewise pointed nullhomotopic. \square

3. PUPPE EXACT SEQUENCE

The main purpose of this section is the proof of Puppe exact sequence in \mathbf{MAP} . (For this sequence, see [9] in the category \mathbf{TOP} and [5] in \mathbf{TOP}_B .) We begin with defining \mathbf{M} -fibrewise pointed mapping cone, \mathbf{M} -fibrewise pointed contractible, \mathbf{M} -fibrewise pointed suspension and exactness of sequence of \mathbf{M} -fibrewise pointed maps.

DEFINITION 3.1. *For an \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$, we call $(CX \cup_\phi X', \tilde{p}, (I \times B) \cup_\alpha B', \tilde{s})$ the \mathbf{M} -fibrewise pointed mapping cone of (ϕ, α) , and denote $\Gamma(\phi, \alpha)$, where CX is the space in Definition 2.8.*

In this definition, the maps

$$\tilde{p} : CX \cup_\phi X' \rightarrow (I \times B) \cup_\alpha B', \quad \tilde{s} : (I \times B) \cup_\alpha B' \rightarrow CX \cup_\phi X'$$

are defined as the \mathbf{M} -fibrewise pointed maps induced by the following maps, respectively:

$$\bar{p} : CX + X' \rightarrow (I \times B) + B', \quad \bar{s} : (I \times B) + B' \rightarrow CX + X'$$

$$\begin{aligned} \bar{p}([t, x]) &= (t, p(x)) && ([t, x] \in CX) \\ \bar{p}(x') &= p'(x') && (x' \in X') \\ \bar{s}(t, b) &= [t, s(b)] && ((t, b) \in I \times B) \\ \bar{s}(b') &= s'(b') && (b' \in B'), \end{aligned}$$

where $CX \cup_\phi X', (I \times B) \cup_\alpha B'$ are adjunction spaces, respectively, determined by

$$\phi : X = 0 \times X \rightarrow X', \quad \alpha : B = 0 \times B \rightarrow B'.$$

For fibrewise adjunction space, see [6].

For an \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ and the \mathbf{M} -fibrewise mapping cone $\Gamma(\phi, \alpha)$, it is easy to see that there is the natural embedding

$$(\phi', \alpha') : (X', p', B', s') \rightarrow \Gamma(\phi, \alpha).$$

For two \mathbf{M} -fibrewise pointed spaces (X, p, B, s) and (X', p', B', s') , we consider the set of all \mathbf{M} -fibrewise pointed homotopy classes of \mathbf{M} -fibrewise pointed maps from (X, p, B, s) to (X', p', B', s') , and denote it by $\pi((X, p, B, s), (X', p', B', s'))$. Then for \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ and any \mathbf{M} -fibrewise pointed space (X'', p'', B'', s'') , we can define an induced map

$(\phi, \alpha)^* : \pi((X', p', B', s'), (X'', p'', B'', s'')) \rightarrow \pi((X, p, B, s), (X'', p'', B'', s''))$ of (ϕ, α) by $(\phi, \alpha)^*([\phi', \alpha']) = [\phi' \phi, \alpha' \alpha]$. It is easy to see that this map is well-defined. Now we shall define exactness of a sequence of \mathbf{M} -fibrewise pointed maps.

DEFINITION 3.2. *A sequence of \mathbf{M} -fibrewise pointed maps*

$$(X_1, p_1, B_1, s_1) \xrightarrow{(\phi_1, \alpha_1)} (X_2, p_2, B_2, s_2) \xrightarrow{(\phi_2, \alpha_2)} (X_3, p_3, B_3, s_3) \xrightarrow{(\phi_3, \alpha_3)} \dots$$

is exact if for any \mathbf{M} -fibrewise pointed space (X', p', B', s') the induced sequence from one in the above

$$\begin{aligned} \pi((X_1, p_1, B_1, s_1), (X', p', B', s')) &\xleftarrow{(\phi_1, \alpha_1)^*} \pi((X_2, p_2, B_2, s_2), (X', p', B', s')) \\ &\xleftarrow{(\phi_2, \alpha_2)^*} \pi((X_3, p_3, B_3, s_3), (X', p', B', s')) \\ &\xleftarrow{(\phi_3, \alpha_3)^*} \dots \end{aligned}$$

is exact.

REMARK 3.3. Note that the latter is exact if

$$\ker(\phi_i, \alpha_i)^* = \text{im}(\phi_{i+1}, \alpha_{i+1})^*$$

where

$$\ker(\phi_i, \alpha_i)^* = \{[\psi_{i+1}, \beta_{i+1}] | (\psi_{i+1} \phi_i, \beta_{i+1} \alpha_i) \text{ is } \mathbf{M}\text{-fibrewise pointed nulhomotopic}\}.$$

We have the following proposition for exactness.

PROPOSITION 3.4. *For \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ and \mathbf{M} -fibrewise pointed space (X'', p'', B'', s'') , the sequence*

$$\begin{aligned} \pi((X, p, B, s), (X'', p'', B'', s'')) &\xleftarrow{(\phi, \alpha)^*} \pi((X', p', B', s'), (X'', p'', B'', s'')) \\ &\xleftarrow{(\phi', \alpha')^*} \pi(\Gamma(\phi, \alpha), (X'', p'', B'', s'')) \end{aligned}$$

is exact.

PROOF. $\text{im}(\phi', \alpha')^* \subset \ker(\phi, \alpha)^*$: It is easy to see that the \mathbf{M} -fibrewise pointed map

$$(\phi', \alpha') \circ (\phi, \alpha) : (X, p, B, s) \rightarrow \Gamma(\phi, \alpha)$$

is extended to an \mathbf{M} -fibrewise pointed map $\Gamma(X, p, B, s) \rightarrow \Gamma(\phi, \alpha)$. Therefore, by Proposition 2.10 $(\phi', \alpha') \circ (\phi, \alpha)$ is \mathbf{M} -fibrewise pointed nulhomotopic.

$\ker(\phi, \alpha)^* \subset \text{im}(\phi', \alpha')^*$: Let

$$(\psi, \beta) : (X', p', B', s') \rightarrow (X'', p'', B'', s'')$$

be an \mathbf{M} -fibrewise pointed map such that $(\psi, \beta) \circ (\phi, \alpha)$ is \mathbf{M} -fibrewise pointed nulhomotopic. Let

$$(\tilde{H}, \tilde{h}) : (I \times X, \text{id} \times p, I \times B, \text{id} \times s) \rightarrow (X'', p'', B'', s'')$$

be the \mathbf{M} -fibrewise pointed nulhomotopy. Then by using \tilde{H}, ϕ, ψ and \tilde{h}, α, β , we can construct

$$(\psi', \beta') : \Gamma(\phi, \alpha) \rightarrow (X'', p'', B'', s'')$$

such that $(\psi', \beta') \circ (\phi', \alpha') = (\psi, \beta)$. \square

We shall define \mathbf{M} -fibrewise pointed contractible and prove two propositions connecting with this concept.

DEFINITION 3.5. *An \mathbf{M} -fibrewise pointed space (X, p, B, s) is \mathbf{M} -fibrewise pointed contractible if there is an \mathbf{M} -fibrewise pointed space (B', p', B', p'^{-1}) (where p' is a homeomorphism) such that*

$$(X, p, B, s) \cong_{(\mathbf{P})}^{\mathbf{M}} (B', p', B', p'^{-1}).$$

PROPOSITION 3.6. *An \mathbf{M} -fibrewise pointed space (X, p, B, s) is \mathbf{M} -fibrewise pointed contractible if and only if $(\text{id}_X, \text{id}_B)$ is \mathbf{M} -fibrewise pointed nulhomotopic.*

PROOF. “Only if” part: Let an \mathbf{M} -fibrewise pointed space (X, p, B, s) be \mathbf{M} -fibrewise pointed contractible. By the definition, there are an \mathbf{M} -fibrewise pointed space (B', p', B', p'^{-1}) and an \mathbf{M} -fibrewise pointed homotopy equivalence

$$(\phi, \alpha) : (X, p, B, s) \rightarrow (B', p', B', p'^{-1})$$

satisfying $\phi = p'^{-1}\alpha p$. Let (ψ, β) be an \mathbf{M} -fibrewise pointed homotopy inverse of (ϕ, α) . Then by the fact

$$\begin{aligned} s(\beta\alpha)p &= (s\beta p')(p'^{-1}\alpha p) \\ &= \psi\phi \end{aligned}$$

and $(\psi\phi, \beta\alpha) \simeq_{(\mathbf{P})}^{\mathbf{M}} (\text{id}_X, \text{id}_B)$, $(\text{id}_X, \text{id}_B)$ is \mathbf{M} -fibrewise pointed nulhomotopic.

“If” part: Let (id_X, id_B) be \mathbf{M} -fibrewise pointed nulhomotopic. There is an \mathbf{M} -fibrewise pointed map

$$(c, \alpha_c) : (X, p, B, s) \rightarrow (X, p, B, s)$$

such that $c = s\alpha_cp$ and $(id_X, id_B) \simeq_{(\mathbf{P})}^{\mathbf{M}} (c, \alpha_c)$. Then we shall prove

$$(X, p, B, s) \cong_{(\mathbf{P})}^{\mathbf{M}} (B, id_B, B, id_B).$$

First, since

$$(pc, \alpha_c) : (X, p, B, s) \rightarrow (B, id_B, B, id_B)$$

is \mathbf{M} -fibrewise pointed, we have

$$\begin{aligned} s(pc) &= sp(s\alpha_cp) \\ &= s\alpha_cp \\ &= c \\ id_B\alpha_c &= \alpha_c \end{aligned}$$

and

$$(s(pc), id_B\alpha_c) = (c, \alpha_c) \simeq_{(\mathbf{P})}^{\mathbf{M}} (id_X, id_B).$$

Next, since

$$\begin{aligned} (pc)s &= p(s\alpha_cp)s \\ &= \alpha_c \\ \alpha_c id_B &= \alpha_c, \end{aligned}$$

we have

$$((pc)s, \alpha_c id_B) = (\alpha_c, \alpha_c) \simeq_{(\mathbf{P})}^{\mathbf{M}} (id_B, id_B).$$

Thus (s, id_B) is an \mathbf{M} -fibrewise pointed homotopy inverse of (pc, α_c) . Therefore, (pc, α_c) is an \mathbf{M} -fibrewise pointed homotopy equivalence. \square

PROPOSITION 3.7. *For an \mathbf{M} -fibrewise pointed space (X, p, B, s) , the \mathbf{M} -fibrewise pointed cone $\Gamma(X, p, B, s)$ is \mathbf{M} -fibrewise pointed contractible.*

PROOF. We define an \mathbf{M} -fibrewise homotopy

$$(H_\tau, h_\tau) : \Gamma(X, p, B, s) \rightarrow \Gamma(X, p, B, s)$$

by

$$\begin{aligned} H_\tau(t, x) &= (t + \tau(1 - t), x) \\ h_\tau(t, b) &= (t + \tau(1 - t), b). \end{aligned}$$

Then this (H_τ, h_τ) is an \mathbf{M} -fibrewise pointed nulhomotopy of (id_X, id_B) . Therefore we complete the proof. \square

From now on, to prove Puppe exact sequence in **MAP**, we shall give some propositions.

PROPOSITION 3.8. *Let $((X, p, B, s), (X_0, p_0, B_0, s_0))$ be a closed \mathbf{M} -fibrewise pointed cofibred pair. Assume that (X_0, p_0, B_0, s_0) is \mathbf{M} -fibrewise pointed contractible. Then the natural projection*

$$(\pi, id_B) : (X, p, B, s) \rightarrow (X, p, B, s)/_{\mathbf{M}}(X_0, p_0, B_0, s_0)$$

is an \mathbf{M} -fibrewise pointed homotopy equivalence.

PROOF. Let

$$(F_t, f_t) : (X_0, p_0, B_0, s_0) \rightarrow (X_0, p_0, B_0, s_0)$$

be an \mathbf{M} -fibrewise pointed nullhomotopy of (id_{X_0}, id_{B_0}) , where (F_1, f_1) is \mathbf{M} -fibrewise pointed constant. By the assumption, for

$$(id_X, id_B) : (X, p, B, s) \rightarrow (X, p, B, s)$$

(F_t, f_t) can be extended to an \mathbf{M} -fibrewise pointed homotopy $(H_t, h_t) : (X, p, B, s) \rightarrow (X, p, B, s)$ of (id_X, id_B) . Let $(H'_1, h'_1) = (H_1, h_1)(\pi, id_B)^{-1}$. Then the \mathbf{M} -fibrewise pointed map

$$(H'_1, h'_1) : (X, p, B, s)/_{\mathbf{M}}(X_0, p_0, B_0, s_0) \rightarrow (X, p, B, s)$$

is induced from (H_t, h_t) . Further $(\pi, id_B) \circ (H_t, h_t)$ induces an \mathbf{M} -fibrewise pointed homotopy

$$(H''_t, h''_t) : (X, p, B, s)/_{\mathbf{M}}(X_0, p_0, B_0, s_0) \rightarrow (X, p, B, s)/_{\mathbf{M}}(X_0, p_0, B_0, s_0).$$

Note that $H''_1 = \pi H'_1, h''_1 = h'_1$. Then since

$$(H'_1, h'_1) \circ (\pi, id_B) = (H_1, h_1) \simeq_{(\mathbf{P})}^{\mathbf{M}} (H_0, h_0) = (id_X, id_B),$$

$$(\pi, id_B) \circ (H'_1, h'_1) = (H''_1, h''_1) \simeq_{(\mathbf{P})}^{\mathbf{M}} (H''_0, h''_0)$$

and (H''_0, h''_0) is the identity of $(X, p, B, s)/_{\mathbf{M}}(X_0, p_0, B_0, s_0)$, (π, id_B) is an \mathbf{M} -fibrewise pointed homotopy equivalence. This completes the proof. \square

PROPOSITION 3.9. *Let $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ be an \mathbf{M} -fibrewise pointed cofibration. Then the natural projection*

$$(\pi, id) : \Gamma(\phi, \alpha) \rightarrow \Gamma(\phi, \alpha)/_{\mathbf{M}}\Gamma(X, p, B, s) \cong_{(\mathbf{P})}^{\mathbf{M}} (X', p', B', s')/_{\mathbf{M}}(X, p, B, s)$$

is an \mathbf{M} -fibrewise pointed homotopy equivalence.

PROOF. We shall construct an \mathbf{M} -fibrewise pointed homotopy of (id, id) of the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(\phi, \alpha)$ which deforms the \mathbf{M} -fibrewise pointed cone $\Gamma(X, p, B, s)$ into its section. Since, if this is done, $\Gamma(X, p, B, s)$ is an \mathbf{M} -fibrewise pointed contractible in $\Gamma(\phi, \alpha)$, we can obtain the result from Proposition 3.8.

We can now define an \mathbf{M} -fibrewise pointed nullhomotopy $(H, h) : I \times \Gamma(X, p, B, s) \rightarrow \Gamma(\phi, \alpha)$ of the inclusion $\Gamma(X, p, B, s) \rightarrow \Gamma(\phi, \alpha)$ by

$$\begin{aligned} H(t, [t', x]) &= [t' + t(1 - t'), x] \\ h(t, t', b) &= (t' + t(1 - t'), b) \end{aligned}$$

Since (ϕ, α) is an \mathbf{M} -fibrewise pointed cofibration, (H, h) can be extended to an \mathbf{M} -fibrewise pointed homotopy

$$(G, g) : I \times (X', p', B', s') \rightarrow \Gamma(\phi, \alpha)$$

of the inclusion $(X', p', B', s') \rightarrow \Gamma(\phi, \alpha)$. Then by using (H, h) and (G, g) we can construct an \mathbf{M} -fibrewise pointed homotopy of (id, id) of the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(\phi, \alpha)$ which deforms the \mathbf{M} -fibrewise pointed cone $\Gamma(X, p, B, s)$ into its section. \square

By using Lemma 2.7 we can prove the next proposition.

PROPOSITION 3.10. *For an \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$, the inclusion $(\phi', \alpha') : (X', p', B', s') \rightarrow \Gamma(\phi, \alpha)$ is an \mathbf{M} -fibrewise pointed cofibration.*

PROOF. Let $M(p, id_B)$ be the \mathbf{M} -fibrewise pointed mapping cylinder of $(p, id_B) : (X, p, B, s) \rightarrow (B, id_B, B, id_B)$. Then it is easy to see that $\Gamma(X, p, B, s)$ and $M(p, id_B)$ are \mathbf{M} -fibrewise pointed equivalent by an \mathbf{M} -fibrewise pointed map of $\Gamma(X, p, B, s)$ to $M(p, id_B)$ defined by

$$[t, x] \mapsto [1 - t, x], \quad (t, b) \mapsto (1 - t, b)$$

Note by Lemma 2.7 that the map $(\sigma_1, \delta_1) : (X, p, B, s) \rightarrow \Gamma(X, p, B, s)$ which maps to 1-level is an \mathbf{M} -fibrewise pointed cofibration.

For any \mathbf{M} -fibrewise pointed homotopy $(H_t, h_t) : (X', p', B', s') \rightarrow (X'', p'', B'', s'')$, $(H_t \circ \phi, h_t \circ \alpha) : (X, p, B, s) \rightarrow (X'', p'', B'', s'')$ is an \mathbf{M} -fibrewise pointed homotopy. Since (σ_1, δ_1) in the above is an \mathbf{M} -fibrewise pointed cofibration, $(H_t \circ \phi, h_t \circ \alpha)$ can be extended to $\Gamma(X, p, B, s)$. Thus by patching the extended homotopy and (H_t, h_t) in $\Gamma(\phi, \alpha)$ we can construct an \mathbf{M} -fibrewise pointed homotopy of $\Gamma(\phi, \alpha)$ to (X'', p'', B'', s'') . This completes the proof. \square

DEFINITION 3.11. *For an \mathbf{M} -fibrewise pointed space (X, p, B, s) , let $\overline{X} = \{0, 1\} \times X$, $\overline{p} = id \times p|_{\{0, 1\} \times X}$, $\overline{B} = \{0, 1\} \times B$ and $\overline{s} = id \times s|_{\{0, 1\} \times B}$. Then the \mathbf{M} -fibrewise pointed collapse*

$$(I \times X, id \times p, I \times B, id \times s) /_{\mathbf{M}} (\overline{X}, \overline{p}, \overline{B}, \overline{s})$$

is called to be \mathbf{M} -fibrewise pointed suspension, and denoted by $\Sigma(X, p, B, s)$. (We denote the total space of $\Sigma(X, p, B, s)$ by ΣX , and the projection of $\Sigma(X, p, B, s)$ by Σp .)

PROPOSITION 3.12. *Let $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ be an \mathbf{M} -fibrewise pointed map, where α is a bijection. Then for the inclusion $(\phi', \alpha') : (X', p', B', s') \rightarrow \Gamma(\phi, \alpha)$, $\Gamma(\phi', \alpha')$ is \mathbf{M} -fibrewise pointed equivalent to the \mathbf{M} -fibrewise pointed suspension $\Sigma(X, p, B, s)$.*

PROOF. We use the same notation of Proposition 3.10. Since (ϕ', α') is an \mathbf{M} -fibrewise pointed cofibration from Proposition 3.10, using Proposition 3.9, $\Gamma(\phi', \alpha')$ is \mathbf{M} -fibrewise pointed homotopy equivalent to

$$\Gamma(\phi', \alpha') /_{\mathbf{M}} \Gamma(X', p', B', s') = \Gamma(\phi, \alpha) /_{\mathbf{M}} (X', p', B', s') = \Sigma(X, p, B, s).$$

This completes the proof. \square

In the process in the above, $((\phi')', (\alpha')')$ is transformed into an \mathbf{M} -fibrewise pointed map

$$(\phi'', \alpha'') : \Gamma(\phi, \alpha) \rightarrow \Sigma(X, p, B, s).$$

Repeating this process we find that $\Gamma((\phi')', (\alpha')')$ is \mathbf{M} -fibrewise pointed equivalent to \mathbf{M} -fibrewise pointed suspension $\Sigma(X', p', B', s')$, and in the process $(((\phi')')', ((\alpha')')')$ is transformed into the \mathbf{M} -fibrewise pointed suspension

$$(\Sigma\phi, id \times \alpha) : \Sigma(X, p, B, s) \rightarrow \Sigma(X', p', B', s'),$$

where $\Sigma\phi$ is the map from ΣX to $\Sigma X'$. Thus we obtain the main theorem of Puppe exact sequence in **MAP**.

THEOREM 3.13. *For an \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ where α is a bijection, the following sequence is exact.*

$$\begin{array}{ccc} (X, p, B, s) & \xrightarrow{(\phi, \alpha)} & (X', p', B', s') \\ & \xrightarrow{(\phi', \alpha')} & \Gamma(\phi, \alpha) \\ & \xrightarrow{(\phi'', \alpha'')} & \Sigma(X, p, B, s) \\ & \xrightarrow{(\phi''', \alpha''')} & \Sigma(X', p', B', s') \\ & \longrightarrow & \dots \end{array}$$

4. AN APPLICATION OF PUPPE EXACT SEQUENCE

In this section, by applying Puppe exact sequence we shall prove the generalized formula for the suspension of \mathbf{M} -fibrewise pointed product spaces, and the proof is simpler than the one in fibrewise version of [5; section 22]. (In our proof, we need not to consider any generalized concept of fibrewise non-degenerate spaces in [5].)

DEFINITION 4.1. (1) *For two \mathbf{M} -fibrewise pointed spaces (X_1, p_1, B_1, s_1) and (X_2, p_2, B_2, s_2) let*

$$\begin{aligned} X_1 \vee^{\mathbf{M}} X_2 &= \cup_{(b, b') \in B_1 \times B_2} (X_{1,b} \times s_2(b') \cup s_1(b) \times X_{2,b'}), \\ p_1 \vee^{\mathbf{M}} p_2 &= p_1 \times p_2 |_{X_1 \vee^{\mathbf{M}} X_2}. \end{aligned}$$

The \mathbf{M} -fibrewise pointed space $(X_1 \vee^{\mathbf{M}} X_2, p_1 \vee^{\mathbf{M}} p_2, B_1 \times B_2, s_1 \times s_2)$ is called the \mathbf{M} -fibrewise pointed coproduct of (X_1, p_1, B_1, s_1) and

(X_2, p_2, B_2, s_2) , and denoted by $(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)$ or $\bigvee_{i \in \{1,2\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i)$. The \mathbf{M} -fibrewise pointed collapse

$$(X_1, p_1, B_1, s_1) \times (X_2, p_2, B_2, s_2) /_{\mathbf{M}} (X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)$$

is called the \mathbf{M} -fibrewise smash product of (X_1, p_1, B_1, s_1) and (X_2, p_2, B_2, s_2) , and denoted by

$$(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2) \text{ or } \bigwedge_{i \in \{1,2\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i).$$

Note that $(X_1, p_1, B_1, s_1) \times (X_2, p_2, B_2, s_2) = (X_1 \times X_2, p_1 \times p_2, B_1 \times B_2, s_1 \times s_2)$.

(2) For $n \geq 3$ and \mathbf{M} -fibrewise pointed spaces (X_i, p_i, B_i, s_i) ($i = 1, \dots, n$), we shall define $\bigvee_{i \in \{1, \dots, n\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i)$ and $\bigwedge_{i \in \{1, \dots, n\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i)$ inductively, as follows:

$$\bigvee_{i \in \{1, \dots, n\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i) = \left(\bigvee_{i \in \{1, \dots, n-1\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i) \right) \vee^{\mathbf{M}} (X_n, p_n, B_n, s_n)$$

$$\bigwedge_{i \in \{1, \dots, n\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i) = \left(\bigwedge_{i \in \{1, \dots, n-1\}}^{\mathbf{M}} (X_i, p_i, B_i, s_i) \right) \wedge^{\mathbf{M}} (X_n, p_n, B_n, s_n).$$

In this paper, we set up the following Hypothesis.

HYPOTHESIS: By Definitions 3.11 and 4.1, the base space of

$$\Sigma\{(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)\}$$

and

$$\Sigma\{(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2)\}$$

is $I \times (B_1 \times B_2)$, but the base space of $\Sigma(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$ and $\Sigma(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$ is $(I \times B_1) \times (I \times B_2)$. So, since $\Sigma\{(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)\}$ and $\Sigma(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$, or $\Sigma\{(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2)\}$ and $\Sigma(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$ have different base spaces, those \mathbf{M} -fibrewise pointed spaces are different, respectively. We want to identify those spaces as \mathbf{M} -fibrewise pointed space, we set the following hypothesis: In $\Sigma(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$ and $\Sigma(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2)$, we always restrict the base space $(I \times B_1) \times (I \times B_2)$ to $\cup\{(t \times B_1) \times (t \times B_2) | t \in I\}$. By this hypothesis, the following equalities always hold.

$$(4.1) \quad \begin{aligned} & \Sigma\{(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)\} \\ &= \Sigma(X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2), \end{aligned}$$

$$(4.2) \quad \begin{aligned} & \Sigma\{(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2)\} \\ &= \Sigma(X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} \Sigma(X_2, p_2, B_2, s_2). \end{aligned}$$

In the rest of this paper, consider automatically these identifications if necessary. The we can prove easily the following lemma by the canonical correspondence.

LEMMA 4.2. *For \mathbf{M} -fibrewise pointed spaces (X_i, p_i, B_i, s_i) ($i = 1, 2, 3$),*

(1)

$$\begin{aligned} & ((X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)) \vee^{\mathbf{M}} (X_3, p_3, B_3, s_3) \\ & \cong_{(\mathbf{P})}^{\mathbf{M}} (X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} ((X_2, p_2, B_2, s_2) \vee^{\mathbf{M}} (X_3, p_3, B_3, s_3)) \end{aligned}$$

(2)

$$\begin{aligned} & ((X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2)) \wedge^{\mathbf{M}} (X_3, p_3, B_3, s_3) \\ & \cong_{(\mathbf{P})}^{\mathbf{M}} (X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} ((X_2, p_2, B_2, s_2) \wedge^{\mathbf{M}} (X_3, p_3, B_3, s_3)) \end{aligned}$$

(3)

$$\begin{aligned} & ((X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_2, p_2, B_2, s_2)) \vee^{\mathbf{M}} (X_3, p_3, B_3, s_3) \\ & \cong_{(\mathbf{P})}^{\mathbf{M}} ((X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_3, p_3, B_3, s_3)) \\ & \quad \wedge^{\mathbf{M}} ((X_2, p_2, B_2, s_2) \vee^{\mathbf{M}} (X_3, p_3, B_3, s_3)) \end{aligned}$$

(4)

$$\begin{aligned} & ((X_1, p_1, B_1, s_1) \vee^{\mathbf{M}} (X_2, p_2, B_2, s_2)) \wedge^{\mathbf{M}} (X_3, p_3, B_3, s_3) \\ & \cong_{(\mathbf{P})}^{\mathbf{M}} ((X_1, p_1, B_1, s_1) \wedge^{\mathbf{M}} (X_3, p_3, B_3, s_3)) \\ & \quad \vee^{\mathbf{M}} ((X_2, p_2, B_2, s_2) \wedge^{\mathbf{M}} (X_3, p_3, B_3, s_3)). \end{aligned}$$

DEFINITION 4.3. *For two sequences of \mathbf{M} -fibrewise pointed maps*

$$\mathcal{F} : (X_1, p_1, B_1, s_1) \rightarrow (X_2, p_2, B_2, s_2) \rightarrow \cdots \rightarrow (X_n, p_n, B_n, s_n) \rightarrow \cdots$$

$$\mathcal{F}' : (X'_1, p'_1, B'_1, s'_1) \rightarrow (X'_2, p'_2, B'_2, s'_2) \rightarrow \cdots \rightarrow (X'_n, p'_n, B'_n, s'_n) \rightarrow \cdots$$

if there are \mathbf{M} -fibrewise pointed homotopy equivalences

$$(\phi_n, \alpha_n) : (X_n, p_n, B_n, s_n) \rightarrow (X'_n, p'_n, B'_n, s'_n)$$

such that all diagrams induced by these maps are commutative, \mathcal{F} and \mathcal{F}' have the same \mathbf{M} -fibrewise pointed homotopy type.

The following proposition is obvious from the construction of Puppe exact sequence.

PROPOSITION 4.4. *Let $\alpha : B \rightarrow B'$ be a bijection. The \mathbf{M} -fibrewise pointed homotopy type of Puppe exact sequence (in the sense of \mathbf{M} -fibrewise pointed) induced from an \mathbf{M} -fibrewise pointed map $(\phi, \alpha) : (X, p, B, s) \rightarrow (X', p', B', s')$ is only depend on the \mathbf{M} -fibrewise pointed homotopy class of (ϕ, α) .*

In particular, if (ϕ, α) is \mathbf{M} -fibrewise pointed nulhomotopic, the \mathbf{M} -fibrewise pointed homotopy type of the sequence induced from (ϕ, α) has the

same \mathbf{M} -fibrewise pointed homotopy type of the sequence induced from an \mathbf{M} -fibrewise constant map (c, α_c)

$$\begin{aligned} (X, p, B, s) &\xrightarrow{(c, \alpha_c)} (X', p', B', s') \\ &\longrightarrow (X', p', B', s') \vee^{\mathbf{M}} \Sigma(X, p, B, s) \\ &\longrightarrow \Sigma(X, p, B, s) \\ &\longrightarrow \dots \end{aligned}$$

As an application of this proposition, we prove the generalized formula for the \mathbf{M} -fibrewise pointed suspension of \mathbf{M} -fibrewise pointed product spaces,
Let

$$(u, id) : (X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s') \rightarrow (X, p, B, s) \times (X', p', B', s')$$

be an \mathbf{M} -fibrewise pointed embedding. We denote the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(u, id)$ of (u, id) by $(X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s')$. The Puppe sequence (in the sense of \mathbf{M} -fibrewise pointed) of (u, id) is as follows:

$$\begin{aligned} (X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s') &\xrightarrow{(u, id)} (X, p, B, s) \times (X', p', B', s') \\ &\xrightarrow{(v, \alpha_v)} (X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s') \\ &\xrightarrow{(w, id)} \Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s') \\ &\longrightarrow \dots \end{aligned}$$

Then we obtain the following.

PROPOSITION 4.5. (w, id) in the above is \mathbf{M} -fibrewise pointed nullhomotopic.

PROOF. Generally, for the inclusion $(f, i) : (X_0, p_0, B_0, s_0) \rightarrow (X_1, p_1, B_1, s_1)$, we can consider the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(f, i)$ of (f, i) as the subspace of $\Gamma(X_1, p_1, B_1, s_1)$. Then the inclusion $(g, \alpha_g) : \Gamma(f, i) \rightarrow \Gamma(X_1, p_1, B_1, s_1)$ is continuous.

By applying this to the inclusion $(X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s') \rightarrow (X, p, B, s) \times (X', p', B', s')$, we find that the inclusion

$$(j, id) : (X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s') \rightarrow \Gamma\{(X, p, B, s) \times (X', p', B', s')\}$$

is continuous. The \mathbf{M} -fibrewise pointed map

$$(w, id) : (X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s') \rightarrow \Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s')$$

can be defined by

$$\begin{aligned} w(t, x, s'(b')) &= ((t, x), (t, s'(b'))) && (x \in X_b, t \in I - \{0\}) \\ w(t, s(b), x') &= ((t, s(b)), (t, x')) && (x' \in X'_{b'}, t \in I - \{0\}) \\ w(0, x, x') &= ((0, s(b)), (0, s'(b'))) && (x \in X_b, x' \in X'_{b'}). \end{aligned}$$

Further, let

$$\begin{aligned}\overline{\Sigma}(X, p, B, s) &= \Sigma(X, p, B, s) /_{\mathbf{M}} \left\{ (t, x) \mid x \in X, t \leq \frac{1}{2} \right\} \\ \underline{\Sigma}(X', p', B', s') &= \Sigma(X', p', B', s') /_{\mathbf{M}} \left\{ (t, x') \mid x' \in X', t \geq \frac{1}{2} \right\}.\end{aligned}$$

Then, since $\{(t, x) \mid x \in X, t \leq \frac{1}{2}\}$ and $\{(t, x) \mid x \in X, t \geq \frac{1}{2}\}$ are \mathbf{M} -fibrewise pointed subspaces of $\Sigma(X, p, B, s)$, those are (\mathbf{M} -fibrewise pointed) homeomorphic to an \mathbf{M} -fibrewise pointed cone, so those are \mathbf{M} -fibrewise pointed contractible from Proposition 3.7. Therefore from Proposition 3.8 we see that the natural projections

$$\Sigma(X, p, B, s) \rightarrow \overline{\Sigma}(X, p, B, s), \quad \Sigma(X', p', B', s') \rightarrow \underline{\Sigma}(X', p', B', s')$$

are \mathbf{M} -fibrewise pointed homotopy equivalences. By patching the projections, the \mathbf{M} -fibrewise pointed map

$$(\rho, id) : \Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s') \rightarrow \overline{\Sigma}(X, p, B, s) \vee^{\mathbf{M}} \underline{\Sigma}(X', p', B', s')$$

is also an \mathbf{M} -fibrewise pointed homotopy equivalence.

Next, we can define an \mathbf{M} -fibrewise pointed map

$$(q, id) : \Gamma\{(X, p, B, s) \times (X', p', B', s')\} \rightarrow \overline{\Sigma}(X, p, B, s) \vee^{\mathbf{M}} \underline{\Sigma}(X', p', B', s')$$

as follows:

$$q(t, x, x') = \begin{cases} [(t, s(p(x))), (t, s'(p'(x')))] & (t \in \{0, \frac{1}{2}, 1\}) \\ ((t, s(p(x))), (t, x')) & (0 < t < \frac{1}{2}) \\ ((t, x), (t, s'(p'(x')))) & (\frac{1}{2} < t < 1). \end{cases}$$

Then it is easy to see that $(q, id) \circ (j, id) = (\rho, id) \circ (w, id)$. On the other hand, since $\Gamma\{(X, p, B, s) \times (X', p', B', s')\}$ is \mathbf{M} -fibrewise pointed contractible from Proposition 3.7, $(\rho, id) \circ (w, id)$ is \mathbf{M} -fibrewise pointed nulhomotopic. Since (ρ, id) is an \mathbf{M} -fibrewise pointed homotopy equivalence, (w, id) is also \mathbf{M} -fibrewise pointed nulhomotopic, which completes the proof. \square

From Propositions 4.4 and 4.5, we find that the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(w, id)$ of (w, id) has the same \mathbf{M} -fibrewise pointed homotopy type of

$$\Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s') \vee^{\mathbf{M}} \Sigma\{(X, p, B, s) \overline{\wedge}^{\mathbf{M}} (X', p', B', s')\}.$$

Since the \mathbf{M} -fibrewise pointed mapping cone $\Gamma(w, id)$ has the same \mathbf{M} -fibrewise pointed homotopy type of $\Sigma\{(X, p, B, s) \times (X', p', B', s')\}$, we have the following.

PROPOSITION 4.6.

$$\begin{aligned}\Sigma\{(X, p, B, s) \times (X', p', B', s')\} &\cong_{(\mathbf{P})}^{\mathbf{M}} \\ \Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s') \vee^{\mathbf{M}} \Sigma\{(X, p, B, s) \overline{\wedge}^{\mathbf{M}} (X', p', B', s')\}.\end{aligned}$$

We now define the following.

DEFINITION 4.7. *An \mathbf{M} -fibrewise pointed space (X, p, B, s) is called \mathbf{M} -fibrewise well-pointed if $(s, id_B) : (B, id_B, B, id_B) \rightarrow (X, p, B, s)$ is an \mathbf{M} -fibrewise pointed cofibration and $s(B)$ is closed in X .*

The next lemma is obvious from Proposition 2.4 in this paper.

LEMMA 4.8. *Assume that \mathbf{M} -fibrewise pointed spaces (X, p, B, s) and (X', p', B', s') are \mathbf{M} -fibrewise well-pointed. Then the \mathbf{M} -fibrewise pointed pair*

$$((X, p, B, s) \times (X', p', B', s'), (X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s'))$$

is an \mathbf{M} -fibrewise pointed cofibred pair.

The next proposition is easily verified from this lemma and Proposition 3.9.

PROPOSITION 4.9. *Assume that two \mathbf{M} -fibrewise pointed spaces (X, p, B, s) and (X', p', B', s') are \mathbf{M} -fibrewise well-pointed. Then the natural projection*

$$\begin{aligned} & (X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s') \\ & \rightarrow (X, p, B, s) \bar{\wedge}^{\mathbf{M}} (X', p', B', s') / \mathbf{M}\Gamma\{(X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s')\} \\ & = (X, p, B, s) \times (X', p', B', s') / \mathbf{M}(X, p, B, s) \vee^{\mathbf{M}} (X', p', B', s') \\ & = (X, p, B, s) \wedge^{\mathbf{M}} (X', p', B', s') \end{aligned}$$

is an \mathbf{M} -fibrewise pointed homotopy equivalence.

We have the following from Propositions 4.6 and 4.9.

COROLLARY 4.10. *Assume that two \mathbf{M} -fibrewise pointed spaces (X, p, B, s) and (X', p', B', s') are \mathbf{M} -fibrewise well-pointed. Then the next formula holds.*

$$\begin{aligned} & \Sigma\{(X, p, B, s) \times (X', p', B', s')\} \cong_{(\mathbf{P})}^{\mathbf{M}} \\ & \Sigma(X, p, B, s) \vee^{\mathbf{M}} \Sigma(X', p', B', s') \vee^{\mathbf{M}} \Sigma\{(X, p, B, s) \wedge^{\mathbf{M}} (X', p', B', s')\}. \end{aligned}$$

By repeatedly using this formula and (4.1),(4.2), we can obtain the following formula.

THEOREM 4.11. *Assume that \mathbf{M} -fibrewise pointed spaces (X_i, p_i, B_i, s_i) , $(i = 1, \dots, n)$ are \mathbf{M} -fibrewise well-pointed. Then the next formula holds.*

$$\sum \left\{ \prod_{i=1}^n (X_i, p_i, B_i, s_i) \right\} \cong_{(\mathbf{P})}^{\mathbf{M}} \bigvee_N^{\mathbf{M}} \sum \left(\bigwedge_{i \in N}^{\mathbf{M}} (X_i, p_i, B_i, s_i) \right)$$

where N runs through all nonempty subsets of $\{1, \dots, n\}$.

5. AN INTERMEDIATE FIBREWISE CATEGORY \mathbf{TOP}_B^H

The category \mathbf{MAP} is an extended one of the category \mathbf{TOP}_B . Further, the proofs of theorems in \mathbf{MAP} in this paper are simpler than those in \mathbf{TOP}_B in a sense; for example, we need not to consider any generalized concept of fibrewise non-degenerate spaces [5; section 22]. But, we have to examine closely whether our theorems and propositions in this paper give another proofs of corresponding theorems and propositions in [5; sections 21 and 22]. For these, we introduce an intermediate fibrewise category \mathbf{TOP}_B^H which combines \mathbf{MAP} with \mathbf{TOP}_B . It is easily verified that Theorem 5.4 in \mathbf{TOP}_B^H is proved by the same methods in \mathbf{MAP} . Further, we can prove in Proposition 5.5 that a fibrewise non-degenerate space is H -fibrewise well pointed. Finally, we can give another proof of [5] Proposition 22.11 as a corollary of Theorem 5.4 using Proposition 5.5.

In this section, for a fixed topological space B we consider in the category \mathbf{TOP}_B . In case considering homotopies in \mathbf{TOP}_B , for a fibrewise pointed space (X, p, B, s) we consider the fibrewise pointed space $(I \times X, id \times p, I \times B, id \times s)$, and the following fibrewise pointed spaces have the base space $I \times B$: fibrewise mapping cylinder, fibrewise pointed cone, fibrewise pointed mapping cone, fibrewise pointed suspension. Further, we naturally identify the diagonal Δ_B (or Δ_{B^n}) with B . Therefore we denote this category by \mathbf{TOP}_B^H . Terminologies “fibrewise \dots ” are in \mathbf{TOP}_B , and “ H -fibrewise \dots ” are in \mathbf{TOP}_B^H . But we use the both in \mathbf{TOP}_B^H . We begin with the following definitions.

DEFINITION 5.1. (1) (cf. Definition 2.1) Let

$$(\phi, id_B), (\theta, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$$

be fibrewise pointed maps. If there exists an \mathbf{M} -fibrewise pointed map $(H, h) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B, s')$ such that (H, h) is an \mathbf{M} -fibrewise homotopy of (ϕ, id_B) into (θ, id_B) with $h\delta_t = id_B$ for $t \in I$, we call it an H -fibrewise pointed homotopy of (ϕ, id_B) into (θ, id_B) . If there exists an \mathbf{M} -fibrewise pointed homotopy of (ϕ, id_B) into (θ, id_B) , we say (ϕ, id_B) is H -fibrewise pointed homotopic to (θ, id_B) and write $(\phi, id_B) \simeq_{(\mathbf{P})}^H (\theta, id_B)$.

A fibrewise pointed map $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$ is called an H -fibrewise pointed homotopy equivalence if there exists a fibrewise pointed map $(\theta, id_B) : (X', p', B, s') \rightarrow (X, p, B, s)$ such that $(\theta\phi, id_B id_B) \simeq_{(\mathbf{P})}^H (id_X, id_B)$, $(\phi\theta, id_B id_B) \simeq_{(\mathbf{P})}^H (id_{X'}, id_B)$. Then we denote $(X, p, B, s) \cong_{(\mathbf{P})}^H (X', p', B, s')$.

(It is obvious that the relations $\simeq_{(\mathbf{P})}^H$ and $\cong_{(\mathbf{P})}^H$ are equivalence relations.)

- (2) (cf. Definition 2.2) A fibrewise pointed map $(u, id_B) : (X_0, p_0, B, s_0) \rightarrow (X, p, B, s)$ is an H -fibrewise pointed cofibration if (u, id_B) has the following H -fibrewise homotopy extension property : Let $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$ be a fibrewise pointed map and $(H, h) : (I \times X_0, id \times p_0, I \times B, id \times s_0) \rightarrow (X', p', B, s')$ an H -fibrewise pointed homotopy such that the following two diagrams

$$\begin{array}{ccc} X_0 & \xrightarrow{\sigma_0} & I \times X_0 & & B & \xrightarrow{\delta_0} & I \times B \\ u \downarrow & & \downarrow H & & id_B \downarrow & & \downarrow h \\ X & \xrightarrow{\phi} & X' & & B & \xrightarrow{id_B} & B \end{array}$$

are commutative. Then there exists an H -fibrewise pointed homotopy $(K, k) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B, s')$ such that $K\kappa_0 = \phi, K(id \times u) = H, k\rho_0 = id_B, k(id \times id_B) = h$, where $\kappa_0 : X \rightarrow I \times X$ and $\rho_0 : B \rightarrow I \times B$ are defined by $\kappa_0(x) = (0, x)$ and $\rho_0(b) = (0, b)$ for $x \in X, b \in B$.

- (3) (cf. Definition 2.3) For a fibrewise pointed subspace (X_0, p_0, B, s_0) of (X, p, B, s) , the pair $((X, p, B, s), (X_0, p_0, B, s_0))$ is called by a closed fibrewise pointed pair if X_0 is closed in X . For a fibrewise pointed pair $((X, p, B, s), (X_0, p_0, B, s_0))$, if the inclusion map $(u, id_B) : (X_0, p_0, B, s_0) \rightarrow (X, p, B, s)$ is an H -fibrewise pointed cofibration, we call the pair $((X, p, B, s), (X_0, p_0, B, s_0))$ an H -fibrewise pointed cofibred pair.
- (4) (cf. Definition 2.5) For a fibrewise pointed map $(u, id_B) : (X_0, p_0, B, s_0) \rightarrow (X_1, p_1, B, s_1)$, we can construct the H -fibrewise pointed push-out $(M, p, I \times B, s)$ of the cotriad

$$(I \times X_0, id \times p_0, I \times B, id \times s_0) \xleftarrow{(\sigma_0, \delta_0)} (X_0, p_0, B, s_0) \xrightarrow{(u, id_B)} (X_1, p_1, B, s_1)$$

where (σ_0, δ_0) is an \mathbf{M} -fibrewise embedding to 0-level, as follows : $M = (I \times X_0 + X_1) / \sim$, where $(0, a) \sim u(a)$ for $a \in X_0$, and $p : M \rightarrow I \times B$ and $s : I \times B \rightarrow M$ are defined, respectively, by

$$p(x) = \begin{cases} [t, p_0(a)] & \text{if } x = [t, a], t \neq 0 \\ [p_0(a)] & \text{if } x = [u(a)], a \in X_0 \\ [p_1(x)] & \text{if } x \in X_1 - u(X_0), \end{cases}$$

$$s(t, b) = \begin{cases} (t, s_0(b)) & \text{if } t \neq 0 \\ (0, s_1(b)) & \text{if } t = 0. \end{cases}$$

where $[*]$ is the equivalence class. Then it is easily verified that p and s are well-defined and continuous. We call the H -fibrewise push-out of the cotriad the H -fibrewise mapping cylinder of (u, id_B) , and denote by $M(u, id_B)$.

- (5) (cf. Definition 2.8) For a fibrewise pointed space (X, p, B, s) , we call the \mathbf{M} -fibrewise pointed collapse (the same space in Definition 2.8)

$$(I \times X, id \times p, I \times B, id \times s) /_{\mathbf{M}} (1 \times X, id \times p | 1 \times X, 1 \times B, id \times s | 1 \times B)$$

the H -fibrewise pointed cone of (X, p, B, s) and denote by $\Gamma_H(X, p, B, s)$ in \mathbf{TOP}_B^H . (We denote the total space of $\Gamma_H(X, p, B, s)$ by $C_H X$.)

- (6) (cf. Definition 2.9) Let $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$ be a fibrewise pointed map. Then we call (ϕ, id_B) to be H -fibrewise pointed nullhomotopic if there is a fibrewise pointed map $(c, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$ such that $c = s'p$ and $(\phi, id_B) \simeq_{(\mathbf{P})}^H (c, id_B)$.
- (7) (cf. Definition 3.1) For a fibrewise pointed map $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$, we call $((C_H X) \cup_{\phi} X', \tilde{p}, (I \times B) \cup_{id_B} B, \tilde{s})$ the H -fibrewise pointed mapping cone of (ϕ, id_B) , and denote $\Gamma_H(\phi, id_B)$, where $C_H X$ is the space in this definition (5).
- (8) (cf. Definition 3.5) A fibrewise pointed space (X, p, B, s) is H -fibrewise pointed contractible if there is a fibrewise pointed space (B, id_B, B, id_B) such that

$$(X, p, B, s) \cong_{(\mathbf{P})}^H (B, id_B, B, id_B).$$

- (9) (cf. Definition 3.11) We use the same notation of Definition 3.11. For a fibrewise pointed space (X, p, B, s) , we call the \mathbf{M} -fibrewise pointed collapse (the same space in Definition 3.10)

$$(I \times X, id \times p, I \times B, id \times s) /_{\mathbf{M}} (\overline{X}, \overline{p}, \overline{B}, \overline{s})$$

H -fibrewise pointed suspension, and denoted by $\Sigma_H(X, p, B, s)$ in \mathbf{TOP}_B^H . (We denote $\Sigma_H(X, p, B, s) = (\Sigma_H X, \Sigma_H p, I \times B, \Sigma_H s)$.)

- (10) (cf. Definition 4.1) For two fibrewise pointed spaces (X_1, p_1, B, s_1) and (X_2, p_2, B, s_2) let

$$\begin{aligned} X_1 \vee^H X_2 &= \cup_{b \in B} (X_{1,b} \times s_2(b) \cup s_1(b) \times X_{2,b}), \\ p_1 \vee^H p_2 &= p_1 \times p_2 | X_1 \vee^H X_2. \end{aligned}$$

The fibrewise pointed space $(X_1 \vee^H X_2, p_1 \vee^H p_2, \Delta_B, s_1 \times s_2 | \Delta_B)$ is called the H -fibrewise pointed coproduct of (X_1, p_1, B, s_1) and (X_2, p_2, B, s_2) , and denoted by $(X_1, p_1, B, s_1) \vee^H (X_2, p_2, B, s_2)$ or $\bigvee_{i \in \{1,2\}}^H (X_i, p_i, B, s_i)$. The fibrewise pointed collapse

$$(X_1 \times_B X_2, p_1 \times_B p_2, \Delta_B, s_1 \times s_2 | \Delta_B) /_B (X_1, p_1, B, s_1) \vee^H (X_2, p_2, B, s_2)$$

is called the H -fibrewise smash product of (X_1, p_1, B, s_1) and (X_2, p_2, B, s_2) , and denoted by $(X_1, p_1, B, s_1) \wedge^H (X_2, p_2, B, s_2)$ or $\bigwedge_{i \in \{1,2\}}^H (X_i, p_i, B, s_i)$, where $p_1 \times_B p_2 = p_1 \times p_2 | X_1 \times_B X_2$. (Note that from $\Delta_B \approx B$, as sets we can put $(X_1, p_1, B, s_1) \vee^H (X_2, p_2, B, s_2) = X_1 \vee_B X_2$ and $(X_1, p_1, B, s_1) \wedge^H (X_2, p_2, B, s_2) = X_1 \wedge_B X_2$.)

For $n \geq 3$ and fibrewise pointed spaces (X_i, p_i, B, s_i) ($i = 1, \dots, n$), we shall define $\bigvee_{i \in \{1, \dots, n\}}^H (X_i, p_i, B, s_i)$ and $\bigwedge_{i \in \{1, \dots, n\}}^H (X_i, p_i, B, s_i)$ inductively, as follows:

$$\begin{aligned} \bigvee_{i \in \{1, \dots, n\}}^H (X_i, p_i, B, s_i) &= \left(\bigvee_{i \in \{1, \dots, n-1\}}^H (X_i, p_i, B, s_i) \right) \vee^H (X_n, p_n, B, s_n) \\ \bigwedge_{i \in \{1, \dots, n\}}^H (X_i, p_i, B, s_i) &= \left(\bigwedge_{i \in \{1, \dots, n-1\}}^H (X_i, p_i, B, s_i) \right) \wedge^H (X_n, p_n, B, s_n). \end{aligned}$$

(11) (cf. Definition 4.3) For two sequences of fibrewise pointed maps

$$\mathcal{F} : (X_1, p_1, B, s_1) \rightarrow (X_2, p_2, B, s_2) \rightarrow \dots \rightarrow (X_n, p_n, B, s_n) \rightarrow \dots$$

$$\mathcal{F}' : (X'_1, p'_1, B, s'_1) \rightarrow (X'_2, p'_2, B, s'_2) \rightarrow \dots \rightarrow (X'_n, p'_n, B, s'_n) \rightarrow \dots$$

if there are H -fibrewise pointed homotopy equivalences

$$(\phi_n, id_B) : (X_n, p_n, B, s_n) \rightarrow (X'_n, p'_n, B, s'_n)$$

such that all diagrams induced by these maps are commutative, \mathcal{F} and \mathcal{F}' have the same H -fibrewise pointed homotopy type.

(12) (cf. Definition 4.7) A fibrewise pointed space (X, p, B, s) is called H -fibrewise well-pointed if $(s, id_B) : (B, id_B, B, id_B) \rightarrow (X, p, B, s)$ is an H -fibrewise pointed cofibration and $s(B)$ is closed in X . (Note that Definition 4.7 is in **MAP** and this definition is in \mathbf{TOP}_B^H .)

REMARK 5.2. By Definition 5.1 (9) and (10), the base space of $\Sigma_H\{(X_1, p_1, B, s_1) \vee^H (X_2, p_2, B, s_2)\}$ and $\Sigma_H\{(X_1, p_1, B, s_1) \wedge^H (X_2, p_2, B, s_2)\}$ is $I \times \Delta_B$, but the base space of $\Sigma_H(X_1, p_1, B, s_1) \vee^H \Sigma_H(X_2, p_2, B, s_2)$ and $\Sigma_H(X_1, p_1, B, s_1) \wedge^H \Sigma_H(X_2, p_2, B, s_2)$ is $\Delta_{I \times B}$. So, we naturally identify $I \times \Delta_B$ with $\Delta_{I \times B}$, and we can have the same formulas (4.1) and (4.2) in \mathbf{TOP}_B^H .

If we consider in \mathbf{TOP}_B^H the theory of sections 2, 3 and 4, it can be verified by slight modifications that we have the same type theorems and propositions in \mathbf{TOP}_B^H . Thus we have the following main theorems.

THEOREM 5.3. For an H -fibrewise pointed map $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$, the following sequence is exact in \mathbf{TOP}_B^H .

$$\begin{array}{ccc} (X, p, B, s) & \xrightarrow{(\phi, id_B)} & (X', p', B, s') \\ & \xrightarrow{(\phi', \delta_0)} & \Gamma_H(\phi, id_B) \\ & \xrightarrow{(\phi'', id \times id_B)} & \Sigma_H(X, p, B, s) \\ & \xrightarrow{(\phi''', id \times id_B)} & \Sigma_H(X', p', B, s') \\ & \longrightarrow & \dots \end{array}$$

THEOREM 5.4. *Assume that fibrewise pointed spaces (X_i, p_i, B, s_i) ($i = 1, \dots, n$) are H -fibrewise well-pointed. Then the next formula holds in \mathbf{TOP}_B^H .*

$$\begin{aligned} \sum_H \left(\prod_{i=1}^n {}_B X_i, \prod_{i=1}^n p_i, \Delta_{B^n}, \prod_{i=1}^n s_i | \Delta_{B^n} \right) \\ \cong_{(\mathbf{P})}^H \bigvee_N^H \sum_H \left(\bigwedge_{i \in N}^H (X_i, p_i, B, s_i) \right) \end{aligned}$$

where N runs through all nonempty subsets of $\{1, \dots, n\}$.

We restate the concept of fibrewise non-degenerate space ([5; Definition 22.2]). Let (X, p, B, s) be a fibrewise pointed space. We regard the fibrewise mapping cylinder $M_B(s)$ of s (cf. [5; section 18]) as a fibrewise pointed space with the section $\check{s} : B \rightarrow M_B(s)$ defined by $\check{s}(b) = (1, b)$, and denote it by $(\check{X}_B, \check{p}, B, \check{s})$. Then the fibrewise pointed space (X, p, B, s) is *fibrewise non-degenerate* if the natural projection $(\rho, id_B) : (\check{X}_B, \check{p}, B, \check{s}) \rightarrow (X, p, B, s)$ is a fibrewise pointed homotopy equivalence.

We now prove the following proposition.

PROPOSITION 5.5. *Let (X, p, B, s) be a fibrewise non-degenerate space. Then (X, p, B, s) is an H -fibrewise well-pointed space.*

PROOF. Since (X, p, B, s) is fibrewise non-degenerate, the natural projection $(\rho, id_B) : (\check{X}_B, \check{p}, B, \check{s}) \rightarrow (X, p, B, s)$ is a fibrewise homotopy equivalence. Therefore there is a fibrewise pointed map $(\eta, id_B) : (X, p, B, s) \rightarrow (\check{X}_B, \check{p}, B, \check{s})$ (with $\eta s = \check{s}$) and a fibrewise pointed homotopy $(G, id_B) : I \tilde{\times} \check{X}_B \rightarrow \check{X}_B$ such that $G_0 = id_{\check{X}_B}$ and $G_1 = \eta\rho$, where $I \tilde{\times} \check{X}_B$ is the reduced fibrewise cylinder of $I \times \check{X}_B$ ([5; section 19]).

To prove that (X, p, B, s) is H -fibrewise well pointed, let $(\phi, id_B) : (X, p, B, s) \rightarrow (X', p', B, s')$ be a fibrewise pointed map, and

$$(H, h) : (I \times B, id \times id_B, I \times B, id \times id_B) \rightarrow (X', p', B, s')$$

an H -fibrewise pointed homotopy such that $H(0, b) = \phi(s(b))$ and $h(t, b) = b$. Then we can construct an H -fibrewise pointed homotopy $(\tilde{H}, \tilde{h}) : (I \times X, id \times p, I \times B, id \times s) \rightarrow (X', p', B, s')$ as follows: For any $(t, x) \in I \times X$, $(t, b) \in I \times B$,

$$\begin{aligned} \tilde{H}(t, x) &= \begin{cases} \phi\rho G(t, (t, x)) & (t = 0) \\ H(t, p\rho G(t, (t, sp(x)))) & (t \neq 0), \end{cases} \\ \tilde{h}(t, b) &= b. \end{aligned}$$

It is easily verified that (\tilde{H}, \tilde{h}) is an H -fibrewise pointed homotopy such that $(\tilde{H}, \tilde{h}) \circ (id \times s, id \times id_B) = (H, h)$ and $(\tilde{H}, \tilde{h}) \circ (i_X, i_B) = (\phi, id_B)$ where $i_X : X \rightarrow I \times X$ defined by $i_X(x) = (0, x)$ and $i_B : B \rightarrow I \times B$ defined by $i_B(b) = (0, b)$. This completes the proof. \square

Using this proposition we can prove [5] Proposition 22.11 as a corollary of Theorem 5.4, which gives an another proof of [5] Proposition 22.11.

PROPOSITION 22.11([5]) *Assume that fibrewise pointed spaces (X_i, p_i, B, s_i) ($i = 1, \dots, n$) are fibrewise non-degenerate spaces. Then the next formula holds in \mathbf{TOP}_B .*

$$\sum_B^B (X_1 \times_B \dots \times_B X_n) \cong_B^B \bigvee_N^B \sum_B^B \bigwedge_{i \in N}^B X_i$$

where N runs through all nonempty subsets of $\{1, \dots, n\}$.

PROOF. First, note from Proposition 5.5 that each fibrewise non-degenerate space (X_i, p_i, B, s_i) ($i = 1, \dots, n$) is H -fibrewise well pointed. Next, we can prove this proposition by the following steps.

(1) For a fibrewise pointed space (X, p, B, s) , the total space (we denote $\Sigma_B^B X$) of the reduced fibrewise suspension $\Sigma_B^B(X)$ in \mathbf{TOP}_B can be obtained from the total space $\Sigma_H X$ of $\Sigma_H(X, p, B, s)$ in \mathbf{TOP}_B^H as follows: the space $\Sigma_B^B X$ is just equal to the fibrewise push-out of the cotriad

$$\Sigma_H X \xleftarrow{i} (id \times s)(I \times B) \xrightarrow{\pi} \{0\} \times s(B)$$

where i is the natural inclusion and π is the natural projection. Further the diagram

$$\begin{array}{ccc} \Sigma_H X & \xrightarrow{c_t} & \Sigma_B^B X \\ \Sigma_H p \downarrow & & \downarrow \Sigma_B^B p \\ I \times B & \xrightarrow{\pi_2} & B \end{array}$$

is commutative, where c_t is the (H -fibrewise pointed) compact map obtaining from the fibrewise push-out of the cotriad in the above, and $\Sigma_B^B p$ is the projection of $\Sigma_B^B(X)$.

(2) From $I \times \Sigma_B^B X$, we construct the reduced fibrewise cylinder $I \tilde{\times} \Sigma_B^B X$ ([5; section 19]), and obtain the natural map $id \tilde{\times} c_t : I \times \Sigma_H X \rightarrow I \tilde{\times} \Sigma_B^B X$ which is an (H -fibrewise pointed) compact map. Further, for an H -fibrewise pointed homotopy

$$(H, h) : I \times \Sigma_H(X, p, B, s) \rightarrow \Sigma_H(X, p, B, s),$$

we can define naturally the fibrewise pointed homotopy $(\tilde{H}, id_B) : I\tilde{\times}\Sigma_B^B(X) \rightarrow \Sigma_B^B(X)$ such that the following diagram is commutative.

$$\begin{array}{ccccc}
 I \times \Sigma_H X & \xrightarrow{H} & \Sigma_H X & & \\
 \downarrow id \times \Sigma_H p & \searrow id \tilde{\times} c_t & \downarrow & \searrow c_t & \\
 I \times (I \times B) & & I \tilde{\times} \Sigma_B^B X & \xrightarrow{\tilde{H}} & \Sigma_B^B X \\
 \downarrow & \downarrow & \downarrow & & \downarrow \\
 I \times B & \xrightarrow{h} & I \times B & & B \\
 \downarrow & & \downarrow & & \downarrow \\
 B & \xrightarrow{id_B} & B & & B
 \end{array}$$

(3) For the fibrewise pointed spaces (X_i, p_i, B, s_i) ($i = 1, \dots, n$), we can consider the formula in Theorem 5.4. Let

$$\begin{aligned}
 (X, p, I \times B, s) &= \sum_H \left(\prod_{i=1}^n {}_B X_i, \prod_{i=1}^n {}_B p_i, \Delta_{B^n}, \prod_{i=1}^n s_i | \Delta_{B^n} \right) \\
 (Y, q, I \times B, t) &= \bigvee_N^H \sum_H \left(\bigwedge_{i \in N}^H (X_i, p_i, B, s_i) \right).
 \end{aligned}$$

Further, let \tilde{X} and \tilde{Y} be the total spaces of

$$\Sigma_B^B(X_1 \times_B \cdots \times_B X_n), \quad \bigvee_N^B \Sigma_B^B \left(\bigwedge_{i \in N} {}_B X_i \right)$$

respectively. Note that Δ_B and Δ_{B^n} are identified with B , and $(X_1, p_1, B, s_1) \vee^H (X_2, p_2, B, s_2) = X_1 \vee_B X_2$, $(X_1, p_1, B, s_1) \wedge^H (X_2, p_2, B, s_2) = X_1 \wedge_B X_2$. By this note and (1) in this proof, there is a compact map $c'_t : Y \rightarrow \tilde{Y}$, and for an H -fibrewise pointed map $(f, id \times id_B) : (X, p, I \times B, s) \rightarrow (Y, q, I \times B, t)$ we can define a fibrewise pointed map $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$ such that the following diagram is commutative.

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 c_t \downarrow & & \downarrow c'_t \\
 \tilde{X} & \xrightarrow{\tilde{f}} & \tilde{Y}
 \end{array}$$

(4) Let $(X, p, I \times B, s)$, $(Y, q, I \times B, t)$, \tilde{X} and \tilde{Y} be the same spaces as those in (3) of this proof. From Theorem 5.4, there are H -fibrewise pointed maps $(f, id \times id_B) : (X, p, I \times B, s) \rightarrow (Y, q, I \times B, t)$ and $(g, id \times id_B) : (Y, q, I \times B, t) \rightarrow (X, p, I \times B, s)$ such that $(gf, (id \times id_B)(id \times id_B)) \simeq_{(\mathbf{P})}^H (id_X, id \times id_B)$, $(fg, (id \times id_B)(id \times id_B)) \simeq_{(\mathbf{P})}^H (id_Y, id \times id_B)$. From these

H -homotopies and H -fibrewise pointed maps $(f, id \times id_B), (g, id \times id_B)$, we can construct, by the same methods of (2) and (3) in this proof, fibrewise pointed homotopies, fibrewise pointed maps $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}, \tilde{g} : \tilde{Y} \rightarrow \tilde{X}$ such that $\tilde{g}\tilde{f} \simeq_B^B id_{\tilde{X}}, \tilde{f}\tilde{g} \simeq_B^B id_{\tilde{Y}}$. Thus, we complete the proof of [5] Proposition 22.11 by using Theorem 5.4. \square

REFERENCES

- [1] D. Buhagiar, *The category \mathcal{MAP}* , Mem. Fac. Sci.& Eng., Shimane Univ. **34** (2001), 1-19.
- [2] M. Crabb and I.M. James, *Fibrewise homotopy theory*, Springer Monographs in Mathematics, 1998.
- [3] R. Engelking, *General topology*, Heldermann, Berlin, rev. ed., 1989.
- [4] T. Hotta and T. Miwa, *A new approach to fibrewise fibrations and cofibrations*, Topology and its Appl. **122** (2002), 205-222.
- [5] I.M. James, *Fibrewise topology*, Cambridge Univ. Press, Cambridge, 1989.
- [6] T. Miwa, *On fibrewise retraction and extension*, Houston J. Math. **26** (2000), 811-831.
- [7] B.A. Pasynkov, *On extensions to mappings of certain notions and assertions concerning spaces*, Mappings and Functions (Moscow), Izdat. MGU, Moscow, 1984, 72-102 (in Russian).
- [8] B.A. Pasynkov, *Elements of the general topology of continuous maps, On Compactness and Completeness Properties of Topological Spaces*, "FAN" Acad. of Science of the Uzbek. Rep., Tashkent, 1994, 50-120 (in Russian).
- [9] D. Puppe, *Homotopiemengen und ihre induzierten Abbildungen*, I, Math. Z. **69** (1958), 299-344.
- [10] E.H. Spanier, *Algebraic topology*, McGraw-Hill, 1966.

Y. Konami
Izumo-Nishi High School
Izumo City 693-0032
Japan

T. Miwa
Department of Mathematics
Shimane University
Matsue 690-8504
Japan
E-mail: miwa@riko.shimane-u.ac.jp

Received: 12.5.2004.

Revised: 30.6.2004.