

Evaluation of the Use and Reasons for Not Wearing a Helmet by Motorcyclists Admitted to the Emergency Ward of Shahid Bahonar Hospital in Kerman

Javad Faryabi¹; Mahboobeh Rajabi^{2,*}; Shahin Alirezaee³

¹Department of Oral and Maxillofacial Surgery, Oral and Dental Diseases Research Center, Kerman University of Medical Sciences, Kerman, IR Iran

²Research Center for Health Services Management, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, IR Iran

³Shahid Bahonar Hospital, Kerman, IR Iran

*Corresponding author: Mahboobeh Rajabi, Research Center for Health Services Management, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, IR Iran. Tel: +98-3432111511, Fax: +98-3432111613, E-mail: ma_rajabi_k@yahoo.com

Received: April 13, 2014; Revised: July 18, 2014; Accepted: August 28, 2014

Background: Motorcycle crashes are the cause of severe morbidity and mortality especially because of head injuries. It seems that wearing a helmet has an effective role in protection against head injuries. Nevertheless, motorcyclists usually have no tendency to wear a helmet when driving in cities and have several reasons for this behavior.

Objectives: This study aimed to evaluate the use and reasons for not using a helmet by motorcyclists admitted to an emergency ward of a trauma hospital due to accident in Kerman, Iran.

Patients and Methods: This study was carried out by recoding the opinions of motorcyclists who had been transferred to the emergency ward of Shahid Bahonar Hospital (Kerman/Iran). Since no data was available on the frequency of the use of helmets, a pilot study was carried out and a sample size of 377 was determined for the main study. Then a researcher-made questionnaire was used to investigate the motorcyclists' reasons for not using a helmet.

Results: Only 21.5% of motorcyclists had been wearing helmets at the time of the accident. The most frequent reasons for not using a helmet were the heavy weight of the helmet (77%), feeling of heat (71.4%), pain in the neck (69.4%), feeling of suffocation (67.7%), limitation of movement of the head and neck (59.6%) and all together, physical discomfort was the main cause of not wearing a helmet during motorcycle rides.

Conclusions: In general, it appears that it is possible to increase the use of helmets by eliminating its physical problems, and increasing the knowledge of community members in relation to the advantages of helmet use, which will result in a significant decrease in traumas resulting from motorcycle accidents.

Keywords: Motorcyclists; Helmet; Use; Reasons

1. Background

Road traffic injuries are a major public health and development crisis and in developing countries the process of modernization, has aggravated this crisis. Thus, about 85% of mortalities, 90% of disability adjusted life years (DALY), and 96% of all children deaths that are caused by road traffic accidents (RTAs) worldwide, occur in these countries. Road traffic trauma is the sixth most common cause of death and will be the third leading cause of mortality and morbidity by 2030 in developing countries according to the world health organization (WHO) (1). Iran has the highest rate of road traffic accidents worldwide and RTAs are considered as the third leading cause of mortality (2). Road traffic injuries cause different issues at different levels. At the national level, they cause considerable financial costs, especially for developing economies. These injuries are estimated to cost 1-2% of the gross national product of low-middle income countries, which is estimated at over 100 billion USD per year (3). Also, in many developing countries, conditions of referral centers, availability of modern equipments

and skilled personnel in medical care centers are great issues. In addition, at the individual level, costs of prolonged and intensive medical care, individual and social problems of victims' disability and loss of income due to death or disability of the head of the household can worsen the families' life conditions (1, 4). Motorcycle has been accepted as a suitable vehicle to transfer the individuals in various communities and the public have welcomed its use. Nevertheless, it is a part of many accidents and it is a source of a large number of traumas and deaths (5). Thus, in many developing countries, the majority of people injured or died in road traffic crashes are motorcyclists. However, the number of road fatalities related to motorcyclists in developed countries, where private cars are more popular, is also disproportionately high (6).

In traffic accidents, motorcyclists are at high risk of head, neck, face and traumatic brain injuries (TBI); TBI being one of the most important causes of neuromotor disabilities and mortality in the world and reports from Europe, Asia, South Pacific, Africa and North Amer-

ica have confirmed this claim (7, 8). In addition, some sources have reported more severe traumas due to motorcycle accidents in comparison with car accidents (5). Literature review reveals a mortality rate of 6.8% to 80.6% due to motorcycle accidents (9-11). For example, mortality rate due to motorcycle accidents in India, Indonesia and Malaysia have been 27%, 42% and 57% of all the road accidents. Furthermore, in two provinces of Iran (Fars & Mazandaran) mortality rates were 23.1% and 25.6%, respectively, achieved from recorded data of forensic medicine of these cities (12-14). In this context, the majority of deaths due to motorcycle accidents occur because of brain traumas (5, 15), and some studies showed a significant relationship between head injury and death in traffic crashes (16, 17). Thus, wearing a helmet significantly decreases the odds of deaths due to brain traumas (5, 15). Also, various studies have proved that the use of a helmet resulted in a significant decrease in injuries, brain traumas and mortality, and in turn lead to lives and financial savings. For example in the United States approximately 14,283 motorcyclists died from 2008 to 2010 due to accidents; 42% were not wearing helmets. In addition, financial savings due to the use of helmets in states implementing the universal law of compulsory use of helmets has been estimated as \$725 for each motorcycle, which was four times higher than that in the states that had not implement such a law (18). Studies showed that wearing a helmet prevented traumas by 69% and decreased the odds of deaths by up to 42% and estimate of effectiveness ranges from OR 0.23 to 0.35 (7, 19). Therefore, recognizing the importance of helmet use in reduction of fatalities increased the implementation of helmet laws in different countries (5, 20-22). Although there is a definite relationship between mortality rate of motorcyclists and brain traumas on one hand and not wearing a helmet on the other, factors affecting wearing or not wearing a helmet by motorcyclists have not been evaluated precisely (23). In spite of the obvious benefits of helmet use by motorcyclists, unfortunately they have no interest in wearing a helmet. The rate of helmet use in different cities of Iran varied from 8.6% to 75% (24-26). In other countries this rate was different. For example this was 56% in Karachi Pakistan, 19.7% in Pamplona Spain, and 90-99% in Vietnam, among adults (19, 27).

2. Objectives

This study, for the first time, aimed to evaluate the frequency of wearing a helmet by motorcyclists in Kerman, Iran. As the reasons for not wearing a helmet by motorcyclists are important, we obtained the motorcyclists' opinions about not wearing helmets. Thus, the results may be used to warn all individuals in the community, especially motorcyclists, through mass media.

3. Patients and Methods

Since no precise data was available on the use of helmets

by motorcyclists in Kerman, we conducted two studies: a pilot study and the main study.

A) Pilot study: this study was carried out for a period of one month in order to determine the sample size. During this one-month, 93 motorcyclists were referred to the emergency ward of Shahid Bahonar Hospital, a Trauma Hospital of Kerman, due to motorcycle crashes. Forty of these motorcyclists had been wearing helmets and 53 had not. Therefore, the sample size was estimated as 377 motorcyclists. Thus, 377 motorcyclists who had referred to the emergency ward of this hospital because of crashes during the time of the study were included and the motorcyclists who had not worn helmets were asked to answer the questionnaire.

B) Main study: Subsequently, a researcher-made questionnaire was used to investigate the motorcyclists' reasons for not using helmets. The first part of this questionnaire consisted of demographic characteristics and history of motorcycle accidents and the second part consisted of 15 questions (Table 1). Injured motorcyclists, under supervision of authors, responded to the questionnaire during all working shifts of the hospital. For assessing the content validity of the questionnaire, we distributed the questionnaire between ten members of Injuries and Disasters Prevention Committee of Kerman University of Medical Sciences to gain their opinion about the questions and with respect to their opinion, some questions were changed and the final questionnaire was prepared for our study. The revised questionnaire was checked by the members of that committee again and regarding their opinions the validity of the final questionnaire, as a whole, was estimated as 0.86. The reliability of the questionnaire based on Cronbach's alpha, was 0.75. Statistical significance was defined at $P < 0.05$.

4. Results

The mean age of motorcyclists who had worn and those who had not worn helmets were 28.36 and 26.33 years, respectively. Most motorcyclists had lower than diploma level of education. Furthermore, 52.7% of the motorcyclists had no previous history of motorcycle accidents and 69.5% of them had not used a helmet; 33.9% of the motorcyclists had a history of one accident and 83.8% of them had not used a helmet. Also, 13.4% of the motorcyclists had a history of more than one accident, 87.2% of them had not used a helmet (Table 2). There was no significant relationship between the motorcyclists' age and educational status, and wearing helmet ($P = 0.12$, $P = 0.42$). However, there was a significant relationship between history of motorcycle accident and wearing a helmet ($P = 0.008$) (Table 2).

Age, educational status and a history of accident were considered in the multiple logistic regression models and the adjusted odds ratio (OR) (95% CI) were estimated. These results also showed no significant relationship between age, educational status and wearing a helmet but there was a significant relationship between a history of

motorcycle accident and wearing a helmet (Table 3). From 377 motorcyclists only 81 motorcyclists (21.5%) had worn a helmet at the time of the accident. The motorcyclists' reasons for not wearing a helmet during the motorcycle ride were the heavy weight of the helmet in 77%; neck pain after wearing a helmet in 69.4%; limitation of movements of the head and neck in 59.6%; visual limitation during wearing a helmet in 57%; feeling of heat in 71.4%; feeling of suffocation in 67.7%; unfavorable appearance of the helmet in 55.4%; difficulty of preserving or holding the helmet before and after the ride in 59.2% and being ridiculed by others due to the use of a helmet in 53.9%. In addition, 64.8%

of motorcyclists believed that summer was not a suitable season for wearing a helmet due to excessive heat and 66.3% believed that winter was a suitable season for wearing a helmet. Furthermore, 53.4% of the cases believed that if they obeyed the rules and rode at low speed, there would be no need to use a helmet. In addition, 41.6% of the subjects believed that they did not need a helmet because they were sure that they could keep their balance if an accident occurred and 49.8% believed that quiet streets were the best places to ride motorcycles without the need of wearing a helmet. Also 59.5% confessed that when they were in a hurry, they did not use their helmet.

Table 1. Questionnaire Investigating Motorcyclist's Reasons of not Using a Helmet

Number	Question details	Completely Agree	Agree	No Opinion	Disagree	Completely Disagree
1	The helmet is heavy and I don't use it for this reason					
2	The helmet causes neck pain					
3	I feel heat on my head when wearing the helmet					
4	I have a feeling of suffocation when wearing the helmet					
5	When I ride slowly there is no need to wear a helmet					
6	I think if an accident occurs I can maintain my balance so there is no need to wear a helmet					
7	It causes limitations in neck and head movement					
8	In summary, I can not use a helmet due to excessive warmth					
9	The winter season is suitable for wearing a helmet					
10	When I am in a hurry, I do not use a helmet					
11	In quiet streets and valleys there is no need to wear a helmet					
12	Helmet leads to restriction of vision					
13	Helmets do not have a suitable appearance					
14	Carrying the helmet before and after use is difficult					
15	Some people ridicule motorcyclists who wear helmet					

Table 2. Demographic Characteristics of Motorcyclist^a

	Helmeted Motorcyclists	Non-Helmeted Motorcyclists	P Value
Age, y	28.36 ± 10.33	26.33 ± 10.19	0.12
Education Level			0.42
Illiterate	3.7	7.9	
Under diploma	44.4	47.9	
Diploma	34.6	31.2	
Academic Education	17.3	13	
History of Accident			0.008
Did not have any	30.5	69.5	
One time	16.2	83.8	
More than one time	12.8	87.2	

^a Data are presented as % or Mean ± SD.

Table 3. The Relationship Between Using a Helmet and Demographic Characteristics of Motorcyclist ^a

Chance for Using Helmet		
Variable	Adjusted OR	P Value
Age	1.43 ± 0.074	> 0.05
Education Level		
Under Diploma	2.88 ± 0.13	> 0.05
Diploma	2.81 ± 0.14	
Academic Education	2.47 ± 0.24	
History of Accident		
One time	0.45 ± 0.016	< 0.05
More than one time	0.38 ± 0.062	

^a Abbreviation: OR, odd ratio.

5. Discussion

Motorcycles crashes are considerable causes of injury-related fatality and disability (22). Head injuries as one of the most common injuries after motorcycle crashes are estimated to be the cause of most deaths and long-term disabilities. Also in some studies, head injuries and deaths had a significant relationship (16, 17). In our study, the mean age of motorcyclists was less than 30 years. This finding was consistent with the results of the Janmohammadi et al. study in Mazandaran (28.1%) but lower than the results of other studies in Yazd (34.8%), Fars (31.4%), Tehran (32.3%) and United State of America (38%) (13, 14, 26, 28, 29). This pattern showed that most motorcyclists' injuries and disabilities occur in the productive age group of the society, which causes enormous social and economic problems. The majority of riders had an education level under diploma, similar to Heydari et al. (13), Vafae-Najar et al. (25) and Amirjamshidi et al. (26) studies in Fars, Mashhad and Tehran, respectively. However, in Mehri et al. (29) study in Yazd 54.7% and in McCartt et al. (30) study in the United States 38% of the participants had college education (13, 25, 26, 29, 30). In the present study, there was no significant relationship between wearing a helmet and age and educational status. Also, in a study carried out by Khan et al. (27) in Pakistan the subjects' awareness of the helmet law, their marital status and their average age had no effect on wearing or not wearing a helmet (27). Brown et al. and Dee studies, did not reveal any significant relationship between the use of a helmet and age or gender (28, 31). Nevertheless, in McCartt's study the frequency of helmet use was related to drivers age (30). In the current study, most participants (52.7%) had no history of accidents by motorcycles, similar to McCartt's study (57%). Furthermore, 33.9% of the participants had a history of one accident, 83.8% of which had not worn a helmet and 13.4% had a history of more than one motorcycle accident, 87.2% of which had not worn a helmet. These findings suggest a pattern of persis-

tent high-risk behavior among people who have the experience of an accident for not using a helmet. In Mehri et al. study, only 10.7% of all motorcyclists with a history of motorcycle crash used a helmet and in Moghisi et al. study 55% of the motorcyclists had at least one accident and only two of them were wearing helmets at the time of the accident (29, 32). Also, there was a significant relationship between a previous history of motorcycle accident and wearing a helmet. Arguments against helmet use for motorcycle riders include the claims of increasing the risk of neck injuries and severity of other body injuries in crashes, decreasing of rider visibility and effectiveness of helmets in reducing mortality (7). However, against these claims the results of different studies in different countries have proved that wearing standard helmets significantly decreases mortality rate due to traumas during riding a motorcycle. Also studies showed a decrease in the rates of fatal injuries and deaths of motorcyclists after the helmet-use laws (33, 34). For example, Dee's (31) study showed that State laws requiring helmet use, reduced motorcyclist fatalities by 27%. Based on the result of the present study, 21.5% of motorcyclists in Kerman wore helmets. This rate was different in other cities of Iran like Tehran during 1999-2000 (8.6%) and 2007 (75%), Mashhad (15.8%) and Ahwaz (10%) (24-26, 35). In Karachi of Pakistan this rate was 56%, in Pamplona of Spain, 19.7%, which increased to 94.8% after the law of compulsory use of a helmet by motorcyclists took effect, in Indonesia 89%, in Texas (1994-2006) 56% and in Ghana 34.2% (27, 28, 36). The reasons for different percentages of using safety helmets by motorcyclists in various studies may be due to cultural behaviors, mandatory motorcycle helmet laws, economic power of people and urban or rural site for riding of motorcycles. Also the type of helmet worn, correct fastening of helmets and cost are secondary issues that are particularly relevant to motorcycle helmet usage in developing countries (7). In the present study the most common reason for not wearing a helmet was the weight of the helmet (77%), and other reasons were the feeling of heat during helmet use (71.4%), neck pain (69.4%), feeling of suffocation (67.7%) and limitations in the movements of the head and neck (59.6%). In Moghisi et al. (32) study looking funny 73.2%, disturbing issues 44.4%, decreased hearing 36.4% and feeling of heat during helmet use 30%, were the most common reasons of not using a helmet. Khan reported that 44% of motorcyclists did not wear a helmet due to physical discomfort and 25% due to visual limitations (27). In a study by Conrad in Indonesia there was differences in time and place where motorcyclists reported wearing helmets. They used helmets rarely at night and physical discomfort and absence of police surveillance was the most common reasons for not wearing helmets (37). Thus, in general it can be concluded that physical discomfort was the most important reason for not wearing a helmet in this study, consistent with the results of other studies. Since in many cases, motorcyclists were the breadwinners of the family and they

were at their productive age, motorcycles' safety devices become of greater importance. Thus manufactures of helmets must consider factors that determine the ease of use and motorcyclists comfort, in construction of standard helmets. Also, motorcyclists training and planning for changing their attitudes and behaviors, should be considered in the future to help increase the usage of helmets and decrease risky behaviors during riding.

Acknowledgements

The authors would like to thank the Vice Chancellor for Research of Kerman University of Medical Sciences and the Kerman Traffic Police Administration for their assistance.

Authors' Contributions

Javad Faryabi: study design, abstraction of data and critical revision of the manuscript for important intellectual content. Mahbobeh Rajabi: data collection, study supervision and manuscript writing. Shahin Alirezaee: data collection.

Funding/Support

This research was part of a project approved and supported by the Vice Chancellor for Research of Kerman University of Medical Sciences, Kerman, Iran.

References

- Peden M, Scurfield R, Sleet D, Mohan D, Hyder AA, Jarawan E, et al. *World Report on Road Traffic Injury Prevention*. Geneva: World Health Organization; 2004. Available from: <http://apps.who.int/bookorders/anglais/detart1.jsp?codlan=1&codcol=15&codch=572>.
- Bhalla K, Naghavi M, Shahrzaz S, Bartels D, Murray CJ. Building national estimates of the burden of road traffic injuries in developing countries from all available data sources: Iran. *Inj Prev*. 2009;**15**(3):150-6.
- World Health Organization. *Global status report on road safety 2013 supporting a decade of action*. Geneva; WHO. 2013.
- Ebrahimzadeh MH. National trauma registry center, as a backbone of trauma management and research. *Arch Trauma Res*. 2012;**1**(3):87-8.
- Rowland J, Rivara F, Salzberg P, Soderberg R, Maier R, Koepsell T. Motorcycle helmet use and injury outcome and hospitalization costs from crashes in Washington State. *Am J Public Health*. 1996;**86**(1):41-5.
- Mohan D. Road safety in less-motorized environments: future concerns. *Int J Epidemiol*. 2002;**31**(3):527-32.
- Liu BC, Ivers R, Norton R, Boufous S, Blows S, Lo SK. Helmets for preventing injury in motorcycle riders. *Cochrane Database Syst Rev*. 2008(1):CD004333.
- Servadei F, Begliomini C, Gardini E, Giustini M, Taggi F, Kraus J. Effect of Italy's motorcycle helmet law on traumatic brain injuries. *Inj Prev*. 2003;**9**(3):257-60.
- Nakahara S, Chadbunchachai W, Ichikawa M, Tipsuntornsak N, Wakai S. Temporal distribution of motorcyclist injuries and risk of fatalities in relation to age, helmet use, and riding while intoxicated in Khon Kaen, Thailand. *Accid Anal Prev*. 2005;**37**(5):833-42.
- Oginni FO, Ugboko VI, Ogundipe O, Adegbehingbe BO. Motorcycle-related maxillofacial injuries among Nigerian intracity road users. *J Oral Maxillofac Surg*. 2006;**64**(1):56-62.

- Tsai MC, Hemenway D. Effect of the mandatory helmet law in Taiwan. *Inj Prev*. 1999;**5**(4):290-1.
- Hyder AA, Waters H, Phillips T, Rehwinkel J. Exploring the economics of motorcycle helmet laws—implications for low and middle-income countries. *Asia Pac J Public Health*. 2007;**19**(2):16-22.
- Heydari ST, Maharlouei N, Foroutan A, Sarikhani Y, Ghaffar-pasand F, Hedjazi A, et al. Fatal motorcycle accidents in Fars Province, Iran: a community-based survey. *Chin J Traumatol*. 2012;**15**(4):222-7.
- Janmohammadi N, Pourhossein M, Hashemi SR. Pattern of motorcyclist's mortality in Mazandran province, Northern Iran. *Iran Red Crescent Med J*. 2009;**11**(1):81-4.
- Rutledge R, Stutts J. The association of helmet use with the outcome of motorcycle crash injury when controlling for crash/injury severity. *Accid Anal Prev*. 1993;**25**(3):347-53.
- Paravar M, Hosseinpour M, Salehi S, Mohammadzadeh M, Shojaee A, Akbari H, et al. Pre-hospital trauma care in road traffic accidents in kashan, iran. *Arch Trauma Res*. 2013;**1**(4):166-71.
- Gennarelli TA, Champion HR, Sacco WJ, Copes WS, Alves WM. Mortality of patients with head injury and extracranial injury treated in trauma centers. *J Trauma*. 1989;**29**(9):1193-201.
- Centers for Disease C, Prevention. Helmet use among motorcyclists who died in crashes and economic cost savings associated with state motorcycle helmet laws—United States, 2008-2010. *MMWR Morb Mortal Wkly Rep*. 2012;**61**(23):425-30.
- Pervin A, Passmore J, Sidik M, McKinley T, Nguyen TH, Nguyen PN. Viet Nam's mandatory motorcycle helmet law and its impact on children. *Bull World Health Organ*. 2009;**87**(5):369-73.
- Keng SH. Helmet use and motorcycle fatalities in Taiwan. *Accid Anal Prev*. 2005;**37**(2):349-55.
- Lardelli-Claret P, Jimenez-Moleon JJ, de Dios Luna-del-Castillo J, Garcia-Martin M, Bueno-Cavanillas A, Galvez-Vargas R. Driver dependent factors and the risk of causing a collision for two wheeled motor vehicles. *Inj Prev*. 2005;**11**(4):225-31.
- MacLeod JB, Digiacomio JC, Tinkoff G. An evidence-based review: helmet efficacy to reduce head injury and mortality in motorcycle crashes: EAST practice management guidelines. *J Trauma*. 2010;**69**(5):1101-11.
- Gkritza K. Modeling motorcycle helmet use in Iowa: evidence from six roadside observational surveys. *Accid Anal Prev*. 2009;**41**(3):479-84.
- Zargar M, Khaji A, Karbakhsh M. Pattern of motorcycle-related injuries in Tehran, 1999 to 2000: a study in 6 hospitals. *East Mediterr Health J*. 2006;**12**(1-2):81-7.
- Vafae-Najar A, Esmaeili H, Ibrahimipour H, Dehnavieh R, Nozadi MS. Motorcycle fatal accidents in khorasan razavi province, iran. *Iran J Public Health*. 2010;**39**(2):95-101.
- Amirjamshidi A, Ardalani A, Nainei KH, Sadeghi S, Pahlevani M, Zarei MR. Comparison of standard and nonstandard helmets and variants influencing the choice of helmets: A preliminary report of cross-sectional prospective analysis of 100 cases. *Surg Neurol Int*. 2011;**2**:49.
- Khan I, Khan A, Aziz F, Islam M, Shafiqat S. Factors associated with helmet use among motorcycle users in Karachi, Pakistan. *Acad Emerg Med*. 2008;**15**(4):384-7.
- Brown CV, Hejl K, Bui E, Tips G, Coopwood B. Risk factors for riding and crashing a motorcycle unhelmeted. *J Emerg Med*. 2011;**41**(4):441-6.
- Mehri A, Saeed MM, Ali MM, Haidar N. Determinants of helmet use behaviour among employed motorcycle riders in Yazd, Iran based on theory of planned behaviour. *Injury*. 2011;**42**(9):864-9.
- McCartt AT, Blunar L, Teoh ER, Strouse LM. Overview of motorcycling in the United States: a national telephone survey. *J Safety Res*. 2011;**42**(3):177-84.
- Dee TS. Motorcycle helmets and traffic safety. *J Health Econ*. 2009;**28**(2):398-412.
- Moghisi A, Mohammadi R, Svanstrom L. Motorcyclists' safety in Iran: implication of haddon matrix in safe community setting. *Med J Islam Repub Iran*. 2014;**28**(37):2-9.
- Sosin DM, Sacks JJ, Holmgren P. Head injury—associated deaths from motorcycle crashes. Relationship to helmet-use laws.

- JAMA. 1990;**264**(18):2395-9.
34. *The National Highway Traffic Safety Administration reports.* Washington; 2010; Research Note Motorcycle Helmet Use in 2010-Overall Results.
 35. Zamani-Alavijeh F, Bazargan M, Shafiei A, Bazargan-Hejazi S. The frequency and predictors of helmet use among Iranian motorcyclists: A quantitative and qualitative study. *Accid Anal Prev.* 2011;**43**(4):1562-9.
 36. Ackaah W, Afukaar FK. Prevalence of helmet use among motorcycle users in Tamale Metropolis, Ghana: an observational study. *Traffic Inj Prev.* 2010;**11**(5):522-5.
 37. Conrad P, Bradshaw YS, Lamsudin R, Kasniyah N, Costello C. Helmets, injuries and cultural definitions: motorcycle injury in urban Indonesia. *Accid Anal Prev.* 1996;**28**(2):193-200.