

Effect of Music on Postoperative Pain in Patients Under Open Heart Surgery

Neda Mirbagher Ajorpaz^{1,*}; Abouzar Mohammadi¹; Hamed Najaran¹; Shala Khazaei¹

¹Department of Nursing, Kashan University of Medical Sciences, Isfahan, Kashan, IR Iran

*Corresponding author: Neda Mirbagher Ajorpaz, Department of Nursing, Kashan University of Medical Sciences, Isfahan, Kashan, IR Iran. Tel: +98-3155540021, Fax: +98-3155546633, E-mail: mirbagher_n@kaums.ac.ir

Received: May 14, 2014; Revised: July 9, 2014; Accepted: July 13, 2014

Background: Music, as a non-pharmacological and inexpensive nursing intervention, can be used easily as complementary technique and be effective in reducing pain with other methods. While some studies have demonstrated pain to decrease after music, others found music to be ineffective on pain.

Objectives: The aim of this study was to investigate the effect of music on postoperative pain in patients under open heart surgery.

Patients and Methods: A quasi-experimental study was performed on 60 patients under open heart surgery referred to ICU of Shahid Beheshti hospital in Kashan city. Patients were randomly divided into two groups including experimental and control groups. Patients in music group listened to nonverbal music for 30 minutes after surgery by headphones. The control group did not receive any intervention. Before and after intervention, pain intensity was measured and recorded by visual analog scale in two groups. Data was analyzed using Chi-Square and t-tests.

Results: Before intervention, the mean of pain intensity was 6.3 ± 0.2 and 6.1 ± 0.2 for experimental and control groups, respectively; and the difference was not significant ($P = 0.2$). After intervention, the mean of pain intensity was 3.1 ± 0.1 and 5.8 ± 0.38 for experimental and control groups, respectively; and the difference was significant ($P = 0.04$).

Conclusions: Hearing the music can reduce postoperative pain. It is suggested that relaxant music be used as a complementary method in patients in order to reduce prospective pain.

Keywords: Music; Pain; Open Heart Surgery

1. Background

Cardiovascular diseases have the highest death rates and will remain the primary cause of death in the world until 2020. Nearly 52% of deaths in the United States and 48% in Europe are related to these diseases (1). A large number of patients with coronary artery diseases, which do not respond to medical therapies have to undergo coronary artery bypass graft surgery (CABG) (2). This common procedure has been done for about 35 years (2). The prevalence of CABG is 26.79% in North America, 0.72% in Asia, 17.94% in Western Europe and 18.14% in the rest of the world. In Iran, 60% of all open-heart surgeries are CABGs (3).

Pain is defined as an unpleasant sensory and emotional experience probably associated with actual or potential tissue damage. Although pain is a predictable part of the postoperative experience, inadequate management of pain is common and can have profound effects. Unrelieved postoperative pain may result in clinical and psychological changes that increase morbidity, mortality, and costs and decrease the quality of life (4). Moreover, negative clinical outcomes resulting from ineffective postoperative pain management include deep vein thrombosis, pulmonary embolism, coronary ischemia, myocardial infarction, pneumonia, poor wound healing, insomnia, and demoralization (5). Moderate to severe

postoperative pain is experienced by more than 80% of patients having surgery (2). Pain has also been reported as one of the primary sources of concern for cardiac surgery patients, and postoperative pain management is important owing to the increasing number of patients undergoing open heart surgery (2, 5, 6). Pain can be caused by incisions, intraoperative tissue retraction and dissection, multiple intravascular cannulations, chest tubes left after surgery, and multiple invasive procedures that patients undergo as part of their therapeutic processes (7). In one study, patients reported chest incision pain as a problem after CABG (8). Because of pain, patients cannot take deep breaths, cough, or start moving around as soon as they should, and this, delays their recovery (9). Associated with these complications are economic and medical implications, such as extended lengths of hospital stay and patient dissatisfaction with medical care (10). It is proposed that effective management of postoperative pain in patients who have undergone CABG can be an important factor in overall recovery. Music is an inexpensive, non-pharmacological, noninvasive nursing interaction that has no side effects and can be effective along with other methods (2, 6). Although music effect mechanism in reducing the pain is not clear completely, increasing of Mio

receptors on the cells surface and increasing of endorphins are suspected as the probable mechanisms in this context (11). In examining the effects of music on patients have undergone CABG, a number of studies have demonstrated decreases in postoperative pain (12, 13), whereas Nilsson *et al.* found no difference in pain (14).

There are limited numbers of published studies on the effect of music in pain reduction after open-heart surgery (12). In addition, music has a cultural implication (15) and insufficient studies have been conducted in countries such as Iran. Meanwhile, the success of music therapy may be greatly enhanced by determining patient's preference, familiarity and cultural context (16). While some studies have demonstrated pain to decrease after music (6, 9, 17-20), others found music to be ineffective on pain (20, 21). Many previous investigations have been limited in a way. For instance, some employed a small sample size (22), some evaluated different types of surgeries and anesthesia in one study (23), and some used a type of music not selected by patients (24, 25). Moreover, while a systematic review recommended a minimum duration of 30 minutes for music therapy to be effective in clinical practice (14), a number of studies played music for less than 30 minutes (12, 26). On the other hand, the constant presence of the researcher during the intervention (25) might have affected patient's response. Therefore, considering the above mentioned facts and according to the cultural, social and economic differences in Iran, we tried to perform this study as well as we could.

2. Objectives

The aim of this study was to investigate the effect of music on postoperative pain in patients under open heart surgery.

3. Patients and Methods

A quasi-experimental study was performed on 60 patients under open heart surgery referred to ICU of Shahid Beheshti hospital in Kashan city during 2012 and 2013. Inclusion criteria were age between 18 to 60 years, complete conciseness, reading and writing ability, moderate and severe pain based on Visual Analog Scale (VAS), not having addiction to sedatives and alcohol, hemodynamic stability, first time open heart surgery, not having mental disorders, not having chronic pain, not having hearing disorders and were on their first postoperative day after CABG or valve replacement. Patients were excluded if they were connected to a ventilator at the time of intervention, used other alternative methods for pain reduction such as massage therapy during the intervention (except of routine pain medication in the ICU).

In the present study, data were collected from the patients undergoing CABG or valve replacement surgery. For those patients undergoing valve replacement surgery, the incision region was in the sternum. However, in patients undergoing CABG surgery, the incision regions

were sternum and leg, where the saphenous vein was removed. Thus, two different pain regions were determined by taking the region of incisional pain into consideration (1 = chest incisional pain region; 2 = chest and leg incisional pain regions).

3.1. Sampling

The sample size in each group was determined based on the following assumptions: power = 0.80, $\alpha = 0.05$, the minimum expected difference in standard deviation = 3.6, and the minimum expected difference in means to be 2.40 (27). According to the formula, the sample size in each group was 30. At first, 82 patients with open heart surgery were assessed for eligibility. Among those, 13 did not have the inclusion criteria, and 9 persons declined to participate. Therefore, 60 patients were randomly assigned into two groups including experimental and control groups (Figure 1). The data collection instrument consisted of two parts; first part was the demographic characteristics including age, gender, marital status, surgery type, education level, and the second part included VAS. According to these criteria, score zero shows that there is no pain and score 10 shows the most level of experienced pain by the person. This scale has extensively been used in the studies related to pain and its validity and reliability have been approved (28, 29).

3.2. Procedures

After explaining the research objectives and obtaining written informed consent from the participants. The data were collected between 3.00 p.m. and 4.00 p.m. when the traffic in ICU was not intense and the patients were not receiving invasive or noninvasive procedures. All of the patients received standard cares under supervision of a cardiovascular specialist. All the patients received 50 mg Pethidine and beta- receptor blocker (metoprolol succinate) at 8 am by the ICU nurse. The patients also received 3-5 ml/min oxygen delivered by nasal cannula to prevent complications and hypoxia after surgery. The second researcher helped the participants lie down in the bed and turn over to a relaxing position on their back, supine position, with the bed head at 20 to 30 degrees. In addition, factors affecting pain intensity, such as incision method, type and extent of incision, having chest tube, and type of analgesic drugs used, were the same for all patients. Demographic and pain intensity scale (VAS) were completed in the first 24 hours after the surgery in the ICU.

The patients in the music group were listening to sedative music by an MP3 player with special headphones for 30 minutes. Sedative music was operationalized as music without lyrics and with a sustained melodic quality, with a rate of 60-80 beats per minute and a general absence of strong rhythms or percussion. Changing the volume was in the control of the patient. Sedative music was selected by a music expert considering the cultural

conditions of the society. During the intervention, for all participants, the environment was enhanced to reduce stimuli and facilitate rest by closing the door and posting a sign to prevent being disturbed by visitors and health care personnel.

During this period, the second researcher stayed in ICU. After the intervention, Pain intensity was again evaluated immediately after the music was completely played in two groups. However, in the control group, headphones were used without playing any music.

3.3. Ethical Considerations

The study was approved by the research deputy and the research ethics committee of Kashan University of Medical Sciences. All participants signed a written informed consent before participation in the study. The respondents were anonymous and all the information was kept confidential in this study. The participants were free to leave the study at any time. The ICU nurses were informed about the results at the end of the study. The researchers observed all ethical issues in accordance with the Helsinki declaration.

3.4. Data Analysis

The data was collected in the specified forms and was

analyzed using SPSS software Version 16 (SPSS, Chicago, IL, USA). The normality the data was analyzed using Kolmogorov-Smirnov test ($P \geq 0.05$). Also, Mann-Whitney-U test was applied where the distribution of the data was not normal. Mean score and standard deviation were calculated. Chi-square and t- tests were used to compare nominal variables between the two groups. T-test was also used to compare the statistical difference between the mean differences of pain in the two groups. $AP < 0.05$ was considered to be statistically significant in all testes.

4. Results

A total number of 60 patients undergoing open heart surgery were enrolled in this study (30 in each group). A proportion of 43.3% of patients were in the age range of 50- 65 years old. Also, 56.6% of the patients were male and all married. No significant difference was observed between the mean of pain intensity in the experimental and control groups before intervention ($P = 0.2$). However, a significant difference was found between the mean of pain intensity in the experimental and control group at 30 minutes after music therapy ($P = 0.04$). Also, a significant difference was observed between the mean of pain intensity in the experimental groups before and after intervention ($p = 0.01$) (Table 1 and 2).




Figure 1. The Sampling Framework of the Study

Table 1. Pain Intensity in the Experimental Groups

Variable	Experimental Group	Control Group	Chi-Square and p-Value
Gender			Chi-square = 3.51 P = 0.1
Female	18 (60%)	13 (43.3%)	
Male	12 (40%)	17 (56.6%)	
Marital status			Chi-square = 2.86 P = 0.09
Married	25 (83.3%)	23 (76.6%)	
Single	5 (16.6%)	7 (23.3%)	
Age Group			Chi-square = 3.11 P = 0.28
20-35	6 (20%)	8 (26.6%)	
35-50	11 (36.6%)	8 (26.6%)	
50-65	13 (43.3%)	14 (46.6%)	
Education level			Chi-square = 3.9 P = 0.14
Secondary education	11 (36.6%)	9 (30%)	
Diploma	9 (30%)	14 (46.6%)	
Academic	10 (33.3%)	7 (23.3%)	
Type of Surgery			Chi-square = 2.91 P = 0.15
Valve replacement	10 (33.3%)	8 (26.6%)	
CABG	20 (66.6%)	22 (73.3%)	

Table 2. Comparing The Mean of Pain Intensity in Two Groups Before and After Intervention

Groups	Before intervention	After Intervention	Independent t-Test
Experimental Group	6.3 ± 0.2	3.1 ± 0.1	T= 4.3/P= 0.01
Control Group	6.1 ± 0.2	5.8 ± 0.3	T= 2.25/P= 0.21
Paired t-Test	T = 2.87/P = 0.2	T = 3.49/P = 0.04	-

5. Discussions

Most of the using methods for reducing postoperative complications such as pain and the amount of analgesic drugs consumption are based on drugs intervention. Our effort in this study was to assess the effect of music as a non-pharmacological and inexpensive intervention in reducing the postoperative pain. Based on the results of this study, music significantly decreased postoperative pain in patients after open heart surgery. Likely to our study, both Sendelbach *et al.* (12) and Hatem *et al.* (30) showed that music reduced patients' pain after cardiac surgery. The finding that listening to music was effective in relieving postoperative pain is consistent with other studies that examined the effects of music on pain for patients undergoing cardiac surgery (18, 31). Voss *et al.* investigated the effect of music on pain during chair rest after open heart surgery and reported that the pain level of the sedative music group was lower than in the scheduled rest or treatment as usual groups (13). This finding is inconsistent with present study. The inconsistencies between different studies may be attributed to differences in type of music, music time, type of disease, patients' culture and music playback time. Aragon *et al.* investigated the effect of harp

music in vascular and thoracic surgical patients by using VAS to measure patients' pain. The VAS was completed just before harp playing 20 minutes after harp playing was started, and 10 minutes after completion. Results indicated that listening to live harp music had a positive effect on patient perception of pain (17). This finding is consistent with findings of our study. However, Aragon *et al.* study used harp music, and the present study used sedative music. It seems that every type of music is effective in reducing pain. Zimmerman *et al.* demonstrated that music intervention decreased postoperative pain in CABG patients (21). Similarly, Nilsson *et al.* stated patients exposed to soft relaxing music intra-operatively, have significantly lower pain scores compared with the control group patients on the first day after the surgery (32). In contrast, Allred *et al.* evaluated patients undergoing knee arthroplasty and did not find any significant pain reduction in the music group as compared with the control group (resting on bed) (2), while it is inconsistent with present study. Allred *et al.* used music for 20 minutes after surgery; whereas, we used music for 30 minutes after surgery in ICU.

The current study is consistent with the findings from

previous studies that sedative music was more effective than scheduled rest or usual treatment in reducing pain in ICU. Studies that have not found significant effects on pain (33) or have found mixed results had small sample sizes with inadequate power to detect significant differences. The lack of significant findings in some studies may have been related to the suddenness of pain experienced during chest tube removal or upon awakening from surgery (33), when patients may have had difficulty focusing on the music intervention. Experts have suggested that sedative music is more effective if the patient is able to concentrate on the intervention (33). It seems that sedative music used in this study, would be effective in decreasing the patients' pain after surgery.

Music ineffectiveness on pain reduction in these studies could be attributed to the type of the music listened by the patients, ie, one type of Spanish guitar music was played for all participants without considering the local and national culture of the patients. However, it has been suggested that preferred music, as opposed to prescribed music, is a critical factor in the effectiveness of music therapy (19). The results of Woldendorp *et al.* showed a significant correlation between the degree of relaxation and liking the music (34). Since the emotional responses differ from culture to culture, a type of music related to the cultural features of the subjects should be selected (3).

The success of music interventions may be greatly enhanced by familiarity and cultural contexts of the patients. The large effects for those who used sedative music may be attributable to having something more pleasant to concentrate on or something to distract their minds from the pain and help them relax their bodies (35). Some researchers believe that music as the senses deviation acts like a mask on the annoying sounds of the ICU and leads to reduction of stress and anxiety and reduction of the pain intensity in the following (36). Music also occupies the patient's mind with something familiar and soothing, which allows the patient to escape into his or her own world (19).

5.1. Study Limitations

This study had several limitations. The study was conducted in only one ICU, and the study sample reflects only one area of Iran. Furthermore, data were collected by one researcher. The available selection of music included only nonverbal musical pieces, which limited participants' choices. Our study only evaluated pain severity, and we did not include vital signs evaluation. Future studies are recommended to include larger samples from different regions of Iran and should also include a larger selection of music pieces.

The findings of the study indicated the significant effect of music on reducing the postoperative pain. Thus, music can be used as a complementary and noninvasive method in relieving postoperative pain. Music, as a nursing intervention, is a noninvasive and safe therapy for

health promotion of patients. However, the music listened by patients should be of a sedative quality, and culturally appropriate selections should be offered. Nurses can use music as an intervention for patients who have undergone open heart surgery to promote nursing autonomy and the idea that nurses are able to affect patients' environment. It is advisable that music should become a part of nursing care offered to patients experiencing pain, because music is a low-cost therapy that has no side effects.

Acknowledgements

The authors would like to thank all patients and physicians who participated in this study, especially Dr. Porabasi (cardiovascular specialist) and all nurses in ICU.

Authors' Contributions

Neda Mirbagher Ajorpaz designed the research and performed data analysis and prepared the first draft of the manuscript and supervised the study. Abouzar Mohammadi contributed in preparing. Hamed Najjaran and Shahla Khazaei collected the data.

Financial Disclosure

There were no conflicts of interest among the authors of the study.

Funding/support

This paper is the result of a research project granted by Kashi University of Medical Sciences and Health Care Services (grant number: 9246). The study was also registered at Iranian registry for clinical trials (IRCT) as IRCT201406028348N7.

References

1. Zand Parsa AF, Ziai H, Haghghi L. The impact of cardiovascular risk factors on the site and extent of coronary artery disease. *Cardiovasc J Afr*. 2012;**23**(4):197-9.
2. Allred KD, Byers JF, Sole ML. The effect of music on postoperative pain and anxiety. *Pain Manag Nurs*. 2010;**11**(1):15-25.
3. Molavi Vardanjani M, Masoudi Alavi N, Sadat Razavi N, Aghajani M, Azizi-Fini E, Vaghefi SM. A Randomized-Controlled Trial Examining the Effects of Reflexology on Anxiety of Patients Undergoing Coronary Angiography. *nursingmidwife*. 2013;**2**(1):3-9.
4. Choiniere M, Watt-Watson J, Victor JC, Baskett RJ, Bussieres JS, Carrier M, et al. Prevalence of and risk factors for persistent postoperative nonanginal pain after cardiac surgery: a 2-year prospective multicentre study. *CMAJ*. 2014;**186**(7):E213-23.
5. Englbrecht JS, Pogatzki-Zahn EM. [Perioperative pain management for abdominal and thoracic surgery]. *Schmerz*. 2014;**28**(3):265-81.
6. Comeaux T. The effect of complementary music therapy on the patient's postoperative state anxiety, pain control, and environmental noise satisfaction. *Medsurg Nurs*. 2013;**22**(5):313-8.
7. Short A, Gibb H, Fildes J, Holmes C. Exploring the role of music therapy in cardiac rehabilitation after cardiothoracic surgery: a qualitative study using the Bonny method of guided imagery and music. *J Cardiovasc Nurs*. 2013;**28**(6):E74-81.
8. Hong SJ, Hong MK, Ko YG, Hong GR, Shim JK, Kwak YL, et al. Migration of calcium and atheromatous plaque in computed tomog-

- raphy: an important mechanism of coronary artery occlusion after transcatheter aortic valve replacement. *J Am Coll Cardiol*. 2014;**63**(12).
9. Bauer BA, Cutshall SA, Anderson PG, Prinsen SK, Wentworth LJ, Olney TJ, et al. Effect of the combination of music and nature sounds on pain and anxiety in cardiac surgical patients: a randomized study. *Altern Ther Health Med*. 2011;**17**(4):16-23.
 10. Trappe HJ. The effects of music on the cardiovascular system and cardiovascular health. *Heart*. 2010;**96**(23):1868-71.
 11. Abrams B. Evidence-based music therapy practice: an integral understanding. *J Music Ther*. 2010;**47**(4):351-79.
 12. Sendelbach SE, Halm MA, Doran KA, Miller EH, Gaillard P. Effects of music therapy on physiological and psychological outcomes for patients undergoing cardiac surgery. *J Cardiovasc Nurs*. 2006;**21**(3):194-200.
 13. Voss JA, Good M, Yates B, Baun MM, Thompson A, Hertzog M. Sedative music reduces anxiety and pain during chair rest after open-heart surgery. *Pain*. 2004;**112**(1-2):197-203.
 14. Nilsson U. The anxiety- and pain-reducing effects of music interventions: a systematic review. *AORN J*. 2008;**87**(4):780-807.
 15. Good M, Ahn S. Korean and American music reduces pain in Korean women after gynecologic surgery. *Pain Manag Nurs*. 2008;**9**(3):96-103.
 16. Stouffer JW, Shirk BJ, Polomano RC. Practice guidelines for music interventions with hospitalized pediatric patients. *J Pediatr Nurs*. 2007;**22**(6):448-56.
 17. Aragon D, Farris C, Byers JF. The effects of harp music in vascular and thoracic surgical patients. *Altern Ther Health Med*. 2002;**8**(5):52-4.
 18. Chan MF. Effects of music on patients undergoing a C-clamp procedure after percutaneous coronary interventions: a randomized controlled trial. *Heart Lung*. 2007;**36**(6):431-9.
 19. Chlan LL, Engeland WC, Savik K. Does music influence stress in mechanically ventilated patients? *Intensive Crit Care Nurs*. 2013;**29**(3):121-7.
 20. Ozer N, Karaman Ozlu Z, Arslan S, Gunes N. Effect of music on postoperative pain and physiologic parameters of patients after open heart surgery. *Pain Manag Nurs*. 2013;**14**(1):20-8.
 21. Jafari H, Emami Zeydi A, Khani S, Esmaili R, Soleimani A. The effects of listening to preferred music on pain intensity after open heart surgery. *Iran J Nurs Midwifery Res*. 2012;**17**(1):1-6.
 22. Ottaviani S, Bernard JL, Bardin T, Richette P. Effect of music on anxiety and pain during joint lavage for knee osteoarthritis. *Clin Rheumatol*. 2012;**31**(3):531-4.
 23. McRee LD, Noble S, Pasvogel A. Using massage and music therapy to improve postoperative outcomes. *AORN J*. 2003;**78**(3):433-42.
 24. Ikonomidou E, Rehnstrom A, Naesh O. Effect of music on vital signs and postoperative pain. *AORN J*. 2004;**80**(2):269-74.
 25. Nilsson U. The effect of music intervention in stress response to cardiac surgery in a randomized clinical trial. *Heart Lung*. 2009;**38**(3):201-7.
 26. Buffum MD, Sasso C, Sands LP, Lanier E, Yellen M, Hayes A. A music intervention to reduce anxiety before vascular angiography procedures. *J Vasc Nurs*. 2006;**24**(3):68-73.
 27. Aghajani M, Mirbagher Ajorpaz N, Kafaei Atrian M, Raofi Z, Abedi F, Naeimi Vartoni S, et al. Effect of Self - Care Education on Quality of Life in Patients With Primary Hypertension: Comparing Lecture and Educational Package. *nursingmidwife*. 2013;**2**(2):71-6.
 28. Etri M, Adib-Hajbaghery M. Effects of Acupressure on Pain and Vital Signs of Patients After Small Abdominal Surgeries: A Clinical Trial. *nursingmidwife*. 2012;**1**(2):67-71.
 29. Xu X, Zhang L, Jiang Y, Huang Y, Huang S, Yang S. [Clinical research of music in relieving orthodontic pain]. *Hua Xi Kou Qiang Yi Xue Za Zhi*. 2013;**31**(4):365-8.
 30. Hatem TP, Lira PI, Mattos SS. The therapeutic effects of music in children following cardiac surgery. *J Pediatr (Rio J)*. 2006;**82**(3):186-92.
 31. Wang Y, Dong Y, Li Y. Perioperative psychological and music interventions in elderly patients undergoing spinal anesthesia: effect on anxiety, heart rate variability, and postoperative pain. *Yonsei Med J*. 2014;**55**(4):1101-5.
 32. Nilsson U, Rawal N, Unestahl LE, Zetterberg C, Unosson M. Improved recovery after music and therapeutic suggestions during general anaesthesia: a double-blind randomised controlled trial. *Acta Anaesthesiol Scand*. 2001;**45**(7):812-7.
 33. Taylor LK, Kuttler KL, Parks TA, Milton D. The effect of music in the postanesthesia care unit on pain levels in women who have had abdominal hysterectomies. *J Perianesth Nurs*. 1998;**13**(2):88-94.
 34. Woldendorp KH, van de Werk P, Boonstra AM, Stewart RE, Otten E. Relation between muscle activation pattern and pain: an explorative study in a bassists population. *Arch Phys Med Rehabil*. 2013;**94**(6):1095-106.
 35. Chen CK, Pei YC, Chen NH, Huang LT, Chou SW, Wu KP, et al. Sedative music facilitates deep sleep in young adults. *J Altern Complement Med*. 2014;**20**(4):312-7.
 36. Gelinas C, Arbour C, Michaud C, Robar L, Cote J. Patients and ICU nurses' perspectives of non-pharmacological interventions for pain management. *Nurs Crit Care*. 2013;**18**(6):307-18.