

ORIGINAL ARTICLE

Iran J Allergy Asthma Immunol
February 2017; 16(1):14-20.

Effects of Submaximal Aerobic Exercise on Regulatory T Cell Markers of Male Patients Suffering from Ischemic Heart Disease

Fariba Raygan¹, Mansour Sayyah², Seyed Mohammad Reza Janesar Qamsari³,
Hassan Nikouejad⁴, and Mojtaba Sehat⁵

¹ Department of Cardiology, Kashan University of Medical Sciences, Kashan, Iran

² Trauma Research Center, Kashan University of Medical Sciences, Kashan, Iran

³ Student Research Center, Kashan University of Medical Sciences, Kashan, Iran

⁴ Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

⁵ Department of Community Medicine, Kashan University of Medical Sciences, Kashan, Iran

Received: 21 May 2016; Received in revised form: 7 January 2017; Accepted: 17 January 2017

ABSTRACT

There are confirmed beneficiary effects of exercise on atherosclerotic inflammation of ischemia-associated heart diseases. The purpose of this study was to evaluate the effect of aerobic exercise on T-regulatory cell markers of IL-35 as well as FoxP3 and T-helper2 marker of IL-33 in patients with ischemic heart disease (IHD).

This research was performed on 44 asymptomatic male patients with ischemic heart disease. The participants were randomly assigned into two groups of submaximal aerobic exercise and control group. Blood samples were collected before and after the termination of the exercise protocol. Serum levels of IL-35 and IL-33 as well as the amount of FoxP3 gene expression in peripheral blood mononuclear cells were measured by Elisa and Real time PCR, respectively.

Serum levels of IL-35 ($p=0.001$) as well as the amount of FoxP3 gene expression increased significantly ($p=0.012$) in exercise group even after controlling the likely confounding effects of age, length of ischemia, duration of the disease, and the amount of such factors before exercise ($p\leq 0.042$).

It seems that exercise may yield a better control of atherosclerotic inflammation in patients with ischemic heart disease through the induction of regulatory T cells.

Keywords: Interleukin-35; Interleukin-33; Regulatory T cells; Sub maximal aerobic exercise

INTRODUCTION

Physical inactivity is a well-documented risk factor for

coronary heart disease and there is evidence showing regular physical activity reduces the risk of hypertension, type 2 diabetes and helps maintain

Corresponding Author: Hassan Nikouejad, MD, PhD;
Nephrology and Urology Research Center, Baqiyatallah University of
Medical Sciences, Baqiyatallah Hospital, P.O.Box: 19395-5487,

Tehran, Iran. Tel: (+98 913) 1615 530, Fax: (+98 21) 8126 2073,
E-mail: hnkuinejad@yahoo.com

Effects of Submaximal Aerobic Exercise on Ischemic Heart Disease

optimal bone mineral density;¹ however, in addition to inactivity, there are other pathophysiologic bases of the disease associated with inflammation. Over the last decades, an abundance of evidence has emerged that inflammation plays a role in all phases of the atherosclerotic disease process, from lesion initiation to progression and, finally, to plaque rupture and the consecutive complications of cardiovascular disease.^{2,3} In fact, the inflammatory reactions are a group of the interfering processes in the development of atherosclerosis and its involved complications.⁴ There are numerous reports that show physical activity has significant beneficiary effects on cardiovascular events⁵⁻⁸ through reducing the risk factors⁹⁻¹¹ as well as processes of atherosclerosis progression.^{12,13} The majority of such studies have examined the changes of inflammatory/anti-inflammatory markers such as CRP, VCAM-1, ICAM-1, fibrinogen, IL-6, TNF- α , IL-10, or TGF- β .¹⁴⁻¹⁸

Regulatory T cells (Tregs), a subset of T cells, suppress different immune cells such as T and B cells, natural killer cells, natural killer T cells, mast cells, and dendritic cells. The main mechanism of such activity is to produce anti-inflammatory cytokines which control various types of inflammation through the mediatory effect of Fork head box P (Foxp3). Foxp3 is the main transcription factor and phenotypic/functional marker of Tregs.^{19,20} Results obtained from animal models of atherosclerosis have shown the protective and fixing role of such cells in atheroma plaque.²¹ It seems that lack of such protection, which is induced by their anti-inflammatory cytokines of IL-10 and TGF- β , may lead to acute coronary events as well as advanced atherosclerosis.^{22,23} IL-35, an IL-12 family cytokine secreted by Tregs may have such anti-inflammatory effects.²⁴

The response of Tregs to various types of endurance exercise has been studied in endurance athletes.²⁵ Perry et al. reported that there was a decrease in both the percentage of Treg lymphocytes and the intensity of FoxP3 expression in CD4+CD25+ Tregs following the participation in endurance exercise in athletes.²⁶ More specifically, the response of FoxP3 in ischemic heart patients to moderate (submaximal) aerobic endurance exercise has not been examined yet.

IL-33, an IL-1 family cytokine secreted by Th2 cells and inducing Th2-associated diseases such as asthma, may have some protective roles in various cardiovascular diseases including atherosclerosis.^{26,27}

It has been demonstrated that mice deprived of IL-33 receptor, namely IL-33Ra, are afflicted to myocardium inflammation, ventricle dilation and cardiac fibrosis. In addition, the increase in extracellular IL-33Ra level is a predictor of clinical myocardial infarction.²⁸

This research was designed, for the first time, to examine the effect of submaximal aerobic instead of intensive aerobic exercise on regulatory T-cells, focusing on two secretory-relevant markers of IL-35 and IL-33 as well as the cell-relevant markers of FoxP3 gene expression in male patients with ischemic heart diseases. Such knowledge may help us to prevent the cardiovascular disorders or to accelerate patients' recovery by promoting anti-inflammatory mechanisms.

MATERIALS AND METHODS

Patients

This semi-experimental research included 44 volunteered asymptomatic male patients with ischemic heart disease diagnosed in Shahid Beheshti Hospital of Kashan University of Medical Sciences. All the patients underwent angiography and at least one of the arteries had 75% or more stenosis. They were randomly assigned into two groups of 21 who participated in prescribed submaximal aerobic exercise and 23 patients served as the control group participating in routine regular physical activities. Men were chosen considering the fact that they are more likely to suffer from such ischemic heart diseases at younger age than women; moreover, men at this age are more likely to be able to participate and complete an exercise program. Both groups received their identical regular treatment. Blood samples were collected before and after the termination of the exercise protocol. The exclusion criteria were any clinical or ECG changes during the exercise, any cardiovascular event within the last 3 months, being candidate for revascularization, ventricular tachycardia, uncontrolled hypertension, malignancy, diabetes, autoimmune diseases, allergy, chronic or acute infectious diseases, kidney or liver failure, use of immune-modulating drugs and erythropoietin, peripheral vascular diseases limiting physical activities, valvular heart diseases, and obstructive pulmonary disease. Both control and exercise groups received their identical conventional medication. The ethical committee of the Kashan University of Medical Sciences approved the

project and informed consent was signed by the participants.

Exercise Protocol

The submaximal aerobic exercise was determined based on formula described in most exercise related texts²⁹ and included three sessions of exercise continued for 12 weeks. An exercise science specialist designed and supervised the exercise protocol. The participants in exercise group performed 5 minutes of mild warm-up exercise prior to performing 40 minutes of submaximal aerobic exercise followed by 5 minutes of cool-down stretching exercises. The target heart rate was monitored by Pollar watch model M300set (China). The health condition of every participant was assessed at the beginning of every exercise session. Weekly general health condition of the participants was evaluated by a cardiologist.

Elisa and Real-time Experiments

24 before and after the termination of the exercise program, 5 cc of venous blood sample was collected from the left arm of the all participants and kept frozen in EDTA-containing tubes until the assays. Serum levels of IL-33 and IL-35 were measured by ELISA sandwich method (eBioscience kit, USA) According to manufacturer's instructions. Total RNA was extracted from peripheral blood mononuclear cells (PBMC) using High Pure RNA Isolation Kit (Cat No: 11828665001, Roche Applied Science, Germany). cDNA synthesized from the extracted RNA using Transcriptor First Strand cDNA Synthesis Kit (Cat No: 04897030001, Roche Applied Science, Germany). The amount of FoxP3 gene expression was measured using Taqman primer probe Comparative CT method) ABI 7300 Real-Time PCR system, USA). β -actin housekeeping gene was used as endogenous control. FoxP3 gene expression in exercise group relative to control group was determined as $2^{\Delta\Delta-CT}$.

Statistical Analysis

Data were analyzed using descriptive statistics including mean and standard deviations. The measurements were compared before and after the exercise using paired and in both groups by independent T-tests. Analysis of covariance

(ANCOVA) was employed to control the confounding variables and baseline differences. The entire significant test was performed by alpha level set to 0.05. SPSS 16.0 (Chicago, USA) and STATA software 11, (Stata Corp., College Station, TX, USA), were used to analyze the data.

RESULTS

Basic and clinical characteristics of the patients have been included in Table 1.

There was no significant difference between the two groups regarding to serum levels of IL-35 and IL-33 as well as the amount of FoxP3 gene expression before exercise.

The changes of both IL-35 serum levels and FoxP3 gene expression in exercise group were significantly higher after the exercise in comparison with those of before exercise ($p \leq 0.012$) (Table 2). The amount of FoxP3 gene expression increased significantly in exercise group compared to control group in post-test condition ($p = 0.012$) (Table 2). There were differences in IL-35 and IL-33 serum levels as well as the amount of FoxP3 gene expression between the groups in the primary condition of the protocol. Therefore, we compared the groups at the post-test state using related factors as the covariates in pre-test state. There was a significant difference in the serum level of IL-35 and the amount of FoxP3 gene expression between both groups at the end of the protocol ($p \leq 0.019$) (Table 3).

Adjusting the effects of total duration of exercise activity, duration of ischemia history, and age of the participants, we performed further multivariate analysis of covariance on IL-35 and IL-33 serum levels and the amount of FoxP3 gene expression to compare the exercise group versus the control group after the termination of the exercise program. The results of analysis indicated that none of those covariates contributed significantly to such difference. This indicates that after the exclusion of the covariate effects, there was still a significant difference of IL-35 and FoxP3 expression between the groups ($p \leq 0.042$). The level of increase after adjustment for those variables remained significant (Table 4).

Effects of Submaximal Aerobic Exercise on Ischemic Heart Disease

Table 1. Basic and clinical characteristics of the ischemic heart disease patients in submaximal exercise and control group

Variants	controls (n=23) (Mean±SD)	exercise (n=21) (Mean±SD)	p value
Age (years)	64.78±12.21	58.5±7.9	0.035
Duration of disease (months)	11.43±4.51	7.71±2.57	0.002
Duration of exercise (hours)	9.61±27.31	27.29±22.84	0.026
Pre-test IL-33 (pg/mL)	75.04±29	89.69±43.75	0.194
Pre-test IL-35(pg/mL)	51.33±22.89	41.53±23.87	0.172
Pre-test FoxP3	12.52±2.85	11.81±2.68	0.4

Table 2. Comparing the levels of IL-35, IL-33, and FoxP3 expression in the ischemic heart disease patients in submaximal exercise and control groups in pre- and post-test condition

Marker	Group	Before	After	Difference	p value
IL-35(pg/mL)	control	51.33 (4.77)	50.42 (4.45)	-0.9 (2.4)	0.714
	exercise	41.53 (5.21)	57.79 (6.86)	16.3 (4.3)	0.001
	Mean (SE)	-9.79	7.36	-	-
	p value	0.172	0.365	-	-
IL-33(pg/mL)	control	75.04 (6.05)	76.84 (7.84)	1.80 (7.33)	0.808
	exercise	89.69 (9.55)	89.95 (7.90)	0.26 (14.70)	0.986
	Mean (SE)	14.65 (10.11)	13.10 (12.37)	-	-
	p value	0.194	0.269	-	-
FoxP3 (pg/mL)	control	12.52 (0.6)	11.87 (0.46)	-0.65 (0.54)	0.243
	exercise	11.81 (0.58)	9.33 (0.87)	-2.48 (0.90)	0.012
	Mean (SE)	0.71	-2.5	-	-
	p value	0.4	0.012	-	-

Table 3. Analysis of covariance (ANCOVA) for the serum levels of IL-35, IL-33, and the amount of FoxP3 gene expression in the ischemic heart disease patients in submaximal exercise and control groups using related covariates

Model		Unstandardized coefficients		Standardized coefficients	t	p value
		B	Standard error	β		
IL-35 (pg/mL)	constant	3.405	6.389		0.353	0.957
	IL-35 before	0.916	0.106	0.814	8.640	0.00
	group	16.335	4.995	0.11	3.279	0.002
IL-33 (pg/mL)	constant	70.633	15.629		4.519	0.00
	IL-33before	0.083	0.174	0.075	0.477	0.636
	group	11.892	12.747	0.146	0.933	0.365
FoxP3 (pg/mL)	constant	6.872	2.196		3.128	0.003
	FoxP3before	0.399	0.168	0.325	2.375	0.22
	group	2.252	0.019	-0.325	-2.450	0.019

Table 4. Multivariate analysis of covariance (ANCOVA) for the serum levels of IL-35 and IL-33 and the amount of FoxP3 gene expression in the ischemic heart disease patients in submaximal exercise and control groups

Model		Unstandardized coefficients		Standardized coefficients	t	p value
		B	Standard error	β		
IL-35(pg/ml)	(constant)	-20.527	19.201		-1.079	0.287
	IL-35before	0.905	0.107	0.803	8.444	0.00
	group	12.176	5.775	0.231	2.108	0.042
	length of ischemia	-1.985	1.114	-0.294	-1.7	0.097
	duration of exercise	0.096	0.097	0.096	0.994	0.327
	age	0.699	0.393	0.0284	1.778	0.083
IL-33(pg/ml)	(constant)	71.360	48.925		1.495	-0.153
	IL-33 before	0.075	0.179	0.067	0.416	0.680
	group	20.302	15.124	0.250	1.342	0.187
	length of ischemia	2.790	2.973	0.280	0.938	0.354
	duration of exercise	-0.066	0.263	-0.043	-0.253	0.802
	age	-0.484	1.057	-0.128	-0.458	0.649
FoxP3(pg/ml)	(constant)	5.366	3.611		1.486	0.146
	FoxP3 before	0.301	0.174	0.245	1.730	0.092
	group	-2.493	1.064	-0.371	-2.343	0.024
	length of ischemia	-0.370	0.218	-0.450	-1.697	0.098
	duration of exercise	-0.026	0.018	-0.202	-1.408	0.167
	age	0.111	0.077	0.355	1.438	0.159

DISCUSSION

This study was conducted to determine the effect of 12 weeks of submaximal aerobic exercise on the serum levels of IL-35 and IL-33 in patients with symptom-free ischemic heart disease. This exercise training program significantly improved the level of regulatory T cell-mediated markers of IL-35 and FoxP3 gene expression.

Our study confirmed the anti-inflammatory role of submaximal aerobic exercise on serum level of IL-35 in patients with ischemic heart disease. These properties of IL-35, at least partly, are mediated through the proliferation of Tregs and inhibition of Th17 cells that play a significant role in recruitment of neutrophils, which in turn, intensify inflammatory processes.²⁹ In the case of inflammation associated with ischemic heart disease, other studies have introduced IL-35 as a significant bio-marker in the incidence of coronary artery disease³⁰ in a way that its decrease may be an indicator of ischemia.³¹ Such protective effect has also been shown in other inflammatory conditions.³²

The findings of this study is in agreement with some other investigations that demonstrated the anti-inflammatory effects of Tregs, where moderate aerobic exercise increased the suppressive function of CD4+CD25+Foxp3+ cells on Th2 inflammatory responses including airway eosinophilia and allergen-induced asthma.^{33,34}

According to our results, sub maximal aerobic does not make any significant changes in serum levels of IL-33. However, one-session of an aerobic exercise made significant increases in the level of serum IL-33 in healthy young men and women.³⁵ These contradictory findings to our study may be attributed to the different types of physical activities and populations employed. Considering blocking effects of Th2 cells on Th1 ones, we emphasize that more physical activity in severity and/or time may also change serum levels of IL-33 and confirm its protective roles in heart ischemia conditions.

The amount of submaximal aerobic exercise applied in this research was sufficient to produce significant changes in regulatory T-cell markers of patients with

Effects of Submaximal Aerobic Exercise on Ischemic Heart Disease

ischemic heart disease. However, the response to moderate level of aerobic physical activity may differ from that of intense and prolonged periods of endurance exercise. A recent report in mice, contradictory to human being, suggested that while intensive exercise increased Treg levels, moderate exercise had no effect on murine Tregs.³⁶ The effects of exercise regarding its intensity have also been the focus of studies in which moderate aerobic exercise reduced the incidence of upper respiratory tract problems, while high intensity and prolonged course of aerobic exercise led to more upper respiratory tract infections in elite athletes.³⁶

It is likely that, other than exercise, some clinical features of our patients affect the production of Tregs. Considering a multivariate regression analysis, we confirmed that exercise was the only explanation for the observed results in our patients.

Our study had some limitations. Firstly, there were no longitudinal follow-up measurements to evaluate the changes of selected markers during the time. This limitation allowed just a cross-sectional analysis with only a limited robustness. Secondly, functional assays which provide further information on the immune-inhibitory effects of the molecules were not performed.

It seems exercise may yield a better control of atherosclerotic inflammation in patients with ischemic heart disease through the induction of regulatory T cells. These findings may be used for both diagnostic approaches of stroke severity and therapeutic strategies.

ACKNOWLEDGEMENTS

This study was supported by Deputy of Research, Kashan University of Medical Sciences (KAUMS), Grant No. 9238. The researchers wish to thank the patients for their sincere and generous contribution to the research protocol.

REFERENCES

1. Paffenbarger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB. The association of changes in physical activity level and other lifestyle characteristics with mortality among men. *N Engl J Med* 1993; 328(8):538–45.
2. Eaton SB, Eaton SB. Physical Inactivity, Obesity, and Type 2 Diabetes: An Evolutionary Perspective. *Res Q Exerc Sport* 2017; 27:1-8. doi: 10.1080/02701367.2016.1268519. PMID: 28129048.
3. Mendis S, Thygesen K, Kuulasmaa K, Giampaoli S, Mähönen M, Ngu Blackett K, et al. World Health Organization definition of myocardial infarction: 2008–09 revision. *Int J Epidemiol* 2011; 40(1):139–46.
4. Elena Galkina and Klaus Ley, Immune and Inflammatory Mechanisms of Atherosclerosis, *Annu Rev Immunol*. *Annu Rev Immunol* 2009; 27:165–97. doi: 10.1146/annurev.immunol.021908.132620.
5. Scott M. Grundy. Rationale for Cholesterol-lowering Strategies Current Problems in Cardiology 1995; 20(5):286-357.
6. Peter W. F. Wilson, Ralph B. D'Agostino, Daniel Levy, Albert M. Belanger, Halit. Prediction of Coronary Heart Disease Using Risk Factor Categories, *Circulation* 1998; 97:1837-47. <https://doi.org/10.1161/01.CIR.97.18.1837>.
7. Ridker PM, Rifai N, Rose L, Buring JE, Cook NR. Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med* 2002; 347(20):1557–65.
8. Ross R. Atherosclerosis –an inflammatory disease. *N Engl J Med* 1999; 340(2):115-26.
9. Zwaka TP, Hombach V, Torzawski J. C-reactive protein-mediated low density lipoprotein uptake by macrophages implications for atherosclerosis. *Circulation* 2001; 103(9):1194–7.
10. Blaschke F, Bruemmer D, Yin F, Takata Y, Wang W, Fishbein MC, et al. C-reactive protein induces apoptosis in human coronary vascular smooth muscle cells. *Circulation* 2004; 110(5):579–87.
11. Grau AJ, Boddy AW, Dukovic DA, Buggle F, Lichy C, Brandt T, et al. Leukocyte count as an independent predictor of recurrent ischemic events. *Stroke* 2004; 35(5):1147–52.
12. Lavie CJ, Thomas RJ, Squires RW, et al: Exercise training and cardiac rehabilitation in primary and secondary prevention of coronary heart diseases. *Mayo Clin Proc* 2009; 84(4):373-83.
13. Hartham SM, Lavie CJ, Milani RV, et al. Benefits of exercise training in secondary prevention of coronary and peripheral arterial disease. *Vasc Dis Prev* 2008; 5:156-68.
14. Lavie CJ, Milani RV: Cardiac rehabilitation update 2008-biological, psychological, and clinical benefits. *US Cardiol* 2008; 5:72-5.
15. Wenger NK. Status of cardiac rehabilitation. *J Am Coll Cardiol* 2008; 51(17):1619-31.
16. Thomas RJ. Cardiac rehabilitation/secondary prevention programs Craft for the rapids: why have we missed the

- boat? *Circulation* 2007; 116(15):1644-6.
17. Milani RV, Lavie CJ. Prevalence and profile of metabolic syndrome in patients following acute coronary events and effects of therapeutic lifestyle change with cardiac rehabilitation. *Am J Cardiol* 2003; 92(1):50-4.
 18. Lavie CJ, Milani RV, Artham SM, Patel DA, Ventura HO. The obesity paradox, weight loss, and coronary disease. *Am J Med* 2009; 122(12):1106-14.
 19. Lavie CJ, Milani RV. Effects of cardiac rehabilitation, exercise training, and weight reduction on exercise capacity, coronary risk factors, behavioral characteristics, and quality of life in obese coronary patients. *Am J Cardiol* 1997; 79(4):397-401.
 20. Lavie CJ, Morshedi-Meibodi A, Milani RV. Impact of cardiac rehabilitation on coronary risk factors, inflammation, and the metabolic syndrome in obese coronary patients. *J Cardiometab Syndr* 2008; 3(3):136-40.
 21. Cardenas GA, Lavie CJ, Cardenas V, Milani RV, McCullough PA. The importance of recognizing and treating low levels of high-density lipoprotein cholesterol: a new era in atherosclerosis management. *Rev Cardiovasc Med* 2008; 9(4):239-58.
 22. Fulya Ilhan and Sevgi Tas Kalkanli, Atherosclerosis and the role of immune cells, *World J Clin Cases*. 2015 Apr 16; 3(4):345–52. doi: 10.12998/wjcc.v3.i4.345 PMID: PMC4391004
 23. Rozanski A, Blumenthal JA, Davidson KW, Saab PG, Kubzansky L. The epidemiology, pathophysiology, and management of psychosocial risk factors in cardiac practice: the emerging field of behavioral cardiology. *J Am Coll Cardiol* 2005; 45(5):637-51.
 24. Lavie CJ, Milani RV. Effects of cardiac rehabilitation and exercise training programs on coronary patients with high levels of hostility. *Mayo Clin Proc* 1999; 74(10):959-66.
 25. Liew FY, Pitman NI, McInnes IB. Disease-associated functions of IL-33: the new kid in the IL-1 family. *Nat Rev Immunol* 2010; 10(2):103-10.
 26. Czyżewska-Buczyńska A, Zuk N, Romanowska-Micherda K, Witkiewicz W. Biological role of Interleukin 33 and its importance in pathophysiology of cardiovascular system]. *Postepy Hig Med Dosw* 2014; 68:834-41.
 27. Milovanovic M, Volarevic V, Radosavljevic G, Jovanovic I, Pejnovic N, Arsenijevic N, et al. IL-33/ST2 axis in inflammation and immunopathology. *Immunol Res* 2012; 52(1-2):89-99.
 28. Milani RV, Lavie CJ, Mehra MR, Ventura HO. Impact of exercise training and depression on survival in heart failure due to coronary heart disease. *Am J Cardiol* 2011; 107(1):64-8.
 29. Dwyer, G.B. and Davis, S.H., *American College of Sport Medicine, health-related physical fitness assessment manual, second edition, 2008*;112-6.
 30. Heeschen C, Dimmeler S, Hamm CW, Fichtlscherer S, Boersma E, Simoons ML, et al. Serum level of the antiinflammatory cytokine interleukin-10 is an important prognostic determinant in patients with acute coronary syndromes. *Circulation* 2003; 107(16):2109–14.
 31. Li X, Mai J, Virtue A, Yin Y, Gong R, Sha X, et al. IL-35 is a novel responsive anti-inflammatory cytokine - a new system of categorizing anti-inflammatory cytokines. *PLoS ONE* 2012; 7(3):e33628.
 32. Milani RV, Lavie CJ. Impact of cardiac rehabilitation on depression and its associated mortality. *Am J Med* 2007; 120(9):799-806.
 33. Ranković G, Milčić B, Savić T, Dindić B, Mancev Z, Pesić G. Effects of physical exercise on inflammatory parameters and risk for repeated acute coronary syndrome in patients with ischemic heart. *Vojnosanit Pregl* 2009; 66(1):44-8.
 34. Pedersen BK. The anti-inflammatory effect of exercise: its role in diabetes and cardiovascular disease control. *Essays Biochem* 2006; 42:105-17.
 35. Khazaei HA, Jalili A, Andarzy S. The effect of one session intense anaerobic exercise (Bruce test) on serum level of IL-6 and IL-33 in volleyball athletes. *Annals of Biological Research*, 2014, 5(2):99-104.
 36. Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, et al. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature* 2010; 463(7280):540-4.