



Pseudomoniasis phytotherapy: a review on most important Iranian medicinal plants effective on *Pseudomonas aeruginosa*

Mahmoud Bahmani^{1,2}, Mahmoud Rafieian-Kopaei³, Hassan Hassanzadazar⁴, Morovat Taherikalani^{5*}

¹Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran

²Razi Herbal Medicine Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

³Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁴Department of Food Safety and Hygiene, Faculty of Health, Zanjan University of Medical Sciences, Zanjan, Iran

⁵Razi Herbal Medicines Research Center and Department of Microbiology, School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

Received: December 2015, Accepted: July 2016

ABSTRACT

Background and Objectives: *Pseudomonas aeruginosa* is a Gram-negative, aerobic bacterium found in water and soil. It is a normal flora in skin and gastrointestinal tract of human beings. *P. aeruginosa* as an opportunistic pathogen involved in nosocomial infections having multiple pathogenic factors and shows high rate of resistance to different antibiotics. The aim of this study was to identify the most important native medicinal plants of Iran effective on *P. aeruginosa*.

Materials and Methods: All required information was obtained by searching keywords such as *P. aeruginosa*, medicinal plant extracts or essential oils in published articles in authentic scientific databases such as Science Direct, Wiley-Blackwell, Springer, Google scholar, Scientific Information Database (SID) and Magiran.

Results: According to the literature review, our results showed 12 different native medicinal plants were effective against *P. aeruginosa* in Iran including *Eucalyptus camadulensis*, *Marticaria chamomilla*, *Ferula gummosa* Boiss, *Lawsonia inermis*, *Ocimumgra tissimum*, *Allium sativum*, *Satureja hortensis* L, *Satureja bachtiarica* Bunge, *Satureja khuzestanica* (Jamzad), *Thymus daenensis* Celak, *Thymus carmanicus* Jalals and *Camellia sinensis*.

Conclusion: Phytochemical analysis has shown that bioactive compounds of medicinal plants with their antioxidant and antimicrobial properties can be good alternatives for the synthetic medicines in food and drug industry.

Keywords: Medicinal plants, *Pseudomonas aeruginosa*, Antimicrobial, Iran

INTRODUCTION

Pseudomonas aeruginosa is a Gram-negative, aerobic bacterium found in water and soil. It is a normal flora of the skin and gastrointestinal tract of human beings (1, 2). This bacterium as an opportunistic pathogen is one of the most important microorganisms in nosocomial infections in immune-compro-

*Corresponding author: Morovat Taherikalani Ph.D, Razi Herbal Medicine Research Center and Department of Microbiology, School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran.
Phone: +98 6633120172
Fax: +98 6633120173
E-mail: taherikalani@gmail.com

mised patients including patients with malignancies, cystic fibrosis and burns with multiple pathogenic factors and high rate of resistance to most of the antibiotics. The presence of glycocalyx in cellular membrane is responsible for easy binding of the bacterium to the host cell, biofilm formation and protection of bacteria against penetration of antimicrobial agents and phagocytic system (3).

In a previous study, *P. aeruginosa* has been isolated from nosocomial pneumonia (16%), nosocomial urinary tract infections (12%), post-operative wound infections (85%) and hospital blood infections (10%) which included 23% of total isolated bacteria from admitted patients (4). Selection of the appropriate treatment approach against *P. aeruginosa* is limited due to its resistance to common recommended antibiotics (5, 6).

Bacterial resistance to antibiotics is increasing which makes humans to apply effective antimicrobial agents with fewer side effects such as medicinal plants instead of medicines with less efficacy and greater side effects (7). Plants have different chemical compounds like secondary metabolites (8, 9) with many biochemical and bioactivity properties showing applications in various industries such as pharmaceutical, chemical, cosmetic and food industry (10-13). The aim of this study was to introduce the most important native medicinal plants of Iran being effective on *P. aeruginosa*.

MATERIALS AND METHODS

All required information was obtained by literature review using keywords including *P. aeruginosa*, medicinal plant extracts or essential oils of published articles in authentic scientific databases such as Science Direct, Wiley-Blackwell, Springer, Google scholar, Scientific Information Database (SID) and Magiran.

RESULTS

According to literature review, our results showed 12 different native medicinal plants were effective against *P. aeruginosa* in Iran including *Eucalyptus camadulensis*, *Marticaria chamomilla*, *Ferula gummosa Boiss*, *Lawsonia inermis*, *Ocimumgra tissimum*, *Allium sativum*, *Satureja hortensis L*, *Satureja*

bachtiarica Bunge, *Satureja khuzestanica (Jamzad)*, *Thymus daenensis Celak*, *Thymus carmanicus Jalals* and *Camellia sinensis*. Effective medicinal plants against *P. aeruginosa* are listed in Table 1.

DISCUSSION

Secondary compounds have attracted particular attention to the plants in recent years. Medicinal plants are good alternatives for synthetic preservatives in food and drug industry due to their antimicrobial compounds. *E. camadulensis* (Eucalyptus) leaves are rich in polyphenols and terpenoids and eucalyptol or cineole (26). Phytochemicals results showed that alfa and beta-pinenes are the main ingredients of *Ferula gummosa Boiss* (galbanum) (27) and *Lawsoniainermis* (Henna) contains manitol, tanic acid, mucilage and galic acid but its most important ingredient is 2-Hydroxy-1,4-naphoquinone (Lawson) recognized as bioactive agent. *Ocimumgra tissimum* (basil) contains essential oils, tannins, glycosides, saponin, anthocyanin, eugenol, linalool, methyl cinnamate, camphor and thymol. Allicin is the major component of *Allium sativum* (28). Different species of *Satureja* (Savory) contain monoterpenes, phenolic compounds especially carvacrol, thymol and eugenol as well as some of the sesquiterpenes. The main phytochemical essential oils of aromatic plants are hydrocarbons, aldehydes, ketones, alcohols, phenols, ethers and esters with phenolic and terpenic sources. The presence of linalool, alpha-pinene, beta-pinene, borneol, carvone, limonene, carvacrol, p-cymene and terpinene essential oils of understudy plants showed antimicrobial properties particularly (28). These compounds with high hydrophobic properties can separate lipids from bacterial cell wall and thereby increase the permeability of the membrane which leads to ion expelling and electron imbalance and eventually cell death. The researchers demonstrated that plant species used in popular medicine are promising resources for antimicrobial treatments (13, 14, 29, 30).

CONCLUSION

Phytochemical analysis has shown that bioactive compounds of medicinal plants with their antioxidant and antimicrobial properties are good substitutions for synthetic drugs in food and drug industry.

Table 1. List of effective Iranian native medicinal plants against *P. aeruginosa* (PA)

Scientific name	Plant Family	Persian name	Discussion
<i>Eucalyptus camadulensis</i>	Myrtaceae	Eucalyptus	Minimum inhibitory concentration (MIC) of alcoholic and aqueous extracts were 3.3 mg/ml and 17.5 mg/l. Increasing concentration of the extract prevented growth more effectively in comparison with control group (13).
<i>Marticaria chamomilla</i>	Asteraceae	Babooneh	At a concentration of 0.5 mg/ml of essential oil the growth of PA was significantly lower than the positive control (14).
<i>Ferula gummosa Boiss</i> (Galbanum)	Umlleliferae	Baryjeh	MIC of methanol and ethanol extracts was 1.25×10^4 and 625×10^3 µg/ml, respectively (15).
<i>Lawsonia inermis</i>	Lythraceae	Henna	MIC of aqueous and ethanol extracts was 10 and 3.5 mg/ml, respectively (16).
<i>Ocimum gratissimum</i> (Basil)	Laminaceae	Reyhan	The inhibition zone diameter of ethanol extract with 100 and 200 mg/ml concentration were 8 and 12 mm (in diameter) but for aqueous extract (with the same concentration), they were 4 and 9 mm, respectively (17).
<i>Allium sativum</i>	Liliaceae	Sir	MIC of the extract at 220 µg/ml conc. had a 16 mm growth inhibition zone (in diameter) (18). In another study MIC and MBC of Allicin were 2.38 and 4.77 mg/ml, respectively (25).
<i>Satureja hortensis L</i>	Labiatae	Marzeh	Growth inhibition zone in disk diffusion method was 19.88 mm (in diameter) (19).
<i>Satureja bachtiarica Bunge</i>	Labiatae	Marzeh bakhtyari	Growth inhibition zone in disk diffusion method was 22.22 mm (in diameter) (19).
<i>Satureja khuzestanica</i> (Jamzad)	Labiatae	Marzeh khoozestani	Growth inhibition zone in disk diffusion method was 18.33 mm (in diameter) (19).
<i>Thymus daenensis Celak</i>	Laminaceae	Avishan denaee	Growth inhibition zone in disk diffusion method was 21.88 mm (in diameter) (19).
<i>Thymus carmanicus</i>	Laminaceae	Avishan kermani	Growth inhibition zone in disk diffusion method was 20.66 mm (in diameter) (19).
<i>Jalals</i>	Laminaceae	Avishan kermani	Growth inhibition zone in disk diffusion method was 20.66 mm (in diameter) (19).
<i>Camellia sinensis</i>	Theaceae	Katchin (Chaysabz)	MIC at 0.01 dilution was 7.24 µg/ml (24).

ACKNOWLEDGEMENT

This paper was prepared with the support of Research Deputy of Shahrekord University of Medical Sciences, Shahrekord, Iran.

REFERENCES

- Jabalamehli F, Mirsalehian A, Sotoudeh N, Jabalamehli L, Aligholi M, Khoramian B, et al. Multiple-locus variable number of tandem repeats (VNTR) fingerprinting (MLVF) and antibacterial resistance profiles of extended spectrum β -lactamase (ESBL) producing *Pseudomonas aeruginosa* among burnt patients in Tehran. *Burns* 2011; 37:1202-1207.
- Jabalamehli F, Mirsalehian A, Khoramian B, Aligholi M, Khoramrooz SS, Asadollahi P, et al. Evaluation of biofilm production and characterization of genes encoding type III secretion system among *Pseudomonas aeruginosa* isolated from burn patients. *Burns* 2012; 38:1192-1197.
- Russell NJ, Gacesa P. Chemistry and biology of the alginate of mucoid *Pseudomonas aeruginosa* in cystic fibrosis. *Mol Aspects Med* 1988; 10: 1-91.
- van Deleden C, Iglewski BH. Cell to cell signaling and *Pseudomonas aeruginosa* infections. *Emerg Infect Dis* 1998; 4: 551-559.
- Duke JA (2002). Handbook of medicinal herbs. CRC Press, Boca Raton.
- Moniri R, Mosayebi Z, Movahedian AH. Increasing trend of antimicrobial drug-resistance in *Pseudomonas aeruginosa* causing septicemia. *Iran J Public Health* 2006; 35: 58-62.
- Rios JL, Recio MC. Medicinal plants and antimicrobial activity. *J Ethnopharmacol* 2005; 100:80-84.
- Gourine N, Yousfi M, Bombarda I, Nadjemi B, Stocker P, Gaydou EM. Anti oxidant activities and chemical composition of essential oil of *Pistacia atlantica* from Algeria. *Ind Crops Prod* 2010; 31: 203-208.
- Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative stress and Parkinson's disease: New

- hopes in treatment with herbal antioxidants. *Curr Pharm Des* 2016; 22: 238-246.
10. Rabiei Z, Rafieian-Kopaei M. Neuroprotective effect of pretreatment with *Lavandula officinalis* ethanolic extract on blood-brain barrier permeability in a rat stroke model. *Asian Pac J Trop Med* 2014; 7S1: S421-S426.
 11. Dangl JL, Jones JD. Plant pathogens and integrated defense responses to infection. *Nature* 2001; 411: 826-833.
 12. Yamane H, Konno K, Sabelis M, Takabayashi J, Sassa T, Oikawa H (2010). Chemical defense and toxins of plants. In: *Comprehensive Natural Products II*. Ed, L Mander, HW Lui. Elsevier, Oxford, pp. 339-385.
 13. Eslami M, Bayat M, Mozaffari Nejad AS, Sabokbar A, Anvar AA. Effect of polymer/nanosilver composite packaging on long-term microbiological status of Iranian saffron (*Crocus sativus* L.). *Saudi J Biol Sci* 2016; 23: 341-347.
 14. Jalali H, Mozaffari Nejad AS, Ebadi AG, Laey G. Ethnobotany and folk pharmaceutical properties of major trees or shrubs in Northeast of Iran. *Asian J Chem* 2009; 21: 5632-2538.
 15. Torkelson AR (1999). The cross name index to medicinal plants, four volume set. CRC Press, Boca Raton.
 16. Aleebrahim-Dehkordy E, Rafieian-Kopaei M, Bahmani M. Study of antibacterial effect of total hydroalcoholic extract of *Thymus daenensis* and *Lavandula officinalis* on *Listeria monocytogenes*, an agent of food spoilage, using microdilution. *Der Pharma Chemica* 2016; 8:238-242.
 17. Mbata TI, Saikia A. Antibacterial activity of essential oil from *Ocimumgra tissimum* on *Listeria monocytogenes*. *Internet J Food Safety* 2006; 7: 15-19.
 18. Rabinkov A, Miron T, Konstantinovski L, Wilchek M, Mirelman D, Weiner L. The mode of action of allicin: trapping of radicals and interaction with thiol containing proteins. *Biochim Biophys Acta* 1998; 1379: 233-244.
 19. Burt S. Essential oils: their antibacterial properties and potential applications in foods- a review. *Int J Food Microbiol* 2004; 94: 223-253.
 20. Dorman HJ, Deans SG. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol* 2000; 88: 308-316.
 21. Duffy CF, Power RF. Antioxidant and antimicrobial properties of some Chinese plant extracts. *Int J Antimicrob Agents* 2001; 17: 527-529.
 22. Farag RS, Daw ZY, Hewedi FM, El-Baroty GSA. Antimicrobial activity of some Egyptian spice essential oils. *J Food Prot* 1989; 52: 665-667.
 23. Carson CF, Mee BJ, Riley TV. Mechanism of action of *Melaleuca alternifolia* (tea tree) oil on *Staphylococcus aureus* determined by time-kill, lysis, leakage and salt tolerance assays and electron microscopy. *Antimicrob Agents Chemother* 2002; 46: 1914-1920.
 24. Cox SD, Mann CM, Markham JL, Bell HC, Gustafson JE, Warmington JR, et al. The mode of antimicrobial action of essential oil of *Melaleuca alternifolia* (tea tree oil). *J Appl Microbiol* 2000; 88: 170-175.
 25. Bahmani MP, Bahmani M, Shahsavari S, Naghdi N, Ezatpour B, Moradniani M, et al. A review of the antiparasitic medicinal plants used in ethnobotany of different regions of Iran. *Der Pharma Chemica* 2016; 8:134-138.
 26. Lambert RJ, Skandamis PN, Coote PJ, Nychas GJ. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *J Appl Microbiol* 2001; 91: 453-462.
 27. Skandamis PN, Nychas GJ. Effect of oregano essential oil on microbiological and physico-chemical attributes of minced meat stored in air and modified atmospheres. *J Appl Microbiol* 2001; 91: 1011-1022.
 28. Thoroski J, Blank G, Biliaderis C. Eugenol induced inhibition of extracellular enzyme production by *Bacillus cereus*. *J Food Prot* 1989; 52: 399-403.
 29. Kamkar A, Shamse Ardekani MR, Shariatifar N, Misagi A, MozaffariNejad AS, Jamshidi AH. Antioxidative effect of Iranian *Pulicaria gnaphalodes* L. extracts in Soybean oil. *S Afr J Bot* 2013; 85: 39-43.
 30. Mozaffari Nejad AS, Shabani S, Bayat M, Hosseini SE. Antibacterial effect of garlic aqueous extract on *Staphylococcus aureus* in Hamburger. *Jundishapur J Microbiol* 2014; 7: e13134.