

***Physalis alkekengi*: A review of Its Therapeutic Effects****Mahmoud Bahmani<sup>1</sup>, Mahmoud Rafeian-Kopaei<sup>2</sup>, Nasrollah Naghdi<sup>1</sup>, Amir Sasan Mozaffari Nejad<sup>3,\*</sup> and Omid Afsordeh<sup>1</sup>**<sup>1</sup> Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran<sup>2</sup> Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran<sup>3</sup> Molecular Medicine Research Center, Hamadan University of Medical Sciences, Hamadan, Iran**\* Corresponding author. Tel.: +98-8138380755. Fax: +98-8138380130****E-mail address: asmozafarinejad@yahoo.in****ABSTRACT**

Medicinal plants have long been used in medicine to treat diseases. There has been a widespread approach to use nature-based, especially plant-based, drugs in recent years. *Physalis alkekengi*, is an herbaceous, perennial plant. The leaves are tailed and paired from family in Solanaceae. This plant can speed up excretion of uric acid and is used for renal and urinary tract diseases, gout, and rheumatism. Other properties of this plant include anti-gout, anti-inflammatory, antibacterial, analgesic, laxative, diuretic, antimetabolic, and malaria-treating properties. Moreover, modern medical investigations have demonstrated that *P. alkekengi* is effective on immunity system, cancer, thyroid hormones, liver enzymes, and sexual and reproductive hormones. Phytochemical investigations indicate that this plant contains alkaloids, glucocorticoids, lycopene, ethanolic compounds, and vitamin C. The therapeutic effects of *P. alkekengi* can be due to these antioxidant compounds. This review article aimed to present the phytochemical and pharmaco-therapeutic properties of this plant.

**KEYWORDS:** Medicinal plants, *Physalis alkekengi*, Therapeutic effects.**1. INTRODUCTION**

There has been a widespread approach to use nature-based drugs in recent years. Medicinal plants have long been used in medicine to treat diseases (Ebrahimie, 2015; Amirmohammadi, 2014; Bahmani, 2012; Eftekhari, 2012; Bahmani, 2013; Gholami-Ahangaran, 2012). Having knowledge about medicinal plants, as a medicinal resource, which have various therapeutic effects dates back to the first human beings (Jalali, 2009; Delfan, 2014; Mozaffari Nejad, 2013). When there were no chemical drugs, human beings used medicinal plants to treat diseases and complications (Asadi Samani, 2014; Delfan, 2014; Bahmani, 2014; Saki, 2014; Bahmani, 2014; Bahmani, 2014; Karamat, 2014; Bahmani, 2014). The medicinal plants are in general cheap and accessible around the world (Bahmani, 2014; Delfan, 2015; Bahmani, 2015; Bahmani, 2015; Ghasemi, 2015; Bahmani, 2015). Implemented as alternative health care treatment for prevent of side effects than chemical drugs (Delfan, 2014; Bahmani, 2013; Bahmani, 2015; Delfan, 2015; Delfan, 2015; Mozaffari Nejad, 2013). However, some parts of medicinal plants are specified as natural products that used for antimicrobial agents. Also, other uses for natural antioxidants contain food additives, bio pharmaceuticals and bioactive nutraceuticals (Mozaffari Nejad, 2014; Eslami, 2016; Bahmani, 2016). *Physalis alkekengi* L. is one of these plants being herbaceous, perennial plant. The leaves are tailed and paired from family Solanaceae. This plant can speed up excretion of uric acid and is used for renal and urinary tract diseases, gout, and rheumatism. Other properties of this plant include anti-gout, anti-inflammatory, antibacterial, analgesic, laxative, diuretic, antimetabolic, and malaria-treating properties. Moreover, modern medical investigations have demonstrated that *P. alkekengi* is effective on immunity system, cancer, thyroid hormones, liver enzymes, and sexual and reproductive hormones. Phytochemical investigations indicate that this plant contains alkaloids, glucocorticoids, lycopene, ethanolic compounds, and vitamin C. The therapeutic effects of *P. alkekengi* L. can be due to these antioxidant compounds. This review article aimed to present the phytochemical and pharmaco-therapeutic properties of this plant (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008; Namjoyan, 2015; Helvaci, 2010).

**Ethnobotany of *Physalis alkekengi*:** Husk tomato, scientifically called *Physalis alkekengi*, is an herbaceous, perennial plant. The leaves are tailed and paired from family Solanaceae (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008).

**Effective compounds of *P. alkekengi*:** Alkaloids and glucocorticoids are the main chemical compounds of this plant. *P. alkekengi* fruit contains lycopene, alkaloids, ethanolic compounds, and a large amount of vitamin C. *P. alkekengi* contains, glucocorticoids, alkaloids, lycopenes, ethanolic compounds, and vitamin C (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008; Namjoyan, 2015).

**Therapeutic effects of *P. alkekengi* according to traditional medicine:** This plant can speed up excretion of uric acid and is used for renal and urinary tract diseases, gout, and rheumatism. Other properties of this plant include anti-gout, anti-inflammatory, antibacterial, analgesic, laxative, diuretic, antimetabolic, and malaria and syphilis treating (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008; Namjoyan, 2015; Helvaci, 2010; Vessal, 1991; Nasimi, 200).

**Therapeutic effects of *P. alkekengi* according to modern medicine:** A review of the findings of different studies indicates that *P. alkekengi* can be effective on immunity system, cancer, thyroid hormones, liver enzymes, and sexual and reproductive hormones shown in Table 1 (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008; Namjoyan, 2015;

**Table.1. Therapeutic effects of *Physalis alkekengi* according to modern medicine.**

Effect of <i>P. alkekengi</i> on immunity system	Findings have demonstrated that physalins <i>P. alkekengi</i> decreased inflammation through decreasing neutrophils infiltration and inhibiting the formation of interleukin-6 and interleukin-12.
Effect of <i>P. alkekengi</i> on cancer	A study demonstrated that <i>P. alkekengi</i> physalins can help to inhibit cancer cells growth by stopping the cells at G2/M phase of cell cycle. In fact, the effects of physalins are exerted through decrease in the expressions and activities of the cyclins A and B and the activity of Cdc2 and increase in the phosphorylation of Cdc2.
Effect of <i>P. alkekengi</i> on thyroid hormones	An experimental study demonstrated that <i>P. alkekengi</i> extract caused increase in the concentrations of T3 and T4 hormones. The increase in thyroid hormones is due to anti-steroidal compounds that cause inhibition of secretion of catecholamines and increase in plasma proteins such as albumin, which causes increase in thyroid hormones in plasma.
Effect of <i>P. alkekengi</i> on liver enzymes and blood biochemical factors	A study demonstrated that aqueous extract of <i>P. alkekengi</i> contains ascorbic acid and polyphenols. Moreover, this extract was able to exert positive antioxidant activity alongside inhibitory activity on lipids peroxidation, and was a free radical scavenger. Intraperitoneal administration of rats with <i>P. alkekengi</i> extract caused increase in alanine aminotransferase aspartate aminotransferase, albumin, and total protein, while no significant changes were seen in the amounts of bilirubin, creatinine, uremia nitrogen, and the activity of alkaline phosphatase.
Effect of <i>P. alkekengi</i> on fertility and sexual hormones	An investigation showed that intraperitoneal administration of pregnant rats with 400 mg/kg of <i>P. alkekengi</i> for eight days caused a significant decrease in progesterone and activity of 3beta-hydroxysteroid dehydrogenase. In this work, the activity of 20alpha-hydroxysteroid dehydrogenase was also measured. Another work investigated the effect of <i>P. alkekengi</i> on fertility in rats. This study demonstrated that intraperitoneal administration of 150 mg/kg <i>P. alkekengi</i> for 56 consecutive days caused contraceptive and antispermatogenic effects.

Modern medical investigations have demonstrated that *P. alkekengi* is effective on immunity system, cancer, thyroid hormones, liver enzymes, and sexual and reproductive hormones. Phytochemical investigations indicate that this plant contains alkaloids, glucocorticoids, physalis, lycopene, ethanolic compounds, and vitamin C. It has antioxidant activity and the some therapeutic effects of *P. alkekengi* have been attributed to its antioxidant compounds (Sadeghi, 2007; Ge, 2009; Gharib Naseri, 2008; Namjoyan, 2015; Zarei, 2011; Helvacı, 2010; Vessal, 1991; Vessal, 2004; Nasimi, 2008).

Antioxidants are substances that are able to prevent harmful chemical reactions in them oxygen is combined with other substances (Rafieian Kopaie and Baradaran, 2013; Baradaran, 2014; Nasri and Rafieian-Kopaei, 2014; Rafieian Kopaie, 2013). Antioxidant can delay undesirable qualities of foods such as rancidity of various foods. They are a wide variety of compounds such as aromatic amines and phenolic compounds. Phenolic compounds are present in numerous medicinal plants (Nasri and Rafieian-Kopaei, 2013; Baharvand Ahmadi, 2016; Nasri, 2014). These phenolic compounds have antioxidant activity and can combat oxidative stress. Hence, they are used for prevention and treatment of numerous diseases (Bahmani, 2016; Sarrafchi, 2016; Nasri, 2015; Akhlaghi, 2011). Other than disease, antioxidants may be used to delay the development of rancidity. For this, organic antioxidants such as propyl gallate, butylated hydroxytoluene (BHT), or butylated hydroxyanisole (BHA) are usually used. Also, these antioxidants used in oil products. However, there are limitations on quantity of antioxidants that may be used. Hence, medicinal plants such as *P. alkekengi* L. which have antioxidant activity might be preferred to synthetic antioxidants. An important subject that should be noted is that this plant is effective in numerous diseases such as diabetes mellitus, atherosclerosis, cancer, cardiovascular diseases, infection and toxicities which most of them are associated with oxidative stress (Baradaran, 2012; Rabiei, 2014; Rahimian, 2013; Rahnama, 2015; Rafieian-Kopaei, 2014; Azizkhani, 2011; Kamkar, 2013). Therefore, patient who have one of these diseases may benefit from antioxidant property of this plant, too. It should be noted that diseases have various mechanism (Baradaran, 2013; Heidarian, 2013; Billah, 2016; Rafieian Kopaie, 2013; Nasri and Rafieian-Kopaei, 2013). This plant also has a wide variety of compounds which have therapeutic effects and antioxidant property of this plant may help them to be treated better.

## 2. CONCLUSION

In the present review, it is suggested that activity of *Physalis alkekengi* investigated against therapeutic properties. The isolation, identification of active principles and pharmacological studies of the active phyto-constituents may be considered and studied elaborately to treat effectively for therapeutic effects.

## REFERENCES

- Akhlaghi M, Shabani Gh, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. Citrus aurantium Blossom and Preoperative Anxiety. *Rev Bras Anesthesiol.*, 61, 2011, 702-712.
- Amirmohammadi M, Khajoenia SH, Bahmani M, Rafieian-Kopaei M, Eftekhari Z, Qorbani M. In vivo evaluation of antiparasitic effects of Artemisia abrotanum and Salvia officinalis extracts on Syphacia obvelata, Aspiculuris tetrapetra and Hymenolepis nana parasites. *Asian Pac J Trop Dis.*, 4, 2014, 250-254.
- Asadi-Samani M, Bahmani M, Rafieian-Kopaei M. The chemical composition, botanical characteristic and biological activities of Borago officinalis: a review. *Asian Pac J Trop Med.*, 7, 2014, 22-28.
- Azizkhani M, Kamkar A, Mozaffari Nejad AS. Effects of Tocopherols on Oxidative Stability of Margarine. *J Chem Soc Pakistan.*, 33, 2011, 134-137.
- Baharvand-Ahmadi B, Bahmani M, Tajeddini P, Naghdi N, Rafieian-Kopaei M. An ethno-medicinal study of medicinal plants used for the treatment of diabetes. *J Nephropathol.*, 5, 2016, 44-50.
- Bahmani M, Banihabib EKH M, Rafieian-Kopaei M and Gholami-Ahangaran M. Comparison of disinfection activities of nicotine with copper sulphate in water containing Limnatis nilotica. *Kafkas Univ Vet Fak Derg.*, 21, 2015, 9-11.
- Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Saki K, Karamati SA, Delfan B. A review on most important herbal and synthetic antihelmintic drugs. *Asian Pac J Trop Med.*, 7, 2014, 29-33.
- Bahmani M, Rafieian-Kopaei M, Jeloudari M, Eftekhari Z, Delfan B, Zargar A, Forouzan SH. A review of the health effects and uses of drugs of plant licorice (*Glycyrrhiza glabra L.*) in Iran. *Asian Pac J Trop Dis.*, 4, 2014, 847-849.
- Bahmani M, Shirzad H, Rafieian S, Rafieian Kopaei M. Silybum marianum: Beyond Hepatoprotection. *Evid J Based Complementary Altern Med.*, 20, 2015, 292-301.
- Bahmani M, Zargar A, Rafieian-Kopaei M. Identification of medicinal plants of Urmia for treatment of gastrointestinal disorders. *Rev Bras Farmacogn.*, 24, 2014, 468-480.
- Baradaran A, Nasri H, Rafieian Kopaei M. Comment on: Anti-oxidative stress activity of Stachys lavandulifolia aqueous extract in humans. *Cell J.*, 15, 2013, 272-273.
- Billah MM, Nawrin K, Ahmed KT, Javed MSU, Islam MN, Uddin MM. GABA mediated response of aqueous, ethanol and ethyl acetate extracts of Dicranopteris linearis leaf in Swiss Albino mice. *J Herbm Pharm.*, 5, 2016, 1-6.
- Changizi Ashtiyani S, Zarei A. The effects of Physalis alkekengi alcoholic extract on certain plasma biochemical factors in rats. *Arak Univ J Med Sci.*, 14, 2011, 18-25.
- Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M. Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran. *Asian Pac J Trop Med.*, 7, 2014, 7(Suppl 1): 376-379.
- Delfan B, Bahmani M, Rafieian-Kopaei M, Delfan M, Saki K. A review study on ethnobotanical study of medicinal plants used in relief of toothache in Lorestan Province, Iran. *Asian Pac J Trop Dis.*, 4, 2014, 879-884.
- Ebrahimi M, Bahmani M, Shirzad H, Rafieian-Kopaei M, Saki K. A Review Study on the Effect of Iranian Herbal Medicines on Opioid Withdrawal Syndrome. *J Evid Based Complementary Altern Med.*, 20, 2015, 302-309.
- Eftekhari Z, Bahmani M, Mohsenzadegan A, Gholami-Ahangaran M, Abbasi J, Alighazi N. Evaluating the anti-leech (*Limnatis nilotica*) activity of methanolic extract of *Allium sativum L.* compared with levamisole and metronidazole. *Comp Clin Path.*, 21, 2012, 1219-1222.
- Eslami M, Bayat M, Mozaffari Nejad AS, Sabokbar A, Anvar AA. Effect of polymer/nanosilver composite packaging on long-term microbiological status of Iranian saffron (*Crocus sativus L.*). *Saudi J Biol Sci.*, 23, 2016, 341-347.
- Ge Y, Duan Y, Fang G, Zhang Y, Wang S. Study on biological activities of Physalis alkekengi var. francheti polysaccharide. *J Sci Food Agric.*, 89, 2009, 1593-1598.
- Gharib-Naseri MK, Mohammadian M, Gharib-Naseri Z. Antispasmodic effect of Physalis alkekengi fruit extract on rat uterus. *Iran J Reprod Med.*, 6, 2008, 193-198.
- Gholami-Ahangaran M, Bahmani M, Zia-Jahromi N. Comparative and evaluation of anti-leech (*Limnatis Nilotica*) effect of Olive (*Olea Europaea L.*) with levamisole and tiabendazole. *Asian Pac J Trop Dis.*, 2, 2012, 101-103.
- Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynara scolymus*) leaf extract against lead toxicity in rat. *Pharm Biol.*, 51, 2013, 1104-1109.
- Helvacı S, Kökdil G, Kawai M, Duran N, Duran G, Güvenç A. Antimicrobial activity of the extracts and physalin D from Physalis alkekengi and evaluation of antioxidant potential of physalin D. *Pharm Biol.*, 48, 2010, 142-150.
- Jalali H, Mozaffari Nejad AS, Ebadi AG, Laey G. Ethnobotany and folk pharmaceutical properties of major trees or shrubs in Northeast of Iran. *Asian J Chem.*, 21, 2009, 5632-5638.
- Javdan N, Estakhr J. Physalis Alkekengi reduces spermatogenesis and camp-responsive element modulator gene expression in rat. *Pharmacologyonline*, 2, 2011, 866-873.

- Kamkar A, Shamse Ardekani MR, Shariatifar N, Misagi A, Mozaffari Nejad AS, Jamshidi AH. Antioxidative effect of Iranian *Pulicaria gnaphalodes* L. extracts in soybean oil. *S Afr J Bot.*, 85, 2013, 39-43.
- Karamati SA, Hassanzadazar H, Bahmani M, Rafieian-Kopaei M. Herbal and chemical drugs effective on malaria. *Asian Pac J Trop Dis.*, 4, 2014, 599-601.
- Mozaffari Nejad AS, Shabani S, Bayat M, Hosseini SE. Antibacterial Effect of Garlic Aqueous Extract on *Staphylococcus aureus* in Hamburger. *Jundishapur J Microb.*, 7, 2014, e13134.
- Namjoyan F, Jahangiri A, Azemi M.E, Arkian E, Mousavi H. Inhibitory Effects of *Physalis alkekengi* L., *Alcea rosea* L., *Bunium persicum* B. Fedtsch. and *Marrubium vulgare* L. on Mushroom Tyrosinase. *Jundishapur J Nat Pharm Prod.*, 10, 2015, e23356.
- Nasimi M, Heydari M, Shiravi A. The Effects of *Physalis alkekengi* fruit's alcoholic extract on placenta growth in Rat. *Journal of Anima Biology*, 1, 2008, 51-60.
- Nasri H, Ahmadi A, Baradaran A, Momeni A, Nasri P, Mardani S, Rafieian-Kopaei M, Mubarak M. Clinicopathological correlations in lupus nephritis; a single center experience. *J Nephrothol.*, 3, 2014, 115-1120.
- Rabiei Z, Rafieian-Kopaei M. Neuroprotective effect of pretreatment with *Lavandula officinalis* ethanolic extract on blood-brain barrier permeability in a rat stroke model. *Asian Pac J Trop Med.*, 7, 2014, 421-426.
- Rahimian GA, Rabiei Z, Tahmasebi B, Rafieian-Kopaei M, Ganji F, Rahimian R. Comparing the combined effect of Garlic and Mint extract with Metronidazole in *Helicobacter Pylori* treatment. *Iran J Pharm Sci.*, 9, 2013, 63-70.
- Rahnama S, Rabiei Z, Alibabaei Z, Mokhtari S, Rafieian-kopaei M, Deris F. Anti-amnesic activity of *Citrus aurantium* flowers extract against scopolamine-induced memory impairments in rats. *Neurol Sci.*, 36, 2015, 553-560.
- Saki K, Bahmani M, Rafieian-Kopaei M. The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review. *Asian Pac J Trop Med.*, 7, 2014, 34-42.
- Sarrafi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative stress and Parkinson's disease: New hopes in treatment with herbal antioxidants. *Curr Pharm Des.*, 22, 2016, 238-246.
- Shekar Forosh S, Changizi-Ashtiyani S AP, Attari MM. The effect of *Physalis alkekengi* alcoholic extract on thyroid hormones concentrations in rats. *Zahedan J Res Med Sci.*, 13, 2012, 1-7.
- Soares MBP, Bellintani MC, Ribeiro IM, Tomassini TCB, Ribeiro dos Santos R. Inhibition of macrophage activation and lipopolysaccharide-induced death by secosteroids purified from *Physalis angulata* L. *Eur J Pharmacol.*, 459, 2003, 107-112.
- Mozaffari Nejad AS, Kamkar A, Giri A, Pourmahmoudi AA. Ethnobotany and folk medicinal uses of major trees and shrubs in Northern Iran. *J Med Plants Res.*, 7, 2013, 284-289.)
- Mozaffari Nejad AS, Bayat M, Ahmadi AA. Investigation of Aflatoxin B1 in Spices Marketed in Hyderabad, India by ELISA Method. *J Pure Appl Microbio.*, 7, 2013, 3219-3223.
- Vessal M, Fathi N, Khoushdel Z. Effect of aqueous extract of *physalis alkekengi* fruits on the activity of ovarian 3beta and 20alpha-hydroxysteroid dehydrogenases in late pregnancy in rat. *Iran J Med Sci.*, 29, 2004, 175-179.
- Vessal M, Mehrani H, Omrani GH. Effects of an aqueous extract of *Physalis alkekengi* fruit on estrus cycle, reproduction and uterine creatine kinase BB-isozyme in rats. *J. Ethnopharmacol.*, 34, 1991, 69- 78.
- Zarei A, Ashtiyani SC, Rasekh F, Mohammadi A, Jabary A. The effect of *physalis Alkekengi* extracts on lipids concentrations in rats. *J Arak Univ Med Sci.*, 14, 2011, 36-42.