



## International Journal of PharmTech Research

CODEN (USA): IJPRIF, ISSN: 0974-4304, ISSN(Online): 2455-9563  
Vol.9, No.6, pp 285-290 2016

### Chemical composition of Elamit *Scrophularia deserti*

Mahnaz Mardani<sup>1</sup>, Sadegh Rezapour<sup>1</sup>, Zohreh Eftekhari<sup>3</sup>,  
Majid Asadi-Samani<sup>4\*</sup>, Marzieh Rashidipour<sup>5</sup>, Omid Afsordeh<sup>2</sup>,  
Farzad Kazemzadeh<sup>2</sup>, Fariba Bahmani<sup>2</sup>

<sup>1</sup>Nutritional Health Research Center, Health and Nutritional Department, Lorestan University of Medical sciences, Khorramabad, Iran

<sup>2</sup>Department of Microbiology, Faculty of Medicine and Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran

<sup>3</sup>Research & Development Department, Research & Production Complex, Pasteur Institute of Iran, Tehran, Iran

<sup>4</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>4</sup>Medical Plants Research Center, Shahrekord University of Medical sciences, Shahrekord, Iran

<sup>5</sup>Young Researchers and Elite Club, Khorramabad Branch, Islamic Azad University, Khorramabad, Iran

**Abstract:** *Scrophularia deserti* is an annual and perennial herb, as well as a genus of shrubs. Flowers have bilateral or rarely radial symmetry reaching height of 10–50 cm. Leaves of plant are thick, hard, brittle with green color. *S. deserti* plant from the family of Scrophulariaceae is mostly grass or shrub and rarely trees. It has five-pointed flowers, corolla with lobes and the fruits usually have a capsule with multiple seeds. Aerial parts of *Scrophularia deserti* were collected during November of 2014 in Dehloran region of Ilam province. After confirming the species and genus of the plant in natural resources research center of Ilam Province, the collected plant was dried in the air and powdered with mixer. Essential oil was extracted and analyzed with chromatography / mass spectrometry (GC-MS), SPME method. Phytochemistry results showed that the most active ingredients of the plant were  $\alpha$ -PINENE,  $\beta$ -Phellandrene and  $\beta$ -Myrcene which respectively formed 24.69%, 20.58% and 11.82% of the essential oils of the plant.

**Keywords:** plant, snapdragon, arid, oil, SPME method,  $\alpha$ -PINENE.

### Introduction

Scrophulariaceae family includes about 268 genera and over 5,100 species. Members of the Scrophulariaceae have a cosmopolitan distribution, with the majority found in high temperate areas, including Arabian Desert and adjacent territory and Touran, including Egypt, Palestine, Jordan, Syria, Iraq, Saudi Arabia, Kuwait, Bahrain and Iran (1). *Scrophularia deserti* is annual and perennial herb, as well as a genus of shrubs. Flowers have bilateral or rarely radial symmetry reaching to 10–50 cm height. Leaves of the plant are thick, hard, brittle with green color. *S. deserti* plant from the family of Scrophulariaceae is mostly grass or shrub and rarely trees, with five-pointed flowers, corolla with lobes and the fruits are usually in a capsule with multiple seeds. (2, 3). This plant is abundant in most parts of Iran and is called *Scrophularia desertisazooei*. This plant

has been used in the west region of Iran especially in Ilam province (4). In traditional medicine, local people experimentally use it in some disorders such as internal infections, mastitis, skin ulcers and episiotomy, inflammation, burns, intestinal pain, inflammation, eye and ear infections and hemorrhoids in the form of decoction by oral or topical uses (5,6). Chemical analysis of snapdragon was performed previously, but the plant essential oil chemical analysis has not been reported. In this study, for the first time chemical analysis of essential oil was examined and reported.

**Table 1. Scrophularia deserti plant**

Species	Plant family	Region	Province	Latitude	Longitude	Altitude (m a.s.l)	Used part
Scrophularia deserti	Scrophulariaceae	Dehloran city	Ilam (West of Iran)	58°26 N	13°33 E	683-790	Aerial organs

## Methods

### Preparation of medicinal plants

Specimens of the species of snapdragon were taken from different parts of Ilam province during 22 to 30 November 2014 (Table 1). Oregon plant shoots of 10 samples were collected. From the aerial parts of the plant ten samples were collected. The collected plant was submitted to Natural Resources Research Center of Ilam province and the plant was identified and the essential oils were determined (table 1)

### Gas chromatography/mass spectrometry (GC–MS) analysis

The essential oils were analyzed using an Agilent 6890N coupled to Agilent 5973 mass detector gas chromatograph (Agilent, USA) with a HP-5MS 5% phenylmethylsiloxane capillary column (HP-5, 30m (length) × 0.25 mm (ID) × 0.25 μm (stationary phase thickness)). Oven temperature was kept at 55 °C for 4 min initially, and then raised at the rate of 4 °C/min to 250 °C. Injector temperature was set at 250 °C, respectively. Helium (99.999%) was used as carrier gas at a flow rate of 0.9 ml/min; samples were injected manually in the split mode. The peaks area percent was used for obtaining quantitative data. The gas chromatograph was coupled to an Agilent 6890N coupled to Agilent 5973 mass detector mass selective detector. Retention indices were calculated for all components using a homologous series of n-alkanes (C5–C24) injected in conditions equal to samples. Identification of oil components was accomplished based on comparison of their retention times with those of authentic standards and by comparison of their mass spectral fragmentation patterns (WILLEY/Chem Station data system)

The characteristics of the instrument were as follow:

Instrument: Gas chromatograph: Agilent 6890N coupled to Agilent 5973 mass detector

Column: HP-5, 30m (length) × 0.25 mm (ID) × 0.25 μm (stationary phase thickness)

Injector type: split/ splitless

### Column temperature program:

Rate(°C/min)	Temperature (°C)	Hold (min)
-	50	-
5	180	-
10	250	0.00

Carrier gas: He (99.999%); Injection type: splitless; Library: Wiley 7n; Injector temperature: 250°C; Flow rate: 0.9 ml/min

Extraction condition:

Extraction mode: Head space solid phase microextraction (HS-SPME)

SMPE fiber: PDMS 100 μm thickness (SUPELCO)

Sample weight: 0.5 gr, Extraction temperature: 60 °C, Extraction time: 15 min, Sonication time: 10 min (Euronada sonication instrument, Italy), Humidity (added water volume): 50  $\mu$ L, Desorption time in injector port of GC-MS: 3 min.

## Results

GC-MS results showed that the *Scrophulariadeserti* plant contains 51 compounds. Phytochemistry laboratory analyses results are shown in Table 2.

**Table 2. Gas chromatography/mass spectrometry (GC–MS) analysis of *Scrophulariadeserti***

No.	Compound	KI	%
1	$\alpha$ -Thujene	920	0.88
2	$\alpha$ -PINENE	931	24.69
3	Sabinene	966	5.81
4	$\beta$ -Myrcene	988	11.82
6	Tricyclene	997	0.94
7	$\beta$ -Phellandrene	1035	20.58
8	3-Carene	1046	0.37
9	$\gamma$ -Terpinene	1054	0.28
10	trans-Sabinene hydrate	1068	0.27
13	Alloocimene	1176	0.75
14	Camphor	1224	0.68
17	Fenchyl acetate	1285	2.25
20	Octadecane	1289	0.18
21	Borneol, acetate	1296	2.98
27	Bicycloelemene	1325	1.14
30	$\alpha$ -Copaene	1378	2.24
31	Calarene	1386	1.48
32	$\beta$ -BOURBONENE	1392	0.26
35	$\alpha$ -Gurjunene	1406	0.29
37	trans-Caryophyllene	1418	9.22
39	$\alpha$ -Bergamotene	1436	1.03
40	Isodene	1449	0.49
41	trans- $\beta$ -Farnesene	1468	3.14
43	$\alpha$ -amorphene	1474	0.40
44	$\beta$ -Cubebene	1485	2.38
46	bicyclogermacrene	1492	1.57
47	$\beta$ -Bisabolene	1512	0.61
48	Germacrene D	1526	1.15
49	$\delta$ -Cadinene	1538	1.88
51	Docosane	1568	0.25

## Discussion

The results of a phytochemical study showed that,  $\alpha$ -PINENE with 24.69% is the most active ingredient of *Scrophularia eserti* plant. Phytochemical analysis revealed that *Scrophularia eserti* extracts containing compounds 3 (zeta) -hydroxy-octadeca-4 (E), 6 (Z) -dienoic acid (1). The known compounds are ajugoside (2), scropolioside B (3), 6-O- $\alpha$ -L-rhamnopyranosylcatalpol (4), buddlejoside A (8) (5), scrospioid A (6), laterioside and 3R-1 -octan-3-yl-3-O-beta-D-glucopyranoside (7). *Scrophularia eserti* is a plant which in traditional medicine, local people experimentally use it in internal infections, mastitis, skin ulcers and episiotomy, inflammation, burns, intestinal pain, inflammation, eye and ear infections and hemorrhoids in the form of decoction by oral or topical uses (5,6). This plant has phenolic compounds and phenolic compounds have antimicrobial activities (8-13). More importantly these compounds have antioxidant activities which

possess various therapeutic effects, especially in diabetes (14-17), hyperlipidemia (18-20), renal toxicities (21-35), pain (36-39) and neurological diseases (40-45). Therefore, this plant might have these properties, too. There are many therapeutic effects of medicinal plants due to pharmaceutical active substances such as phenols, flavonoids, tannins, anthocyanin and etc (46-55).

## References

1. Mabberley DJ, editor. The Plant-Book. 2nd ed. Cambridge: Cambridge University Press; 1997.
2. Mozafarian VL. Flora of Ilam. 2008; 15(5): 902.
3. Azadbakht M. Classification of medical plants. Tehran: Teimorzadeh Pub. 2000; p: 7-276. Persian
4. Amin GH. Traditional Medicinal Plants of Iran. 1th ed. Tehran: Research assistance , Ministry of Health and Medical Education; 1991; 57.
5. Shohani F. People Journalism of Ivan. Ilam Cultural Heritageorg 2003; 56-7.
6. Bahmani M, Saki K, Rafieian-Kopaei M. Medicinal Plants of Thyme Land in Iran. LAP Lambert Academic Publishing: Germany. 2014; 1-3.
7. Stavri M, Mathew KT, Gibbons S. Antimicrobial constituents of *Scrophularia deserti*. Phytochemistry. 2006; 67(14): 1530-3.
8. Gupta A, Chaphalkar SR. Anti-inflammatory and anti-microbial activities of aqueous leaves extract of *Butea frondosa*. J Herbmed Pharmacol. 2016;5(2):85-88.
9. Rahimian GA, Rabiei Z, Tahmasebi B, Rafieian-Kopaei M, Ganji F, Rahimian R. Comparing the combined effect of garlic and mint extract with metronidazole in helicobacter pylori Treatment. Iranian Journal of Pharmaceutical Sciences. 2013;9(3):63-70.
10. Asadi-Samani M, Bahmani M, Rafieian-Kopaei M. The chemical composition, botanical characteristic and biological activities of *Borago officinalis*: a review. Asian Pac J Trop Med 2014; 7(Suppl 1): 22-28.
11. Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Saki K, Karamati SA, Delfan B. A review on most important herbal and synthetic antihelmintic drugs. Asian Pac J Trop Med 2014; 7(Suppl 1): 29-33.
12. Karamati SA, Hassanzadazar H, Bahmani M, Rafieian-Kopaei M. Herbal and chemical drugs effective on malaria. Asian Pac J Trop Dis 2014; 4(Suppl 2): 599-601.
13. Bahmani M, Karamati SA, Hassanzadazar H, Forouzan SH, Rafieian-Kopaei M, Kazemi-Ghoshchi B, Asadzadeh J, Kheiri AGh, Ehsan Bahmani E. Ethnobotanic study of medicinal plants in Urmia city: identification and traditional using of antiparasites plants. Asian Pac J Trop Dis 2014; 4(Suppl 2): 906-910.
14. Nasri H, Shirzad H, Baradaran A, Rafieian-Kopaei M. Antioxidant plants and diabetes mellitus. Journal of Research in Medical Sciences. 2015; 20:491-502.
15. Baharvand-Ahmadi B, Bahmani M, Tajeddini P, Naghdi N, Rafieian-Kopaei M. An ethno-medicinal study of medicinal plants used for the treatment of diabetes. J Nephropathol. 2016; 5(1):44-50.
16. Nasri H, Shirzad H, Baradaran A. Rafieian-kopaei M. Antioxidant plants and diabetes mellitus. J Res Med Sci 2015; 20:491-50.
17. Bahmani M, Zargarani A, Rafieian-Kopaei M, Saki M. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran. Asian Pac J Trop Med 2014; 7(Suppl 1): 348-354.
18. Bahmani M, Mirhoseini M, Shirzad H, Sedighi M, Shahinfard N, Rafieian-Kopaei M. A review on promising natural agents effective on hyperlipidemia. J Evid Based Complementary Altern Med. 2015 Jul;20(3):228-38. doi: 10.1177/2156587214568457. Epub 2015 Jan 28.
19. Mirhosseini M, Baradaran A, Rafieian-Kopaei M. *Anethum graveolens* and hyperlipidemia: A randomized clinical trial. J Res Med Sci 2014;19:758-61.
20. Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulago angulata* extract on serum lipids and lipid peroxidation. Evidence-Based Complementary and Alternative Medicine; 2014 (2014), Article ID 680856, 4 pages <http://dx.doi.org/10.1155/2014/680856>
21. Rafieian-Kopaei M, Hosseini M, Shirzad H. Comment on: Effect of pomegranate flower extract on cisplatin-induced nephrotoxicity in rats. Journal of Nephropathology. 2014;3(4):121-123.
22. Bahmani M, Shirzad H, Rafieian S, Rafieian-Kopaei M. *Silybum marianum*: Beyond Hepatoprotection. Journal of Evidence-Based Complementary and Alternative Medicine. 2015; 20(4):292-301

23. Nasri H, Tavakoli M, Ahmadi A, Baradaran A, Nematbakhsh M, Rafieian-Kopaei M. Ameliorative effect of melatonin against contrast media induced renal tubular cell injury. *Pak J Med Sci.* 2014 Mar;30(2):261-5.
24. Nasri, H., Tavakoli, M., Ahmadi, A., Baradaran, A., Nematbakhsh, M., Rafieian-Kopaei, M. Ameliorative effect of melatonin against contrast media induced renal tubular cell injury. *Pak J Med Sci.* 2014; 30(2): 261-265
25. Nasri H., Rafieian-Kopaei M. Tubular kidney protection by antioxidants. *Iranian J Publ Health.* 2013; 42(10): 1194-1196 IF=0.405
26. Rafieian-Kopaei M, Nasri H. Re: Erythropoietin ameliorates oxidative stress and tissue injury following renal ischemia/reperfusion in rat kidney and lung. *Med Princ Pract.* 2014;23(1):95. doi: 10.1159/000350842. Epub 2013 May 23.
27. Nasri H, Abedi-Gheshlaghi Z, Rafieian-Kopaei M. Curcumin and kidney protection; current findings and new concepts. *Acta Persica Pathophysiol.* 2016; 1(1):e01.
28. Khodadadi S, Rafieian-Kopaei M. Herbs, health and hazards; a nephrology viewpoint on current concepts and new trends. *Ann Res Antioxid.* 2016; 1(1):e05.
29. Rafieian-Kopaei M. Medicinal plants for renal injury prevention. *J Renal Inj Prev.* 2013 Jun 1; 2(2):63-5.
30. Rafieian-Kopaei M. Nasri H. Herbal antioxidant therapy in dialysis patients. *Ann Res Dial.* 2016; 1(1):e02.
31. Nasri H. Herbal drugs and new concepts on its use. *J Prev Epidemiol.* 2016; 1(1):e01.
32. Kafeshani M. Diet and immune system. *Immunopathol Persa.* 2015; 1(1):e04.
33. Mardani, S, Nasri H, Rafieian-Kopaei M, Hajian S. Herbal medicine and diabetic kidney disease. *J Nephropharmacol.* 2015; 2(1): 1-2.
34. Kafeshani M. Ginger, micro-inflammation and kidney disease. *J Renal Endocrinol.*2015; 1:e04.
35. Tamadon MR, Zahmatkesh M. World kidney day 2015. *J Parathy Dis* 2015; 3(2):34-36.
36. Dehghan Shahreza F. Renal tubular cell injury and its protection by antioxidants; new trends. *J Inj Inflamm.*2016; 1(1):e01.
37. Bahmani M, Shirzad HA, Majlesi M, Shahinfard N, Rafieian-Kopaei M. A review study on analgesic applications of Iranian medicinal plants. *Asian Pac J Trop Med* 2014; 7(Suppl 1):
38. Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M. Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran. *Asian Pac J Trop Med* 2014; 7(Suppl 1): 376-379.
39. Shirani M, Alibabaei Z, Kheiri S, Shirzad H, Taji F, Asgari A, Rafieian M. Effect of *Euphorbia helioscopia* extract on acute and chronic pain in mice *Journal of Babol University of Medical Sciences.* 2011; 13(4):14-18.
40. Shayganni E, Bahmani M, Asgary S, Rafieian-Kopaei M. 2015. Inflammaging and cardiovascular disease: management by medicinal plants. *Phytomedicine*,
41. <http://dx.doi.org/10.1016/j.phymed.2015.11.004>, pii: S0944-7113(15)00356-6.[Epub ahead of print].
42. Bahmani M, Sarrafchi A, Shirzad H, Rafieian-Kopaei M. Autism: Pathophysiology and promising herbal remedies. *Curr Pharm Des.* 2016; 22(3):277–285. DOI: 10.2174/1381612822666151112151529
43. Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative stress and Parkinson's disease: New hopes in treatment with herbal antioxidants. *Curr Pharm Des.* 2016; 22(2): 238 – 246. DOI: 10.2174/1381612822666151112151653
44. Saki K, Bahmani M, Rafieian-Kopaei M. The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review. *Asian Pac J Trop Med* 2014; 7(Suppl 1): 34-42.
45. Rahnama S, Rabiei Z, Alibabaei Z, Mokhtari S, Rafieian-kopaei M, Deris F. Anti-amnesic activity of *Citrus aurantium* flowers extract against scopolamine-induced memory impairments in rats. *Neurological Sciences.* 2015 Apr;36(4):553-60. doi: 10.1007/s10072-014-1991-2. IF=2.4
46. Rabiei Z, Rafieian-kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of meynert in rat. *Neurochemical research.* 2014;39(2):353-60
47. Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M. The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception. *Biomedicine & Aging Pathology.* 2014;4(1):71-6.

48. Nikfarjam M, Bahmani M, Naimi A. Native medicinal plants of Iran effective on Memory and Learning: A Review. International Journal of PharmTech Research 2016; Vol.9, No.5, pp 466-473.
49. Asadi-Samani M, Moradi MT, Bahmani M, Shahrani M. Antiviral medicinal plants of Iran: A Review of Ethnobotanical evidence. International Journal of PharmTech Research 2016; Vol.9, No.5, pp 427-434.
50. Moradi MT, Asadi-Samani M, Bahmani M. Hypotensive medicinal plants according to Ethnobotanical evidence of Iran: A Systematic Review. International Journal of PharmTech Research 2016. Vol.9, No.5, pp 416-426.
51. Moradi MT, Asadi-Samani M, Bahmani M, Shahrani M. Medicinal plants used for liver disorders based on the Ethnobotanical documents of Iran: A Review. International Journal of PharmTech Research 2016; Vol.9, No.5, pp 407-415.
52. Bahaa El-Din Mekki, Hebat-Allah Hussien, Hanaa Salem(2016). Role of Glutathione, Ascorbic Acid and  $\alpha$ -Tocopherol in Alleviation of Drought Stress in Cotton Plants, International Journal of ChemTech Research, Vol.8, No.4, pp 1573-1581.
53. Helmina Br. Sembiring, TonelBarus, LamekMarpaung, and Partomuan Simanjuntak (2015), International Journal of PharmTech Research, Vol.8, No.9, pp 24-30.
54. Kartini Zailanie, Hartati Kartikaningsih, Umi Kalsum, Yushinta Aristina Sanjaya(2015), Fucoxanthin Effects of Pure Sargassum filipendula Extract Toward HeLa Cell Damage, International Journal of PharmTech Research, Vol.8, No.3, pp 402-407.
55. Hashish Kh.I., Rawia A. Eid, Magda M. Kandil and Azza A.M. Mazher(2015), Study on Various Level of Salinity on Some Morphological and Chemical composition of gladiolus Plants by Foliar Spray with Glutathione and Thiamine, International Journal of ChemTech Research, Vol.8, No.9, pp 334-341.

\*\*\*\*\*

**PAGE Number = 291**



**Extra page: not to be printed**

**International Journal of PharmTech Research**

**log on to - [www.sphinxesai.com](http://www.sphinxesai.com)**


**Indexed/Abstracted/  
Ranked by**

**Elsevier SCOPUS-  
scimagojr.**

Scimago  
Lab

Powered by  
**Scopus**



**International Journal of PharmTech Research** is an open access Bimonthly Journal, 9 Years old. It contains more than 3500 published papers since 2009.

**Subject areas:** This journal publishes the Research and Review papers of the following subject/areas. Pharmaceutics, Pharmaceutical Chemistry, Biopharma, Pharmacology, Pharmacy Practice, Pharmacognosy, Analytical Chemistry, Biotechnology, Microbiology, Biochemistry, , Medicinal Science, Clinical Pharmacy, Medichem, and applied related subject areas.

PAGE Number = 292

## Extra page: not to be printed

### [1] RANKING:

It has been ranked from India (subject: Pharma Sciences) from India at International platform, by SCOPUS- scimagojr.

It has topped in total number of CITES AND CITABLE DOCUMENTS.

Find more by clicking on SCOPUS-scimagojr SITE....AS BELOW....

[http://www.scimagojr.com/journalrank.php?area=3000&category=0&country=IN&year=2013&order=tc&min=0&min\\_type=tc](http://www.scimagojr.com/journalrank.php?area=3000&category=0&country=IN&year=2013&order=tc&min=0&min_type=tc)

Please log on to - [www.sphinxesai.com](http://www.sphinxesai.com)

### [2] Indexing and Abstracting.

International Journal of PharmTech Research is selected by -

CABI, CAS(USA), SCOPUS, MAPA (India), ISA(India),DOAJ(USA),Index Copernicus, Embase database, EVISA, DATA BASE(Europe), Birmingham Public Library, Birmingham, Alabama,Worldcat, RGATE Databases/organizations, Beardslee Library Journals, Holland.

UNIVERSITY LIBRARY OF University of SASKATCHEWAN, ResearchBible/Journal Seeker,

AYUSH India, [ersa.lib.sjtu.edu.cn](http://ersa.lib.sjtu.edu.cn), many libraries for Indexing and Abstracting.

It is also in process for inclusion in various other databases/libraries.

### [3] Editorial across the world.

### [4] Authors across the world:

### [5] It has good SJR [SCImago Journal Rank] =

<http://www.scimagojr.com/journalsearch.php?q=19700175060&tip=sid&clean=0>

Please log on to - [www.sphinxesai.com](http://www.sphinxesai.com)

Search for the best References and submit your papers for publication.

[www.sphinxesai.com](http://www.sphinxesai.com)

\*\*\*\*\*