See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/286922214

Ruta graveolens plant: A plant with a range of high therapeutic effect called cardiac plant

Article in Der Pharmacia Lettre · December 2015		
CITATIONS		READS
0		186
1 author:		
	Mahmoud Bahmani	
	213 PUBLICATIONS 1,621 CITATIONS	
	SEE PROFILE	

Some of the authors of this publication are also working on these related projects:



The effect of metoclopramide addition to lidocaine on pain of patients with grades II and III postepisiotomy repair View project

All content following this page was uploaded by Mahmoud Bahmani on 15 December 2015.

Available online at www.scholarsresearchlibrary.com



Scholars Research Library

Der Pharmacia Lettre, 2015, 7 (11):172-173 (http://scholarsresearchlibrary.com/archive.html)



Ruta graveolens plant: A plant with a range of high therapeutic effect called cardiac plant

 1 Babak Baharvand-Ahmadi, 2 Mahmoud Bahmani, 3 Arman Zargaran, 4 Zohreh Eftekhari, 5 Kourosh Saki, 6 Sara Baharvand-Ahmadi and 7* Mahmoud Rafieian-Kopaei

¹Lorestan University of Medical Sciences, Khorramabad, Iran
²Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran
³Pharmaceutical Sciences Research Center and Department of Phytopharmaceuticals (Traditional Pharmacy),
School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran
⁴Pasteur Institute of Iran, Iran
⁵Department of Pharmacology, Debrescen University, Hungary
⁶Shahid Beheshti University of Medical Sciences, Tehran, Iran
⁷Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

INTRODUCTION

Belonging to the family Rutaceae, Rue or *Ruta graveolens* L. (RG) is an evergreen shrub herb and is distributed all around the world. From the ancient times, the herb is applied for various types of disorders [1]. In Iran, RG is known as *Pim* or *Sodab* and generally grows in the northern parts as Gilan province and Rasht city [2]. Concurrent with other traditional systems of medicine, RG has been administered for numerous types of diseases in Traditional Persian Medicine (TPM). Avicenna, the outstanding Persian physician, believed that RG have antispasmodic, anti-inflammatory, analgesic and antihelmintic effects and also exhibit aphrodisiac activity. In addition, this natural medicament has been administered in gastrointestinal, pulmonary and genecology disorders [3]. Later Persian scholars have recommended the herb as a vasodilator, antiflatulent, diuretic and a herb to treat amenorrhea and dysmenorrhea. Moreover, leaves of RG have been applied as an appetizer, a medicament for skin disorders, sciatalgia, chill and fever, kidney stones, toxins and poisons, neurological disorders and also a flavor to control the taste and odor of syrup dosage forms [4].

The herb is also widely used in other medical systems. Chinese and Indian traditional physicians applied the herb for syncope, neuralgia, rheumatalgia and ascites. However the application has been contraindicated in abortions and pregnancies [5]. Traditional healers in some Latin America countries such as Uruguay and Mexico applied the herb for abortion and GI disorders and cardiovascular system [6-7]. The abortive effect, menstruation function and management of dyspepsia in veterinary medicines are also other applications of RG in some countries or states such as Trinidad, Tobago and British Colombia of Canada [8]. Furthermore, African traditional practitioners have used RG dry leaves for urinary ailments, diabetes, headache, and GI disorders and cardiovascular system [9]. British and Denmark traditional medicines report the antirheumatic, tranquilizing and memory enhancing effects and cardiovascular disorders of the herb [10-11].

With reference to current knowledge, RG showed antitumor activity against Dalton's lymphoma ascites (DLA), Ehrlich ascites carcinoma (EAC) and L929 cells (IC100 = 16 mg/ml) which may be related to prooxidant and hydroxyl radicals scavenging activities of the extract [12]. The extract has also evaluated in some neurological disorders such as multiple sclerosis which was found effective [13]. Additionally, the genus, *Ruta* has shown

hypnotic, anticonvulsant and anti-anxiety properties [14] and also could inhibit and destroy the growth of cancer cells [15]. The herb also possessed analgesic, anti-inflammatory and moderate antimicrobial activities [16-17]. In the field of cardiology, RG revealed to have hypotensive activity as well as inotropic and chronotropic effects in animal models [18]. The antifertility effects of RG roots, leaves and stem were assessed intragastrically and resulted in the respective activity [19]. In an animal study and compared to the control group, RG aqueous extract reduced the count of spermatocytes and spermatids at a dose of 500 mg/kg for 60 days [20].

Different parts of RG plant contain numerous secondary metabolites such as glycosides, quinoline alkaloids, coumarins, lignans, flavonoids, furanocoumarins, hydroxycoumarins, and hydrocarbons [21-23].

Despite all these pharmacological and clinical properties, RG may cause photosensitization and photodermatosis due to furanocoumarins as well as gastric inflammation, hypothermia and metritis [2]. The herb also is forbidden in pregnancy due to epigastric pain, vomiting, liver and kidney damage, tremor and spasm [24]. Medicinal herbs are always a source to treat diseases such as cardiovascular, nervous and ect[25-27].

REFERENCES

- [1] R Kannan, UV Babu. Anc Sci Life. 2012; 32(1): 16-19.
- [2] A Zargari. Tehran University Press. 1996: 464 7.
- [3] A Avicenna. Soroush press. 1997: 249 51.
- [4] MH Aghili Khorasani Shirazi. Tehran University of Medical Sciences. Sahba Press. 1992: 450-451.
- [5] H Mirheydar. Islamic Cultuer Publishing office. 1992: 203-207.
- [6] C Ciganda, A Laborde. J Toxicol Clin Toxicol. 2003; 41(3): 235-239.
- [7] F Calzada, L Yepez-Mulia, A Aguilar. J Ethnopharmacol. 2006; 108(3): 367-370.
- [8] Lans C, Turner N, Khan T, Brauer G. Vet Parasitol. 2007; 148(3-4): 325-340.
- [9] TS Thring, FM Weitz. J Ethnopharmacol. 2006; 103(2): 261-275.
- [10] A Waldstein. J Ethnopharmacol. 2006; 108(2): 299-310.
- [11] A Adsersen, B Gauguin, L Gudiksen, AK Jager. J Ethnopharmacol. 2006; 104(3): 418-422.
- [12] KC Preethi, G Kuttan, R Kuttan. Asian Pac J Cancer Prev. 2006; 7(3): 439-443.
- [13] EW Bethge, KH Bohuslavizki, W Hansel, A Kneip, E Koppenhofer. Gen Physiol Biophys. 1991; 10(3): 225-244.
- [14] ME Gonzalez-Trujano, D Carrera, R Ventura-Martinez, E Cedillo-Portugal, A Navarrete. *J Ethnopharmacol*. **2006**: 106(1): 129-135.
- [15] S Pathak, AS Multani, P Banerji. Int J Oncol. 2003; 23(4): 975-982.
- [16] NS Alzoreky, K Nakahara. Int J Food Microbiol. 2003; 80(3): 223-230.
- [17] AH Atta, A Alkofahi. J Ethnopharmacol. 1998; 60(2): 117-124.
- [18] KW Chiu, AY Fung. Gen Pharmacol. 1997; 29(5): 859-862.
- [19] YC Kong, CP Lau, KH Wat, KH Ng, PP But, KF Cheng, PG Waterman. Planta Med. 1989; 55(2): 176-178.
- [20] NA Khouri, Z El-Akawi. Neuro Endocrinol Lett. 2005; 26(6): 823-829.
- [21] EE Stashenko, R Acosta, JR Martinez. J Biochem Biophys Methods. 2000; 43(1-3): 379-390.
- [22] RF Weiss, V Fintelman. Thieme. 2000: 181-330.
- [23] S Milesi, B Massot, E Gontier, F Bourgaud, A Guckert. Plant Science. 2001; 161(1): 189-199.
- [24] Staff TP. PDR for Herbal Medicines: Thomson PDR. 2004.
- [25] B Baharvand, M Esmailidehaj, J Alihosaini, SH Bajoovand, S Esmailidehaj. *Iranian Biomedical Journal* **2016**; 20(1): 41-48.
- [26] M Bahmani, A Sarrafchi, H Shirzad, M Rafieian-Kopaei. Curr Pharm Des. 2015 Nov 12.
- [27] A Sarrafchi, M Bahmani, H Shirzad, M Rafieian-Kopaei. Curr Pharm Des. 2015 Nov 12.