

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd

Document heading

doi: 10.1016/S2222-1808(14)60449-7

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

In vivo evaluation of antiparasitic effects of *Artemisia abrotanum* and *Salvia officinalis* extracts on *Syphacia obvelata*, *Aspiculoris tetrapetra* and *Hymenolepis nana* parasites

Mahdi Amirmohammadi¹, Shole Khajoenia², Mahmoud Bahmani^{3*}, Mahmoud Rafeian-Kopaei⁴, Zohre Eftekhari⁵, Mostafa Qorbani^{6,7}

¹Jiroft Branch, Islamic Azad University, Jiroft, Iran

²Department of Clinical Science, Faculty of Medicine, Medical Science University Jiroft, Iran

³Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran

⁴Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁵Veterinary Medicine Faculty, Tehran University, Tehran, Iran

⁶Alborz University of Medical Sciences, Karaj, Iran

⁷Department of Epidemiology, Tehran University of Medical Sciences, Tehran, Iran

PEER REVIEW

Peer reviewer

Dr. Hassan hassanzadazar, PhD of food hygiene, food control expert of food and drug deputy, Urmia university of medical science, Urmia, Iran.

Tel: +989144412026

Fax: +984413455839

E-mail: h.hassanzadazar@yahoo.com

Comments

This is a good research in which the authors determined effects of two local plant extracts on three various parasites in mice. The results are interesting that *S. officinalis* and *A. abrotanum* extracts had good antiparasitic effects on three various parasite.

Details on Page S253

ABSTRACT

Objective: To evaluate the effects of *Salvia officinalis* and *Artemisia abrotanum* extracts against digestive system parasites of mice.

Methods: The ethanol extract was prepared and dissolved in distilled water. The mebendazole was used as positive control and distilled water as negative control. After counting eggs per gram feces, infected mice with 16 eggs per gram feces contained two to three parasites of *Syphacia obvelata*, *Aspiculoris tereptra* and *Hymenolipis nana* designated in 4 groups. The first group was given extracts of *Artemisia* (150 mg/kg), the second group was given *Salvia* extract (150 mg/kg), the third group was given mebendazole (10 mg/kg) and finally the fourth group was given distilled water (2 mL/kg).

Results: The ethanol extracts of *Artemisia* and *Salvia* plants reduced the number of parasite eggs per gram of feces. Results showed significant reduction (P -value<0.001) in the number of eggs excreted by *Hymenolepis nana*, *Aspiculoris tetrapetra*, *Syphacia obvelata* in mice.

Conclusions: These results revealed that antiparasitic effects of *Artemisia* and *Salvia* are reasonable and these two plants might be used as antiparasitic natural products.

KEYWORDS

Salvia officinalis, *Artemisia abrotanum*, Mice, Parasite, *Syphacia obvelata*, *Aspiculoris tereptra*, *Hymenolipis nana*

1. Introduction

Parasitic infections cause major public health problems leading to morbidity and sometimes mortality of the

victims. Chemical drug against parasites is beneficial, but this method has some problems such as drug resistance, residue and unwanted side effects associated. Therefore, study on alternative therapies are important^[1]. For

*Corresponding author: Mahmoud Bahmani, Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran.

Tel: 984412772023

Fax: 984412787814

E-mail: mahmood.bahmani@gmail.com

Foundation Project: Supported by Jiroft Branch, Islamic Azad University of Jiroft, Iran (Grant No. 1713864).

Article history:

Received 28 Nov 2013

Received in revised form 5 Dec, 2nd revised form 11 Dec, 3rd revised form 16 Dec 2013

Accepted 20 Jan 2014

Available online 28 Jan 2014

centuries, medicinal plants have been used to eliminate parasites in many parts of the world, and humans used the plant to treat various diseases. Herbal medicines are used in recent years due to the high reliability and effectiveness, and cost is also taken into consideration[2,3]. In ethnoveterinary medicine, which draws inspiration from traditional practice, there seems to be various plants suitable for treating almost every parasitic disorder of livestock. Although the majority of the evidence on antiparasitic activity of plants has been customarily based on anecdotal observations, there are currently an increasing number of controlled experimental studies aimed to verify, validate and quantify such plant activity in a scientific manner. Currently, there are 119 drugs composed of plants, and only 90 of the 250 000 known plant species, has been obtained[2,3]. In Chaharmahal va Bakhtiari and Kerman provinces in Iran, *Artemisia abrotanum* (*A. abrotanum*) and *Salvia officinalis* (*S. officinalis*) are used as medicinal plants against parasites.

Cultivars of *S. officinalis* are quite variable in size, leaf and flower color and foliage pattern with many variegated leaf types. The Old World type grows approximately 0.61 m tall and wide with lavender flowers, the most common color. And the flower can also be white, pink, or purple. The plant flowers bloom in late spring or summer. The leaves are oblong, ranging in size up to 6.4 cm long by 2.5 cm wide. Leaves are grey–green, rugose on the upper side, and nearly white underneath due to many short soft hairs. Modern cultivars include leaves with purple, rose, cream, and yellow in many variegated combinations. Sage leaf contains tannic acid, oleic acid, ursolic acid, ursolic acid, cornsole, cornsolic acid, fumaric acid, chlorogenic acid, caffeic acid, niacin, nicotinamide, flavones, flavonoid glycosides, and estrogenic substance[3]. Its leaves contain essential oils and tannins that can help to facilitate digestion with anticonvulsant, anti–fever, antiseptic and anti–diabetic effect. Modern evidence shows possible uses as an anti–sweating, antibiotic, antifungal, astringent, antispasmodic, estrogenic, hypoglycemic and tonic agent. This plant is also used for the treatment of gout, chronic rheumatism, dizziness and headache. Investigations have be performed to use sage as a medication for Alzheimer’s disease.

Wormwood is used to treat liver, spleen and stomach disorders. It is seldom used for medicinal purpose today, except in Germany, where poultices are placed on wounds, splinters and skin conditions and it is employed occasionally to treat frostbite[4].

In the present study, the antiparasitic effects of *Artemisia* and *S. officinalis* plants were investigated on the most common gastrointestinal tract parasites such as *Hymenolepis nana* (*H. nana*), *Aspicularis tetraoptera* (*A. tetraoptera*) and *Syphacia obvelata* (*S. obvelata*).

2. Materials and methods

The methanolic extracts of *A. abrotanum* and *S. officinalis* were prepared by adding a 1:3 ratio of garlic and methanol, respectively, and this was subjected to Soxhlet extraction for 72 h according to the prescribed method of Eidi *et al*[5]. After extraction, the solvent was filtered[5]. The information about the studied plants is presented in Table 1.

Table 1

Scientific name, English name, part used, extract type, and traditional treatment of *A. abrotanum* and *S. officinalis*.

Scientific name	English name	Part used	Extract	Traditional use[6]
<i>A. abrotanum</i>	Wormwood	Leaf	Ethanol extract	Antiparasite
<i>S. officinalis</i>	Garden sage	Leaf	Ethanol extract	Antiparasite

S. officinalis (garden sage, common sage) is a small, perennial, evergreen subshrub, with woody stems, grayish leaves, and blue to purplish flowers. It is a family member of Lamiaceae and is native to the Mediterranean region, and it has been naturalized in many places throughout the world. It has a long history of medicinal and culinary use, and in modern times as an ornamental garden plant.

Wormwood or southernwood (*A. abrotanum*) is a flowering plant. Southernwood is antiseptic and kills intestinal worms.

First, a few infected mice were randomly divided into four groups. The mice were treated with doses of 20, 50, 100 and 150 mg/kg body weight of ethanol extracts or normal saline (2 mL/kg), orally. All animals were tested during 48 h for symptoms of poisoning and death. After that, 16 mice with equal weight, nutritional conditions, infected with *H. nana*, *A. tetraoptera* and *S. obvelata* were tested for four groups that each group consisted of four mice. The Lynn Clayton test was employed for detecting parasite eggs per gram feces. These groups were treated as following:

Group 1: treated with 150 mg/kg *Salvia* extract for 5 d.

Group 2: treated with 150 mg/kg *Artemisia* extract for 5 d.

Group 3: treated with 10 mg/kg mebendazole for 5 d as positive control .

Group 4: treated with 2 mL/kg normal saline as negative control.

The parasite eggs were counted on the first, second, third and fifth day. On the seventh day, one mouse from each group was killed and the intestine was checked regarding the presence of parasites. The differences between the control and treated groups were analyzed using M–stat program.

3. Results

In the present study, the ethanol extracts of wormwood, and garden sage, and mebendazole was examined for the effect on digestive tract parasites of mice. The results

showed that in all groups which received drug or herbal extracts, but not in control group, the number of eggs excreted in per gram of feces was reduced one day after treatment. Results showed significant differences (P -value<0.001) in the number of eggs excreted by the *H. nana*, *A. tetraptera* and *S. obvelata* in mice, among the groups which had used the wormwood or garden sage and control group.

This study showed a noticeable effect for wormwood and garden sage over time against parasites. So that on the seventh day, the number of parasite eggs in feces was at the lowest amount (P <0.001). The garden sage and mebendazole revealed better effects on *H. nana* and *A. tetraptera* in comparison to Wormwood. Wormwood, however, had better effect on *S. obvelata* in comparison to garden sage. Average number of eggs in per gram of feces excreted by *S. obvelata*, *A. tetrapetra* and *H. nana* on Day 0, 1, 3, 5 and 7 after treatment in four groups is specified in Figure 1, 2 and 3. The mean and standard deviation of the different treatment groups is shown in Figure 4.




Figure 1. Average number of eggs in per gram of feces excreted by *H. nana* on Day 0, 1, 3, 5 and 7 after treatment in four groups.




Figure 2. Average number of eggs in per gram of feces excreted by *A. tetrapetra* on Day 0, 1, 3, 5 and 7 after treatment in four groups.




Figure 3. Average number of eggs in per gram of feces excreted by *S. obvelata* on Day 0, 1, 3, 5 and 7 after treatment in four groups.




Figure 4. The mean and standard deviation of the different treatment groups.

4. Discussion

In the present study, the ethanol extracts of wormwood, garden sage and mebendazole for treatment of the digestive tract parasites of mice was examined. The results showed significant differences (P -value<0.001) in the number of eggs excreted by the *H. nana*, *A. tetrapetra* and *S. obvelata* in mice among the groups which had used the wormwood or garden sage and control group. This study found a noticeable effect of wormwood and garden sage over time, so that the number of parasite eggs in the feces was at the lowest amount on the seventh day.

In Iranian traditional medicine, one species of *Artemisia*, namely *Artemisia aucheri* Bioss, is known as antiparasitic plant. Ghasemi pirbalouti showed that *Artemisia salina* had antiparasite and anti-pesticide effect, *Artemisia inculata* had anti-parasite and laxative effects, and *Saliva verbeneca* L. had good effect for skin disorders[6].

The essence of *S. officinalis* can help to repair cell, stimulate, and strengthen the skin and hair, and decrease depression. Sharifi *et al.* revealed that the extraction of

Artemisia aucheri Bioss has an antiparasitic effect on the *Leishmania major*[7].

Mahrami Poor and Neghaban in 2007 showed that essential oils of *Artemisia scoparia* and *Artemisia sieberi* dependently reduced spawning, hatching eggs, and mortality of first instars larvae of the beetle (parasite of grains)[8]. Jalali *et al.* revealed the insecticide effect of the *Artemisia annua* L[9].

S. officinalis essential oil contains compounds such as thujine, cis humulen- α , 1,8-cineole, (E)-Caryophyllene, borneole, monoterpenes, diterpenes and diterpenes. The artemisinin is one of the most important components of *Artemisia*[9].

Results of study by Mirza showed that 32 compound has been identified on sage. Among the compounds identified, using gas chromatograph coupled with mass spectrometer, β -caryophyllene (16.3%), sclareol (13.3%), hexyl octanoate (12.2%) and bicyclogermacrene (10%) had the highest share, respectively[10].

The results of Bahmani *et al.* showed that high doses of the plant extract on antiparasitic effect against leeches (*Limnatis nilotica*) was found[11]. Santonin is antiparasitic herb drug that is produced from the *Artemisia* plant and its active ingredient is artemisinin[6]. Perhaps the powerful antiparasitic effects of garden sage and wormwood plants are due to these compounds.

The antiparasitic effect of *Artemisia* on *Eimeria* has been previously shown and it has been introduced as an anti-coccidian drug[12–14]. Bruce had demonstrated the effects of *Artemisia* extract on malarial protozoa[13].

Esfandiari *et al.* has shown effect of different concentrations of *Artemisia* extract on *Syphacia nematode*[15]. Khosraviead and Allen *et al.* showed that the artemisinin, a component of *Artemisia* (different species) has appropriate antiparasite effect on Coccidia[16,17].

Artemisia absinthium Linn. is used traditionally by people as a vermifuge in addition to its other livestock uses. The results of the study of Tariq *et al.* showed that *Artemisia absinthium* extracts are a promising alternative to the commercially available anthelmintics for the treatment of GI nematodes of sheep[18]. Study of Ezzat-zadeh *et al.* showed that the plant *Artemisia kulbadica* had antimicrobials and antioxidants effects[19]. Studies of Arokiyaraj showed that *Artemisia nilagirica* showed good antibacterial effects[20]. In studies of Suresh it was found that phytochemical contents of *Artemisia parviflora* Roxb were alkaloids, sterols/terpenoids, flavonoids, tannins, phenols and coumarins[21]. Results of two studies showed

that traditional effects of antiparasitic on intestinal worms are introduced for *Artemisia herba-alba*[22,23].

This study and other studies revealed appropriate antiparasite effect of *Artemisia* and *S. officinalis* which indicate that they might be used instead of chemical drugs in parasites control programs.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This work was supported by Jiroft Branch, Islamic Azad University of Jiroft, Iran. Grant number of this study is 1713864.

Comments

Background

Parasitic infections in animals and humans are one of the medical problems. In recent decades the use of local medicinal herbs and non-chemical remedies instead of chemical drugs to treat diseases has been regarded. Herbal plants in different areas based on species and amount of active ingredient have different therapeutic effects.

Research frontiers

Studies are performed in order to determine effects of two local plant extracts on three various parasites in mice. Certain dose of two plant extracts were fed to ternary test groups of infected mice.

Related reports

The data about the effects of two plant extracts on three parasites are different. This difference is probably due to extracts' main ingredients and drug resistance of parasites that cause various results during one week consuming of plant extracts.

Innovations & breakthroughs

Data regarding effects of different plant extracts on various parasites are scarce. This study showed that *S. officinalis* and *A. abrotanum* extracts can be use for

progressive study on human and domestic animals.

Applications

The results of this study showed that *S. officinalis* and *A. abrotanum* extracts had good antiparasitic effects on three various parasite, one of which is common in human. So we can test those extracts on other parasites in human and domestic animals.

Peer review

This is a good research in which the authors determined effects of two local plant extracts on three various parasites in mice. The results are interesting, and *S. officinalis* and *A. abrotanum* extracts had good antiparasitic effects on three various parasite.

References

- [1] Tyler YE, Brady LR, Robbers JE. *Pharmacognosy*. 9th ed. Philadelphia, USA: Lea & Febiger; 1988, p. 492.
- [2] Fabrican DS, Farnsworth NR. The value of plants used in traditional medicine for drug discovery. *Environ Health Perspect* 2001; **109**(Suppl 1): 69–75.
- [3] Saleem TS, Chetty SM, Ramkanth S, Rajan VS, Kumar MK, Gauthaman K. Hepatoprotective herbs—a review. *Int J Res Pharm Sci* 2010; **1**: 1–5.
- [4] Grando SA, Kist DA, Qi M, Dahl MV. Human keratinocytes synthesize, secrete and degrade acetylcholine. *J Invest Dermatol* 1993; **101**: 32–36.
- [5] Eidi A, Eidi M, Esmaeili E. Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin-induced diabetic rats. *Phytomedicine* 2006; **13**: 624–629.
- [6] Ghasemi-pirbalouti A. *Third listen: plants, traditional medicine and ethnoveterinary, medicinal and aromatic plant*. 1. Shahrekord: Saman–Danesh; 2008, p. 158–190.
- [7] Sharif M, Ziaei H, Azadbakht M, Daryani A, Ebadattalab A, Rostami M. Effects of methanolic extracts of *Artemisia aucheri* and *Camellia sinensis* on *Leishmania major* (in vitro). *Turk J Med Sci* 2006; **36**(6): 365–369.
- [8] Negahban N, Moharrampour S. Efficiency of *Artemisia sieberi* Besser and *Artemisia scoparia* waldst et kit essential oils on biological activity of *Callosobruchus maculatus* F. (Col.: Bruchidae). *Iran J Med Aromat Plants* 2007; **23**(2): 146–156.
- [9] Jalali A, Arbab A, Aliakbar A. The efficacy of aqueous plant extracts of wormwood and dwarf elder against elm leaf beetle *Xanthogaleruca luteola* Mull. (Coleoptera: Chrysomelidae). *Agric Sci* 2005; **15**(1): 115–120.
- [10] Mirza M, Ahmadi L. Identification of essential oil and extraction of *Salvia sclareae*. *Iran J Med Aromat Plants Res* 1999; **216**: 115–136.
- [11] Bahmani M, Avijgan M, Gholami–Ahangaran M, Rafieian M. The comparison of the anti *Limnatis nilotica* effects of albendazole and some of the Iranian medicinal plants. *Iran South Med J* 2012; **15**: 25–34.
- [12] Arab HA, Rahbari S, Rassouli A, Moslemi MH, Khosravirad F. Determination of artemisinin in *Artemisia sieberi* and anticoccidial effects of the plant extract in broiler chickens. *Trop Anim Health Prod* 2006; **38**(6): 497–503.
- [13] Bruce–Chwatt LJ. Qinghaosu: a new antimalarial. *Br Med J* 1982; **284**: 767–768.
- [14] Youn HJ, Noh JW. Screening of the anticoccidial effects of herb extract against *Eimeria tenella*. *Vet parasitol* 2001; **96**(4): 257–263.
- [15] Esfandiari B, Youssefi M, Keighobadi M, Nahrevanian H. *In vivo* evaluation of anti-parasitic effects of *Artemisia absinthium* extracts on *Syphacia parasite*. *Int J Parasitic Dis* 2007; **2**(2): 1–5.
- [16] Allen PC, Lyndon J, Danforth HD. Effects of components of *Artemisia annua* on coccidia infections in chickens. *Poult Sci* 1997; **76**: 1156–1163.
- [17] Khosravi Rad. Effect of of *Artemisia* plant on poultry eimeria[D]. Tehran: Tehran University; 2001.
- [18] Tariqa KA, Chishtia MZ, Ahmada F, Shawl AS. Anthelmintic activity of extracts of *Artemisia absinthium* against ovine nematodes. *Vet Parasitol* 2009; **160**(1–2): 83–88.
- [19] Ezzatzadeh E, Farjam MH, Rustaiyan A. Comparative evaluation of antioxidant and antimicrobial activity of crude extract and secondary metabolites isolated from *Artemisia kulbadica*. *Asian Pac J Trop Dis* 2012; **2**: S431–S434.
- [20] Arokiyaraj S, Sripriyaa N, Bhagyaa R, Radhikaa B, Prameelaa L, Udayakash NK. Phytochemical screening, antibacterial and free radical scavenging effects of *Artemisia nilagirica*, *Mimosa pudica* and *Clerodendrum siphonanthus*—an *in-vitro* study. *Asian Pac J Trop Biomed* 2012; **2**(2): S601–S604.
- [21] Suresh J, Ahuja J, Paramakrishnan N. Pharmacognostical investigation of *Artemisia parviflora* Roxb. *Asian Pac J Trop Biomed* 2012; **2**(2): S532–S535.
- [22] Bahmani M, Rafieian–Kopaei M, Avijgan M, Hosseini S, Golshahi H, Eftekhari Z, et al. Ethnobotanical studies of medicinal plants used by Kurdish owner’s in south range of Ilam Province, west of Iran. *Am Eurasian J Agric Environ Sci* 2012; **12**(9): 1128–1133.
- [23] Bahmani M, Eftekhari Z. An ethnoveterinary study of medicinal plants in treatment of diseases and syndromes of herd dog in southern regions of Ilam province, Iran. *Comp Clin Pathol* 2013; **22**: 403–407.