



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Medicine

journal homepage: www.elsevier.com/locate/apjtm



Document heading doi: 10.1016/S1995-7645(14)60257-1

## Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran

Mahmoud Bahmani<sup>1</sup>, Arman Zargaran<sup>2</sup>, Mahmoud Rafieian-Kopaei<sup>3\*</sup>, Kourosh Saki<sup>4</sup><sup>1</sup>Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran<sup>2</sup>Pharmaceutical Sciences Research Center and Department of Traditional Pharmacy, Faculty of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran<sup>3</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran<sup>4</sup>Shahid Beheshti University of Medical Sciences, Tehran, Iran

## ARTICLE INFO

## Article history:

Received 3 Jul 2014

Received in revised form 25 Aug 2014

Accepted 31 Aug 2014

Available online 29 Sep 2014

## Keywords:

Ethnomedicine

Diabetes

Urmia

Iran

## ABSTRACT

**Objective:** To collect and document information on anti-diabetic plants traditionally used in the treating of diabetes in Urmia at Northwest Iran because ethnomedicines are considered as valuable sources to find new potential drugs.

**Methods:** We used the method of direct observation and interview (35 traditional healers) along with gathering herbarium specimens mentioned plants in site.

**Results:** There were 30 medicinal plants from 17 families for the treatment of diabetes. The family with most plants was Lamiaceae (20%). Leaves (20%) are often used and its form is decoction (70%). It was also found that *Citrullus colocynthis* has the most frequency of use among traditional healers.

**Conclusions:** Furthermore, base on current findings many of the mentioned plants have potential active ingredients to influence diabetes.

## 1. Introduction

Diabetes mellitus is one of the most prevalent diseases in endocrine gland system with an increasing incidence in human community[1]. Type I diabetes is caused by insulin secretion deficit while type II diabetes is accompanied with progressive rate of insulin resistance in liver and peripheral tissues, reducing  $\beta$ -cells mass, and deficient insulin secretion[2,3]. This disease brings about acute metabolic side effects including ketoacidosis, hyperosmolar coma accompanied with chronic disorders and long term, adverse side effects such as retinopathy, renal failure, neuropathy, skin complications, as well as increasing cardiovascular complication risks[4,5]. Also, common symptoms of diabetes

are frequent urine, thirsty, and overeating[6].

Diabetes inflicts 100 million people yearly and is recognized as the seventh cause of death in the world[7]. It has been estimated that the number of diabetic people will increase from 150 million individuals in 2003 to 300 million by 2025[8].

The essential and effective drugs for diabetes mellitus are insulin injection and hypoglycemic agents, but these compounds possess several adverse effects and have no effects on diabetes complications in long term. Regarding the human increased knowledge about this disease and its complications, it is necessary to find effective compounds with lower side effects in treating diabetes[9]. Medicinal plants are good sources as alternative or complementary treatments for this and other diseases[10–12].

Although various plants have been traditionally used throughout history to reduce blood glucose and improve diabetes complications, there is not enough scientific information about some of them. Herbal medicines are

\*Corresponding author: Prof. Mahmoud Rafieian-Kopaei, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran.

Tel: 0098 381 3349509, 0098 913 1811842

Fax: 0098 381 3349509

E-mail: rafieian@yahoo.com

Foundation Project: Supported by the Deputy for Research and Technology of Lorestan University of Medical Sciences (Grant No. 0207).

commonly prescribed throughout the world because of low side effects, availability, roughly low cost, and also its effectiveness<sup>[13,14]</sup>.

In the present situation, herbal medicines' usage has significantly increased and published studies from developed countries emphasize that a paramount proportion of medicines supplied by them have herbal origins, so growing and producing the herbal medicines could be helpful to both economic development and community's health<sup>[15,16]</sup>. World Health Organization has announced that 80% of people around the world use herbal medicines at initial level of health remedy in the same way and they need to be evaluated scientifically for their efficacies<sup>[17]</sup>.

The plants as a major source of food supplements are effective in controlling blood glucose and preventing long term complications in type II diabetes<sup>[18]</sup>. The positive effects of many herbal medicines in reducing blood glucose and its complications have been already recognized<sup>[19–21]</sup>.

Iran as an old civilized country has a valuable traditional system of medicine called traditional Persian medicine and rooted in thousands years ago<sup>[22–25]</sup>. During early medieval age, Persian physicians such as Akhawayni, Rhazes, Avicenna, Haly Abbas, Jorjani, *etc.* have flourished medical sciences while some of their manuscripts like Canon of Medicine were added in main textbooks in eastern and western universities until 18th century<sup>[26–30]</sup>. Regarding, current ethno–medicine in Iran with such great background it is popular among people and is a potentially good source for finding new natural remedies. Twenty–three Persian physicians knew and diagnosed diabetes for centuries and had potentially good medicines (based on current knowledge) to manage and treat it<sup>[31]</sup>.

Urmia, a big city in Northwest Iran, is one of the places that ethno–medicine is popular and has good mountain weather to growth medicinal plants. In this study, we tried to obtain and analyze ethno pharmacological data of medicinal plants used to control and treatment of diabetes mellitus by traditional healers.

## 2. Methods

### 2.1. Study area

Forest landscape of Darre Ghasemloo and its adjacent regions with an area of 57 700 km<sup>2</sup> is located in southern part of Urmia City in Iran. It covers 45° 5' to 45° 10' E and 37° 15' to 37° 20' N geographical coordinates. It is located in 30 km right part of Urmia–Oshnaviyeh Road. This region is mountainous with the lowest altitude of 1420 m and the highest altitude of 2280 m<sup>[32]</sup>. With regard to the meteorological information

of region and annual humidity condition of soils, the soil moisture in this area under study were xeric and mesic, respectively<sup>[33]</sup>. According to climatic divisions, this region is cold and semi–arid. The mean annual rain fall is 367.5 mm, the highest and the lowest annual mean temperature is 33.1 °C and –15.5 °C, respectively. In this area, February and August are the coldest and the warmest months of year, respectively<sup>[34]</sup>.

### 2.2. Method of collecting data

This study was conducted via questionnaire and interview during November, 2013 to December, 2013 by traditional healers who are considered as unwritten knowledge.

Firstly, a full list of Urmia county herbalists (traditional healers) was obtained from Food and Medicine Assistance Department of Urmia University of Medical Sciences. Observation and interview accompanied (face to face) by gathering herbarium specimens of medicinal herbs and typical and effective impacts in treating diabetes were used. The questionnaires consisted of personal information and a list of native plants' names, the used organ parts, its usage, and traditional remedy effect.

There are 47 traditional healers in Urmia that only 35 of them cooperate with us. They announced that they referred to people with diabetes who had been diagnosed with a blood test lab.


The herbarium specimens obtained based on the native herbalists' information were collected from study area. The herbarium samples obtained from data of local traditional physicians in the questionnaire were collected from the region and then they were authenticated by a botanist (Mortaza Rafieian). A herbarium specimen from each plant (whole or the used part) was prepared and deposited in the herbarium unit of Shahrekord University of Medical Sciences. Finally, the data were analyzed using Excel 2010 software.

Prepared ethno–medicinal plants were listed and compared with current knowledge about them. Also, the frequency of ethno–medicinal use in Urmia was calculated via data mentioned by traditional healers (as a percentage among 35 healers). Finally, it is tried to present potential medicinal herbs for diabetes mellitus treatment via these data.

## 3. Results

Thirty medicinal plants from 17 families for diabetes treatment were identified. The families with most antidiabetic plants included Lamiaceae (6%), Fabaceae (4%),

and Rosaceae (4%) (Figure 1).




**Figure 1.** The families and percentage of plants with antidiabetic activity included.

Ast: Asteraceae; Bra: Brassicaceae; Poa: Poaceae; Ber: Berberidaceae; Ros: Rosaceae; Fab: Fabaceae; Equ: Equisetaceae; Jug: Juglandaceae; Lam: Lamiaceae; Cuc: Cucurbitaceae; Pap: Papaveraceae; Ann: Anacardiaceae; Rub: Rubiaceae; Pol: Polygonaceae; Urt: Urticaceae; Cru: Cruciferae; Lau: Lauraceae.

Leaves (20%) were frequently used and its form was decoction (70%). It was also found that *Citrullus colocynthis* (*C. colocynthis*) is frequently used by traditional healers.

The most used organ parts are shown in Figure 2. As it is illustrated in Figure 2, the most used organ parts in diabetes remedy in present study was 20% leaf, 16% flowering shoot, 13% fruit, 11% crust stem and 11% seed. Urmia traditional

healers prescribed 70% of antidiabetic herbal medicines in the decoction forms (Figure 3). The percentage of different used organ parts and of used type are indicated in Figures 2 and 3, respectively.



**Figure 2.** The percentage of different organ parts used.

The list and information of ethnobotanical remedies for diabetes treatment mentioned by Urmia, traditional healers are presented in Table 1. It shows that *C. colocynthis*, *U. dioica*, *L. album*, *Rosa foetida*, *Sanguisorba minor*, *Sophora alopecuroides*, *Trifolium pratense*, *Salvia nemorosa*, *Teucrium orientale*, *T. polium* were the most commonly used species for management of diabetes (frequency of use) by traditional healers.

**Table 1**

Ethnobotanical remedies for the treatment of diabetes.

Scientific name	Family	English name	C u r a t i v e effect	Biological form	Growth form	Used part	Frequency of use*	Way of using
<i>Achillea millefolium</i> L.	Asteraceae	Milfoil	Antidiabetic	He	Herbaceous perennial	Inflorescence	9 (25.71%)	Boiled, steamed
<i>Alyssum desertorum</i> Stapf.	Brassicaceae	Hedge mustards	Antidiabetic	Th	Herbaceous annuals	Seed	10 (28.57%)	Boiled, herbal fumigation
<i>Arctium lappa</i> L.	Asteraceae	Burdock	Antidiabetic	He	Two-year herbaceous	Root, leaf	8 (22.85%)	Boiled, steamed
<i>Avena sativa</i> L.	Poaceae	Oat	Blood refining	Th	Herbaceous annuals	Seed, glumelle	10 (28.57%)	Boiled
<i>Berberis integrifolia</i> Bunge.	Berberidaceae	Barberry	Antidiabetic	Ph	Shrub	Fruit, leaf, skin	9 (25.71%)	Boiled, steamed
<i>Cerasus microcarpa</i>	Rosaceae	Sour cherry	Blood refining	Ph	A bushes	Fruit	10 (28.57%)	Boiled, raw use
<i>Cinnamomum verum</i>	Lauraceae	Cinnamon	Antidiabetic	Th	Herbaceous perennial	Skin	8 (22.85%)	Boiled
<i>C. colocynthis</i> (L.) Schrad.	Cucurbitaceae	(Common caper-bush) Colocynthis	Antidiabetic		Herbaceous annuals	Fruit	28 (80%)	Boiled
<i>Coronilla varia</i> L.	Fabaceae	Garden alfalfa	Antidiabetic	He	Herbaceous perennial	Leaf	8 (22.85%)	Raw use, boiled
<i>Crataegus aronia</i> (L.) Bosc ex DC.	Rosaceae	Hawthorn	Antidiabetic	Ph	Shrub	Fruit and skin	9 (25.71%)	Raw use, boiled
<i>Crataegus oxyacantha</i> L.	Rosaceae	Hawthorn	Antidiabetic	Ph	Shrub	Fruit, flower root, skin	9 (25.71%)	Raw use, boiled
<i>Equisetum arvense</i> L.	Equisetaceae	Horse pipe	Antidiabetic	Cr	Herbaceous perennial	Aeration organ	9 (25.71%)	Boiled
<i>Juglans regia</i>	Juglandaceae	Walnut tree	Antidiabetic	Ph	Arboreal	Fruit, leaf and skin	9 (25.71%)	Boiled
<i>L. album</i> L.	Lamiaceae	White nettle	Antidiabetic	Th	Herbaceous annuals	Flowering offshoot	25 (71.42%)	Boiled
<i>Nasturtium officinale</i> (L.) R. Br.	Cruciferae	Water cress	Antidiabetic	He	Herbaceous perennial	Leaf, root	8 (22.85%)	Boiled
<i>Nepeta bracteata</i> Benth.	Lamiaceae	Horsemint	Antidiabetic	Th	Herbaceous annuals	Flowering offshoot	8 (22.85%)	Boiled, steamed
<i>Nepeta meyeri</i> Benth.	Lamiaceae	Horsemint	Antidiabetic	Th	Herbaceous annuals	Flowering offshoot	8 (22.85%)	Boiled, steamed
<i>Papaver rhoeas</i> L.	Papaveraceae	Copse	Antidiabetic	Th	Herbaceous annuals	Seed, capsule	8 (22.85%)	Boiled
<i>Polygonum aviculare</i> L.	Polygonaceae	Polygonum	Antidiabetic	Th	Herbaceous annuals	Aeration organ	8 (22.85%)	Boiled
<i>Rhus coriaria</i> L.	Anacardiaceae	Sumach	Blood refining	Ph	Arboreal	Fruit, leaf, resin	8 (22.85%)	Boiled
<i>Rosa foetida</i> Heram.	Rubiaceae	Yellow eglantine	Antidiabetic	Ph	A bushes	Petal	25 (71.42%)	Boiled
<i>Rumex sculantus</i> L.	Polygonaceae	Sorrel	Blood refining	He	Herbaceous perennial	Fruit, leaf	9 (25.71%)	Raw use, boiled
<i>Salvia nemorosa</i> L.	Lamiaceae	Farm sage	Antidiabetic	He	Herbaceous perennial	Flowering offshoot	22 (62.85%)	Boiled
<i>Sanguisorba minor</i> Scop.	Rosaceae	Poterium	Antidiabetic	He	Herbaceous perennial	Fruit, leaf	24 (68.57%)	Raw use, boiled
<i>Sophora alopecuroides</i>	Fabaceae	Pagoda tree	Antidiabetic	He	Herbaceous perennial	Inflorescence	24 (68.57%)	Boiled
<i>Teucrium orientale</i> L.	Lamiaceae	Germander	Antidiabetic	He	Herbaceous perennial	Leaf	22 (62.85%)	Boiled
<i>T. polium</i> L.	Lamiaceae	Germander	Antidiabetic	He	Herbaceous perennial	Flowering offshoot	22 (62.85%)	Boiled
<i>Trifolium pratense</i> L.	Fabaceae	Clover	Antidiabetic	He	Herbaceous perennial	Flowering offshoot	22 (62.85%)	Boiled
<i>Trifolium purpureum</i> Loisel.	Fabaceae	Clover	Antidiabetic	He	Herbaceous perennial	Flowering offshoot	11 (31.42%)	Boiled
<i>U. dioica</i> L.	Urticaceae	Nettle	Antidiabetic	He	Herbaceous perennial	Seed, aeration organ	25 (71.42%)	Boiled

\*: Calculated by 35 healers, n(%); *Lamium album* L.: *L. album*; *Teucrium polium* L.: *T. polium*; *Urtica dioica*: *U. dioica*.




Figure 3. The percentage of type used.

#### 4. Discussion

Current findings show many using medicinal herbs by Urmia traditional healers have antidiabetic effects. *C. colocynthis* could be applied in controlling blood glucose increment[35]. *Colocynthis* fruits have been prescribed in order to reduce blood glucose in traditional systems of medicine of many countries[36]. *In-vivo* studies show *colocynthis* juice can lead to reducing blood glucose[37]. An experimental study indicates that *U. dioica* possesses glucose reduction effect[38]. The fractions of nettle (*U. dioica*) cause an increase in insulin secretion in Langerhans Island taken from hyperglycemic mouse[39]. *U. dioica* and *L. album* can enhance insulin secretion in Langerhans Island and cause antidiabetic and blood glucose-reducing effects[39]. Current findings also showed that anti-diabetic property of nettle was due to its  $\alpha$ -glycosidase inhibitor effect[40]. Furthermore, flavonoids can be effective in improving the blood glucose indexes via their antioxidant activity[41]. Regarding, hydro alcoholic extraction of nettle could lead to rebuild  $\beta$ -cells in pancreas via its antioxidant characteristics. On the other hand, tannin and carotenoids, as nettle's constituents could be effective in improving blood glucose indexes[42].

The fruit of a kind of dog rose (*Rosa canina* L.) is conventionally used in order to prevent and treat sweet diabetes[43].

Investigations show barberry could lead to induce glucose uptake in a time- and dose-dependent manner. Several inhibitors indicated that this effect was not through insulin signaling but by means of adenosine monophosphate-activated protein kinase, which its downstream molecule is P38 mitogen-activated protein kinase[44].

Burdock contains inulin. In a study, sugar was not obviously seen in diabetes-afflicted people's urine who took 120 g inulin daily[45,46]. Taking inulin led to reduce the risk of cardiovascular disease because of triglycerides synthesis and fatty acids in liver. Nowadays, inulin is used successfully because of enjoying lower calorie content, food fiber enrichment, and other nutritional characteristics[47].

A kind of *Salvia officinalis* has glucose reducing effect[48]. Some investigations confirmed that *Salvia officinalis* could showed hypoglycemic and antidiabetic feature[49,50].

In herbal medicine, walnut is considered as a plant effective in treating diabetes[51]. The compounds of walnut tree leaves like phenolic acids and flavonoids (including quercetin, galactoside, quercetin pentoxide derivatives, quercetin arabinoside, quercetin glycosid, and quercetin rhamnoside) inhibit *in vitro* human plasma and low density lipoprotein oxidation[52,53]. Generally, flavonoids can lead to glucose plasma decline[54]; therefore, walnut antidiabetic quality may be due to the flavonoids involved.

Flavonoids are as key compounds for treating and managing diabetes in plants, vegetables, and fruits. Flavonoids, such as quercetin, cause insulin secretion and are also considered as a strong inhibitor in sorbitol accumulation throughout the body[55,56]. It may be the reason for positive effect of traditional herbal medicines effective in treating diabetes. The useful effect of flavonoids could be the result of increasing inter-cellular antioxidants levels, preventing capillaries rupture, and boosting the body immune system, all of which are totally effective in diabetes improvement[57,58]. Taking 1–2 g flavonoids mixture is helpful to cure diabetes disease and its complications[59].

Mulberry plant reduces blood glucose due to enhancing glucose peripheral re-absorption[60]. Antidiabetic activities of mentioned plants may also have been attributed to the following components. Barberry owns some effective substances such as chelidonic acid, resin, tannin, citric acid, and malic acid. Watercress contains gluconasturtiin. Milfoil has achillin. Sorrel dock owns oxalate salts. Copse includes *Papaver* and resins and also clover (trifolium) has phytoestrogen[61].

*Abrus precatorius* includes trigonelline which is a phytohormone and reduce blood glucose through its inhibitory effect on glucose-6-phosphatase and glycogen phosphorylase[62]. Trigonelline leads to enhance insulin receiver contents in blood red globules, hence improving glucose uptake in body tissues. Alleviating glucose level and inhibitory effect on two main enzymes of glucose metabolism by trigonelline could be promising for producing new medicines in treating diabetes[63].

In Arasbaran area (northwest of Iran), *Equisetum arvense* L. and *Rubus caesius* L. are used as antidiabetic drugs[64]. In the traditional medicine of Shiraz (Southern Iran), plants *Glaucium oxylobum* Boiss and Buhse, *Glaucium grandiflorum* Boiss and Huet, *L. album* Ky. ex Boiss., *Scorzonera canada* (C.A.Mey.) O. Hoffm., *T. polium* and *Solanum nigrum* are the antidiabetic herbs[65,66]. Plants in Kerman Province, located in east of Iran, *Coriandrum sativum*, *Hordeum vulgare* and *Peganum harmala* are antidiabetic herbs in that region[67]. In Isfahan (Central Iran), plants such as *Juglans regia* L., *Hordeum vulgare* L., *Cucurbita pepo*, *Coriandrum sativum* L. and *Mentha spicata* L. are used as antidiabetic plant sources[68]. In Ilam Province

(west of Iran), plants such as *C. colocynthis* L., *Glycyrrhiza glabra* L., *Gundelia tournefortii* L. and *Myrtus communis* L. are used in antidiabetic medicine in this region<sup>[69]</sup>. Comparison of the antidiabetic plants in Ghasemloo Valley with other parts of Iran shows that many plants are indigenous which has been reported for the first time.

Currently, there is no data on herbal plants used to treat diabetes in northwest of Iran. Therefore, these findings are so important in the management of diabetes and future study on traditional medicine in drug development.

It is important to find curative medicines effective in treating diabetes and also disease improving drugs with lower side effects. Many studies have been carried out about herbal medicines and plants' effective substances could be applicable to diabetes remedy. One of the most significant research scopes is to exam the disease changing effect, *i.e.* the efficiency of herbal medicines in preventing diabetes progress, which requires doing experiments on mechanisms, herbal medicines efficiency and safety as well as useful potential botanical substances in diabetes disease remedy.

It should be noted that diabetes mellitus is accompanied with severe increases in oxidative stress<sup>[70–73]</sup>. Oxidative stress, other than diabetes, is also involved in a lot of curable hard diseases such as atherosclerosis<sup>[74,75]</sup>, cardiovascular diseases<sup>[76,21]</sup>, cancer<sup>[55,77]</sup>, Alzheimer<sup>[78]</sup>, gastrointestinal<sup>[79,80]</sup> and infectious diseases<sup>[81,82]</sup>. Most of medicinal plants possess antioxidant activity<sup>[83,84]</sup>, thanks to polyphenolic compounds such as flavonoids, flavonols and anthocyanins which have been seen in a lot of medicinal plants reported in this paper<sup>[83,85,86]</sup>. Medicinal plants with antioxidant activity have been able to counteract with the development of these complications. The antioxidants usually scavenge the free radicals which cause oxidative stress<sup>[73,87]</sup>. Therefore, plants' effectiveness in diabetes mellitus should counteract, at least in part, with oxidative stress induced through diabetes, reducing its symptoms.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

The authors appreciate scientific supports of Urmia University of Medical Sciences and Shahrekord University of Medical Sciences. This research was supported by the Deputy for Research and Technology of Lorestan University of Medical Sciences (Grant No. 0207).

### References

- [1] Morey SC. ADA recommends a lower fasting glucose value in the diagnosis of diabetes mellitus. *Am Fam Physician* 1997; **56**(8): 2128, 2130.
- [2] Srinivasan K, Ramarao P. Animal models in type 2 diabetes research: an overview. *Indian J Medical Res* 2007; **125**(3): 451–472.
- [3] Kazemi S, Asgary S, Moshtaghian J, Rafieian M, Adelnia A, Shamsi F. Liver-protective effects of hydroalcoholic extract of *Allium hirtifolium* Boiss. In rats with alloxan-induced diabetes mellitus. *ARYA Atheroscler* 2010; **6**(1): 11–15.
- [4] Gleckman R, Morr J. Diabetes-related foot infections. *Contemp Intern Med* 1994; **6**(8): 57–64.
- [5] Behradmanesh S, Derees F, Rafieian-kopaei M. Effect of *Salvia officinalis* on diabetic patients. *J Ren Inj Prev* 2013; **2**(2): 51–54.
- [6] Meigs JB. Epidemiology of the metabolic syndrome, 2002. *Am J Manag Care* 2002; **8**(11 Suppl): S283–S292; quiz S293–S296.
- [7] Boulton AJ, Vileikyte L, Ragnarson-Tennvall G, Apelqvist J. The global burden of diabetic foot disease. *Lancet* 2005; **366**(9498): 1719–1724.
- [8] Williams RH, Larsen PR. *Textbook of endocrinology*. 10th ed. Philadelphia: Saunders; 2003.
- [9] Swanston-Flatt SK, Day C, Bailey CJ, Flatt PR. Evaluation of traditional plant treatments for diabetes: studies in streptozotocin diabetic mice. *Acta Diabetol Lat* 1989; **26**(1): 51–55.
- [10] Nasri H, Rafieian-Kopaei M. Herbal medicine and diabetic kidney disease. *J Nephropharmacol* 2013; **2**(1): 1–2.
- [11] Eddouks M, Chattopadhyay D, De Feo V, Cho VC. Medicinal plants in the prevention and treatment of chronic diseases 2013. *Evid Based Complement Alternat Med* 2014; doi: 10.1155/2014/180981.
- [12] Jamila F, Mostafa E. Ethnobotanical survey of medicinal plants used by people in Oriental Morocco to manage various ailments. *J Ethnopharmacol* 2014; **154**(1): 76–87.
- [13] Venkatesh S, Reddy GD, Reddy BM, Ramesh M, Rao AV. Antihyperglycemic activity of *Caralluma attenuate*. *Fitoterapia* 2003; **74**(3): 274–279.
- [14] Nasri H, Shirzad H. Toxicity and safety of medicinal plants. *J HerbMed Pharmacol* 2013; **2**(2): 21–22.
- [15] Singh VK, Govil JN, Singh G. *Recent progress in medicinal plants, volume 1: ethnomedicine and pharmacognosy*. New York: Sci Tech Publishing Inc; 2002.
- [16] Behradmanesh M, Ahmadi M, Rafieian-kopaei M. Effect of glycol on blood glucose level of patients with type II diabetes. *Iran J Endocrinol Metab* 2012; **14**(2): 163–168.
- [17] Kennedy J. Herb and supplement use in the US adult population. *Clin Ther* 2005; **27**(11): 1847–1858.
- [18] Gallagher AM, Flatt PR, Duffy G, Abdel-Wahab YHA. The effects of traditional antidiabetic plant on *in vitro* glucose diffusion. *Nutr Res* 2003; **23**(3): 413–424.
- [19] Marles RJ, Farnsworth NR. Antidiabetic plants and their active constituents. *Phytomedicine* 1995; **2**(2): 137–189.

- [20] Akbari F, Ansari–Samani R, Karimi A, Mortazaei S, Shahinfard N, Rafieian–kopaei M. Effect of turnip on glucose and lipid profiles of alloxan–induced diabetic rats. *Iran J Endocrinol Metab* 2013; **14**(5): 492–497.
- [21] Khosravi–Boroujeni H, Mohammadifard N, Sarrafzadegan N, Sajjadi F, Maghroun M, Khosravi A, et al. Potato consumption and cardiovascular disease risk factors among Iranian population. *Int J Food Sci Nutr* 2012; **63**(8): 913–920.
- [22] Zargaran A. Ancient Persian medical views on the heart and blood in the Sassanid era (224–637 AD). *Int J Cardiol* 2014; **172**(2): 307–312.
- [23] Hamed A, Zarshenas MM, Sohrabpour M, Zargaran A. Herbal medicinal oils in traditional Persian medicine. *Pharm Biol* 2013; **51**(9): 1208–1218.
- [24] Mohagheghzadeh A, Zargaran A, Daneshamuz S. Cosmetic sciences from ancient Persia. *Pharm Hist (Lond)* 2011; **41**(2): 18–23.
- [25] Zargaran A, Fazlzadeh A, Mohagheghzadeh A. Surgeons and surgery from ancient Persia (5,000 years of surgical history). *World J Surg* 2013; **37**(8): 2002–2004.
- [26] Zargaran A, Kordafshari G, Hosseini SR, Mehdizadeh A. Akhawayni (?–983 AD): a Persian neuropsychiatrist in the early medieval era (9th–12th Century AD). *J Med Biogr* 2014; doi: 10.1177/0967772014525105.
- [27] Zarshenas MM, Mehdizadeh A, Zargaran A, Mohagheghzadeh A. Rhazes (865–925 AD). *J Neurol* 2012; **259**(5): 1001–1002.
- [28] Zargaran A, Mehdizadeh A, Zarshenas MM, Mohagheghzadeh A. Avicenna (980–1037 AD). *J Neurol* 2012; **259**(2): 389–390.
- [29] Zargaran A, Zarshenas MM, Ahmadi SA, Vessal K. Haly Abbas (949–982 AD). *J Neurol* 2013; **260**(8): 2196–2197.
- [30] Zarshenas MM, Zargaran A, Abolhassanzadeh Z, Vessal K. Jorjani (1042–1137). *J Neurol* 2012; **259**(12): 2764–2765.
- [31] Zarshenas MM, Khademian S, Moein M. Diabetes and related remedies in medieval Persian medicine. *Indian J Endocrinol Metab* 2014; **18**(2): 142–149.
- [32] Heidari Rikan M, Malekmoohamadi L. Medicinal plants in Ghasemloo Valley of Uromieh. *Iran J Med Aromat Plants* 2007; **23**(2): 234–250.
- [33] Mokhber Dezfuli MR, Hajiakhoondi A, Bahmani M, Fazeli R, Eftekhari Z, et al. *Biological and pharmacological on natural products and medicinal plants extracts, general methods*. 1st ed. Tehran: University of Tehran Press (No. 3374). 1th ed. 2013, p. 1.
- [34] Bahmani M, Sepahvand R, Delfan B, Saki K, Abdi F, Bahmani A. *Alternative medicine beliefs of martyrs Valley (Valley Ghasemloo) in Western Azerbaijan*. 1st ed. Tehran: Grove Press; 2014, p. 15–18.
- [35] Rahman MA, Mossa JS, Al–Said MS, Al–Yahya MA. Medicinal plant diversity in the flora of Saudi Arabia 1: a report on seven plant families. *Fitoterapia* 2004; **75**(2): 149–161.
- [36] Behradmanesh S, Ahmadi M, Rafieian–kopaei M. Effect of diabetan on blood glucose, glycosylated hemoglobin, lipid profile, liver and kidney function tests of diabetic patients: a clinical, double blind, randomized trial. *Afr J Pharm Pharmacol* 2013; **7**(2): 50–53.
- [37] Craig WJ. Health–promoting properties of common herbs. *Am J Clin Nutr* 1999; **70**(3 Suppl): 491S–499S.
- [38] Bnouham M, Merhfour FZ, Ziyat A, Mekhfi H, Aziz M, Legssyer A. Antihyperglycemic activity of the aqueous extract of *Urtica dioica*. *Fitoterapia* 2003; **74**: 677–681.
- [39] Farzami B, Ahmadvand D, Vardasbi S, Majin FJ, Khaghani Sh. Induction of insulin secretion by a component of *Urtica dioica* leave extract in perfused Islets of Langerhans and its *in vivo* effects in normal and streptozotocin diabetic rats. *J Ethnopharmacol* 2003; **89**(1): 47–53.
- [40] Onal S, Timur S, Okutucu B, Zihnioğlu F. Inhibition of alpha–glucosidase by aqueous extracts of some potent antidiabetic medicinal herbs. *Prep Biochem Biotechnol* 2005; **35**(1): 29–36.
- [41] Song Y, Manson JE, Buring JE, Sesso HD, Liu S. Associations of dietary flavonoids with risk of type 2 diabetes, and markers of insulin resistance and systemic inflammation in women: a prospective study and cross–sectional analysis. *J Am Coll Nutr* 2005; **24**(5): 376–384.
- [42] Mangels AR, Holden JM, Beecher GR, Forman MR, Lanza E. Carotenoid content of fruits and vegetables: an evaluation of analytic data. *J Am Diet Assoc* 1993; **93**(3): 284–296.
- [43] Wenzig EM, Widowitz U, Kunert O, Chrubasik S, Bucar F, Knauder E, et al. Phytochemical composition and *in vitro* pharmacological activity of two rose hip (*Rosa canina* L.) preparations. *Phytomedicine* 2008; **15**(10): 826–835.
- [44] Cheng Z, Pang T, Gu M, Gao AH, Xie CM, Li JY, et al. Berberine–stimulated glucose uptake in L6 myotubes involves both AMPK and p38 MAPK. *Biochim Biophys Acta* 2006; **1760**(11): 1682–1689.
- [45] Van den Ende W. Fructan metabolism in chicory roots [dissertation]. Belgium: University of Leuven, 1996.
- [46] Roberfroid MB. Inulin–type fructans: functional food ingredients. *J Nutr* 2007; **137**(11 Suppl): 2493S–2502S.
- [47] Myers RH, Montgomery DC, Anderson–Cook CM. *Response surface methodology: process and product optimization using designed experiments*. 2nd ed. New Jersey: Wiley; 2002, p. 115–124.
- [48] Eidi M, Eidi A, Zamanizadeh H. Effect of *Salvia officinalis* L. leaves on serum glucose and insulin in healthy and streptozotocin–induced diabetic rats. *J Ethnopharmacol* 2005; **100**(3): 310–313.
- [49] Geller SE, Studee L. Contemporary alternatives to plant estrogens for menopause. *Maturitas* 2006; **55**(Suppl 1): S3–S13.
- [50] Grzegorzczak I, Matkowski A, Wysokinska H. Antioxidant activity of extracts from *in vitro* cultures of *Salvia officinalis* L. *Food Chem* 2007; **104**(2): 536–541.
- [51] Zargari A. *Medicinal plants*. Tehran: Tehran Univ.; 1997, p. 106–111.
- [52] Anderson KJ, Teuber SS, Gobeille A, Cremin P, Waterhouse AL, Steinberg FM. Walnut polyphenolics inhibit *in vitro* human plasma and LDL oxidation. *J Nutr* 2001; **131**(11): 2837–2842.
- [53] Sharafati–Chaleshtori R, Sharafati–Chaleshtori F, Rafieian–Kopaei M. Biological characterization of Iranian walnut (*Juglans regia*) leaves. *Turk J Biol* 2011; **35**(5): 635–639.
- [54] Li WL, Zheng HC, Bukuru J, De Kimphe N. Natural medicines used in the traditional Chinese medical system for therapy of diabetes mellitus. *J Ethnopharmacol* 2004; **92**(1): 1–21.
- [55] Shirzad H, Taji F, Rafieian–Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI–164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food* 2011; **14**(9): 969–974.
- [56] Hii CS, Howell SL. Effects of flavonoids on insulin secretion and  $^{45}\text{Ca}^{2+}$  handling in rat islets of Langerhans. *J Endocrinol* 1985; **107**(1): 1–8.

- [57] Sakai I, Izumi SI, Murano T, Okuwaki S, Makino T, Suzuki T. Presence of aldose reductase inhibitors in tea leaves. *Jpn J Pharmacol* 2001; **85**(3): 322–326.
- [58] Asgary S, Kelishadi R, Rafieian-Kopaei M, Najafi S, Najafi M, Sahebkar A. Investigation of the lipid-modifying and antiinflammatory effects of *Cornus mas* L. supplementation on dyslipidemic children and adolescents. *Pediatr Cardiol* 2013; **34**(7): 1729–1735.
- [59] Tamadon MR, Baradaran A, Rafieian-Kopaei M. Antioxidant and kidney protection; differential impacts of single and whole natural antioxidants. *J Renal Inj Prev* 2014; **3**(2): 41–42.
- [60] Chen F, Nakashima N, Kimura I, Kimura M. [Hypoglycemic activity and mechanisms of extracts from mulberry leaves (*Folium mori*) and cortex mori radices in streptozotocin-induced diabetic mice]. *Yakugaku Zasshi* 1995; **115**(6): 476–482. Japanese.
- [61] Ghasemi-pirbalouti A. [*Third listen: plants, traditional medicine and ethnoveterinary, medicinal and aromatic plant*]. 1st ed. Shahrekord: Saman-Danesh Pub; 2009, p. 158–190. Persian.
- [62] Zeiger E. Trigonelline: review of toxicological literature. [dissertation]. North Carolina: National Institute of Environmental Health Sciences; 1997.
- [63] Monago Comfort C, Nwodo O Fred C. Antidiabetic effects of crude flavanoid and alkaloid of *Abrus precatorius* Linn. seed in alloxan diabetic rabbits. *Res J Pharmacogn Phytochem* 2010; **2**(4): 331–335.
- [64] Zolfaghari E, Adeli I, Mozafarian V, Babaiy S, Habibi Bibalani GH. Identification of Mrasbaran medicinal plants and ethnobotanical study of rural people knowledge (case study: Arasbaran Forest, Mardanaghom Watershed). *Iran J Med Arum Plants* 2012; **28**(3): 534–550.
- [65] Sadeghi E, Borjian A. [Medicinal plants of *Cheshme anger* region (Shiraz)]. *J Res Plants Sci* 2013; **1**(7): 25, 42–59. Persian.
- [66] Dolatkhaei M, Ghorbani Nohooji M, Mehrafarin E, Amini Nejad GR, Dolatkhaei A. Ethnobotanical study of medicinal plants in Kazeroon, Iran: identification, distribution and traditional usage. *J Med Plants* 2012; **11**(42): 163–178.
- [67] Rajaei P, Mohamadi N. Ethnobotanical study of medicinal plants of Hezar Mountain allocated in South East of Iran. *Iran J Pharma Res* 2012; **11**(4): 1153–1167.
- [68] Shaahin MN, Mansoureh V. Ethno-botany of medicinal plants by Mobarakeh's people (Isfahan). *J Herb Drugs* 2012; **3**(2): 111–129.
- [69] Ghasemi Pirbalouti A, Momeni M, Bahmani M. Ethnobotanical study of medicinal plants used by Kurd Tribe in Dehloran and Abdanan Districts, Ilam Province, Iran. *Afr J Tradit Complement Altern Med* 2012; **10**(2): 368–385.
- [70] Baradaran A, Madihi Y, Merrikhi A, Rafieian-Kopaei M, Nasri H. Serum lipoprotein (a) in diabetic patients with various renal function not yet on dialysis. *Pak J Med Sci* 2013; **29**: 354–357.
- [71] Behradmanesh S, Horestani MK, Baradaran A, Nasri H. Association of serum uric acid with proteinuria in type 2 diabetic patients. *J Res Med Sci* 2013; **18**(1): 44–46.
- [72] Rafieian-Kopaei M, Behradmanesh S, Kheiri S, Nasri H. Association of serum uric acid with level of blood pressure in type 2 diabetic patients. *Iran J Kidney Dis* 2014; **8**(2): 152–154.
- [73] Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014; **19**(1): 82–83.
- [74] Madihi Y, Merrikhi A, Baradaran A, Rafieian-Kopaei M, Shahinfard N, Ansari R, et al. Impact of sumac on postprandial high-fat oxidative stress. *Pak J Med Sci* 2013; **29**: 340–345.
- [75] Madihi Y, Merrikhi A, Setorki M, Baradaran A, Ghobadi S, Shahinfard N, et al. Bioactive components and the effect of hydroalcoholic extract of *Vaccinium myrtillus* on postprandial atherosclerosis risk factors in rabbits. *Pak J Med Sci* 2013; **29**: 384–389.
- [76] Sarrafzadegan N, Khosravi-Boroujeni H, Esmailzadeh A, Sadeghi M, Rafieian-Kopaei M, Asgary S. The association between hypertriglyceridemic waist phenotype, menopause, and cardiovascular risk factors. *Arch Iran Med* 2013; **16**(3): 161–166.
- [77] Shirzad H, Shahrani M, Rafieian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes *in vivo*. *Int Immunopharmacol* 2009; **9**(7–8): 968–970.
- [78] Rabiei Z, Rafieian-Kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of Meynert in rat. *Neurochem Res* 2014; **39**(2): 353–360.
- [79] Sedighi M, Rafieian-kopaei M, Noori-Ahmadabadi M. *Kelussia odoratissima* Mozaffarian inhibits ileum contractions through voltage dependent and beta adrenergic receptors. *Life Sci J* 2012; **9**(4): 1033–1038.
- [80] Rahimian G, Sanei MH, Shirzad H, Azadegan-Dehkordi F, Taghikhani A, Salimzadeh L, et al. Virulence factors of *Helicobacter pylori* vacA increase markedly gastric mucosal TGF- $\beta$ 1 mRNA expression in gastritis patients. *Microb Pathog* 2014; **67–68**: 1–7.
- [81] Rafieian-Kopaei M, Baradaran A, Maghsoudi AR, Ghobadi S, Nasri H. *Helicobacter pylori* infection and serum homocysteine in hemodialysis patient. *Life Sci J* 2012; **9**(4): 3696–3702.
- [82] Bagheri N, Rahimian G, Salimzadeh L, Azadegan F, Rafieian-Kopaei M, Taghikhani A, et al. Association of the virulence factors of *Helicobacter pylori* and gastric mucosal interleukin-17/23 mRNA expression in dyspeptic patients. *EXCLI J* 2013; **12**: 5–14.
- [83] Nasri H, Rafieian-Kopaei M. Medicinal plants and antioxidants: why they are not always beneficial? *Iran J Public Health* 2014; **43**(2): 255–257.
- [84] Rafieian-Kopaei M, Baradaran A, Rafieian M. Oxidative stress and the paradoxical effects of antioxidants. *J Res Med Sci* 2013; **18**(7): 629.
- [85] Baradaran A, Nasri H, Rafieian-Kopaei M. Oxidative stress and hypertension: possibility of hypertension therapy with antioxidants. *J Res Med Sci* 2014; **19**(4): 358–367.
- [86] Kafash-Farkhad N, Asadi-Samani M, Rafieian-Kopaei M. A review on phytochemistry and pharmacological effects of *Prangos ferulacea* (L.) Lindl. *Life Sci J* 2013; **10**(8s): 360–367.
- [87] Asgary S1, Sahebkar A, Afshani MR, Keshvari M, Haghjooyjavanmard S, Rafieian-Kopaei M. Clinical evaluation of blood pressure lowering, endothelial function improving, hypolipidemic and anti-inflammatory effects of pomegranate juice in hypertensive subjects. *Phytother Res* 2014; **28**(2): 193–199.