

Clinical investigation of the acute effects of pomegranate juice on blood pressure and endothelial function in hypertensive individuals

**Sedigheh Asgary⁽¹⁾, Mahtab Keshvari⁽²⁾, Amirhossein Sahebkar⁽³⁾,
Mohamad Hashemi⁽⁴⁾, Mahmoud Rafieian-Kopaei⁽⁵⁾**

Original Article

Abstract

BACKGROUND: Pomegranate juice (PJ) is rich in bioactive phytochemicals with antioxidant, and anti-inflammatory and cardioprotective functions. The present trial investigated the acute effects of PJ consumption on blood pressure and markers of endothelial function.

METHODS: In this single-arm study, thirteen hypertensive men aged 39–68 years were recruited. Included subjects were assigned to natural PJ (150 ml/day) following a 12 hour fast. Systolic blood pressure (SBP), diastolic blood pressure (DBP), and flow-mediated dilation (FMD), along with serum concentrations of C-reactive protein (CRP), intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), E-selectin and interleukin-6 (IL-6) were measured at baseline and 4-6 hours after PJ consumption.

RESULTS: Comparison of pre- vs. post-trial values revealed a significant reduction in both SBP (7%; $P = 0.013$) and DBP (6%; $P < 0.010$). However, changes in FMD (20%) as well as circulating levels of CRP, ICAM-1, VCAM-1, E-selectin, and IL-6 did not reach statistical significance ($P = 0.172$).

CONCLUSION: PJ has promising acute hypotensive properties. Consumption of PJ could be considered in the context of both dietary and pharmacological interventions for hypertension.

Keywords: Punica Granatum L., Cardiovascular Disease, Hypertension, Inflammation, Endothelium-Dependent Dilation

Date of submission: 11 Apr 2013, *Date of acceptance:* 20 Jul 2013

Introduction

Cardiovascular disorders are among the leading causes of death and disability in the world.^{1,2} Hypertension is a major risk factor for cardiovascular and cerebrovascular disease, end stage renal disease, type 2 diabetes, and metabolic syndrome. Hypertension has a high global prevalence of about 15% that is estimated to reach as high as 30% by 2025.^{3,4} Controlled studies have indicated that each 5 mmHg decrease in diastolic blood pressure (DBP) is associated with 15% and 40% reductions in the risk of cardiovascular disease and stroke, respectively.⁵ In spite the introduction of several classes of anti-hypertensive agents with different mechanisms of action, uncontrolled hypertension resistant to drug therapy still remains a frequent medical problem.

Pomegranate (*Punica granatum* L.; Family Punicaceae) is a popular edible fruit with wide applications in traditional medicine.^{6,7} Several lines of modern scientific evidence have also indicated the therapeutic efficacy of pomegranate against different types of disorders.⁸⁻¹¹ The pomegranate is characterized by considerable amounts of biologically active phytochemicals including flavonoids (e.g. anthocyanins, catechins, quercetin, and rutin), other types of polyphenols, ellagitannins, and antioxidant vitamins.¹²⁻¹⁵ Many of these phytochemicals have been shown to possess antioxidant and anti-inflammatory properties plus additional biological activities such as inhibition of angiotensin converting enzyme.¹⁶⁻²⁰ All these activities of the pomegranate are potentially beneficial for the treatment of hypertension and improvement of

1-Professor, Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

2- Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

3- Assistant Professor, Biotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

4- Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

5- Professor, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

Correspondence to: Sedigheh Asgary, Email: sasgary@yahoo.com

endothelial function.¹¹ However, clinical studies investigating the hypotensive and cardioprotective effects of pomegranate have been scarce.²⁰⁻²⁶

According to the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC7), prehypertension is defined as 120 mmHg \leq systolic blood pressure (SBP) $<$ 140 mmHg 12 and/or 80 mmHg \leq DBP $<$ 90 mmHg.²⁷ Prehypertension is a clinical stage where subjects are at increased risk (2 folds higher) of developing hypertension in the near future.²⁸ Emerging findings suggest that interventions at the prehypertension stage can prevent or delay the progression of disease into established hypertension and subsequent detrimental outcomes.²⁹ The present trial investigated the acute effects of pomegranate juice (PJ) on blood pressure and endothelial function in subjects with diagnosed prehypertension.

Materials and Methods

Subjects

Thirteen hypertensive men aged 39–68 years were recruited for this trial.³⁰ The Ethics Committee at the Shahrekord University of Medical Sciences (Iran) approved the study protocol (code: 92-3-16) and written informed consents were obtained from all participants.

The inclusion criteria were body mass index (BMI) \leq 30, and diagnosed hypertension defined as SBP $>$ 120 mmHg and/or DBP $>$ 80 mmHg. Exclusion criteria were type 1 or 2 diabetes, chronic pancreatitis, liver cirrhosis, kidney stones, renal failure, use of non-steroidal anti-inflammatory drugs, use of antioxidant or vitamin supplements, intense physical activity ($>$ 5 h/week), smoking habit, being vegetarian or having any restrictive dietary requirements, and pregnancy.

Study design

The present study was designed as a single-arm clinical trial. The included subjects were assigned to natural PJ (150 ml/day) following a 12-h fast. SBP, DBP, and flow-mediated dilation (FMD), along with serum concentrations of C-reactive protein (CRP), intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), E-selectin, and interleukin-6 (IL-6) were measured at baseline and 4-6 h after PJ consumption. Participants were asked not to eat or drink anything during the interval between PJ consumption and final measurements.

Total anthocyanin assay

A pH differential method was carried out to

determine the total anthocyanin content of PJ. In this method, the difference in the absorbance of sample at pHs 1.0 and 4.5 is proportional to the total anthocyanin content. Details of this method have been published previously.^{24,31}

Anthropometric and BP measurements

Measurement of weight and height was carried out using a standard procedure as described previously.^{24,32} Body mass index (BMI) was calculated as weight in kg divided by height in meters squared (m²). BP was measured by a single operator at baseline and 4-6 h after intake of PJ, according to a standard protocol. BP recordings were performed after rest, employing a stethoscope and calibrated sphygmomanometer (Accurtorr 1A; Datascope, Japan). Systolic blood pressure was defined as the appearance of the first sound (Korotkoff phase 1) and diastolic blood pressure was defined as the disappearance of the sound (Korotkoff phase 5) during deflating of the cuff.^{24,32}

FMD measurement

Endothelium-dependent FMD was measured on the right brachial artery as described previously.³³ All measurements were carried out by a single operator following a 5-min rest and employing a GE vivid 3 ultrasound apparatus (AtCor Medical, Solingen, Germany). A BP cuff was inflated around the forearm to 200 mmHg for 5 min. Images were recorded at baseline (before inflation), 30 s before cuff release, and then every 15 sec after cuff release for 3 min. Arterial diameter was measured at the end of end-diastolic phase, coinciding with R-wave on the electrocardiogram (ECG). Brachial FMD was expressed as the percentage change in arterial diameter from baseline.³⁴

Blood sampling and biochemical analyses

Twelve-hour fasted blood samples were taken from the left antecubital vein. After being allowed to clot for 2-3 h, serum was isolated by centrifugation at 3500-4000 rpm for 10 min. Serum samples were kept at -80°C prior to biochemical analyses. Biochemical analyses were performed using an automated enzymatic assay (Pars Azmoon, Tehran, Iran) on a Hitachi 902 autoanalyzer (for CRP), or enzyme-linked immunosorbent assay (ELISA) with commercial kits (Boster Biological Technology Ltd., Wuhan, China) (for ICAM-1, VCAM-1, E-selectin, and IL-6). Inter-assay coefficients of variation for ICAM-1, VCAM-1, E-selectin and IL-6 were 4.1-6.4%, 6.1-7.7%, 6.6-8.1%, and 3.1-5.5%, respectively. Intra-assay coefficients of variation for ICAM-1, VCAM-1, E-selectin, and IL-6 were 3.4-5.1%, 2.3-3.7%, 5.2-6.9%, and 2.3-4.9%, respectively.

Statistical analysis

All statistical analyses were performed using SPSS for Windows (version 17; SPSS Inc., Chicago, IL, USA). Data were expressed as mean \pm SD. Group comparisons were made using paired t-test (in case of normally distributed data) or Wilcoxon signed-ranks test (in case of non-normally distributed data). A two-sided P-value of < 0.05 was considered to be statistically significant.

Results

This trial was comprised of 13 hypertensive male adolescents with a mean age, weight, and BMI of 55.92 ± 7.92 yrs, 80.42 ± 11.01 kg, and 27.34 ± 3.82 kg/m², respectively. All 13 subjects completed the study. Total anthocyanin content of PJ was determined to be 5.8 mg per 100 ml of the administered juice.

Comparison of pre- vs. post-trial values revealed a significant reduction in both SBP ($P = 0.013$) and DBP ($P = 0.010$), amounting to an approximate reduction by 7% and 6%, respectively. However, percentage changes in FMD (20%) was not found to be statistically significant ($P = 0.172$). In the same manner, there was no significant difference in the circulating concentrations of inflammatory biomarkers namely hsCRP ($P = 0.263$), ICAM-1 ($P = 0.248$), VCAM-1 ($P = 0.657$), E-selectin ($P = 0.182$), and IL-6 ($P = 0.763$) following consumption of PJ. Baseline and post-trial values for the evaluated parameters are summarized in table 1.

Discussion

The present pilot trial is one of the few clinical evidences on the acute hypotensive and vascular effects of PJ. The results indicated amelioration of both SBP and DBP following consumption of a single dose of PJ. In a previous study, Aviram and Dornfeld investigated the effects of 2-week

supplementation with PJ (50 ml/day) on the SBP of hypertensive patients.²⁰ The findings revealed a significant decrease in SBP amounting to 5%. In the same study, a 36% decrement in the activity of serum angiotensin converting enzyme (ACE) was reported from PJ.²⁰

The same group also investigated the effect of chronic supplementation with PJ (50 ml/day) in patients with carotid artery stenosis. Their results indicated a significant reduction in SBP, but not DBP, starting from 1 month after starting supplementation and generally increasing up to month 12 (equivalent to 12% decrement). However, no further reduction was observed when supplementation was continued for another 2 years.²¹ Another study conducted by Lynn et al. indicated a significant reduction in both SBP and DBP following consumption of PJ (330 ml/day) by healthy subjects for 4 weeks.²² Mathew et al. reported that consumption of pomegranate extract, either during or 15 min before a high-fat meal, can effectively prevent postprandial SBP rise at 2 and 4 h postprandial. Nevertheless, no significant effect was found on DBP.²³ In a recent trial conducted by our group, the effects of a 2-week intake of PJ (from the same source as that used in the present study) were evaluated on BP of hypertensive subjects. Our results implied a significant reduction of both SBP (5%) and DBP (4.5%) compared to the control group who consumed water instead of PJ.²⁴ Although most of the findings by previous trials infer the hypotensive impact of PJ consumption, there are some contrasting findings that are worth attention. Kelishadi et al. investigated the acute (4 h) and chronic (1 month) effects of PJ (240 ml/day) on BP of adolescents with metabolic syndrome and could not find any significant effect.²⁵ In another trial, administration of PJ (240 ml/day) for 3 months was not found to significantly affect SBP or DBP in patients with stable coronary heart disease (CHD), as reported by Sumner et al.²⁶ These negative

Table 1. Acute effects of pomegranate juice on blood pressure, flow-mediated dilation, and biochemical parameters

Parameters	Before	After	P
SBP (mmHg)	125.38 \pm 11.80	116.15 \pm 7.94	0.013
DBP (mmHg)	82.69 \pm 5.25	78.08 \pm 3.25	0.010
FMD (%)	0.30 \pm 0.17	0.36 \pm 0.17	0.172
Hs-CRP (mg/L)	1.39 \pm 1.14	1.20 \pm 0.94	0.263
ICAM-1 (ng/mL)	297.00 \pm 122.30	265.23 \pm 125.35	0.248
VCAM-1 (ng/mL)	1008.31 \pm 450.99	993.92 \pm 447.42	0.657
E-Selectin (ng/mL)	27.77 \pm 15.89	25.49 \pm 14.19	0.182
IL-6 (ng/mL)	0.45 \pm 0.16	0.46 \pm 0.15	0.763

SBP: Systolic blood pressure; DBP: Diastolic blood pressure; FMD: Flow-mediated dilation; hsCRP: High-sensitivity C-reactive protein; ICAM-1: Intracellular adhesion molecule-1; VCAM-1: Vascular cell adhesion molecule-1; IL-6: Interleukin-6; Group comparisons were made using paired t-test (in case of normally distributed data) or Wilcoxon signed-ranks test (in case of non-normally distributed data).

findings might be attributed to the difference in the inclusion criteria applied by these two latter trials. It appears that hypotensive effects of PJ are more likely to be elicited in hypertensive patients, rather than patients with established CHD or metabolic syndrome. Moreover, the overall findings of the trials conducted so far weigh in favor of the beneficial effect of PJ consumption on BP.

The hypotensive properties of PJ could be ascribed to the promising antioxidant properties of phytochemicals present in this complex juice. Oxidative stress is known to play a key role in the pathogenesis of hypertension.³⁵ Increased levels of oxidants have been shown in different experimental models of hypertension.³⁶⁻³⁹ Detrimental effects of oxidative stress are mainly due to the interaction of reactive oxygen species (ROS) (in particular superoxide anion) with vital cellular components, lipids, and proteins, which leads to endothelial dysfunction and vascular resistance. In addition, ROS interfere with the production and vasodilatory actions of endothelium-derived nitric oxide (NO) via attenuating NO synthase (NOS) activity and enhancing NO breakdown.^{40,41} Epidemiological evidence has indicated that diets rich in natural antioxidants are associated with a reduced risk of developing hypertension and cardiovascular events.^{42,43} A plethora of studies have confirmed the considerable antioxidant and radical scavenging effects of PJ, which are mainly due to the anthocyanins and hydrolysable tannins present in the fruit. Interestingly, it has been suggested that some 50% of the total antioxidant activity of PJ is exerted by a specific ellagitannin, named punicalagin.¹⁶ Intestinal hydrolysis of punicalagin yields ellagic acid; the latter being a strong antioxidant compound.⁴⁴

Apart from antioxidant properties, PJ may lower BP through a direct interaction with ACE. As referred above, a significant reduction in the activity of serum ACE has been observed in hypertensive patients following PJ consumption.²⁰ Besides, in an animal study by Mohan et al. PJ administration attenuated angiotensin II-induced hypertension in diabetic rats, and also blocked the effects of different catecholamines on arterial BP and vasoreactivity. Furthermore, chronic administration of PJ counterbalanced the increased ACE activity in diabetic hypertensive rats.⁴⁵ In spite of these positive reports, the ACE inhibitory effect of PJ needs to be further explored as Lynn et al. failed to report any change in serum ACE concentration following PJ consumption for 4 weeks.²²

Another primary outcome measure that was evaluated by the present trial was changes in endothelium-dependent flow-mediated dilation. Our results did not indicate any improvement in FMD and this is in agreement with our recent report on the effects of 2-week PJ intake.²⁴ However, findings from another trial indicated significant improvement in both endothelium-dependent and -independent (nitroglycerin-induced) dilation after 4 h of PJ consumption. In addition, this increased vasodilation was persisted until the end of supplementation period (1 month).²⁵ In the present study, biomarkers of endothelial function, namely ICAM-1, VCAM-1, and E-selectin, remained statistically unaltered compared to baseline levels. It should be noticed that alterations in the circulating levels of these biomarkers is more likely to be exerted by chronic, rather than single dose, consumption of PJ and needs a longer term evaluation to allow the turnover of previously released proteins and observation of possible changes in the expression and subsequent release of these markers due to PJ consumption. This notion is corroborated by our previous study which showed a decreasing trend in serum levels of the aforementioned biomarkers following consumption of PJ.²⁴

The present study has certain limitations that need to be acknowledged. This study did not include a control group. Therefore, our findings might have been confounded by bias. In addition, the present trial was conducted in pilot scale and with a small population size. This small size could potentially account for lack of detecting significant difference despite the increasing trend in FMD and decreasing trends in serum CRP, ICAM-1, VCAM-1, and E-selectin levels. With respect to these limitations, findings of the present trial may not be generalizable to the general population and should be interpreted with caution.

In conclusion, the key finding to emerge from the present study is the acute hypotensive effect of PJ in hypertensive patients. While this trial is not a substitution for well-designed randomized controlled trials, it generates a hypothesis and motivates further research on this topic. Future large-scale investigations are indeed warranted in order to obtain a mechanistic understanding on this observed hypotensive activity.

Acknowledgements

This study was conducted with financial support

that was provided by the Shahrekord University of Medical Sciences, Shahrekord, Iran.

Conflict of Interests

Authors have no conflict of interests.

References

- Kochanek KD, Xu J, Murphy SL, Minino AM, Kung HC. Deaths: Final Data for 2009. *National Vital Statistics Reports* 2011; 60(3): 1-110.
- Whitworth JA. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003; 21(11): 1983-92.
- Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet* 2005; 365(9455): 217-23.
- Hajjar I, Kotchen JM, Kotchen TA. Hypertension: trends in prevalence, incidence, and control. *Annu Rev Public Health* 2006; 27: 465-90.
- Fazizi F, Esmailzadeh A, Mirmiran FP. Obesity and cardiovascular disease risk factors in Tehran adults: a population-based study. *East Mediterr Health J* 2004; 10(6): 887-97.
- Lansky EP, Newman RA. Punicagranatum (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *J Ethnopharmacol* 2007; 109(2): 177-206.
- Amruthesh S. Dentistry and ayurveda v-an evidence based approach. *International Journal of Clinical Dental Science* 2011; 2(1): 3-9.
- Ismail T, Sestili P, Akhtar S. Pomegranate peel and fruit extracts: a review of potential anti-inflammatory and anti-infective effects. *J Ethnopharmacol* 2012; 143(2): 397-405.
- Johanningsmeier SD, Harris GK. Pomegranate as a functional food and nutraceutical source. *Annu Rev Food Sci Technol* 2011; 2: 181-201.
- Haber SL, Joy JK, Largent R. Antioxidant and antiatherogenic effects of pomegranate. *Am J Health Syst Pharm* 2011; 68(14): 1302-5.
- Stowe CB. The effects of pomegranate juice consumption on blood pressure and cardiovascular health. *Complement TherClinPract* 2011; 17(2): 113-5.
- Jurenka JS. Therapeutic applications of pomegranate (*Punicagranatum L.*): a review. *Altern Med Rev* 2008; 13(2): 128-44.
- Li Y, Guo C, Yang J, Wei J, Xu J, Cheng S. Evaluation of antioxidant properties of pomegranate peel extract in comparison with pomegranate pulp extract. *Food Chemistry* 2006; 96(2): 254-60.
- Tezcan F, Gültekin-Ozgiiven M, Diken T, Ozcelik B, Erim FB. Antioxidant activity and total phenolic, organic acid and sugar content in commercial pomegranate juices. *Food Chemistry* 2009; 115(3): 873-7.
- Afaq F, Saleem M, Krueger CG, Reed JD, Mukhtar H. Anthocyanin- and hydrolyzable tannin-rich pomegranate fruit extract modulates MAPK and NF-kappaB pathways and inhibits skin tumorigenesis in CD-1 mice. *Int J Cancer* 2005; 113(3): 423-33.
- Gil MI, Tomas-Barberan FA, Hess-Pierce B, Holcroft DM, Kader AA. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J Agric Food Chem* 2000; 48(10): 4581-9.
- Shukla M, Gupta K, Rasheed Z, Khan KA, Haqqi TM. Bioavailable constituents/metabolites of pomegranate (*Punicagranatum L.*) preferentially inhibit COX2 activity ex vivo and IL-1beta-induced PGE2 production in human chondrocytes in vitro. *J Inflamm (Lond)* 2008; 5: 9.
- Akpınar-Bayizit A, Ozcan T, Yilmaz-Ersan L. The Therapeutic Potential of Pomegranate and Its Products for Prevention of Cancer. In: Georgakilas A, Editor. *Cancer Prevention-From Mechanisms to Translational Benefits*. Winchester, UK: INTECH; 2012.
- Chong MF, Macdonald R, Lovegrove JA. Fruit polyphenols and CVD risk: a review of human intervention studies. *Br J Nutr* 2010; 104 (Suppl 3): S28-S39.
- Aviram M, Dornfeld L. Pomegranate juice consumption inhibits serum angiotensin converting enzyme activity and reduces systolic blood pressure. *Atherosclerosis* 2001; 158(1): 195-8.
- Aviram M, Rosenblat M, Gaitini D, Nitecki S, Hoffman A, Dornfeld L, et al. Pomegranate juice consumption for 3 years by patients with carotid artery stenosis reduces common carotid intima-media thickness, blood pressure and LDL oxidation. *Clin Nutr* 2004; 23(3): 423-33.
- Lynn A, Hamadeh H, Leung WC, Russell JM, Barker ME. Effects of pomegranate juice supplementation on pulse wave velocity and blood pressure in healthy young and middle-aged men and women. *Plant Foods Hum Nutr* 2012; 67(3): 309-14.
- Mathew AS, Capel-Williams GM, Berry SE, Hall WL. Acute effects of pomegranate extract on postprandial lipaemia, vascular function and blood pressure. *Plant Foods Hum Nutr* 2012; 67(4): 351-7.
- Asgary S, Sahebkar A, Afshani MR, Keshvari M, Haghjooyjavanmard S, Rafieian-Kopaei M. Clinical Evaluation of Blood Pressure Lowering, Endothelial Function Improving, Hypolipidemic and Anti-Inflammatory Effects of Pomegranate Juice in Hypertensive Subjects. *Phytother Res* 2013.
- Kelishadi R, Gidding SS, Hashemi M,

- Hashemipour M, Zakerameli A, Poursafa P. Acute and long term effects of grape and pomegranate juice consumption on endothelial dysfunction in pediatric metabolic syndrome. *J Res Med Sci* 2011; 16(3): 245-53.
26. Sumner MD, Elliott-Eller M, Weidner G, Daubenmier JJ, Chew MH, Marlin R, et al. Effects of pomegranate juice consumption on myocardial perfusion in patients with coronary heart disease. *Am J Cardiol* 2005; 96(6): 810-4.
 27. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003; 42(6): 1206-52.
 28. Vasan RS, Larson MG, Leip EP, Kannel WB, Levy D. Assessment of frequency of progression to hypertension in non-hypertensive participants in the Framingham Heart Study: a cohort study. *Lancet* 2001; 358(9294): 1682-6.
 29. Savica V, Bellinghieri G, Kopple JD. The effect of nutrition on blood pressure. *Annu Rev Nutr* 2010; 30: 365-401.
 30. Amr M, El-Mogy A, Shams T, Vieira K, Lakhan SE. Efficacy of vitamin C as an adjunct to fluoxetine therapy in pediatric major depressive disorder: a randomized, double-blind, placebo-controlled pilot study. *Nutr J* 2013; 12: 31.
 31. Lee J, Durst RW, Wrolstad RE. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: collaborative study. *J AOAC Int* 2005; 88(5): 1269-78.
 32. Moohebati M, Bidmeshgi S, Azarpazhooh MR, Daloe MH, Ghayour-Mobarhan M, Tavallaie S, et al. Simvastatin treatment reduces heat shock protein 60, 65, and 70 antibody titers in dyslipidemic patients: A randomized, double-blind, placebo-controlled, cross-over trial. *Clin Biochem* 2011; 44(2-3): 192-7.
 33. Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, et al. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol* 2002; 39(2): 257-65.
 34. Dickinson KM, Keogh JB, Clifton PM. Effects of a low-salt diet on flow-mediated dilatation in humans. *Am J Clin Nutr* 2009; 89(2): 485-90.
 35. Griendling KK, FitzGerald GA. Oxidative stress and cardiovascular injury: Part II: animal and human studies. *Circulation* 2003; 108(17): 2034-40.
 36. Friedman J, Peleg E, Kagan T, Shnizer S, Rosenthal T. Oxidative stress in hypertensive, diabetic, and diabetic hypertensive rats. *Am J Hypertens* 2003; 16(12): 1049-52.
 37. Trollet MR, Rudd MA, Loscalzo J. Oxidative stress and renal dysfunction in salt-sensitive hypertension. *Kidney Blood Press Res* 2001; 24(2): 116-23.
 38. Lerman LO, Nath KA, Rodriguez-Porcel M, Krier JD, Schwartz RS, Napoli C, et al. Increased oxidative stress in experimental renovascular hypertension. *Hypertension* 2001; 37(2 Pt 2): 541-6.
 39. Dobrian AD, Davies MJ, Schriver SD, Lauterio TJ, Prewitt RL. Oxidative stress in a rat model of obesity-induced hypertension. *Hypertension* 2001; 37(2 Pt 2): 554-60.
 40. Laursen JB, Somers M, Kurz S, McCann L, Warnholtz A, Freeman BA, et al. Endothelial regulation of vasomotion in apoE-deficient mice: implications for interactions between peroxynitrite and tetrahydrobiopterin. *Circulation* 2001; 103(9): 1282-8.
 41. Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol* 1996; 271(5 Pt 1): C1424-C1437.
 42. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 1997; 336(16): 1117-24.
 43. Esposito K, Marfella R, Ciotola M, Di PC, Giugliano F, Giugliano G, et al. Effect of a mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA* 2004; 292(12): 1440-6.
 44. Seeram NP, Henning SM, Zhang Y, Suchard M, Li Z, Heber D. Pomegranate juice ellagitannin metabolites are present in human plasma and some persist in urine for up to 48 hours. *J Nutr* 2006; 136(10): 2481-5.
 45. Mohan M, Waghulde H, Kasture S. Effect of pomegranate juice on Angiotensin II-induced hypertension in diabetic Wistar rats. *Phytother Res* 2010; 24(Suppl 2): S196-S203.

How to cite this article: Asgari S, Keshvari M, Sahebkar A, Hashemi M, Rafieian-Kopaei M. **Clinical investigation of the acute effects of pomegranate juice on blood pressure and endothelial function in hypertensive individuals.** *ARYA Atheroscler* 2013; 9(6): 326-31.