

Depression and Demographical Features in Diabetic Patients

Farzaneh Sheikholeslami¹, Rohangiz Norozi Nia², Zinat Alsadat Mirpoor³, Abolazim Tavakoly Vardanjani⁴
Safar Ali Esmaili Vardanjani⁵

¹Faculty member, Guilan University of medical sciences, Guilan, Iran.

²Ms in Nursing, Alborz University of medical sciences, Alborz, Iran.

³Ms Student in psychology, Disable and Elderly Hospice Guilan, Iran.

⁴Ms in Intensive Nursing, Tehran University of Medical Science, Tehran, Iran

⁵Ms in Nursing Education, ShahreKord University of medical sciences, ShahreKord, Iran.

safaraliesmaili@yahoo.com

Abstract: Depression is one of the psychiatric prevalent illnesses across the world. Depression is an illness which has a direct relationship with diabetes. The purpose of determining the level of depression among the diabetic persons referring to Rasht diabetes center. This research is a correlation – descriptive study which was conducted using the simple random sampling method within 3 months on 144 subjects of the social workers afflicted with diabetes referring to Rasht diabetes center in 2008. Data gathering tool was a questionnaire consisting of two parts. The first and second parts of the questionnaire consisted of demographic data and Beck depression standard, respectively. Finally, the data out of this research were analyzed by using the SPSS version 15 statistical software and T-test statistical tests. Results showed that out of 144 subjects, there were 79.86 % female and 20.14% male, 16 people (11.1%) forty years old and younger, 18 people (12.5%) older than 65 years, and 2.1% single and 6.9% divorced. In one hand, in terms of education level, 47.9% and 6.9% were illiterate and educated, respectively. On the other hand, in terms of the illness type, 25 people (17.4%) and 75 people (82.6%) were afflicted with diabetes types 1 and 2, respectively. In addition, statistical tests showed that 35.41% of tester were afflicted with acute depression of which 88.23%, 84.31%, and 84.33% were female, married, and afflicted with diabetes type 2, respectively. It was also shown that there is a significant relationship between depression and gender ($p=0.004$), marital status ($p<0.05$), education level ($p<0.05$), and jobs of the afflicted ($p<0.05$). The present study shows that there is a significant relationship between the depression in people afflicted with diabetes and their gender, marital status, education level, and jobs and also 22.22%, 22.22%, 19.47% and 35.41% of the people were afflicted with weak, low, average, and acute depression.

[Sheikholeslami F, Norozi Nia R, Mirpoor ZS, Tavakoly Vardanjani A, Esmaili Vardanjani SA. **Depression and Demographical Features in Diabetic Patient.** *Life Sci J* 2012;9(4):5542-5547] (ISSN:1097-8135).

<http://www.lifesciencesite.com> 822

Keywords: Depression, Diabetes, Beck test

1. Introduction

One of the problems which threaten human being from the childhood to the life end has to do with Endocrine illnesses among which diabetes is the most prevalent (Fips, 2008). The number of diabetic people has increased to more than 70 million ones during the ten previous years and so World Health Organization (WHO) has called this event a tocsin for developing countries (Shahbazian et al., 2004). Diabetes is one of the prevalent illnesses in human being societies that unfortunately in spite of the medical sciences progresses, not only its prevalence has not decreased, but also it is ever on the rise (Shahbazian et al., 2004). Diabetes is among the medical disorders it can have psychiatric symptoms. Mode disorders with depression manifestation which are made by the presence of common medical problems are equal in both genders, while the acute depression disorder is dominant among the female (Sadock and Sadock, 2007). Diabetes is an illness

which can be treated in spite of its chronicity, in such a way that the person survive and continue his / her life in the case the illness is controlled. Diabetic patients are generally aware of the fact that they are prone to the late coming effects of diabetes and their life time is shortened (Peyrol, 1997).

Recent studies show that depression increases the danger of death in diabetic patients very much (Black, 2003). Depression has since long been mentioned in writings, and Boqrat has also applied the term "hypocondria" (melancholy) to refer to a kind of depression about 450 B.C (Kalpan and sadock, 2007). Depression symptoms consist of fluctuations in weight, appetite, and sleep hours which if are along with the symptoms resulting from the diabetes, then the depression arising from the diabetes is diagnosed clinically (Ludman et al., 2004). Of course, it is worth mentioning that the researches have shown just 50% of the patients afflicted with depression are identified in the initial

stages (Egede, 2004). Therefore, it is not surprising that emotional reactions towards the diabetes often make problems in treatment (Debra, 1996).

Depression is one of the psychiatric prevalent illnesses across the world with which 15 and 25 percents of the men and women in the society are respectively afflicted in their life spans (Fips, 2008). In the united States, 9.5% of the population, or 19 million people in other words, are yearly afflicted with depression (Regier et al., 1993). Depression is a dangerous disease because it leads to suicide, increases the danger of affliction and mortality resulting from the disease. Decreases the life quality, and is known as a cost taking disorder. Studies show that about 20% of the patients who refer to out clinics suffer from depression. Mentally, the first symptom of depression is reduction in motivation and enthusiasm after which reduction in activity and efficiency, hope in life, self-caring, and also tendency to death are emerged in the person (Kaplan and Sadock, 2002). For the first time, the relationship between diabetes and depression was determined by an English physician called Dr. Tomas Vilis in 1684 (Willis, 1971). In another study done by Grandinei et al, On 574 patients indigenous to Havaei region, it was known that the increase in depression prevalence is observed among the diabetic patients (Grandinetti and Kaholokula, 2000). In addition, there is some relationship between depression and blood sugar controlling in diabetic patients (Gory et al., 2000). It has been mentioned in Lostman's study that depression is along with the increase in blood sugar in the patients afflicted with diabetes of types 1 and 2 (Grandinetti and Kaholokula, 2000). In researches conducted by Gardeo et al. in Mexico in 1998 on diabetic patients of type 2, it was concluded that 46% of diabetic persons are afflicted with depression and also being female and time duration of affliction with diabetes are among the danger factors in depression occurrence (Graduno and Teles, 1998). On the whole, depression is one of the mental disorders which has the most accompaniment with diabetes. The issue

whether this relationship has to do with either beginning of progress and deterioration of diabetes or preparation and reaction towards it, is under discussion (Talbot et al., 2000). Accordingly, considering the daily increase in the number of the people afflicted with diabetes and also the role of depression in making and controlling this disease, a study was conducted with the purpose of determining the degree of depression among the diabetic persons referring to Rasht diabetes center.

2. Material and Methods

This research is a correlation – descriptive study which was conducted using the simple random sampling method within 3 months on 144 subjects of the social workers afflicted with diabetes referring to Rasht diabetes center in (2008). Data gathering tool was a questionnaire consisting of two parts. The first and second parts of the questionnaire consisted of demographic data and Beck depression Questionnaire, respectively. Finally, the data out of this research were analyzed by using the SPSS 15 statistical software and T-test statistical tests.

3. Results

Results showed that out of 144 subjects (tested), there were 16 persons (11.1%) forty years old and younger, 18 persons (12.5%) older than 65 years, 66.4% between 41 to 65 years old, 2.1% single, 6.9% divorced, and 91% married. In addition, in terms of education level, 47.9% and 6.9% were illiterate and educated, respectively. On the other hand, in terms of the illness type, 25 people (17.4%) and 75 people (82.6%) were afflicted with diabetes types 1 and 2, respectively. Besides, statistical tests showed that 35.41% of tester were afflicted with acute depression of which 88.23%, 84.31%, 60.78%, 80.39%, and 84.33% were female, married, illiterate, in age interval of 41-65 years old, and afflicted with diabetes type 2, respectively. It was also shown that there is a significant relationship between depression and gender ($p=0.004$), marital status ($p<0.05$), education level ($p<0.05$), and jobs of the afflicted ($p<0.05$). (Table 1)

Table 1: Frequency distribution and relationship between personal identifications and depression in Rasht diabetic patients.

	Percent	Number	Percent	Number	Percent	Number	Percent	Number		
Not Significant	9.80%	5	10.71%	3		0	19.51%	8	40 years old and younger	Age
	80.39%	41	78.57%	22	87.5%	21	63.41%	26	Between 41 to 65 years old	
	9.80%	5	10.71%	3	12.5%	3	17.7%	7	Older than 65 years	
Significant $p=0.004$	88.23%	45	82.14%	23	75%	18	70.73%	29	Female	Gender
	11.76%	6	17.85%	5	25%	6	29.26%	12	Male	
Significant $p<0.005$		0		0	4.16%	1	4.87%	2	Single	Marital Status
	84.31%	43	100%	28	87.5%	21	92.68%	38	Married	
	15.68%	8		0	8.33%	2	2.43%	1	Divorced	
	17.64%	9	10.71%	3	4.16%	1	9.75%	4	Free	

Significant p <0.005	7.84%	4	14.28%	4	16.66%	4	39.02%	16	Employee	Job
		0	3.57%	1	8.33%	2	4.87%	2	Jobless	
	74.50%	38	71.42%	20	70.83%	17	46.34%	19	Housekeeper	
Significant p <0.005	60.78%	31	42.87%	12	50%	12	34.14%	14	Illiterate	Education level
	25.49%	13	46.42%	13	33.33%	8	34.14%	14	Secondary	
	11.76%	6	3.57%	1	8.33%	2	19.51%	8	Diploma	
Not Significant	1.96%	1	7.14%	2	8.33%	2	12.19%	5	University	Living place
	6%	3	7.40%	2	4.16%	1		0	Village	
	94%	47	92.59%	25	95.83%	23	100%	39	City	
Not Significant	29.44%	15	17.85%	5	29.16%	7	21.95%	9	Rental	Living Status
	70.58%	36	82.14%	23	70.83%	17	78.04%	32	Personal	
Not Significant	15.68%	8	17.85%	5	16.66%	4	19.51%	8	Diabetes 1	Type of disease
	84.33%	43	82.14%	23	83.33%	20	80.48%	33	Diabetes 2	

4. Discussions

The present study shows that there is a significant relationship between the depression in people afflicted with diabetes and their gender, marital status, educations level, and jobs. According to the researchers conducted by Sepehrmanesh et al. Regarding the prevalence and types of depression on 300 diabetic persons in 1382, it was shown that men and women were 71% and 57.7% depressed, respectively, and also out of 30% patients afflicted with acute depression, 25 (32%) and 105(47.3%) persons were men and women, respectively. Researches showed that depression in diabetic women was almost 1.4 times as equal as diabetic men. Reference books proclaim acute depression in

women about two times as equal as men (Kaplan H & Sadock, 2000). Reached also to the result in their research in 2001 based on which depression in diabetic women was more seen than in diabetic men (Anderson et al., 2001). Soinkok et al., (2001) noticed in their research that depression in diabetic people has just to do with the female gender (Soinkok et al., 2001). In researches conducted by Gardeo et al in Mexico in 1998 on diabetic patients of type 2, it was concluded that 46% of diabetic persons are afflicted with depression and also being female and time duration of affliction with diabetes are among the danger factors in depression occurrence (Graduno and Teles, 1998).




Figure 1: Distribution of individuals' depression types degrees in terms of education level

In another research conducted by Toziki et al., the sexual distribution of the patients under study was as follows: 35 people (34%) of the women were in the group of the undepressed and the other left 68

people (66%) were depressed. In the men's group as well, there were 23 people (48.9%0 undepressed and 24 ones (51.1%) depressed. The data obtained from the study indicate that depression is not under the

influence of gender. A significant relationship was also observed in our research ($p < 0.05$) (Talbot et al.,

2000). (Figure 2).




Figure 2: Distribution of individuals' depression types degrees in terms of their gender

In the United States, 6% of the people are afflicted with the depression arisen from diabetes. 1.5% of the people between 20 to 40 years old as well as 20% of the people older than 75 years are afflicted with depression whose 90% are among the diabetic type 2. In the present research, results showed that there were 16 people (11.1%) forty years old and younger as well as 18 ones (12.5%) older than 65 years out of the total 144 subjects (tester). In addition, in terms of the illness type, there were 25 (17.4%) and 75 (82.6%) people afflicted with diabetes types 1 and 2, respectively. In addition, statistical tests showed that 35.41% of tester were afflicted with acute depression of which 88.23%, 84.31%, and 84.33% were female, married, and afflicted with diabetes type 2, respectively (Figure 3). According to the psychiatric references, the average age of depression initiation is about 40 years old (Kaplan and Sadock, 1998). In our research as well, the people between 41 to 60 years old made up the most distribution of the diabetic. Some researches show the degree of depression occurrence in the people under 20 years old is on the rise (Talbot et al., 2000). Based on the reference books, the acute

depression is more seen in the people who do not usually have any close interpersonal relationships (Kaplan and Sadock, 1999). Fisher et al (2001) proclaimed in their research that life stresses such as education level are effective on occurring depression in the diabetic as the added and independent agents (Fisher et al., 2001). The married made up the most percentage of patients (84.31%) in the study conducted by us. It is while a high percentage of these people were ranked in low levels in terms of the education (Figures 1 – 4). In the research conducted in a clinic in California, it was seen a significant relationship between diabetes and the participants' employment status as well as between the regression analysis of depression variable and the employment status, respectively (Robert and Nanji, 1986). In addition, it was shown in the study carried by ru-Ling Baia et al, that most of the participants were financially independent ($n=113$; 72.4%), and also the results have not shown a significant difference in terms of the financial status and income level (Yu and Chou, 2008).




Figure 3: Distribution of individuals' depression types degrees in terms of marital status




Figure 4: Distribution of individuals' depression types degrees in terms of their jobs

Acknowledgements:

The authors would like to thank the Dean School of Nursing and Midwifery, Guilan University of medical sciences and all of the patients and clients who participated in our study.

Corresponding Author:

Safar Ali Esmaili Vardanjani
 Ms in Nursing Education,
 ShahreKord University of medical sciences,
 ShahreKord, Iran.
 E-mail: safaraliesmaili@yahoo.com

References

1. Fips, Lngman. Medical Surgical Nursing 2008.
2. Mohamadi Mohamadhasan et al. Rahavard Danesh 2001: 4 (1), pp 25.
3. Shahbazian H, Amani R, Shahbazian H, Shahbazian N, Shahbazian M. What the diabetic patients must beknow. Qom: Darolnashre Eslam press 2004: 10 (in Persian)
4. Sadock BJ, Sadock VA. Kaplan & Sadock's synopsis of psychiatry. 10th ed. Philadelphia: Lipincott Williams & wilkins; 2007
5. Mr. Peyrol levels and risks of depression and anxiently symptoms among diabetic adults:diabetes care 199:20 (4):585-90

6. Black SA; Markides KS; Ray LA. Depression predicts increased incidence of adverse health outcomes in older Mexican Americans with type 2 diabetes. *Diabetes Care* 2003; 26(10):2822–8.
7. Kalpan H sadock synopsis of psychiatry . 8h ed.maryland.williams and wilkins.1998: p: 363-282
8. Ludman EJ; Katon W; Russo J. et al. Depression and diabetes symptom burden. *Gen Hosp Psychiatry* 2004; 26(6):430–6.
9. Egede LE. Diabetes, major depression, and functional disability among US adults. *Diabetes Care* 2004 27(2):421–8.
10. Debra h.g.,management of diabetes mellitus,London,mosby,how to LEADA healthy lifa despite diabetes,ub.spd 1996: p:15
11. Regier DA; Narrow WE; Rae DS; et al. The de facto US mental and addictive disorders service system. Epidemiologic catchment area prospective 1-year prevalence rates of disorders and services. *Arch Gen Psychiatry* 1993; 50(2):85–94.
12. Kaplan H & Sadock Synopsis of psychiatry 9th ed. Baltimore: Williams and Wilkins 2002: 15:534-590.28:822-843
13. Willis T. Diabetes: a medical odyssey. New York: Tuckahoe; 1971
14. Grandinetti a. Kaholokula ik.relationship between depressive symptoms and diabetes among native hawailas psychoneuro endocrinology 2000: 25(3)239-49.
15. Gory.tl. CrumRM. Cooper PL depressive symptoms and metabolic control in African amricans with type 2.Diabetes care 2000: 23(1):23-9
16. Grandinetti a. Kaholokula ik.relationship between depressive symptoms and diabetes among native hawailas psychoneuro endocrinology 2000: 25(3)239-46
17. Graduno-Espinosaj. Teles-zenteno Jf.frequency of depression in pations with diabetes mellitus type 2. 1998: 50(4):287-91
18. Talbot F & et al. A review of the relationship between depression and diabetes in adults: is there a link? *Diabetes Care* 2000: oct; 23 (10): 1556-62.
19. Anderson - RJ & et al. The Prevalence of comorbid depression in adults with diabetes meta analysis - diabetes care 2001: Jun; 24(6): 1069-78.
20. Sevincok L & et al. Depression in a sample of Turkish type 2 diabetes patients, Turkey: *Eur Psychiatry* 2001: jun; 16 (4): 229-31.
21. Kaplan H Sadock B.Comprehensive text book of psychiatry, Philadelphia: LippincottWilliams and wilkins 1999: 1: 1284-1441
22. Fisher L & et al. Contributors to depression in latino and European American patients with type 2 diabetes; *Diabetes Care* 2001: oct; 24 (10) : 1751-70.
23. Robert Friis, G. Nanji Jndappa. DIABETES, DEPRESSION AND EMPLOYMENT STATUS, SW. *Sci. Med* 1996: Vol. 23. No. 5.
24. Yu-Ling Baia, Chou-Ping Chioub,Correlates of depression in type 2 diabetic elderly patients:A correlational study, *International Journal of Nursing Studies* 2008: 45, 571–579

12/21/2012