



NIH PUBLIC ACCESS

Author Manuscript

Nat Neurosci. Author manuscript; available in PMC 2010 January 10.

Published in final edited form as:

Nat Neurosci. 2007 September ; 10(9): 1125–1127. doi:10.1038/nn1946.

Kinase activity is not required for α CaMKII-dependent presynaptic plasticity at CA3-CA1 synapses

Mohammad Reza Hojjati^{1,2}, Geeske M van Woerden¹, William J Tyler^{3,6}, Karl Peter Giese⁴, Alcino J Silva⁵, Lucas Pozzo-Miller³, and Ype Elgersma^{1,5}

¹Department of Neuroscience, Erasmus University Medical Center, Dr. Molewaterplein 50, 3015 GE Rotterdam, The Netherlands. ²Department of Physiology, Shahrekord University of Medical Sciences, Kashani Blvd, Shahrekord 88155, Iran. ³Department of Neurobiology, McKnight Brain Institute, Civitan International Research Center, University of Alabama at Birmingham, Birmingham, Alabama 35294-2182, USA. ⁴Institute of Psychiatry, King's College London, 125 Coldharbour Lane, London SE59NU, UK. ⁵Departments of Neurobiology, Psychiatry and Psychology, Brain Research Institute, 695 Charles Young Drive South, University of California, Los Angeles, California 90095-1761, USA.

Abstract

Using targeted mouse mutants and pharmacologic inhibition of α CaMKII, we demonstrate that the α CaMKII protein, but not its activation, autophosphorylation or its ability to phosphorylate synapsin I, is required for normal short-term presynaptic plasticity. Furthermore, α CaMKII regulates the number of docked vesicles independent of its ability to be activated. These results indicate that α CaMKII has a nonenzymatic role in short-term presynaptic plasticity at hippocampal CA3-CA1 synapses.

The α isoform of Ca^{2+} /calmodulin-dependent protein kinase II (α CaMKII) was originally identified as synapsin I kinase¹. Subsequent studies showed that α CaMKII is abundantly associated with presynaptic vesicles by binding to synapsin I (ref. 2). Together with the observation that α CaMKII is one of the most abundant proteins of the hippocampus³, these results suggest that α CaMKII also has a nonenzymatic function, but such a function has not directly been demonstrated yet. Analysis of α CaMKII knockout (KO) mice confirmed a presynaptic role for α CaMKII in short-term presynaptic plasticity^{4,5}, but these experiments did not address whether this role is mediated by α CaMKII as a kinase, as a structural protein, or as both.

To study the enzymatic requirements of α CaMKII in presynaptic plasticity, we used four different lines of α CaMKII mutants. Autophosphorylation at the threonine 286 (T286) and T305/T306 sites was prevented by using α CaMKII-T286A (T286 is mutated to alanine) mice, which lack α CaMKII autonomous (Ca^{2+} /CaM independent) activity⁶, and α CaMKII-T305V/

© 2007 Nature Publishing Group

Correspondence should be addressed to Y.E. (y.elgersma@erasmusmc.nl).

⁶Present address: Arizona State University, School of Life Sciences, 1711 South Rural Road, Tempe, Arizona 85287, USA.

Note: Supplementary information is available on the Nature Neuroscience website.

COMPETING INTERESTS STATEMENT

The authors declare no competing financial interests.

Published online at <http://www.nature.com/natureneuroscience>

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions>

T306A (T305 and T306 are mutated to valine and alanine, respectively) mice, which lack α CaMKII inhibitory autophosphorylation⁷. Furthermore, we used α CaMKII-T305D (T305 is mutated to an aspartic acid) mice, in which constitutive autophosphorylation at the T305 site in the Ca²⁺/CaM domain is mimicked, preventing α CaMKII in these mice from becoming activated⁷. The fourth line (α CaMKII-KO) lacks the entire α CaMKII protein⁷ (for an overview of all phenotypes see⁸).

Because these mutants were backcrossed in C57BL/6, we first tested whether the originally reported long-term potentiation (LTP) deficits (in hybrid 129Sv/C57BL/6 mice) were still present^{6,7,9}. We confirmed that α CaMKII activation and its subsequent autophosphorylation at T286 are absolute requirements for LTP, but that loss of α CaMKII can partially be compensated for (Supplementary Fig. 1 online). In contrast, loss of inhibitory phosphorylation in α CaMKII-T305V/T306A mice reduced the threshold for LTP induction as reported previously⁷ (Supplementary Fig. 1). Western blots of isolated synaptosomes of all the mutants did not reveal changes in the levels of the β , γ and δ CaMKII isoforms, nor was the amount of calmodulin affected in these mutants (Supplementary Fig. 2 online).

We looked at the ability of these mutants to phosphorylate synapsin I at serine 603 (S603, site 3), which is an exclusive CaMKII site. α CaMKII-KO mice showed a marked decrease of synapsin I phosphorylation compared to wild-type mice ($P < 0.001$, ANOVA, Fig. 1), suggesting that none of the other CaMKII isoforms could efficiently compensate for the loss of α CaMKII phosphorylation of synapsin I *in vivo*. Notably, steady-state levels of phosphorylated synapsin I were not affected in α CaMKII-T286A ($P = 0.2$) mice and in α CaMKII-T305V/306A mice ($P = 0.3$), indicating that loss of autonomous activity or self-inhibition does not have a large effect on synapsin I phosphorylation *in vivo*. In contrast, activation of α CaMKII is an absolute requirement for synapsin I phosphorylation, as S603 phosphorylation in the α CaMKII-T305D mutant was not significantly above background level ($P = 0.2$; Fig. 1a). The dominant-negative nature of the α CaMKII-T305D mutation was further illustrated by the fact that this was also the only mutant in which autophosphorylation at both α CaMKII-T286 and β CaMKII-T287 was indistinguishable from background level (both $P > 0.8$; Fig. 1b), further suggesting that α CaMKII is inactive in this mutant. Taken together, these results show that these mutants provide an ideal tool for dissecting the requirements of α CaMKII activation, α CaMKII autophosphorylation and synapsin I S603 phosphorylation in short-term presynaptic plasticity.

Previous studies from multiple laboratories^{4,5,8}, using independently generated targeted deletions of α CaMKII^{5,9}, have demonstrated that the loss of α CaMKII results in enhanced augmentation and decreased synaptic fatigue, which are both measures of presynaptic plasticity. Augmentation is an increase in the evoked postsynaptic response that is observed several seconds after high-frequency afferent stimulation, caused by facilitated exocytosis¹⁰. We measured augmentation at hippocampal CA3-CA1 synapses using extracellular field recording (Supplementary Methods online). α CaMKII-KO mice showed a striking increase in synaptic augmentation (during 3–11 s: $F_{4,92} = 13$, $P < 0.0001$, repeated measures ANOVA; Fig. 2), confirming that α CaMKII critically regulates this form of presynaptic plasticity⁴. Activation of presynaptic α CaMKII may be important in synaptic augmentation, as transiently elevated presynaptic calcium is thought to be a major factor underlying this process. However, although α CaMKII activity is regulated by autophosphorylation, augmentation was not affected in the T286A and T305V/T306A autophosphorylation-deficient mutants (Fig. 2c). The lack of a phenotype in these mutants could reflect the short-term nature of this kind of plasticity, and/or the fact that phosphorylation of synapsin I S603 was unaffected in these mutants (Fig. 1). Unexpectedly however, augmentation was also unaffected in α CaMKII-T305D mutants ($P = 0.5$, ANOVA; Fig. 2b,c), where α CaMKII activation was blocked and synapsin I phosphorylation was absent (Fig. 1). These results strongly suggest that synaptic

augmentation does not depend on the ability of α CaMKII to phosphorylate synapsin I, nor on the activation of α CaMKII.

Because activation of both α CaMKII and β CaMKII seem to be impaired in the α CaMKII-T305D mutant (Fig. 1b), we would expect that a similar result should be obtained if the augmentation experiment is carried out in the presence of the membrane-permeable CaMKII inhibitor KN-93 (ref. 11). This inhibitor competes with Ca^{2+} /calmodulin binding and therefore prevents activation of both α CaMKII and β CaMKII. LTP experiments in the presence of this inhibitor showed that KN-93 was indeed able to block LTP (Supplementary Fig. 1). In contrast, we observed similar levels of augmentation in the presence of KN-93 or its inactive analog KN-92 ($P = 0.5$, ANOVA; Fig. 2c), confirming that CaMKII activation is not required for normal augmentation.

Our biochemical analyses (Fig. 1) indicated that CaMKII activity in the α CaMKII-T305D mutant was reduced to such an extent that phosphorylation of CaMKII substrates was undetectable. If CaMKII activity is indeed not required for augmentation, the potent CaMKII inhibitor AIP-II (autocamtide-2-related inhibitory peptide II) should not affect augmentation either. This inhibitor is 500 times more potent than KN-93, and it is noncompetitive for Ca^{2+} /calmodulin and exogenous substrates, thus also blocking basal and autonomously active CaMKII¹². Efficient penetration was achieved by making use of AIP that was fused to the antennapedia transport peptide, and by preincubating slices for 1 h with AIP (see Supplementary Methods). Indeed, LTP was fully blocked, indicating that the drug was able to penetrate the slice (Supplementary Fig. 1), and notably, Ant-AIP-II showed no discernable effect on synaptic transmission (Supplementary Fig. 3 online), which makes it suitable for use in these experiments. However, like KN-93, Ant-AIP-II did not affect augmentation ($P = 0.1$, ANOVA; Fig. 2c). Taken together, these results indicate that α CaMKII protein, but not α CaMKII activity, is required for normal augmentation.

Previous whole-cell patch-clamp recordings in CA3-restricted α CaMKII-KO mice showed a substantial enhancement of the excitatory postsynaptic current amplitude during repetitive stimulation of CA3-CA1 synapses⁵, suggesting that the fatigue rate of neurotransmitter release is regulated by α CaMKII. Likewise, extracellular field recordings from our global α CaMKII-KO mice also demonstrated reduced synaptic fatigue (Fig. 2d). The responses to repetitive 10-Hz stimulation reveal the competing processes of facilitation, vesicle depletion and vesicle mobilization¹⁰. α CaMKII-KO mutants had a similarly shaped curve as wild-type mice; however, there was a significant effect of genotype ($F_{1,45} = 4.5$, $P < 0.05$) and a significant interaction between genotype and stimulus number ($F_{99,4455} = 4.5$, $P < 0.0001$). α CaMKII-KO mutants were only significantly different from wild-type mice after 20 stimuli (first 20: $F_{19,855} = 1.2$, $P = 0.3$; last 80: $F_{79,3555} = 2.6$, $P < 0.001$), suggesting a differential rate of vesicle depletion and/or mobilization, the cellular processes primarily responsible for the maintenance of excitatory postsynaptic potential (EPSP) amplitude during prolonged stimulation¹⁰.

To test the α CaMKII autophosphorylation and synapsin I phosphorylation requirements in this presynaptic measure, we repeated this experiment in the α CaMKII point mutants. No significant effect of genotype was observed in either α CaMKII-T286A or α CaMKII-T305V/T306A autophosphorylation-defective mutants (both $P > 0.3$, ANOVA at stimulus number 100; Fig. 2f). Notably, the depletion rate was also not affected in α CaMKII-T305D mutants ($P > 0.7$; Fig. 2e,f), in which activation of α CaMKII was prevented and phosphorylation of synapsin I was absent. In addition, there was no discernable effect on depression in slices treated with KN-93 or Ant-AIP-II ($P = 0.9$ and $P = 0.7$, respectively; Fig. 2f). Taken together, these data strongly suggest that the α CaMKII protein plays a structural role rather than an enzymatic role, during this form of short-term presynaptic plasticity.

Mechanistically, enhanced synaptic augmentation and reduced synaptic depression could reflect changes in the pool sizes of the synaptic vesicles, in particular the size of the readily releasable pool (RRP). Therefore, we used electron microscopy to measure the number of docked vesicles, a morphological correlate of the RRP^{13,14}. We obtained measurements at the active zone of excitatory synapses on CA1 spines of wild-type, α CaMKII-KO and α CaMKII-T305D mice. Indeed, there was a significant effect of genotype ($F_{2,41} = 4.7$, $P < 0.05$), with synapses in α CaMKII-KO mice showing a 20% increase in the total number of docked vesicles compared with wild-type (Fisher's PLSD, $P < 0.05$) or α CaMKII-T305D mice ($P < 0.01$) (Fig. 3a). In contrast, no significant difference in the number of docked vesicles was observed between the α CaMKII-T305D mutant and wild-type mice (Fisher's PLSD, $P = 0.24$). Additional measurements of the number of reserve pool vesicles, active zone length and presynaptic terminal area were similar between the mutants and wild-type mice (all measures $P > 0.2$; Supplementary Fig. 4 online). Thus, the absence of α CaMKII protein results in an increased number of docked vesicles, whereas the loss of α CaMKII activation and synapsin I S603 phosphorylation does not affect vesicle docking.

Together, these results suggest a model in which α CaMKII functions nonenzymatically to limit the size of the RRP, thereby modulating short-term presynaptic plasticity. If the observed increase in the size of EPSPs during repetitive stimulation in α CaMKII-KO mice is indeed mediated by a larger RRP, then presynaptic function should be normal under conditions that minimize depletion of the RRP. Accordingly, we decreased the extracellular calcium concentration, which limits the rate of depletion from the RRP¹⁵. Decreasing the extracellular calcium concentration from 2.5 to 0.8 mM normalized the responses during 10-Hz stimulation in α CaMKII-KO mice ($F_{1,31} < 0.0001$, $P = 1$, Fig. 3b), supporting the idea that α CaMKII limits the available pool of readily-releasable neurotransmitter vesicles.

Taken together, our experiments suggest that synapsin I S603 phosphorylation and α CaMKII activity are not required for short-term presynaptic plasticity measures such as augmentation and synaptic fatigue during repetitive stimulation. Specifically, we have demonstrated that, at hippocampal CA3-CA1 synapses, α CaMKII functions independently of its kinase activity to modulate short-term presynaptic plasticity by limiting the number of presynaptic docked neurotransmitter vesicles.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

We greatly appreciate the help of H. Beck, M. Merkens, R. Anwyl, K. Wang and R. Colbran for their (hands-on) advice on the use of CaMKII inhibitors, and from G. Borst and S. Kushner and members of the Silva and Elgersma lab for stimulating discussions and critically reading the manuscript. We thank M. Elgersma, H. van der Burg and E. Phillips for technical support. This work was supported by grants from NWO-ZonMW (TOP, VIDI) and Neuro-BSIK to Y.E. We also thank the University of Alabama, Birmingham Neuroscience Cores (P30-NS47466, P30-HD38985, P30-NS57098).


References

1. Bennett MK, Erondou NE, Kennedy MB. *J. Biol. Chem* 1983;258:12735–12744. [PubMed: 6313675]
2. Benfenati F, et al. *Nature* 1992;359:417–420. [PubMed: 1328883]
3. Erondou NE, Kennedy MB. *J. Neurosci* 1985;5:3270–3277. [PubMed: 4078628]
4. Chapman PF, Frenguelli BG, Smith A, Chen CM, Silva AJ. *Neuron* 1995;14:591–597. [PubMed: 7695905]

5. Hinds HL, Goussakov I, Nakazawa K, Tonegawa S, Bolshakov VY. *Proc. Natl. Acad. Sci. USA* 2003;100:4275–4280. [PubMed: 12629219]
6. Giese KP, Fedorov NB, Filipkowski RK, Silva AJ. *Science* 1998;279:870–873. [PubMed: 9452388]
7. Elgersma Y, et al. *Neuron* 2002;36:493–505. [PubMed: 12408851]
8. Elgersma Y, Sweatt JD, Giese KP. *J. Neurosci* 2004;24:8410–8415. [PubMed: 15456813]
9. Silva AJ, Stevens CF, Tonegawa S, Wang Y. *Science* 1992;257:201–206. [PubMed: 1378648]
10. Zucker RS, Regehr WG. *Annu. Rev. Physiol* 2002;64:355–405. [PubMed: 11826273]
11. Sumi M, et al. *Biochem. Biophys. Res. Commun* 1991;181:968–975. [PubMed: 1662507]
12. Ishida A, Kameshita I, Okuno S, Kitani T, Fujisawa H. *Biochem. Biophys. Res. Commun* 1995;212:806–812. [PubMed: 7626114]
13. Schikorski T, Stevens CF. *Nat. Neurosci* 2001;4:391–395. [PubMed: 11276229]
14. Tyler WJ, Pozzo-Miller LD. *J. Neurosci* 2001;21:4249–4258. [PubMed: 11404410]
15. Borst JG, Sakmann B. *Nature* 1996;383:431–434. [PubMed: 8837774]

**Figure 1.**

Phosphorylation of synapsin I and CaMKII-T286/T287 in synaptosomes obtained from α CaMKII mutants. **(a)** Phosphorylation of synapsin I S603 was not affected by impaired α CaMKII autophosphorylation, but required α CaMKII protein and its activation by Ca^{2+} /calmodulin. **(b)** Phosphorylation of α CaMKII-T286 and β CaMKII-T287 was absent in α CaMKII-T305D mice. Graph represents data from β CaMKII-T287 only. Error bars indicate s.e.m. Each sample contains pooled fractions from four independent isolations.

**Figure 2.**

Presynaptic short-term plasticity requires α CaMKII protein, but not its autophosphorylation, activation or activity. (**a–c**) Increased synaptic augmentation in α CaMKII-KO mutant mice was not caused by the lack of CaMKII kinase activity. fEPSP responses (normalized to pretetanus baseline) of CaMKII-KO (**a**) and CaMKII-T305D (**b**) mice were recorded at the indicated time after a 10 theta-burst tetanus. Traces are from baseline response (gray) and the response 5 s post-tetanzation (black). Augmentation summary of responses obtained 5 s post-tetanus normalized to baseline is shown in **c**. Black bars represent mutants or drug-treated slices, white bars represent control slices as indicated. (**d–f**) Decreased synaptic fatigue during repetitive stimulation in α CaMKII-KO mice was not caused by the lack of α CaMKII kinase activity. (**d,e**) fEPSP responses (normalized against baseline) of CaMKII-KO (**d**) and CaMKII-T305D (**e**) mice were recorded during a 10-Hz tetanus. Only the first and even numbered stimuli are shown for clarity. Traces are from wild-type (gray) and α CaMKII-KO slices (black) recorded from stimulus number 21–30. Depletion summary of the last (100) stimulus of the 10-Hz train is shown in **f**. Black bars represent mutant or drug-treated slices, normalized against the controls as indicated (white bars, set at 100%). Numbers between brackets indicate the number of slices. Error bars indicate s.e.m.




Figure 3. α CaMKII protein regulates the number of docked vesicles. **(a)** Quantitative electron microscopy of asymmetric synapses on dendritic spines of CA1 pyramidal neurons showed a 20% increase in the number of docked vesicles in α CaMKII-KO mice. **(b)** Decreasing the depletion rate by lowering extracellular calcium reversed the phenotype of the α CaMKII-KO mice during repetitive stimulation. Error bars indicate s.e.m.