



Identification of chemical compounds of *Nardostachys Jatamansi* essence available in Iran

Nasrolah Ghassemi-Dehkordi¹, Ebrahim Sajjadi¹, Hamed Shafiei-Koojani^{1*}, Mahtab Keshvari², Seyyed-Masih Hoseini³

¹Department of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

²Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

³Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

ARTICLE INFO

Article Type:
Original Article

Article History:
Received: 12 July 2014
Accepted: 3 October 2014
ePublished: 1 December 2014

Keywords:
Essence
Nardostachys Jatamansi
Iran
Retention index

ABSTRACT

Introduction: With regard to using drugs with plant origin and with the aim of suitable use of these types of drugs and preventing them to be abused, it is necessary to determine the standards of these plants. The aim of the present study was to identify and study chemical compounds of *Nardostachys Jatamansi* essence in Iran and define monograph of this plant for the Iranian plant pharmacopeia.

Methods: In an experimental study the *Nardostachys Jatamansi* specimen was prepared from the market in Iran. The essence of the plant was prepared by the hydro-distillation in *Clevenger apparatus*. Essence was obtained as a greenish yellow oil layer with the 0.07 % yield. The essence compounds were identified quantitatively by Gas Chromatography-Mass Spectroscopy (GC/MS) method.

Results: Totally, 29 compounds were identified in *Nardostachys Jatamansi* essence. The retention indexes (RI) were only similar with overall standard values in two compounds like mesitylene and P-cymene. In this line the RI values about three compounds of valerenic acid, palmitic acid, and valeranyl isovalerate were determined significantly higher than standard values of RI.

Conclusion: The essence prepared from the *Nardostachys Jatamansi* plant in Iran was different in terms of some compounds and components including valerenic acid, palmitic acid, and valeranyl isovalerate and so it is necessary to identify and register quality and quantity characteristics of compounds available in this plant in the Iranian medicinal plants pharmacopeia.

Implication for health policy/practice/research/medical education:

The essence prepared from *Nardostachys Jatamansi* plant in Iran seems to be different in terms of some compounds and components. So it is necessary to identify and register quality and quantity characteristics of compounds available in this plant in the Iranian medicinal plants pharmacopeia.

Please cite this paper as: Ghassemi-Dehkordi N, Sajjadi E, Shafiei-Koojani H, Mahtab Keshvari, Hoseini SM. Identification of chemical compounds of *Nardostachys Jatamansi* essence available in Iran. J HerbMed Pharmacol. 2014; 3(2):83-86.

Introduction

Today although most of consumed drugs are synthetic, but it has been estimated that at least one third of all medicinal products have plant origin or have been transformed after they are extracted from the plants. Using medicinal plants for diseases has an old history (1). Medicinal plants have less side effects and risks compared to chemical drugs due to having natural origin and also contiguity

and compatibility with the human body physiology. This property is one of the main reasons of returned approach and tendency of people toward medicinal plants and using it compared to chemical drugs (1,2). Therefore, pharmaceutical industries and research groups in many countries have focused their attention on cultivating and producing medicinal plants (2). Citronella or Indian nard (with the scientific name of *Valeriana sisymbriifolium*

*Corresponding author: Hamed Shafiei-Koojani,
Email: hamedshafiei1366@gmail.com

from Valerianaceae family) is a plant that its height reaches to 10-60 cm and mainly grows in summer. This plant has been used as an herbal medicine from the sixteen century (3). It has been used traditionally in different regions of Europe and Asia and today it is accepted as a medicinal plant with various benefits even in North America (4). Accordingly, the *Valeriana sisymbriifolia* has been paid attention by experts in pharmacology and also in the traditional and plant medicine and is used as a medicinal or food complement (5). According to clinical observations, the extract prepared from root of this plant has sedative property and so it is used as an anti-anxiety drug (6-8). Due to the antioxidant property of the extract of this plant, its inhibitive effects on the brain ischemia in animal models of study have been confirmed (9). Also antibacterial effects of this plant in preventing and treating infectious diseases have been proved (10). With regard to the fact that it is economical to use a medicinal plant with high effective materials in pharmaceutical industries, identification of chemical compounds present in plants essence and their effect is very important (11). It is fundamentally possible to understand the drug effect and also to determine effective mechanisms of the plant extract based on identifying compounds and components of the plant extract. Some main extract components of the plant root of *Valeriana sisymbriifolium* include: alkaloids (actinidin, creatinine, valerianine, and valerine), isoalderamide, gamma amino butiric acid (GABA), isoalderic acid, iridoids (isoaltrate).

In the present study the aim was to identify quality and quantity of chemicals in the plant essence of Iran.

Materials and Methods

In a experimental study, *Valeriana sisymbriifolia* specimen was prepared from the Iran market. The dried plant was grinded by the crusher device. Then the essence was collected by the Hydro-distillation method in a Clevenger apparatus for 3 hours. Essence weight ratio of the plant was measured to be 0.07 w/w. The essence was maintained in colored glasses with closed lid in the refrigerator. Compounds available in the essence were determined by the a Gas chromatography device connected to the GC/MS (Hewlett, Packard 5792A, USA), with the ion-trap system, column with the length of 60 m and the diameter of 0.25 mm that the static phase thickness was 0.25 micrometer. The column temperature was set at 60 to 280 °C with gradient of 4 °C per minute, the injection chamber temperature of 280 °C and the ion resource temperature of 250 °C. The helium gas with the purity degree of 99/999, with gas speed of 2 ml/min and the PSI pressure of 17.7 and Electron ionization energy of 70 eV were used.

Results

The greenish yellow essence with the 0.07% yield in terms of the weight (w/w) was obtained from this plant. Totally

29 compounds were extracted in evaluating different components of the *Valeriana sisymbriifolia* extract that retention time values of these compounds were variable between 4.96 related to mesitylene and 34.32 related to valereryl isovalerate. In result of comparing the RI of each of the *Valeriana sisymbriifolia* essence components in Iran with standard cases (Table 1), It was determined in this study that the RI values were similar to world standard values in only two compounds including mesitylene and p-cymene. In this line, the RI values were a little higher from standard values about some compounds including n-octanol, linalool, naphthalene, α -terpineole, cuminal, carvone, trans-anethole, thymole, carvacrol and valeranal in the compound presented in our study. However, the RI value was determined significantly higher than the standard values of RI about some compounds like valerenic acid, palmitic acid and valereryl isovalerate.

Discussion

In evaluating compounds of the *Valeriana sisymbriifolia* essence and materials by the GC/MS method the index of retention time is used. However, this index is potentially affected by other variables and conditions like temperature, chromatography column length and the column diameter, so other index named retention index or RI was used. In this study based on determining the RI index about each component of *Valeriana sisymbriifolia* and comparing it with standard RI values we found that the main difference between compounds presented in our study and cases mentioned in standard pharmacopeias was mainly in three compounds of valerenic acid, palmitic acid and valereryl isovalerate that RI related to each three cases was determined significantly higher than standard values. In other words, *Valeriana sisymbriifolia* specimens in Iran can be differentiated from other specimens in other countries in terms of values of the above three compounds.

Valerenic acid has been identified as GABA-A receptor agonist that plays an important role as a sedative. Also, its role as an anti-inflammation is also proposed (14,15). About palmitic acid also recent researches have suggested sedative effects of this fatty acid and also its effect on modulating glucose metabolism (16). Based on above findings and with regard to obvious differences in above compounds between Iranian specimens and standard in the *Valeriana sisymbriifolia* essence, the therapeutic effects of this essence in Iranian specimens can be different from specimens in other points and so it is necessary to add mentioned characteristics in pharmacopeia of Iranian medicinal plants. Totally, it can be concluded that the essence prepared from the *Valeriana sisymbriifolia* plant in Iran is different in terms of amount of some compounds like valerenic acid, palmitic acid and valereryl isovalerate and with regard to clinical effects of these components specially sedative effects, anti-inflammation and also its effects in regulating the glucose metabolism, it is necessary

Table 1. Comparing the RI of each of the *Valeriana sisymbriifolia* essence components in Iran with standard cases

| No | RT | Compound | % | RI | Book-R |
|----|-------|-----------------------------------|-----|------|--------|
| 1 | 4.96 | mesitylene | 0.1 | 994 | 994 |
| 2 | 5.66 | p-cymene | 0.2 | 1026 | 1026 |
| 3 | 6.84 | n-octanol | 0.1 | 1072 | 1070 |
| 4 | 7.65 | linalool | 0.4 | 1099 | 1098 |
| 5 | 10.04 | Naphthalene | 0.7 | 1181 | 1179 |
| 6 | 10.33 | α -terpineole | 0.2 | 1190 | 1189 |
| 7 | 11.84 | Cuminal | 0.6 | 1240 | 1239 |
| 8 | 11.97 | carvone | 0.3 | 1244 | 1243 |
| 9 | 13.27 | Trans-anethole | 0.9 | 1284 | 1283 |
| 10 | 13.58 | thymol | 1.0 | 1292 | 1290 |
| 11 | 13.89 | Carvacrol | 1.1 | 1302 | 1298 |
| 12 | 16.19 | β -patchoulene | 0.5 | 1377 | 1380 |
| 13 | 17.10 | α -gurjunene | 0.2 | 1405 | 1410 |
| 14 | 17.75 | β - gurjunene | 0.6 | 1428 | 1434 |
| 15 | 18.02 | aromadendrene | 0.5 | 1437 | 1441 |
| 16 | 18.44 | α - patchoulene | 0.4 | 1451 | 1456 |
| 17 | 18.61 | allo-aromadendrene | 0.8 | 1457 | 1461 |
| 18 | 19.60 | valencene | 0.3 | 1489 | 1491 |
| 19 | 20.31 | 7- epi- α -selinene | 0.4 | 1513 | 1517 |
| 20 | 21.04 | α -calacorene | 0.2 | 1539 | 1542 |
| 21 | 22.18 | Caryophyllene oxide | 3.9 | 1578 | 1583 |
| 22 | 22.31 | Globulol | 1.5 | 1582 | 1585 |
| 23 | 24.37 | Patchouli alcohol | 6.1 | 1656 | 1658 |
| 24 | 24.75 | 14-hydroxy-9-epi-(E) caryophyllen | 5.2 | 1669 | 1670 |
| 25 | 24.88 | valeranone | 6.0 | 1674 | 1675 |
| 26 | 26.07 | Valerenal | 3.4 | 1717 | 1715 |
| 27 | 30.15 | Valerenic acid | 0.9 | 1880 | 1862 |
| 28 | 32.44 | Palmitic acid | 0.4 | 1976 | 1969 |
| 29 | 34.32 | Valerenyl isovalerate | 0.2 | 2058 | 2052 |

Retention indices on HP-5 capillary column t: trace ($\leq 0.05\%$), %: calculated from TIC data

to identify and register qualitative and quantitative properties of *Valeriana sisymbriifolia* components in Iranian medicinal plant pharmacopeia.

Authors' contributions

All the authors wrote the manuscript equally.

Conflict of interests

The authors declared no competing interests.

Ethical considerations

Ethical issues (including plagiarism, misconduct, data fabrication, falsification, double publication or submission, redundancy) have been completely observed by the authors.

Funding/Support

None.

References

1. Sewell RD, Rafieian-Kopaei M. The history and ups and downs of herbal medicine usage. *J HerbMed Pharmacol* 2014; 3(1):1-3.
2. Nasri H, Shirzad H. Toxicity and safety of medicinal plants. *J HerbMed Pharmacol* 2013; 2(2): 21-2.
3. Houghton PJ. The scientific basis for the reputed activity of Valerian. *J Pharm Pharmacol* 1999; 51:505–12.
4. Hendriks H, Bos R, Allersma DP, Malingre TM, Koster AS. Pharmacological screening of valeranal and some other components of essential oil of *Valeriana officinalis*. *Planta Med* 1981; 42:62–8.
5. Ortiz JG, Nieves-Natal J, Chavez P. Effects of *Valeriana officinalis* extracts on [3H] flunitrazepam binding, synaptosomal [3H] GABA uptake, and hippocampal [3H] GABA release. *Neurochem Res* 1999; 24:1373–8.
6. Leathwood PD, Chauffard F, Heck E, Munoz-Box R.

- Aqueous extract of valerian root (*Valeriana officinalis* L.) improves sleep quality in man. *Pharmacol Biochem Behav* 1982; 17:65–71.
7. Vorbach EU, Gortelmeyer R, Bruning J. Therapy of insomnia. The efficacy and tolerability of valerian. *Psychopharmakotherapie* 1996; 3:109–15.
 8. Leathwood PD, Chauffard F. Quantifying the effects of mild sedatives. *J Psychiatr Res* 1982; 83:17:115–22.
 9. Ortiz JG, Nieves-Natal J, Chaves P. Effects of *Valeriana officinalis* extracts on [3H] flunitrazepam binding, synaptosomal [3H] GABA uptake, and hippocampal [3H] GABA release. *Neurochem Res* 1999; 24:1373–8.
 10. Watanabe K, Takatsuki H, Sonoda M, Tamura S, Murakami N, Kobayashi N. Anti-influenza A effects of novel nuclear export inhibitors from *Valeriana Radix* and *Alpinia galanga*. *Drug Discoveries and Therapeutics* 2011; 5(1):26–31.
 11. Radford AE. Quantitation analysis of polysaccharids and glycoprotein fractions in *Echinacea purpurea* and *Echinacea anngustifolia* by HPLC-ELSD for quality control of raw material. *J Pharmacol Biomed* 2007; 45(4): 115-20.
 12. Das J, Mao AA, Handique PJ. Terpenoid compositions and antioxidant activities of two Indian valerian oils from the Khasi Hills of north-east. *India Nat Prod Commun* 2011; 6(1):129-32.
 13. Verma RS, Verma RK, Padalia RC, Chauhan A, Singh A, Singh HP. Chemical diversity in the essential oil of Indian valerian (*Valeriana jatamansi* Jones). *Chem Biodivers* 2011; 8(10):1921-9. doi: 10.1002/cbdv.201100059.
 14. Yuan CS, Mehendale S, Xiao Y, Aung HH, Xie, JT, Ang-Lee, MK. The gamma-aminobutyric acid effects of valerian and valerenic acid on rat brainstem neuronal activity. *Anesth Analg* 2004; 98(2): 353–8.
 15. Khom S, Baburin I, Timin E, Hohaus A, Trauner G, Kopp B, *et al.* Valerenic acid potentiates and inhibits GABAA receptors: Molecular mechanism and subunit specificity. *Neuropharmacology* 2007; 53 (1): 178–87.
 16. Benoit SC, Kemp CJ, Elias CF, Abplanalp W, Herman JB, Migrenne S, *et al.* Palmitic acid mediates hypothalamic insulin resistance by altering PKC- θ subcellular localization in rodents. *J Clin Invest* 2009; 119(9):2577–87