

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/237042430>

Effect of exercise training on health-related physical fitness factors and blood lipids profile of former addicted persons

Article · January 2010

CITATIONS

2

READS

36

5 authors, including:



Mohammad Faramarzi

Shahrekord University

38 PUBLICATIONS 52 CITATIONS

SEE PROFILE



Ebrahim Banitalebi

Tarbiat Modares University

15 PUBLICATIONS 13 CITATIONS

SEE PROFILE



Reza Nuri

26 PUBLICATIONS 74 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Effect of aerobic exercise on leptin and ghrelin in patients with colorectal cancer [View project](#)



The relationship between obesity indexes with vascular endothelial function after a period of circuit combined training in older women [View project](#)

ORIGINAL PAPER (ARTIGO ORIGINAL)

EFFECT OF EXERCISE TRAINING ON HEALTH-RELATED PHYSICAL FITNESS FACTORS AND BLOOD LIPIDS PROFILE OF FORMER ADDICTED PERSONS

Ebrahim Banitalebi¹, Mohammad Faramarzi², Reza Nuri², Jahanbakhsh Khosrozadeh³, Mohsen Ghafoorian³

¹ Islamic Azad University, Shahrekord Branch. Shahrekord, Iran.

² Department of Physical Education and Sport Sciences, Faculty of Human Sciences, Shahrekord University, Shahrekord, Iran.

³ Sama organization (affiliated with Islamic Azad university) - shiraz branch, Shiraz, Iran.

Corresponding author:

Ebrahim Banitalebi

Islamic Azad University, Shahrekord Branch. Shahrekord, Iran

Email: Ebi_wushu58@yahoo.com

Tel: +989132818216

Fax: +983814424241

Submitted for publication: Apr 2010

Accepted for publication: Aug 2010

ABSTRACT

BANITALEBI, E.; FARAMARZI, M.; NURI, R.; KHOSROZADEH, J.; GHAFOORIAN, M. Effect of exercise training on health-related physical fitness factors and blood lipids profile of former addicted persons. *Brazilian Journal of Biomotricity*, v. 4, n. 3, p. 190-197, 2010. Dysfunctional eating patterns and excessive weight gains have been observed during recovery from drug and alcohol addictions. The purpose of this study was to determine the effect of exercise training on health-related physical fitness factors and blood lipids profile of former addicted persons. Thirty seven males who were 23-49 years old, and had one-year quitting history were selected and randomized (exercise group, n= 18 and control, n= 19). Thirty eight individuals completed the entire study; 16 persons were in exercise group and 15 persons were in control group. Exercise training was consisted primarily of some game-based aerobic exercise. Exercise training duration progressed from 20 minutes at the baseline to 45 minutes at the end of weeks 12th, and intensity of exercise progressed from 50% of heart rate reserve of baseline to 70 % at 12 weeks. Weight, BMI and WHR were measured. Muscle endurance, flexibility and Vo₂Peak were measured using by pull up, Sit -and -Rich test and one-mile Rockport walk test, respectively. Body composition was assessed using the sum of three skin-fold measurement specific for males (chest, abdomen, and thigh). Total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C) and triglyceride (TG) were measured enzymatically using diagnostic Pars kits. All variables were measured at baseline. Data analyzed by using ANCOVA analysis. There were no significant differences in weight (p=0.208), BMI (P=0.2631), CT (P=0.428), HDL (0.833), LDL (0.396), VLDL (P=0.169), TG (P=0.283), Vo₂peak (p=0.884), flexibility (P=0.923) and Pull-up (P=0.44) after 12 weeks exercise training between two groups, but there was significant difference in WHR (p=0.044). It appears that, exercise training can prevent weight gain after quitting drugs and substances.

Key words: Exercise training, Heart risk factor, Lipids, Quit drug.



INTRODUCTION

There is a meaningful increase in the number of persons admitted to recovery programs for drug abuse. Overweight and dysfunctional eating patterns have been observed among people in recovery from drug and alcohol addictions. Six progressive recovery stages has been suggested: (1) transition or pretreatment phase, in which the addicted individual recognizes the addiction problem; (2) stabilization, marked by detoxification and recuperation from acute withdrawal; (3) early recovery, in which the compulsion to use chemicals is reduced and the sober individual becomes more comfortable with being abstinent; (4) middle recovery, in which the individual begins to experience a balance and productive daily life; (5) late recovery, in which the addicted individual begins to move towards healthy living; and (6) maintenance, marked by an awareness for continued growth and development (COWAN and DEVINE, 2008). This study emphasized the differences in weight changes, meaningful use of food, and eating behaviors at different stages of recovery and the opportunities for nutrition interventions at each stage. The excessive weight gain that usually occurs in early recovery and the struggles to lose weight later on in the recovery process further highlight the importance of early nutrition and weight interventions in drug and alcohol treatment facilities. Clark et al (2006) illustrated that approximately half of females who are seeking smoking cessation treatment are weight concerned and that one quarter of male smokers are weight concerned (CLARK et al, 2006). Levine et al (2001) reported that although weight-concerned women smokers expected to gain large amounts of weight after quitting, they expressed a willingness to tolerate only minimal weight gain (LEVINE et al, 2001). The discrepancy between expected and tolerable weight gain may undermine efforts to quit smoking in this group of women.

Abrantes (2006) illustrated that there is some concern that substance abuse treatment may actually pose a risk toward unhealthy eating and weight gain. In a study examining the relationship between supervised drug abstinence and increased weight gain among 215 adolescents in a residential substance abuse treatment center, Hodgkins and colleagues (2004) found that neurotransmitter release in the nucleus accumbens use has been linked to self-administration and learning following drug use. This endogenous reward system is also activated following food intake or sex. Therefore, rebound hyperphagia following abstinence may be a mechanism to replenish the release of neurotransmitters in this reward system, leading to increased weight gain and a rise in Body Mass Index during recovery from substance abuse.

One area that has not been sufficiently explored among currently available adolescent drug use treatment programs is the effect of promoting healthy living integrating structured physical activity and nutritional intervention programs. Overweight and obesity are important lifestyle-related public health problems in the world. Overweight and obesity increase the risks of hypertension, type 2 diabetes, dyslipidemia, coronary heart disease, gout, osteoarthritis, gallbladder disease, endometrial and colon cancer, psychosocial problems, sleep apnea, disability and premature mortality (TIAN et al, 2009). There is growing evidence that a weight loss of between 5 and 10% of baseline weight will substantially reduce health risk. Exercise that expends at least 2000 kcal per week is effective to evoke a weight loss of at least 5% for men and will likely prevent weight gain (DONNELLY et al, 2003).

Endurance exercise training improves plasma lipoprotein and lipid profiles and thus reduces CVD risk. Fifteen to 20 miles of brisk walking or jogging (energy expenditure of 5021-9205 kJ/wk) was associated with HDL-C increases of 2 to 8 mg/dl (HALVESRSTADT et al, 2007). A relatively high amount of regular exercise improved overall lipoprotein profiles, including an increase in total HDL concentration, large HDL particle concentration,



and HDL particle size, even without clinically significant weight loss (TIAN et al, 2009). Long-term restriction of energy intake has proven difficult and generally results in weight re-gain over time. Exercise has been shown to provide an energy expenditure that is not fully compensated with other components of the energy balance equation and thus provides a stimulus for weight loss and maintenance (DONNELLY et al, 2003).

However, few well-controlled studies are available to determine whether regular exercise is indeed an effective weight management strategy for former addicted persons. Thus, the purpose of this study is to determine the effect of exercise training on health-related physical fitness factors and blood lipids profile of former addicted persons.

MATERIALS AND METHODS

Subjects

Subjects were recruited from the Behzisti Institution (a rehabilitation centre in Iran). They were referred there for quitting drugs or had quit drugs. Sixty-two subjects who were 23–49 years old, and had more than one year of quitting history were recruited for the present study. They were invited to an informational meeting where they received written and verbal information about the study protocol that was approved by exercise psychologist experts from Islamic Azad University of Shahrekord. The subjects were randomly allocated into either the exercise group (32 persons) or the control group (30 persons). Thirty one individuals completed the entire study with 16 persons in the exercise group and 15 persons in the control group. All subjects completed informed consent form. Before participating in the study, subjects completed a medical history questionnaire.

Procedures

Endurance exercise training consisted of 12 weeks of supervised endurance exercise including 3 sessions per week. Subjects were allowed to use various types of Game-based aerobic exercise training: badminton, walking, and jogging. Exercise training began with 20 minutes at 50% of heart rate reserve and progressed to 70% of heart rate reserve for 60 minutes where it remained for the final 12 weeks. The control group participated in psychological consulting sessions only. All exercise training was performed under the direct supervision of research personnel throughout the exercise training sessions. Because exercise training intensity can not be controlled by observation, target heart rate was verified during each session by using pulse-meters that was used randomly by ten subjects at any time.

Measurements

Body mass index (BMI) was calculated by measuring height and weight at baseline and the end of 12 weeks. Body composition was assessed using the sum of three skin-fold measurements specific for males (chest, abdomen, and thigh) (ACSM, 2000). 10 ml of blood was drawn for each subject after 12 hours fasting from superficial forearm vein. Total cholesterol, high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C) and triglyceride (TG) were measured enzymatically using Diagnostic Pars Kits. Each subject performed a middle intensity Vo_2 Peak test on a treadmill to determine the cardiovascular fitness (one-mile Rockport walking test) (ACSM, 2000). Flexibility and muscular endurance were assessed by Sit-and-Reach Test (HOEGER and HOEGER, 2007) and modified Pull-up Test (ACSM, 2000), respectively. Heart rate (HR) was recorded by using a Polar heart rate monitor belt. All variables were measured at baseline and end of weeks 12th.

Statistical analysis

Data were analyzed using SPSS (Version 13.0). For the description of data, mean and



standard deviation were used. Also, mean values of two groups in pre and post tests were compared by analysis of covariance (ANCOVA analysis) for measurements. The significance levels of this study was set at $p < 0.05$.

RESULTS

Table 1 indicated that there were no significant differences between the exercise and control group for age, duration of without abusing drugs, duration of using drugs and number of cigarettes at baseline ($p > 0.05$).

Table1 - Baseline characteristics of participants in the exercise group (EG) and control group (CG)

Variable	pretest		F	Sig	P-value
	CG	EG			
Number	15	16	-	-	-
Age(years)	35.60 ± 7.53	35.38 ± 6.71	0.456	0.505	0.931
Duration of without abusing (months)	21.46 ± 13.24	21 ± 1.18	0.90	0.351	0.916
Duration of using drugs (years)	13.40 ± 6.52	15.06 ± 7.14	0.000	0.993	0.505
Number of cigarettes per day	9 ± 10.36	11.25 ± 11.61	0.681	0.416	0.575
Height(cm)	177.40 ± 6.31	174.06 ± 4.71	1.737	0.198	0.105

Table2 - Effect of 12 weeks exercise training on variables in subjects

Variable	Changes		F	Sig
	CG	EG		
Weight (kg)	0.9 ± 3.18	-2.31 ± 2.44	1.655	0.208
BMI (Kg/m ²)	0.29 ± 1.04	-0.78 ± 0.80	1.303	0.263
WHR	0.013 ± 0.020	-0.04 ± 0.04	4.433	0.044*
CT (mg.dl ⁻¹)	-1.40 ± 15.03	-9.87 ± 13.17	0.647	0.428
TG mg.dl ⁻¹	0.93 ± 22.21	-1.81 ± 11.66	1.195	0.283
HDL- C (mg.dl ⁻¹)	-1.133 ± 2.06	1.12 ± 2.06	0.045	0.833
LDL-C (mg.dl ⁻¹)	0.80 ± 2.01	-2.62 ± 2.89	0.744	0.396
VLDL (mg.dl ⁻¹)	0.93 ± 1.28	-0.75 ± 1.65	1.99	0.169
Pull- up (Modified test)	0.47 ± 2.23	3.06 ± 2.90	0.613	0.44
Sit- up	1.06 ± 2.76	6.62 ± 3.34	0.376	0.545
Flexibility (cm)	0.533 ± 2.06	3.125 ± 2.47	0.010	0.923
Vo ₂ peak(ml/kg/min)	-0.74 ± 1.55	1.49 ± 1.67	0.022	0.884

Furthermore, as indicated in table 2, there were no significant differences between post tests for weight ($P=0.208$), BMI ($P=0.263$), CT ($P=0.428$), TG ($P=0.283$), LDL-C ($P=0.396$), VLDL ($P=0.169$), HDL-C ($P=0.833$), Pull- up ($P=0.44$), Sit- up ($P=0.545$), Flexibility ($P=0.923$) and Vo₂peak ($p=0.884$) after 12 weeks. Statistically significant differences were found for WHR between pre and post test. Weight was decreased 1.781 kg and 0.8 kg in EG and CG, respectively

DISCUSSION

The aim of present study was effects of 12-week exercise training on health-related physical fitness factors and blood lipids profile of former addicted persons. The findings of this study showed that 12 week exercise training can not decrease weight. Cowan and Devine (2008) reported that there were three main stages (early, mid and later) following recovery from substance addiction. Men in early recovery (1-6months) described dysfunctional eating practices such as binge eating, the use of food as a substitute for drug use, and the use of food to satisfy cravings (COWAN and DEVINE 2008). The findings about weight gain (CG: $0.9 \pm 3.18\text{kg}$), increasing BMI (CG: 0.29 ± 1.04), WHR (CG: 0.013 ± 0.020) after quitting substances were similar to that of Williams (2003), Cowan & Devine (2008), Trinko et al (2007). Other studies have focused more on weight gain after quitting smoking (CLARK et al, 2006; COOPER et al, 200; KNAUSS et al, 2005; PISINGER and JO RGENSEN, 2007; PRAPAVESSIS et al, 2007; SCHOELLER and SAY, 1997; TOLL et al, 2008a; TOLL et al, 2008b, USSHER et al, 2007; and WHITE et al, 2007). One reason for the weight gain is that nicotine and drugs speed up body metabolism. After quitting, body metabolism slows down and it causes weight gain even if persons do not eat more. It is postulated that for these measurements (weight and BMI), increase is not evitable (TALCOTT et al, 1995). Exercise training had no significant effect on body weight, BMI, CT, TG, HDL, LDL and VLDL ($p < 0.05$). Exercise training (180 minutes) of moderate intensity per week could not reduce the weight in compared to control group. As noted, dysfunctional eating patterns and weight gain have been observed during recovery from drug and alcohol addiction. In addition, Williams 2002 found that dieting to prevent this weight gain is ineffective and may actually interfere with quit efforts. Diet does not provide a long-term solution and over 50% of individuals who lose weights through diet eventually regain the weight they lost. Dietary restraint is difficult across time; it represents a state of deprivation, and runs counter to the current environment that provides enormous amounts of palatable, high energy, low cost foods, and beverages available at almost any location (DONNELLY et al, 2004). Therefore, physical activity and exercise can prevent or manage overweight after quitting drugs, substances and smoking. ACSM recommended that individuals decided lose weight should participate in at least 150 minutes of moderate intensity exercise training per week, and to accumulate >200 minutes of moderate intensity exercise training per week when possible. Marcus et al (2006) reported that vigorous exercise facilitates short- and longer-term smoking cessation in women when combined with a cognitive-behavioral smoking cessation program (MARCUS et al 2006). Vigorous exercise improves exercise capacity and delays weight gain following smoking cessation. Data from the National Weight Control Centre (JAKICI et al, 1999; KLEM et al., 1997; SCHOELLER and SHAY, 1997) illustrated that prevention of weight regain is associated with 280 to 450 minutes of moderate intensity exercise training per week, while 180 minutes per week was used in the present study. In the EG, 180 minutes exercise per week result in no significant weight reduction. Thus, 180 minutes moderate-intensity exercise training per week can be as a guideline for the prevention of weight gain in persons who have quitted drugs and substances for more than one year.

In the present study, there was a significant increase in health-related physical fitness factors such as muscular endurance, flexibility and Vo_2peak in the EG after 12 week exercise training, but there was no significant change in the CG. Wilmore et al (1996) reported that physical activity and exercise training result in increased muscular strength and endurance through neuromuscular adaptations in early weeks, and in later weeks it was done by muscular mechanism (WILMORE et al., 1996). The finding of present study showed no significant increase in Vo_2pak after 12 weeks of exercise training, that similar to

others (Power and HOWLEY 199, WILMOR 1996, BABB et al (1997).

Also the findings of this study indicated significant increase in VLDL ($p=0.014$), and a significant decrease in HDL-C ($p=0.025$) in CG. There were no significant differences in CT, LDL, HDL and VLDL ($p>0.05$) in post test between two groups. As noted, there was no significant decrease in TG between post tests. Exercise training should be enough intense and long to decrease TG. In this study, because of condition of subjects, the exercise was not long enough and intense was lesser ($\sim 70\%$ Vo_2 peak). Enough intensity to decrease TG is 85% Vo_2 max (HOGGER and HOGGER, 2007). Despite of only 1.78 kg weight lost, this is important because it demonstrates that individuals can acquire positive health benefits from moderate exercise (HALVERSTTADT et al, 2007).

In conclusion, our data can not provide evidence for the beneficial effects of moderate intensity endurance exercise training, independent of diet on weight, BMI, and plasma lipoprotein and lipid profiles in former addicted persons at risk for CVD. However, confirm this findings need to more studies with different intensity and duration of exercise training.

PRACTICAL APPLICATION

The findings of present study demonstrated that moderate intensity exercise training is a good training to manage weight gain in former addicted persons who have quitted drugs more than one year and to have loss weight they should exercise more intensity and long duration.

ACKNOWLEDGEMENTS

The authors would like to acknowledge and thank the necrotic annoyant (NA) group in the study for their commitment and effort throughout study.

REFERENCES

- ABBRANTES, A. M. Promoting Wellness among Adolescents in Substance Abuse Treatment Programs. A Review of the Empirical Literature. Butler Hospital / Brown Medical School. 2006.
- ACSM. ACSM'S guidelines for testing and prescription. 6th Edn. Lippincott, Williams & Wilkins, Baltimore, v. 24, p. 63-66, 2000.
- BABB, T. G.; LONG, K. A.; RODRATE, I. R. The relationship between maximal expiratory flow and increases of maximal exercise capacity with exercise training. *Am-T-Respir-Crit-Care-Med*, v. 156, n. 1, p. 116-121, 1997.
- CLARK, M. M.; HURT, R. D.; CROGHAN, I. T.; PATTEN, C. A.; NOVOTNY, P.; SLOAN, J. A.; DAKHIL, S. R.; CROGHAN G. A.; WOS, E. J.; ROWLAND, K. M.; BERNATH, A.; MORTON, R. F.; HOMAS S. P.; TSCHETTER, L. K.; GARNEAU, S.; STELLA, P. J.; EBBERT, L. P.; WENDER, D. B.; LOPRINI, C. L. The prevalence of weight concerns in a smoking abstinence clinical trial. *Addicted Behaviors*, v. 31, n. 4 ,p. 1144-1152, 2006.
- COOPER, T.; DUNDON, M.; HOFFMAN, B.; STOEVEER, C. J. General and smoking cessation related weight concerns in veterans. *Addicted Behaviors* v. 31, p. 722-725, 2006.
- COWAN, J.; DEVINE, C. Food, eating, and weight concerns of men in recovery from substance addition. *Appetite Journal*, v. 50, n. 1, P. 33- 42, 2007.
- DONNELLY, E. J.; HILL, J.; JACOBSEN, D. Effect of a 16-month Randomized Controlled

Exercise Trial on body Weight and Composition in young, Overweight Men and Women, *Achieves of International Medicine*, v. 163, n. 11, p. 1343-1350, 2003.

DONNELLY, E. J.; SMITH, B.; JACOBSEN, D. J.; KIRK, E.; DUBOSE, K.; HYDER, M.; BAILEY, B. The role of exercise for weight loss and maintenance. *Best Practice & Research Clinical Gastroenterology*, v. 18, n. 6, p. 1009–1029, 2004.

EATON, B. C. Traditional and Emerging Risk Factors for Cardiovascular Disease. *Primary Care Clinical Office Practition*, v. 32, p. 963–976, 2005.

HALVERSTADT, A.; PHARES, D. A.; WILUND, K. R.; GOLDBERG, A. P.; HAGBERG, J. M. Endurance exercise training raises high-density lipoprotein cholesterol and lowers small low-density lipoprotein and very low-density lipoprotein independent of body fat phenotypes in older men and women. *Metabolism Clinical and Experimental*, v. 56, p. 444–450, 2007.

HODGKINS, C. C., CAHILL, K. S., SERAPHINE, A. E., FROST-PINEDA, K., GOLD, M. S. Adolescent drug addiction treatment and weight gain. *Journal of Addiction Disease*, v. 23, n. 3, p.55-65, 2004.

HOEGER, W.; HOEGER, S. *Fitness and Wellness*, seventh edition. TOMSON WADSWORTH; 36-37, 2007.

HU, G.; TUOMILEHTO, J.; SILVENTOINEN, K.; BARENGO, N.; JOUSILAHT, P. Joint effects of physical activity, body mass index, waist circumference and waist-to-hip ratio with the risk of cardiovascular disease among middle-aged Finnish men and women. *European Heart Journal*, v. 25, p. 2212–2219, 2004.

JAKICI, J.; WINTERS, C.; LANG, W.; WING, P. R. Effect of intermittent exercise and use of home exercise equipment on adherence, weight loss. *American Journal of Clinical Nutrition*, v. 282, p. 1544-1560, 1999.

KLEM, M. L.; WING, P. R.; MCGUIRE, M. T.; SEAGLE, H. M.; HILL, J. O. A descriptive study of individuals successful at long-term maintenance of substantial weight loss. *American Journal of Clinical Nutrition*, v. 66, p. 239-246, 1997.

KOKKINOS, P.; NARAYAN, P.; COLLERAN, J.; FLETCHER, R.; LAKSHMAN, R.; PAPADEMETRIOU, V. Effects of Moderate Intensity Exercise on Serum Lipids in African-American Men with Severe Systemic Hypertension. *American Journal of Cardiology*, v. 81, p. 732–735, 1998.

KNAUSS, F.; MATZ, D.; PARKER, A.; CUTIN, L.; PAI, S. Women and cigarette smoking: Dose amount of weight gain following a failed quit attempt affect social disapproval? *Addicted Behaviors*, v. 30, n. 4, p. 643-651, 2005.

LEVINE, M.; PERKINS, K.; MARCUS, M. The characteristics of women Smoker concerned about post cessation weight gain. *Addictive Behaviors Journal*, v. 26, n. 5, p. 749-756, 2001.

MARCUS, B. H.; ALBRECH, A. E.; KING, T. K.; PARISI, A. F.; INTO B. M.; ROBETS, M.; NIAURA, R. S.; ABRAMS, D. B. The Efficacy of Exercise as an Aid for Smoking Cessation in Women. *Archives of Internal Medicine*, v. 159, p. 1229-1234, 1999.

OUNIS, O.; ELLOUMI, M.; AMRI, M.; TRABELSI, Y.; LAC, G.; TABKA, Z. Impact of training and hypocaloric diet on fat oxidation and body composition in obese adolescents, *Science & Sports*, v. 24, p. 178-185, 2008.

PISINGE, C.; JORGENSEN, T. Weight concerns and smoking in a general population: The Inter99 study. *Preventive Medicine*, v. 44, n. 4, p. 283-289, 2007.

POWERS, S. K.; HOWLEY, E. T. Exercise theory and application to fitness and performance. University of Knoxville W.M.C. Brown, 1990.

PRAPAVESSIS, H.; CAMERON, L.; BALDI, J. C, ROBINSON, S.; BORRIE, K.; HARPER, T.; GROVE, J. The effect of exercise and nicotine replacement therapy in smoking rates in women. *Addicted Behaviors*, v. 32, n. 7, p. 1416-1432, 2007.

ROBERGS, R. A.; ROBERTS, S. O. Fundamental Principle of Exercise Physiology: for Fitness, Performance and Health. Human Kinetic. 2000.

SCHNEIDER, K.; SPRING, B.; L, PAGOTO, S. Affective Benefits of Exercise While Quitting Smoking: Influence of Smoking-Specific Weight Concern. *Psychology Addicted Behaviors*, v. 21, n. 2, p. 255-260, 2007.

SCHOELLER, D.; SHAY, K. R. How much Physical activity is needed to minimize weight gain in previously obese women? *American Journal of Clinical Nutrition*, v. 66, p. 551-556, 1997.

STEFANICK, M. L. Nutrition in the Prevention and Treatment of Disease, Chapter32, Obesity: Role of Physical Activity. 2001.

TALCOTT, G. W.; FIEDLER, E. R.; PASCALE, R. W. Is weight gain after smoking cessation inevitable? *Journal of Consulting and Clinical Psychology*, v. 63, n. 2, p. 313-316, 1995.

TIAN, H.; XIE, H.; SONG, G.; ZHANG, H.; HU, G. Prevalence of overweight and obesity among 2.6 million rural Chinese adults. *Preventive Medicine*, v. 48, p. 59-63, 2009.

TOLL, B.; KATULAK, N.; WILLIAMS-PIEHOTA, P.; O'MALLEY, S. Validation of a scale for the assessment of food cravings among smoker. *Appetite Journal*, v. 50, n. 1, p. 25-32, 2008.

TOLL, B.; LEARY, V.; WU, R.; SALOVEY, P.; MEANDZIJA, B.; O'MALLEY, S. A Preliminary investigation of naltrexone augmentation of bupropion o stops smoking with less weight gain. *Addicted Behaviors*, v. 33, n. 1, p. 73-179, 2008.

TRINKO, R.; SEARS, R.M.; GUARNIERI, D.J.; DILEONE, R.J. Neural mechanisms underlying obesity and drug addiction. *Physiology & Behavior*. Is weight gain after smoking cessation inevitable?. v. 91, p. 499-505, 2007.

USSHER, M.; WEST, R.; MCEWEN, A.; TAYLOR, A.; STEPTONE, A. Randomized controlled trial of physical activity counseling as an aid to smoking cession: 12 month follow up. *Addicted Behaviors*, v. 32, n. 12, p. 3060-3064, 2007.

WHITE, M.; MCKEE, S.; O'MALLEY, S. Smoke and mirrors: Magnified believes that cigarette smoking suppresses weight. *Addicted Behaviors*, v. 32, n. 10, p. 2200-2210, 2007.

WILLIAMS, J. Women and Smoking: Sensory factors, Attitudes about Weight, Phase of Menstrual Cycle All Key to Quitting. *NIDA NOTES*, v. 17, n. 4, 2003.

WILMORE, J. H.; STANFORTH, P. R.; GAGNON, J.; LEONE, A. S.; RAO, D. C.; SKINNER, J. S.; BOUCHARD, C. Endurance exercise training has a minimal effect on resting heart rate: the HERITAGE Study, *Medicine and Science in Sports and Exercise*, v. 28, n. 7, p. 829-35, 1996.