

Cephalometric Comparison of Position of the Hyoid Bone in Class I and Class II Patients

Sanaz Soheilifar,¹ and Mohammad Ali Momeni^{1*}

¹Orthodontic Department, Hamadan University of Medical Sciences, Hamadan, IR Iran

*Corresponding author: Mohammad Ali Momeni, Orthodontic Department, School of Dentistry, Shahid Fahmide Blvd., Hamadan, IR Iran. Tel: +98-9194886113, E-mail: alimomeni362@gmail.com

Received 2016 May 08; Accepted 2016 June 28.

Abstract

Background: According to the close proximity of hyoid bone with dentofacial structures and its muscular attachments, a probable relationship between it and different types of skeletal patterns is suspected.

Objectives: The aim of this study is compare the position of hyoid bone in skeletal class I and class II patients.

Methods: In this study 50 cephalograms were divided into two groups, skeletal class I ($1 \leq ANB \leq 4$) and skeletal class II ($ANB > 4$), with 24 and 26 patients in each group, respectively. Horizontal and vertical position of hyoid bone were evaluated. SPSS software and student t-test were used to analyze the data.

Results: According to the results of our study, there is no statistically significant difference between the hyoid bone position in skeletal class I and skeletal class II patients.

Conclusions: Since the hyoid bone position is similar in skeletal class I and class II patients, the skeletal pattern is not the only determinant of the position of hyoid bone.

Keywords: Cephalometry, Hyoid Bone, Angle Class II Malocclusion, Angle Class I Malocclusion

1. Background

The hyoid bone is a u-shaped (horse shoe shaped) bone, situated on the anterior midline of the neck between symphysis and larynx, suspended from the tip of stylohyoid process of temporal bone by stylohyoid ligaments (1). Unlike the other bones of head and neck, it has no osseous joint and is only distantly articulated to mandible, cranium and pharynx by muscles and ligaments (2). Function of this bone is to maintain equilibrium of respiratory way by serving as anchoring structure for the tongue and also is to keep normal head posture (3-5).

The hyoid bone position could be a good diagnostic guide to malocclusions elicited by destructive oral habits such as atypic deglutition or mouth breathing (6, 7). Changes in the anteroposterior head posture and mandibular inclination can affect the hyoid bone position (3, 8, 9). Treatment with functional appliances that position mandible downward and forward, may also displace the hyoid bone (10-13). Recent orthodontic studies indicate that evaluation of the hyoid bone position may play an important role in assessment of dentofacial structures especially in evaluation of relapse prevention after orthodontic treatment and orthognathic surgeries (14). Furthermore, in several studies, the hyoid bone position and pharyn-

geal airway space are mentioned in relation. Jose et al, who evaluated this relation in different types of malocclusions, found a positive relationship between lower airway and horizontal distance from the hyoid bone to the retrognathion only in class I skeletal patient with normal growth pattern (15). Guven et al. reported changes in the hyoid bone position and pharyngeal airway space after mandibular body osteotomy (16). In addition, Parkkinen observed lower hyoid bone position in children with breathing difficulties during sleep. Although several studies reported a positive correlation between the hyoid bone position and mandible (17-19), the other studies concluded that there is no significant correlation between them (7, 20).

2. Objectives

The aim of this study was to evaluate the cephalometric hyoid bone position in patients with skeletal class I and skeletal class II pattern.

3. Methods

Based on sample size in similar studies (21), 24 cephalograms were selected in class I group and 26 cephalograms

for class II group from the archive of dentistry faculty of Hamedan University of medical sciences between “1389-92”.

Inclusion criteria was as follows: being Iranian, “neither” adenoidectomy “nor” tonsillectomy history, lack of presence of pharyngeal lesions such as tonsil hypertrophy or tonsillitis, having normal growth pattern (FMA = 22-28) (22), patients in the age group of 9-11 years who didn't attain the age of maturity (in the stage of CS1, CS2 and CS3) (23) and having comfortable nasal breathing. The patients' files were checked for inclusion criteria. Any patient who didn't meet these criteria was excluded. All the cephalograms were taken using the Soredex Digora machine; used in Hamedan dentistry school with standard method in normal head posture and centric occlusion.

Our comparing groups were skeletal class I ($1 \leq ANB \leq 4$) and skeletal class II ($ANB > 4$) malocclusions including 24 and 26 patients, respectively. These two groups were matched regarding the stage of cervical vertebra and sex.

Table 1 shows the frequency of patients in each group.

Horizontal (24) and vertical (25) measurements of the hyoid bone position were conducted according to the previous studies (Figure 1).

All cephalograms were traced manually and 10 cephalograms were randomly selected and traced again after a week to evaluate systematic error.

SPSS19 software was used for data analysis. Paired t-test was used for estimation of systematic error and independent t-test was used for analysis of difference between two groups' variables. Significance level was considered as $P < 0.05$.

4. Results

Independent t-test, shows no statistically significant difference between two groups of skeletal class I and II malocclusions, considering horizontal dimension measurements. Table 2 illustrates the information related to this evaluation.

Independent t-test, shows no statistically significant difference between two groups of skeletal class I and II malocclusions, considering vertical dimension measurements. Table 3 illustrates the information related to this evaluation. Pearson correlation showed that there was no statistical correlation between ANB and variables.

5. Discussion

The position of the hyoid bone in class I, II and III malocclusions has been evaluated in several studies and various results are achieved (17-22). The position of the hyoid

bone is different, not only from an individual to the other one, but also in different time intervals in the same patient (15). Evaluation of the position, shape and function of the hyoid bone seems to be important due to its effect on equilibrium maintenance of surrounding tissues.

In present study, we found no statistically significant difference between class I and II malocclusions, regarding the horizontal and vertical position of hyoid bone. Similar to our study, Carvalho et al. reported no statistically significant difference in the hyoid bone position in class I and II malocclusions and statistically significant difference only was shown with ANS-PNS measurement for the evaluation of vertical position (7). Furthermore, in Jose et al., who evaluated vertical and horizontal position of the hyoid bone in class I, II and III malocclusions using Bibby and Preston analysis (3), no statistically significant difference was found between these three groups (15).

Unlike our study, in the study of Kuroda et al., there was statistically significant difference in hyoid bone position in skeletal class I, II and III patients. The hyoid bone in the class II patients was situated closer to the skull base in comparison with class III patients (17). The controversial results are due to the fact that Kuroda et al. found difference in hyoid position in class II and class III subjects, but in the present study class I and class II subjects are compared.

Adamidis and Spyropoulos evaluated the effect of hypertrophic adenoid tissue on the hyoid bone position in children with mean age of 9 years. The results suggested that more downward inclination of mandible in the mouth breathers as compared to nasal breathers (without hypertrophy of tonsils), can be seen. This study also indicates that the hyoid bone position follows the mandibular posture in the mouth breathing patients and hence, the hyoid bone shows downward inclination, too. The authors said that the reason is indirect effect of craniofacial complex and function of suprahyoid and infrahyoid muscles on the position of hyoid bone (18).

Galvao et al. reported that the hyoid bone is located closer to the rim of mandible in class II patients and is situated farther in class III patients and difference was statistically significant (19). The reason of this finding may be due to investigation of skeletal class II and III patients “not” skeletal class I and II ones.

Haralabakis et al. reported that the distance from the hyoid bone to the palatal plane was found to be statistically greater in the open bite class II group than class I normal patients. They stated that the palatal plane plays a role in the development of skeletal malocclusions, and also the findings strongly suggest that the hyoid bone moves only in close relation with pharynx, cervical spine and mandibular plane in patients with entirely different skeletal pattern. Most of horizontal measurements revealed no

Table 3. Mean, Standard Deviation and P-Value Obtained from Independent Test for Evaluating the Variables of Vertical Position of the Hyoid Bone

Variable	Mean \pm SD		P Value
	Class I	Class II	
H-SN	87.500 \pm 14.6140	85.381 \pm 18.1645	0.647
H-FH	72.033 \pm 5.3722	71.190 \pm 6.0632	0.603
H-PP	51.000 \pm 5.2850	51.333 \pm 9.1013	0.870
H-MP	11.967 \pm 4.9234	10.214 \pm 4.5403	0.203

ference was observed only in the horizontal position of the hyoid bone after three series of cephalometric evaluation relative to T0 (26). In the present study all of the patients were in the same age group (9-11 years), therefore it does not seem that the age of patients in two groups would affect the hyoid bone position.

Valenzuela et al. didn't find any statistically significant difference between anteroposterior head posture and the hyoid bone (27). Smith et al. suggested that the hyoid bone position varies with change in the posture from upright to supine due to change in larynx posture and reduction in oropharyngeal airway (28). Since in the present study the cephalograms were taken in natural head position, no difference is expected in the position of hyoid bone due to change in postures.

The results of our study suggest that horizontal and vertical dimensions of the hyoid bone position have no significant difference in skeletal class I and class II patients. Some studies suggest that compensating function of surrounding muscles and soft tissues might result in similarity of the hyoid bone position in different skeletal groups (29).

5.1. Conclusion

Hyoid bone position is not significantly different in skeletal class I and class II patients. Some compensational mechanisms in surrounding structures of dentofacial complex may be responsible for maintaining the position of hyoid bone, despite the difference in the position of maxilla and mandible in skeletal class I and class II subjects.

References

- Williams PL, Warwick R, Dyson M, Bannister LH. Gray's anatomy. 378. Churchill livingstone Edinburgh; 1989.
- Romanes GJ. Cunningham's Manual of Practical Anatomy: Volume 3: Head and Neck and Brain. Oxford University Press; 1986.
- Bibby RE, Preston CB. The hyoid triangle. *Am J Orthod.* 1981;**80**(1):92-7. [PubMed: 6942659].
- Brodie AG. Anatomy and physiology of head and neck musculature. *Am J Orthod.* 1950;**36**(11):831-44. [PubMed: 14783190].
- Atik E, Kocadereli I. . Craniofacial morphology, head posture and hyoid bone position. *Turkish J Orthodon.* 2011;**24**:237-46.
- Bibby RE. The hyoid bone position in mouth breathers and tongue-thrusters. *Am J Orthod.* 1984;**85**(5):431-3. [PubMed: 6586082].
- Carvalho E, Nouer DF, Garbui IU, Nouer PRA. Relationship of the hyoid bone and posterior surface of the tongue for class II division 1 orthodontic patients. *Rev Gaucha Odontol.* 2013;**61**(1):61-8.
- Gustavsson U, Hansson G, Holmqvist A, Lundberg M. Hyoid bone position in relation to head posture. *Sven Tandlak Tidsskr.* 1972;**65**(8):423-30. [PubMed: 4517205].
- Valk JWP, Zonnenberg AJJ, van Maanen CJ, Onno GT. The biomechanical effects of a sagittal split ramus osteotomy on the relationship of the mandible, the hyoid bone, and the cervical spine. *Am J Orthod Dentofacial Orthop.* 1992;**102**(2):99-108.
- Rocabado M. Biomechanical relationship of the cranial, cervical, and hyoid regions. *J Craniomandibular Pract.* 1983;**1**(3):61-6. [PubMed: 6586872].
- Wenzel A, Williams S, Ritzau M. Changes in head posture and nasopharyngeal airway following surgical correction of mandibular prognathism. *Eur J Orthod.* 1989;**11**(1):37-42. [PubMed: 2714390].
- Winnberg A, Pancherz H. Head posture and masticatory muscle function. An EMG investigation. *Eur J Orthod.* 1983;**5**(3):209-17. [PubMed: 6578043].
- Erdinc AM, Dincer B, Sabah ME. Evaluation of the position of the hyoid bone in relation to vertical facial development. *J Clin Pediatr Dent.* 2003;**27**(4):347-52. [PubMed: 12924735].
- Gunduz Arslan S, Dildes N, Devecioglu Kama J. Cephalometric investigation of first cervical vertebrae morphology and hyoid position in young adults with different sagittal skeletal patterns. *Scientific World J.* 2014;**2014**.
- Jose NP, Shetty S, Mogra S, Shetty VS, Rangarajan S, Mary L. Evaluation of hyoid bone position and its correlation with pharyngeal airway space in different types of skeletal malocclusion. *Contemp Clin Dent.* 2014;**5**(2):187-9. doi:10.4103/0976-237X.132313. [PubMed: 24963244].
- Güven O, Saracoglu U. Changes in pharyngeal airway space and hyoid bone positions after body osteotomies and sagittal split ramus osteotomies. *J Craniofac Surg.* 2005;**16**(1):23-30. [PubMed: 15699641].
- Kuroda T, Nunota E, Hanada K, Ito G, Shimizu K, Nakamura S, et al. [A roentgenocephalometric study on the position of the hyoid bone]. *Nihon Kyosei Shika Gakkai Zasshi.* 1966;**25**(1):31-8. [PubMed: 5225031].
- Adamidis IP, Spyropoulos MN. The effects of lymphadenoid hypertrophy on the position of the tongue, the mandible and the hyoid bone. *Eur J Orthod.* 1983;**5**(4):287-94. [PubMed: 6581051].
- Galvao C. Estudo cefalométrico da posição do osso hioide em indivíduos com oclusão normal e portadores de maloclusão. *Rev Odont Unesp.* 1983;**12**(1/2):143-52.
- Stepovich ML. A cephalometric positional study of the hyoid bone. *Am J Orthod.* 1965;**51**(12):882-900. [PubMed: 5214894].
- Mergen DC, Jacobs RM. The size of nasopharynx associated with normal occlusion and Class II malocclusion. *Angle Orthod.* 1970;**40**(4):342-6. doi: 10.1043/0003-3219(1970)040<0342:TSONAW>2.0.CO;2. [PubMed: 5272410].

22. Jacobson A, Jacobson RL, Khoury F, Antoun H, Missika P, Sclar AG. Radiographic cephalometry: From basics to 3-d imaging, (Book/CD-ROM set). 2006.; 2007.
23. Proffit WR, Fields Jr HW, Sarver DM. Contemporary orthodontics. Elsevier Health Sciences; 2014.
24. Yamaoka M, Furusawa K, Uematsu T, Okafuji N, Kayamoto D, Kurihara S. Relationship of the hyoid bone and posterior surface of the tongue in prognathism and micrognathia. *J Oral Rehabil.* 2003;**30**(9):914-20. [PubMed: [12950973](#)].
25. Haralabakis NB, Toutountzakis NM, Yiagtzis SC. The hyoid bone position in adult individuals with open bite and normal occlusion. *Eur J Orthod.* 1993;**15**(4):265-71. [PubMed: [8405131](#)].
26. Kollias I, Krogstad O. Adult craniocervical and pharyngeal changes- a longitudinal cephalometric study between 22 and 42 years of age. Part I: Morphological craniocervical and hyoid bone changes. *Eur J Orthod.* 1999;**21**(4):333-44. [PubMed: [10502896](#)].
27. Valenzuela S, Miralles R, Ravera MJ, Zuniga C, Santander H, Ferrer M, et al. Does head posture have a significant effect on the hyoid bone position and sternocleidomastoid electromyographic activity in young adults?. *Cranio.* 2005;**23**(3):204-11. doi: [10.1179/crn.2005.029](#). [PubMed: [16128355](#)].
28. Smith AM, Battagel JM. Non-apneic snoring and the orthodontist: radiographic pharyngeal dimension changes with supine posture and mandibular protrusion. *J Orthod.* 2004;**31**(2):124-31. doi: [10.1179/146531204225020418](#). [PubMed: [15210928](#)].
29. Ravanmehr H, Abdollahi D. The position of hyoid bone in skeletal class I, II and III patients. *J Dental Med.* 2000;**13**(3):27-37.