

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/272842885>

Effect of hope therapy on the hope of diabetic patients

Article in *Iranian journal of nursing and midwifery research* · February 2015

Source: PubMed

CITATIONS

3

READS

46

4 authors, including:



Zahra Ghazavi

Isfahan University of Medical Sciences

18 PUBLICATIONS 28 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Environmental medicine: social and medical aspects [View project](#)



Scientific Hypothesis as a source of innovation and discovery [View project](#)

All content following this page was uploaded by [Zahra Ghazavi](#) on 12 April 2015.

The user has requested enhancement of the downloaded file.

Effect of hope therapy on the hope of diabetic patients

Zahra Ghazavi¹, Firouz Khaledi-Sardashti², Mohamad Bagher Kajbaf¹, Mojtaba Esmailzadeh²

ABSTRACT

Background: Hope is the most important factor in diabetic patients' life. The level of hope may be changing among these individuals as a result of chronic nature of diabetes and its complications. When the level of hope increases among these patients, they can resist against physical and psychological complications of diabetes more, accept the treatment better, enjoy life more, and adapt with their situations more efficiently. This study aimed to define the efficacy of hope therapy on hope among diabetic patients.

Materials and Methods: This was a quasi-experimental study conducted on 38 diabetic patients referring to Sedigheh Tahereh Research and Treatment Center affiliated to Isfahan University of Medical Sciences in Iran in 2012. The subjects were selected based on the goals and inclusion criteria of the study and then were randomly assigned to study and control groups. Herth Hope Index (HHI) was completed by both groups before, after, and 1 month after intervention. In the study group, 120-min sessions of hope therapy were held twice a week for 4 weeks. Descriptive and inferential statistical tests were adopted to analyze the data through SPSS version 12.

Results: Comparison of the results showed that hope therapy significantly increased hope in diabetic patients after intervention in the study group compared to control ($P < 0.001$).

Conclusions: The results showed that hope therapy increased hope among diabetic patients. This method is suggested to be conducted for diabetic patients.

Key words: Diabetics, group therapy, hope, Iran

INTRODUCTION

Diabetes refers to a group of metabolic diseases with a common characteristic of increased blood sugar resulting from a defect in secretion of insulin or a defect in its function and/or both.^[1] One of the reasons for the health and treatment system focusing on diabetes is its prevalence. Based on the International Federation for Diabetes report, there are 189 million diabetic patients in the world^[2] and it is also predicted to reach 360 million people in 2030.^[3,4]

The second reason to focus on diabetes is the complications and mortality associated with this disease. Blood sugar increase results in short- and long-term complications in diabetic patients.^[1] Treatment and the costs associated with the care are very high for diabetes. The costs related

to diabetes are divided into two groups: Direct costs which include treatment and medical services and the indirect ones which include short-term or long-term disability, job loss, and early mortality.^[4] The imposed burden of diabetes is about 44 billion dollars as direct and 54 billion dollars as indirect costs. In Iran, yearly burden (physicians' visit fee) accounts to 10 billion dollars.

On the other hand, per capita cost of a diabetic patient is 250 dollars.^[5] Although chronic patients have longer life span compared to the past due to medical advancements, they simultaneously face adaptation problems more than before; therefore, despite the fact that the main concern in health and treatment was individuals' life preservation, the main challenge of the present century is to care about general health and a qualified and joyful life.^[6] Diabetic patients' mental health is a specific concern.^[7] Research shows that psychological, social, and disabling problems such as feeling of tiredness, irritability, anger, depression, and anxiety are seen more in diabetic patients than other individuals.^[8] Concurrency of emotional problems such as depression and anxiety among diabetic patients is accompanied with reduction of quality of life, self-care defective behaviors, and reduction of blood sugar control among these patients. Management and control of diabetes can be disturbed by existence of diabetes-related emotional problems such as fear of hypoglycemia, worrying about complications, or not accepting the disease. Psychological and social problems

¹Department of Psychiatry Nursing, Faculty of Nursing and Midwifery, Isfahan University of Medical Sciences, ²Department of Psychology, Faculty of Psychology and Educational Sciences, Isfahan University, Isfahan, Iran

Address for correspondence: Ms. Zahra Ghazavi, Department of Psychiatry Nursing, Faculty of Nursing and Midwifery, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: Zahra_ghazavi@nm.mui.ac.ir

Submitted: 15-Nov-12; Accepted: 05-Aug-13

1 may be caused by diet and activity limitations, the need
2 for precise and constant self-care, and the possibility of
3 serious physical complications like renal, ocular, cardiac,
4 and medullary problems. Individuals' acceptance of the
5 disease to change their lifestyle based on that is not always
6 easy.^[9] There are numerous non-medication methods and
7 treatments to help diabetic patients to cope and adapt with
8 these problems, such as complementary treatments. Hope
9 therapy is one of the complementary treatments.

10
11 Various researches have shown that hope is a meaningful
12 element in life and helps individuals to adapt with the
13 disease and its problems, decrease their mental suffering,
14 and enhance their quality of life and mental and social
15 health.^[10] It can also physiologically and emotionally help
16 patients to tolerate the crisis of the disease.^[11] Hope is an
17 essential element in chronic patients' life and has high effects
18 on their adaptation with disease. Hope plays a major role in
19 patients' quality of life and affects their various stages of the
20 disease. Hope has been defined as an internal power that
21 can enrich life and enable the patients to have a perspective
22 farther than their present state.^[10] It is also an important issue
23 in Islam as God always invites the human beings to hope and
24 show optimism toward life and designs a brighter future for
25 them. On the contrary, disappointment has been counted
26 as hideous and is named as the second great sin as it leads
27 to disappointment from mercy and compassion of God
28 and not believing in His endless power and kindness.^[11] In
29 Quran, it is said, "who gets hopeless from Lord's compassion
30 except the perverse"^[12] and "keep hopeful to God's mercy
31 as only the perverse give up hope."^[12] Positive effects of
32 helpful psychological structures (like optimism and hope) on
33 physical and mental health have been emphasized in various
34 studies.^[13] Positive individuals are healthier and happier
35 and their immunity system functions better. They can cope
36 with psychological tensions better through application
37 of more efficient coping strategies like reevaluation and
38 problem solving. They also actively avoid stressful events
39 of life and make a better supportive social network around
40 themselves. They have healthier biological styles which
41 protect them against diseases, and if they are diseased, they
42 follow medical recommendations better through behavioral
43 patterns, which accelerate recovery.^[14] Lack of hope and a
44 goal in life leads to patients' lower quality of life and hopeless
45 beliefs. People who deal with chronic diseases have different
46 physiologic, psychological, social, and emotional needs
47 compared to healthy individuals.

48
49 Therefore, finding some methods to fulfill these needs is
50 a part of the process of coping with the disease. During
51 fulfillment of patients' physical needs, their emotional
52 and social needs may be ignored. So, the best option
53 in either patients' recovery or in relation with fulfillment
54 of their needs is the group of interventions which not
55
56


1 only consider physical treatments but also psychological
2 and social treatments, for which hope therapy can play
3 this role well.^[10] Hope can be considered as a symbol of
4 mental health in nursing investigations.^[15] On the one
5 hand, nurses spend more time with diabetic patients
6 compared to physicians and have different responsibilities
7 in diabetes team, and on the other hand, they play a
8 helpful role in diagnosis and treatment of these patients'
9 emotional problems.^[16] Through working with chronic
10 patients, nurses are in relation with the whole of a
11 person (regarding physical, mental, and social aspects)
12 in addition to dealing with his/her physical problems and
13 this holistic approach needs a vast field of knowledge in
14 behavioral and psychological sciences.^[17] Meanwhile,
15 psychiatric nurses have numerous and efficient roles in
16 complementary medicine related interventions whose
17 education and application is among their duties in relation
18 with promotion of individuals' health. Psychiatric nurses
19 are also counted as the practitioners, performers, and
20 prescribers of this type of medicine.^[18,19] Johnswin, a
21 complementary medicine physician, has emphasized on
22 the importance of psychiatric nurses' role in conducting
23 and performing complementary skills and medicine and
24 highlights complementary interventions as nurses' role.
25
26

27
28 He argues that psychiatric nurses often conduct those
29 interventions that focus on mental, social, and biological
30 aspects.^[20] Application of group hope therapy, which is
31 one among the complementary treatments, has a specific
32 position, and despite the importance of hope therapy as
33 a complementary treatment in motivating and causing
34 behavior changes, it has been ignored in health-related
35 domains, so poor research is observed in this field.^[21]
36 Research shows that promotion of hope is an efficient
37 way to promote chronic patients' quality of life. Increased
38 hope consequently leads to an increase in self-care level,
39 quality of life, and general health among this group of
40 patients.^[22] The highest volume of research on hope therapy
41 in recent decades has been conducted by Seligman, the
42 father of positive psychology (2000), and Snyder.^[23] They
43 believe that disappointment leads to physical and mental
44 diseases.^[21,24,25] Snyder's studies revealed that chronic
45 patients undergoing hope therapy show a more appropriate
46 response after receiving hope therapy intervention when
47 faced with disease-related stress and tensions, resist more
48 during treatment, and accept and follow the suggested
49 treatment better.^[26] Hope therapy in the present study is a
50 combination of Snyder's hope therapy and hope program
51 in Islam. As few studies have been conducted on the effect
52 of hope therapy on the level of hope in diabetic patients,
53 the present study aimed to investigate the effect of group
54 hope therapy on the level of diabetic patients' hope.
55
56

MATERIALS AND METHODS

This was a quasi-experimental two-group (study and control) two-step study conducted on 38 diabetic patients in 2012. Study population included all diabetic patients having a medical record and referring to Sedigheh Tahereh Research and Treatment Center in Isfahan, Iran. Inclusion criteria were the interest to attend the study and Muslim diabetic patients aged 30-50 years residing in Isfahan, who could express their experiences and memories. On the other hand, the subjects were excluded in case of attending other psychotherapy programs during the study, taking psychotic medications, being involved in acute depression, and finally, in the event of death of any of their immediate relatives. Out of 250 patients' medical files studied, 38 diabetic patients who met the inclusion criteria were selected through random convenient sampling. After getting their written informed consent, they were randomly assigned to two groups of study and control. Data collection tool was a two-section questionnaire. The first section was about personal characteristics including age, sex, marital status, education level, length of diabetes, types of complications due to diabetes, and types of diabetes medication taken or insulin. The second section included Herth Hope Index (HHI) containing 12 questions based on a 4-point Likert scale scored as score 1 (absolutely disagree), 2 (disagree), 3 (agree), and 4 (absolutely agree). Questions 3 and 6 were scored inversely. HHI scores in the range 12-48 and higher scores show a better hopefulness status. Scores 12-24 show low level of hope (hopelessness), 25-32 indicate moderate level of hope, and 37-48 indicate high level of hope. Herth confirmed the validity of this index through Cronbach's alpha (0.97) and by test-re-test and calculation of correlation coefficient of 0.91. In Iran, after extraction and translation of Herth's questionnaire, Pourghaznein used both the above-mentioned methods to calculate test validity based on Cronbach's alpha of 0.76 and Pearson correlation coefficient of 0.84. He also confirmed the reliability of the tool through some expert psychologists.^[27] The questionnaire was filled by both study and control groups before intervention. As in psychotherapy, the number of group members is 7-12 members,^[28] the study group was divided into two groups of 9 and 10 members, and each group separately attended the hope therapy sessions based on an already made schedule on different days twice a week for eight sessions of 2 h duration. The location of the sessions was a classroom in Sedigheh Tahereh Center. The educational program comprised a combination of Snyder hope therapy and hope program in Islam (hope giving Hadis and verses and Quran stories like those of Joseph and Jonah). Hope therapy is a narrative and story-based approach on individuals' life, in which every person tells his/her story of life based on his/her own culture. As this

study was conducted in Iran which is an Islamic country, the researcher tried to make a combination of hope program in Islam and Snyder's hope therapy method. This model has already been localized concerning Iranian and Islamic culture.^[22] After researcher's explanation about hope program in Islam and Snyder hope therapy model, every member framed his/her story of life and retold that. This program had been already localized for patients with basic hypertension,^[22] but no localized study was conducted on diabetic patients.

Each session included four sections. In the first section, activities and assignments of the former session were discussed for 25 min, and the members were encouraged to help one another to solve problems and do their assignments like getting hopeful through telling their story of life, hope domination, and empowerment. In the second section, the subjects underwent education of psychological and hope-related skills through a treatment unity between the subjects and the therapist, lead to increase of hope through different techniques such as goal expanding optimizing techniques. They included providing a structure for uncovering goals, coming up with clear and workable goals, finding the silver lining, and Selgman optimism model, as well as hope maintaining skills for about 30 min. In this section, the subjects learned a new hope-related skill. In the third section, which lasted for 50 min, the methods of application for these skills in daily life were discussed and the subjects practically conducted them. The subjects were encouraged to go over the problems objectively and clearly and help one another to solve them by use of hope skills. In the last 15 min of the session, the assignments of the future session were determined. No intervention was conducted for the control group. Immediately after and month after the intervention, HHI questionnaire was completed by the study and control groups again. Finally, the collected data were analyzed by descriptive and inferential statistical tests such as independent *t*-test, Chi-square test, repeated measure analysis of variance (ANOVA), Mann-Whitney, and Fischer's exact test. 

RESULTS

The obtained results in both groups showed that there was no significant difference in subjects' baseline demographic characteristics such as age, number of children, length of diabetes, sex, marital status, types of complications that resulted from diabetes, types of medication consumed to control diabetes, and level of education ($P > 0.05$).

The results also showed that all subjects suffered from type 2 diabetes. Independent *t*-test showed no significant difference

in the mean scores of hope before intervention in the study and control groups ($P = 0.92$), but immediately after and 1 month after the intervention, the mean scores of hope were significantly higher in the study group compared to the control group ($P < 0.001$) [Table 1]. Repeated measure ANOVA showed no significant difference in the mean scores of hope in the control group before, immediately after, and 1 month after intervention ($P = 0.26$), but a significant difference was observed in the mean scores of hope in the study group before, immediately after, and 1 month after intervention ($P < 0.001$). Since no intervention was conducted in the control group, the mean score of hope had no significant difference immediately after and 1 month after intervention compared to before intervention. Meanwhile, in the study group in which hope therapy intervention was conducted, the mean score of hope was notably increased after intervention compared to before intervention, and this increase continued until 1 month after intervention. Therefore, it can be concluded that hope therapy has been effective on increase of hope in diabetic patients [Table 2]. Least significant difference (LSD) *post-hoc* test showed a significant increase in the scores of hope before and after intervention ($P < 0.001$) and before and 1 month after intervention ($P = 0.003$) in the study group, but not before and 1 month after intervention in control group ($P = 0.28$), which showed that the effect of hope therapy continued until 1 month of follow-up.

DISCUSSION

The results of the study showed that hope therapy could affect the level of hope among diabetic patients in the study group. It can be reasoned that hope is a strong adaptation

Table 1: Comparison of mean scores of hope in the two groups of study and control

Mean score	Study		Control		Independent t-test	
	M	SD	M	SD	t	P value
Before intervention	34.5	6.4	34.3	5.4	0.1	0.92
After intervention	40.9	4.6	34.5	4.7	4.23	<0.001
One month after intervention	39.8	5.1	33.7	4.1	4.02	<0.001

SD: Standard deviation

Table 2: Comparison of mean scores of hope at three time periods of intervention in the two groups

Group	Before		After		One month after		Repeated measures ANOVA	
	M	SD	M	SD	M	SD	F	P value
Study	34.5	6.4	40.9	4.6	39.8	5.1	13.8	<0.001
Control	34.3	5.4	34.5	4.7	33.7	4.1	1.46	0.26

SD: Standard deviation, ANOVA: Analysis of variance

mechanism among chronic patients including diabetic patients, so that hopeful individuals can tolerate the crisis of the disease more conveniently.^[29] This finding is in line with that of some other studies. Irving *et al.*,^[30] Klausner *et al.*,^[31,32] and Hankins,^[33] in their studies during the investigation on hope-based interventions among adults, showed that this therapy led to an increase in hope and a decrease in depression. Shekarabi *et al.* showed that group hope therapy significantly increased hope among mothers of children with cancer and reduced their depression, and the results of a 2-month follow-up showed no significant change in that high level of hope in this group of mothers compared to after intervention.^[34] Cheavens *et al.*^[35] in a community sample showed that group hope therapy based on Snyder and McDermott hope approach increased hope in the study group, but the difference was not significant, possibly due to lack of identical and homogenous subjects. Ghasemi *et al.*^[36] showed that group education intervention based on Snyder hope theory style led to an increase in joyfulness of the elderly. Bijari *et al.*^[37] showed that group therapy based on hope therapy approach significantly increased hope and decreased depression among women with cancer compared to the control group.

Although all of the aforementioned studies have emphasized that hope therapy can increase hope, they cannot be compared with the present study as none of them have been conducted among diabetic patients and used hope program in Islam; thus, the research in this field is poor.

Sotoudeh Asl *et al.*^[22] showed that hope therapy promoted quality of life inpatients with hypertension more than medication, so that the effect of treatment remained until 3 months post intervention. The present study, like the study of Sotoudeh Asl *et al.*, was a combination of Snyder hope therapy and hope program in Islam, with a difference in the length of follow-up (1 month in the present study).

This study cannot be compared with the study of Sotoudeh Asl *et al.* as the subjects were hypertensive patients and the variable of hope was not measured directly in their study. Ebadi *et al.*^[38] showed that application of positivism with emphasis on Quran verses increased hope among widowed women in Ahwaz and the effect of intervention with regard to follow-up test results had the essential stability. Their results were similar to the results obtained by us to some extent, as they used the effect of Quran verses on the level of hope, which is a positive psychological component; this similarity is reinforced. No studies (even conducted on non-diabetic patients) were found to report that hope therapy had negative effects on hope and lowered that. The following factors have been effective in description and explanation of the efficacy

and constant effect of hope therapy on increase of hope among diabetic patients:

Encouragement of the subjects to tell the story of their life is one of the principles in hope therapy. These stories let the therapist highlight the sparkles of hope, especially those that may disappear due to some memories and thoughts.^[31]

Determination of realistic and measurable goals and changing the great goals to some minor and more accessible goals is a characteristic of hopeful individuals.^[39] The diabetic subjects started this step by selection of minor and logical goals, so as to achieve them in 1 month.

Limitations of the study

Low sample size can be mentioned as one of our limitations. It is suggested to recruit a higher sample size in future studies. Another limitation was use of a self-report data collection tool. In future studies, objective behavior indexes or semi-structured interviews can be adopted. As the sessions were managed by the researcher, unexpected bias may have affected approval of the hypotheses, which can be counted as another limitation for the present study.

CONCLUSION

With concentration on hope-bringing thoughts, hope therapy helps individuals to use their problem-solving abilities better, and consequently, they use problem-solving skills more when faced with problems and challenges, instead of problem-avoiding behaviors.^[40] Hope therapy used positive psychology instead of concentration on disabilities. Positive self-regard, hopeful imagination, healthy diet, sport, and being a member of social support networks are some characteristics of hopeful individuals which were focused in the present study.^[41]

With regard to the results of the present study, it can be noted that group hope therapy can enhance hope among diabetic patients and it can be concluded that the efficacy of hope therapy can last during follow-up period.

ACKNOWLEDGMENT

Hereby, many appreciations go to all those who assisted us in conducting the present project and also other esteemed staff of Sedighe research center affiliated to Isfahan University of Medical Sciences, Isfahan, Iran. This article was derived from a master thesis of ??? with project number ??? Isfahan University of Medical Sciences, Isfahan, Iran.

REFERENCES

1. Fauci A, Braunwald E, Braunwald E, Kasper D, Hauser S,

Longo D, et al. Harrison's Principles of Internal Medicine (endocrine and metabolic diseases). Translated by: Sotodehnia AH and Asareh MH. 18th ed. Tehran: Ketab Arjmand Publication; 2012. p. 192-3.

2. Schram MT, Baan CA, Pouwer F. Depression and quality of life in patients with diabetes: A systematic review from the European depression in diabetes (EDID) research consortium. *Curr Diabetes Rev* 2009;5:112-9.
3. Brunner LS, Smeltzer SC, Bare BG, Hinkle JL, Cheever KH. Brunner and Suddarth's textbook of medicalsurgical. Nursing. 12th ed. Philadelphia: Lippincott Williams and Wilkins; 2010.
4. American Diabetes Association. National diabetes fact sheet. Available from: <http://www.diabetes.org/udeocumentations/nationaldiabetesfactsheetRev.pdf> [Last accessed on 2005 ??].
5. Soltanian AR, Bahreini F, Afkhami-Ardekani M. People awareness about diabetes disease and its complications among aged 18 years and older in Bushehr port inhabitants (Iran), Diabetes and Metabolic Syndrome: Clinical Research and Reviews (2007) 1, 245-249.
6. Strine TW, Chapman DP, Balluz L, Mokdad AH. Health-related quality of life and health behaviors by social and emotional support: Their relevance to psychiatry and medicine. *Soc Psychiatry Psychiatr Epidemiol* 2008;43:151-9.
7. Zhao W, Chen Y, Lin M, Sigal RJ. Association between diabetes and depression: Sex and age differences. *Public Health* 2006;120:696-704.
8. Mahmodi A, Sharifi A. Comparison of Prevalence and factors associated with depression in patients with diabetes and non-diabetic individuals. *Journal of Urmia Nursing and Midwifery Faculty* 2008;2:87-93.
9. Moser A, van der Bruggen H, Widdershoven G. Competency in shaping one life: Autonomy of people with type 2 diabetes mellitus in a nurse-led, shared-care setting: A qualitative study. *Int J Nurs Stud* 2006;43:417-27.
10. Abdi N, Taghdisi M, Naghdi S. Efficacy of increasing hope interventions in cancer patient of Sanandag. *Armaghane-danesh, Journal of Yasuj University of Medical Sciences* 2009;14:13-21.
11. Ebadi N, Faghihi A. Efficacy of positivism (optimism) on increase hope of widow Ahvaz women with emphasis on Koran. *J Psychol Relig* 2010;2:61-74.
12. Qoran.hojar, verse 56. Josef, verse 87.
13. Scheier MF, Carver CS, Bridges MW. Optimism, pessimism, and psychological well-being». *Change optimism and pessimism*. Washington, Dc: American Psychological Association; 2001. p. 182-216.
14. Schram MT, Baan CA, Pouwer F. Depression and Quality of life in patients with diabetes: A Systematic Re-view from the European Depression in Diabetes (EDID).Research Consortium, National Institute for Public Health and the Environment. Centre for prevention and health services research, Bilthoven. The Netherlands, *Curr Diabetes Rev*. May 2009;5 (2):112-119.
15. Dufault, K., and Martocchio, B. C. (1985). Hope: Its spheres and dimensions. *Nursing Clinics of North America*, 20 (2), 379-391.
16. Pouwer F, Beekman AT, Lubach C, Snoek FJ. Nurses' recognition and registration of depression, anxiety and diabetes-specific emotional problems in outpatients with diabetes mellitus. *Patient Educ Couns* 2006;60:235-40.
17. Brunner LS, Smeltzer SC, Bare BG. Brunne and Suddarth's textbook of medical-surgical nursing. 12th ed. Philadelphia: Lippincott Williams and Wilkins; 2009.
18. Kneisl CR, Trigoboff E. Contemporary psychiatric mental health

- nursing. 2nd ed. Pearson Education, Inc. Upper Saddle River, New Jersey; 2009.
19. Sturt GW. Principle and practice of psychiatric nursing. 9th ed. Mosby, Elsevier publication :2009;??
20. Fortinash K. Psychiatric mental health nursing. 4th ed. Mosby: Elsevier; 2008.
21. Seligman ME, Csikszentmihalyi M. Positive psychology: An introduction. *Am Psychol* 2000;55:5-14.
22. Sotodehasl N, Taherneshat dost H, Kalantari M, Talebi H, Khosravi A. Comparison of effectiveness of two methods of hope therapy and drug therapy on the quality of life in the Patients with Essential Hypertension. *Journal of clinical psychology* 2010, 2 (1): 27-34.
23. Snyder CR. Handbook of hope: Theory, measures, and applications. USA: Academic Press; 2000.
24. Herth K. Development and implementation of a Hope intervention program. *Oncology Nursing Forum* 1998;28 (6): 1009-1017.
25. Snyder CR. The psychology of hope: You can get there from here. New York: Free Press; 1994.
26. Snyder CR, Lopez S. Striking a balance: A complementary focus on human weakness and strength. In: Lopez SJ, Snyder CR, editors. Models and measures of positive assessment. Washington, DC: American Psychological Association; 2003.
27. Porghaznin T, Ghafari F. Relationship between Hope and Self Steam of Clients with kidney transplantation in Emam Reza Mashhad Hospital. *J Yazd Med Sci Univ* 2005;1:52-61.
28. Snyder KM, Jerald K. Group therapy. Translated by In: Bahari S, Ranjgar BA, Hossinshahi H, Mirhashemi M, Naghshbandi S, editors. 7th ed. Tehran. Ravan Publications; 2011.
29. Baljani E, Kazemi M, Amanpour E, Tizfahm T. A survey on relationship between religion, spiritual wellbeing, hope and quality of life in patients with cancer. *Medical Sciences Journal of Islamic Azad University, Uremia Medical Branch* 2011;1:51-61.
30. Irving ML, Snyder CR, Gravel L, Hanke J, Hilberg P, Nelson N. Hope and the effectiveness of a pre- therapy orientation group from community mental health center clients. Seattle, WA: Western psychological Association Convention; 1997.
31. Klausner E, Snyder CR, Cheavens J. A hope-based group treatment for depressed older adult outpatients. In: Williamson GM, Parmlee PA, Shaffer DR, editors. *Physical Illness and Depression in older adults: A handbook of Theory, Research, and Practice*. New York: Plenum; 2000. p. 295-310.
32. Klausner EJ, Clarkin JF, Spielman L, Pupo C, Abrams R, Alexopoulos GS. Late-life depression and functional disability: The role of goal-focused group psychotherapy. *Int J Geriatr Psychiatry* 1998;13:707-16.
33. Hankins SJ. Measuring the efficacy of the Snyder hope theory as an intervention with an inpatient population. A dissertation presented for the doctorate of philosophy. Mississippi, United States: The University of Mississippi; 2004.
34. Shekarabi Ahari GH, Younesi J, Borjali A, Ansari Damavandi SH. The Effectiveness of group hope therapy on hope and depression of mothers with children suffering from Cancer in Tehran. *Iran J Cancer Prev* 2012;4:183-8
35. Cheavens SJ, Gum A, Feldman BD, Micheal ST, Snyder CR. A group intervention to increase hope in a community sample. San Francisco: American Psychological Association; 2001.
36. Ghasemi A, Baghban EI. Effectiveness of Group Hope Therapy Based Snyder hope theory on well-bing (fulfillment) of old people. *Khorasghan Islamic Azad Univ* 2009;44:17-40.
37. Bijari H, Ghanbari Hashemabadi B, Aghamohamadian H, Homaii SH. Effectiveness of Group therapy based on hope therapy on increase hope of women with cancer. *Psychol Educ Stud Mashhad Ferdooosi Univ* 2009;1:171-84.
38. Ebadi N, Sodani M, Faghihi A, Hossinpor M. Effectiveness education of optimism based on Koran verses on increase Hope of Ahvaz Divorced Women. *Khozestan: Khozestan Islamic Azad university publication*. 2009. p. 71-84.
39. Snyder CR, Irving L, Anderson JR. Hope and health: Measuring the will and the ways. In: Snyder CR, editor. *Handbook of social and clinical psychology: The health perspective*. New York: Pergamon Press; 2000.
40. Tsevat J. Spirituality/religion and quality of life in patients with HIV/AIDS. *J Gen Intern Med* 2006;21:S1-2.
41. Benzein EG, Berg AC. The level of and relation between hope, hopelessness and fatigue in patients and family members in palliative care. *Palliat Med* 2005;19:234-40.

How to site: ???

Source of Support: Grant No. 391027, **Conflict of Interest:** Nil.

Author Queries???

AQ1: Kindly confirm authors name

AQ2: Kindly provide page range

AQ3: Kindly confirm last name of author

AQ4: Kindly provide access date and month

EQ1: Please mention the ethical considerations

EQ2: Kindly provide research number of your project