

## Diarrhea-associated micronutrient deficiencies and risk of subsequent diarrhea in admitted children to Hajar hospital in Shahrekord, Iran

Bahman Khalili<sup>1</sup>, Antony Hart<sup>2</sup>, Masoud Mardani<sup>3</sup>, Mehdi Khalili<sup>1</sup>, Frank Mcardle<sup>2</sup>, Luis Cuevas<sup>4</sup>

<sup>1</sup> Department of Infectious Disease, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>2</sup> Department of Medical Microbiology and Genitourinary Medicine, Liverpool

<sup>3</sup> Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti Medical University, Tehran, Iran

<sup>4</sup> Liverpool School of Tropical Medicine, Liverpool, England

### ABSTRACT

**Background:** Acute infectious diarrhea is still one of the most important causes of death in childhood and malnutrition increases its morbidity and mortality. There is a strong correlation between the nutritional status of the child and the risk of subsequent diarrhea. Micronutrient deficiencies also increase the child's susceptibility to diarrhea and vitamin A and zinc supplementation has been shown to reduce the incidence and hasten recovery from acute diarrhea episodes.

**Materials and methods:** This study describes the association of nutritional deficiencies and other factors on the risk of subsequent diarrhea in children in Shahrekord, Iran. A cohort of 211 children less than 5 years old admitted with acute diarrhea to Hajar Hospital in Shahrekord, were followed for 14 weeks after hospital discharge.

**Results:** Fifty-eight (27%) of these children developed a new diarrhea episode during the follow up period. Children who were vitamin A and zinc deficient at the time of admission, above 12 months of age, kept animals at home or had weight-for-age and weight-for-height z scores <-1 during the univariate analysis had a higher risk of experiencing subsequent diarrhea. Vitamin A and zinc deficiencies, keeping animals at home, diarrhea duration  $\leq 4$  days on enrolment and weight-for-age z score <-1 remained as independent risk factors during multivariate analysis. The aggregation of these factors had a synergistic effect on the risk of subsequent diarrhea.

**Conclusion:** Children with micronutrient deficiencies and in contact with animal had the highest risk of suffering subsequent diarrhea. Our findings support the current approach of providing multiple micronutrient supplements for the prevention of infection in order to reduce mortality in children.

**Keywords:** *Micronutrient deficiencies, Children, Diarrhea, Iran.*

*(Iranian Journal of Clinical Infectious Diseases 2007;2(3):121-128).*

### INTRODUCTION

Acute infectious diarrhea is still one of the most important causes of death in childhood (1). Malnutrition increases the morbidity and mortality of diarrhea through its negative impact on non-

specific defence mechanisms and by altering the host immune mechanisms (2) and several studies have reported a strong correlation between the nutritional status of the child and the risk of subsequent diarrhea (3-5). Micronutrient deficiencies of vitamin A and zinc have also been shown to increase the child's susceptibility to acute and persistent diarrhea and supplementation of

Received: 21 November 2005 Accepted: 14 June 2006

Reprint or Correspondence: Bahman Khalili, MD.

Department of Infectious Disease and Tropical Medicine,  
Shahrekord University of Medical Sciences.

E-mail: bahman55\_khalili@yahoo.com

children with single micronutrients has been shown to reduce the incidence and hasten recovery from acute diarrhea episodes (6-8).

Although there is ample evidence that malnutrition and individual micronutrient deficiencies are associated with increased morbidity, most studies investigating these associations have focused on single micronutrients. There is thus a need to investigate if the presence of concurrent nutritional deficiencies increases the risk of subsequent diarrhea episodes in children.

## PATIENTS and METHODS

Children less than 5 years old admitted to Hajar Hospital with a clinical diagnosis of acute diarrhea were enrolled systematically during working days from October 2001 to August 2002. Hajar Hospital is a reference hospital serving the population of Chaharmahal va Bakhtiari province. Acute diarrhea was defined as the presence of three or more liquid or semi-liquid stools or a single watery stool per day of less than 14 days duration. After informed parental consent, parents were interviewed using standard questionnaires to obtain their socio-economic background, family characteristics and medical history. A total of 64 variables identified as risk factors in previous studies or considered to be potential risk factors within the local context were explored. The investigators did not modify the routine management of the children in the hospital and patients with clinically significant vitamin A deficiency should have received vitamin supplements. None of the children participating in the study, however, were deemed to require micronutrient supplements by the treating clinicians and the result of micronutrient concentration measurements were only available at the end of study.

After hospital discharge, children were followed up fortnightly by telephone, post or household visits for 14 consecutive weeks to assess if the child had experienced further diarrhea episodes.

Parents not available during a scheduled visit were given new appointments and new attempts were made to contact them.

A child was classified as having recurrent diarrhea if the mother indicated that the child had a new episode 14 or more days after the initial acute diarrhea episode.

Blood samples (3 ml) were collected at the time of enrolment during the initial diarrhea episode to assess the micronutrient status of the children. Blood samples were left to clot, centrifuged at 1500 rpm for 5 minutes and the sera were stored at -20°C. Serum samples were kept in dark-coloured safety tubes for the detection of vitamin A and E using the High-Performance Liquid Chromatography (HPLC) method, as described by Catignani et al (1983). Zinc, selenium and copper concentrations were measured by the Inductively Coupled Plasma–Mass Spectrometry (ICP –MS) method, as described by Delves et al (1997). Micronutrient measurements were performed at the end of the follow up period, once samples had been transferred to the U.K (Department of Medicine, University of Liverpool). Cut off values for each micronutrient were selected following internationally accepted cut off points to classify children as deficient or based on the statistical dispersion of the sample if reference cut off values were not available. Children with vitamin A and E concentrations <70 µmol/l and <16 µmol/l were defined as vitamin A and E deficient, respectively (9-11). Zinc deficiency was defined as a plasma serum concentration <9.94 µmol/l (12). Copper and selenium deficiencies were defined as values of <1.5 µmol/l and <0.90 µmol/l. These cut off values were selected as there are no internationally recognised cut off values for these micronutrients (13,14). Stool samples were collected at the time of enrolment and tested for the presence of known pathogens.

The characteristics of the children and their micronutritional status at the time of admission

were correlated to the presence or absence of subsequent diarrhea episode using univariate analysis. A conservative approach was used to select variables for multivariate analysis and variables with  $p$  values  $< 0.20$  were entered into a multiple logistic regression using the logistic regression programme of Epi 2002 to obtain adjusted odds ratios and 95% confidence intervals. Ethical issues in relation to all aspects were respected. Ethical approval has obtained from Research Ethic Committees of the Liverpool School of Tropical Medicine and the Shahrekord University of Medical Sciences.

## RESULTS

Two hundred and eleven children were enrolled. Eighty-nine (42%) of the participants were female and 122 (58%) male. The acute diarrhea episodes of the children at the time of enrolment had lasted for a median of 3 days (range 1 to 13 days). Most children had diarrhea for one (14%), two (20%) or three days (18%) and only 10 (5%) children had diarrhea for 13 days. One hundred and thirty eight (65%) participants had vomited the week before enrolment.

One hundred and eighty four (87%) children had complete follow up and 27 (13%) had missed one or more appointments. Three out of the 27 children with incomplete follow up had missed one visit, 6 two visits, 7 three visits and 11 had missed  $\geq 4$  visits. The mean (SD) age on enrolment was 15.6 (12.5) with a range from 2–59 months and the mean (SD) birth weight was 3.11 (0.56) Kgs. Only 19 (9%) children had a history of low birth weight. Children with complete follow up were compared to children who had missed one or more visits. Their medical history, clinical characteristics of the diarrhea episodes, laboratory results and family background were similar to those with incomplete follow up.

Fifty-eight (27%) children had subsequent diarrhea episodes during the follow up. Most

children (44) had one episode, 12 had 2 episodes and 2 children had three or more episodes as shown in table 1.

**Table 1.** Number of episode of subsequent diarrhea days with diarrhea and number of stools per day observed during the follow up period

	Number of	Number of children (%) (n=211)
Subsequent diarrhea episodes	0	153 (72%)
	1	44 (21%)
	2	12 (6%)
	3	1 (0.5%)
	4	1 (0.5%)
Days with diarrhea	0	153 (72%)
	1	34 (16%)
	2	12 (6%)
	3	8 (4%)
	4	1 (0.5%)
	$\geq 5$	3 (1.5%)
Watery stool per day	0	153 (72%)
	1	6 (3%)
	2	6 (3%)
	3	15 (7%)
	4	18 (9%)
	$\geq 5$	6 (3%)
Unknown		7 (3%)

The mean (SD) duration of the initial acute diarrhea episode was 4.3 (3.3) days compared to 1.9 (1.6) days for subsequent episodes ( $p < 0.01$ ). The mean (SD) number of stools per day in the children with subsequent diarrhea episodes was 3.2 (1.2) compared to 7.1 (3.8) stools per day for the initial acute episodes ( $p < 0.01$ ). Thirty-four (59%) of the subsequent episodes lasted for one day, 12 (21%) for 2 days and 12 (21%) for 3 or more days. None of the children developed persistent diarrhea and the longest duration of subsequent diarrhea episodes was 10 days. Children with subsequent diarrhea were older than children without further episodes with a mean (SD) of 9.6 (15.3) versus 14.1 (10.9) months, respectively ( $p < 0.05$ ). Forty-eight (83%) of the children with subsequent diarrhea episodes returned to the health services compared to 51 (32%) of the children without subsequent diarrhea ( $p < 0.01$ ).

**Table 2.** Characteristics of children with and without subsequent diarrhea

Characteristics	Followed (n= 211)	Subsequent diarrhea		OR (95% CI)	P value
		With (n=58)	Without(n=153)		
Male/female (% female)	122/89 (42%)	35/23(40%)	87/66 (43%)	0.87 (0. 5-1.7)	0.76
Breast fed for ≤ 6 months	34 (11%)	9 (16%)	25 (16%)	0.96 (0.4- 2.4)	0.90
Age >12 months	118 (56%)	23 (40%)	95 (62%)	2.48 (1.3-4.9)	0.005*
Birth weight <2500	18 (9%)	4 (7%)	14 (9%)	1.34 (0.4- 5.2)	0.78
Weight-for-height z score < -1	87 (42%)	33 (57%)	54 (35%)	2.50 (1.3-4.9)	0.005*
Weight-for-age z score < -1	121 (57%)	41 (71%)	80 (52%)	2.20 (1.1-4.4)	0.02*
<b>Family background</b>					
Keep animals at home	56 (27%)	23 (38%)	33 (22%)	2.39 (1.2-4.8)	0.01*
Mother <20 years	5 (10%)	1 (2%)	14 (9%)	0.18 (0.1-1.3)	0.07*
Father has manual job	99 (47%)	34 (59%)	24 (16%)	1.90 (1- 3.7)	0.06*
<b>Previous medical history</b>					
Diarrhoea in the previous year	92(44%)	30 (52%)	62(41%)	0.64 (0.3-1.2)	0.20
Hospitalized in previous year	71 (34%)	25 (43%)	46 (30%)	1.80 (0.9-3.5)	0.09*
<b>Initial diarrhea episode</b>					
Stool ≤ 4 times	67 (32%)	16 (28%)	51 (33%)	0.77 (0.4-1.6)	0.55
With blood in stools	23 (11%)	4 (7%)	19 (12%)	0.52 (0.1- 1.7)	0.35
Duration ≤ 4 days	139 (66%)	35 (60%)	73 (48%)	1.65 (0.9-3.2)	0.15*
Fever (previous 5 days)	162 (77%)	48 (83%)	114 (75%)	1.60 (0.7- 2.4)	0.31
Vomiting (previous week)	138 (65%)	41 (71%)	97 (63%)	1.37 (0.7- 2.8)	0.43
Mild dehydration	108 (51%)	36 (62%)	87 (57%)	1.24 (0.6-2.4)	0.60
Presence of pathogens in stool	142 (67%)	22 (38%)	60 (39%)	1.06 (0.56-2.1)	0.98
<b>Micronutrient status</b>					
Selenium <0.90 µmol/l	131 (69%)	35 (60%)	96 (63%)	0.87 (0.4-1.7)	0.78
Vitamin A <0.70 µmol/l	57 (27%)	25 (43%)	32 (21%)	2.91 (1.4-5.9)	0.001*
Zinc <9.94 µmol/l	158 (75%)	49 (84%)	109 (71%)	2.92 (0.9-10.7)	0.07*
Copper <1.5 µmol/l	89 (45%)	61 (43%)	23 (55%)	0.62 (0.29-1.3)	0.24
Vitamin E <16 µmol/l	165 (87%)	124 (89%)	41 (82%)	1.7 (0.64- 4.5)	0.34

\* Selected for multivariate analysis

**Table 3.** Multivariate analysis and adjusted odds ratios of the variables selected

Variable	Subsequent diarrhoea		OR (95%CI)	P value	AOR (95%CI)	P value
	With	Without				
Age <12 months	23 (40%)	95 (62%)	2.48 (1.3-4.9)	0.005		
Weight-for-height z score	33 (57%)	54 (35%)	2.50 (1.3-4.9)	0.005		
Weight-for-age z score	41 (71%)	80 (52%)	2.20 (1.1-4.4)	0.02	2.5 (1.3-5.3)	0.01
<b>Family background</b>						
Keep animals at home	23 (38%)	33 (22%)	2.39 (1.2-4.8)	0.01	2.7 (1.3- 5.5)	0.01
Mother age <20 years	1 (2%)	14 (9%)	0.18 (0.1-1.3)	0.07		
Father has manual job	34 (59%)	24 (16%)	1.90 (1-3.7)	0.06		
<b>Previous medical history</b>						
Hospitalised in previous year	25 (41%)	46 (31%)	1.80 (0.9-3.5)	0.09		
Diarrhoea in previous year	32 (55%)	60 (38%)	0.64 (0.3-1.2)	0.20		
<b>Index diarrhea episode</b>						
Duration ≤ 4 days	35 (60%)	73 (48%)	1.65 (0.9-3.2)	0.15	2.7 (1.2-6.1)	0.01
<b>Nutrition status</b>						
Vitamin A <0.70 µmol/l	25 (43%)	32 (21%)	2.91 (1.4-5.9)	0.001	2.8 (1.3-5.8)	0.006
Zinc <9.94 µmol/l	49 (84%)	109 (71%)	2.92 (0.9-10)	0.07	3.9 (1.2-12.8)	0.04

A summary of the univariate analysis of the risk factors investigated is shown in table 2. Children with anthropometric z scores <-1 (weight-for-height and weight-for-age) had an increased risk of subsequent diarrhea. Age <12 months had an inverse risk association, with younger infants experiencing less subsequent diarrhea than older children. Among the family background, children whose families kept animals at home and those who had a father working in manual activities had a higher risk of subsequent diarrhea, while children with young mothers (<20 years) had a lower risk. Children with a history of hospitalisation in the previous year were also more likely to experience subsequent diarrhea episodes following discharge from hospital. Seventy-six (36%) of the 211 children had rotavirus and 78 (37%) had other pathogens. Although the presence of rotavirus was associated with an increased risk of hospitalisation during the acute episodes on enrolment (15), the aetiology of the initial episodes was not associated with an increased risk of subsequent diarrhea. Among the micronutrients measured, vitamin A and zinc deficiency (<0.70  $\mu\text{mol/l}$  and <9.94 $\mu\text{mol/l}$  respectively) were associated with an increased risk of subsequent diarrhea, while selenium, copper, and vitamin E deficiency were not associated.

Vitamin A<0.70  $\mu\text{mol/l}$ , zinc<9.94  $\mu\text{mol/l}$  and weight-for-age zscore<-1 were nutritional variables independently associated with recurrent diarrhea after the multivariate analysis, as shown in table 3 (AOR = 2.8, 3.9, and 2.5 respectively). There was also a negative association between the duration of the initial diarrhea episode and the risk of subsequent diarrhea and children with shorter diarrhea episodes on enrolment were more likely to have subsequent diarrhea (AOR =2.7). Of the variables related to the environment and family background, only ownership of animals (AOR= 2.7) remained significant. The combination of the risk factors had synergistic effect. Children who had zinc and vitamin A deficiencies and kept

animals at home had the highest risk of subsequent diarrhea (AOR of 9.2).

## DISCUSSION

Despite the development of effective strategies for the management of acute diarrhea, this disease remains as one of the most important causes of death in childhood. Our study revealed that in this population, more than a quarter of the children with episodes of acute diarrhea severe enough to require hospitalisation, experience further diarrhea episodes after hospital discharge. The increased risk of recurrent diarrhea soon after an acute diarrhea episode has been recognised before, although there are surprisingly few studies that systemically investigate the risk factors for subsequent diarrhea. In Australia, aboriginal children and those who were severely dehydrated during the initial episodes, were more likely to suffer further episodes (16) and In the U.K, children attending health centres for acute diarrhea were six times more likely than population controls to have subsequent diarrhea episodes, with 10% of cases consulting their doctor in the 3 months following an initial episode and 2% being referred to the hospital (17). Information on the incidence and risk factors for recurrent diarrhea after an initial diarrhea episode in immunocompetent children in Africa or the Middle East is very scanty.

Diarrhea and other infections are often associated with malnutrition in a cycle that feeds each other. It has been estimated that 12 million children <5 years old die annually due to infection and malnutrition and about 70% these deaths are due to malnutrition(18-21). Malnutrition, particularly wasting, is a strong predictor of diarrheal disease duration, exacerbating nutritional faltering, thereby increasing the subsequent risk of death (22,23). The prevalence of micronutrient deficiencies in Iran, including zinc, vitamin A and

iodine deficiencies is high even though malnutrition, as assessed by anthropometry, is less prevalent and few children have weight-for-height z scores  $< -1$ . This may be a reflection of the poor quality of their diets more than a shortage of food (24). There is however a body of evidence that supplementation with micronutrients reduces the incidence and duration of diarrhea in children (25,2). Community-based trials have been shown that zinc or vitamin A supplementation of children reduces the incidence of childhood diarrhea (8,27-30). In most, but not all studies, vitamin A supplementation has a synergistic effect when applied in combination with zinc supplements (31).

One limitation of our study is that micronutrient concentrations were measured at the time of the acute episode not at the time of discharge from hospital and plasma zinc and vitamin A concentrations change vary rapidly during the acute phase of infection. The micronutrient concentrations measured therefore might not reflect long term deficiencies before or after the acute diarrhea. Despite this limitation, our study revealed that low vitamin A and zinc concentrations at the time of admission to the hospital were still independent and synergistic factors increasing the child's risk of subsequent diarrhea, suggesting that the low concentrations had a prolonged effect that extended to the follow up period. In addition to the low micronutrient concentrations, malnutrition defined as a weight-for-age z score  $< -1$ , keeping animals at home and experiencing a short initial episode of acute diarrhea were independent risk factors.

An association between diarrhea and the presence of animals in the close environment has been reported for *Campylobacter* (26), *Cryptosporidium* (32), *Salmonella typhimurium* (33), *Cyclospora* (34) and other pathogens, and outbreaks of diarrhea have been described in visitors to farms (35,36). It is therefore likely that contact with animals in our setting is associated with a higher frequency of exposure to pathogens

and to the lower hygiene standard of families residing in urban farms.

Although it was surprising that the aetiology of the initial episode was not a risk factor for subsequent diarrhea in our setting, this is in agreement with a report of Guinean children who had an increased incidence of subsequent diarrhea if the episode was caused by another pathogen other than rotavirus (37). Our children were not protected even if the initial diarrhea episode was due to rotavirus and further studies should be conducted in our setting. It is also interesting that children with shorter diarrhea episodes had a higher risk of subsequent diarrhea, independently of nutritional factors. This could be related to the infective dose of the organisms causing the diarrhea and children with severe episodes may have developed stronger immunological responses than children with mild infections.

Few studies to date have described the effect of a combination of factors on the risk of subsequent diarrhea. The aggregation of the factors had a synergistic effect on the risk of subsequent diarrhea and the children at the highest risk were those with a combination of micronutrient deficiencies and in contact with animals. Our findings support the current approach of providing multiple micronutrient supplementation for the prevention of infection and reduction of mortality in children.

## REFERENCES

---

1. Thapar N, Sanderson IR. Diarrhea in children: an interface between developing and developed countries. *Lancet* 2004;363:641-53.
2. Brown KH. Diarrhea and malnutrition. *J Nutr* 2003;133:328S-32S.
3. Kossman J, Nestel P, Herrera MG, et al. Undernutrition in relation to childhood infections: a prospective study in the Sudan. *Eur J Clin Nutr* 2000;54:463-72.
4. Kossman J, Nestel P, Herrera MG, et al. Undernutrition and childhood infections: a prospective study of childhood infections in relation to growth in the Sudan. *Acta Paediatr* 2000;89:1122-28.

5. Wierzba TF, El-Yazeed RA, Savarino SJ, et al. The interrelationship of malnutrition and diarrhea in a periurban area outside Alexandria, Egypt. *J Pediatr Gastroenterol Nutr* 2001;32:189-96.
6. Black RE, Sazawal S. Zinc and childhood infectious disease morbidity and mortality. *Br J Nutr* 2001;85 Suppl 2:S125-29.
7. Black RE. Zinc deficiency, infectious disease and mortality in the developing world. *J Nutr* 2003;133:1485S-89S.
8. Fawzi WW, Mise R, Spiegelman D, et al. Vitamin A supplements and diarrheal and respiratory tract infections among children in Dar es Salaam, Tanzania. *J Pediatr* 2000;137:660-67.
9. Agarwal DK, Singh SV, Gupta V, et al. Vitamin A status in early childhood diarrhea, respiratory infection and in maternal and cord blood. *J Trop Pediatr* 1996;42:12-14.
10. Hughes K, New AL, Lee BL, et al. Plasma vitamins A, C and E in the general population of Singapore, 1993 to 1995. *Ann Acad Med Singapore* 1998;27:149-53.
11. Wetherilt H, Ackurt F, Brubacher G, et al. Blood vitamin and mineral levels in 7-17 years old Turkish children. *Int J Vitam Nutr Res* 1992;62:21-29.
12. Hotz C, Peerson JM, Brown KH. Suggested lower cutoffs of serum zinc concentrations for assessing zinc status: reanalysis of the second National Health and Nutrition Examination Survey data (1976-1980). *Am J Clin Nutr* 2003;78:756-64.
13. Fisberg M, Castillo Duran C, Egana, J, et al. Plasma zinc and copper in infants with protein-calorie malnutrition. *Arch Latinoam Nutr* 1984;34:568-77.
14. Mutanda LN, Kinoti SN, Gemert W, et al. Age distribution and seasonal pattern of rotavirus infection in children in Kenya. *J Diarrheal Dis Res* 1984;2:147-50.
15. Khalili B, Cuevas LE, Reisi N, et al. Epidemiology of rotavirus diarrhea in Iranian children. *J Med Virol* 2004;73:309-12.
16. Lee AH, Flexman J, Wang K, et al. Recurrent gastroenteritis among infants in Western Australia: a seven-year hospital-based cohort study. *Ann Epidemiol* 2004;14:137-42.
17. Cumberland P, Sethi D, Roderick PJ, et al. The infectious intestinal disease study of England: a prospective evaluation of symptoms and health care use after an acute episode. *Epidemiol Infect* 2003;130:453-60.
18. Rice AL, Sacco L, Hyder A, et al. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ* 2000;78:1207-21.
19. Roy SK, Tomkins AM, Akramuzzaman SM, et al. Randomised controlled trial of zinc supplementation in malnourished Bangladeshi children with acute diarrhea. *Arch Dis Child* 1997;77:196-200.
20. Umeta M, West CE, Verhoef H, et al. Factors associated with stunting in infants aged 5-11 months in the Dodota-Sire District, rural Ethiopia. *J Nutr* 2003;133:1064-69.
21. Wekell P, Hakanson A, Krantz I, et al. Children in poor countries also have right to good health care. A new care program will reduce child mortality. *Lakartidningen* 1997;94:3637-41.
22. Patwari AK. Diarrhea and malnutrition interaction. *Indian J Pediatr* 1999;66:S124-34.
23. Stephenson LS, Latham MC, Ottesen EA. Global malnutrition. *Parasitology* 2000;121 Suppl:S5-22.
24. Mahmoodi MR, Kimiagar SM. Prevalence of zinc deficiency in junior high school students of Tehran City. *Biol Trace Elem Res* 2001;81:93-103.
25. Anand K, Sundaram KR, Lobo J, et al. Are diarrheal incidence and malnutrition related in under five children? A longitudinal study in an area of poor sanitary conditions. *Indian Pediatr* 1994;31:943-48.
26. Butzler JP. *Campylobacter*, from obscurity to celebrity. *Clin Microbiol Infect* 2004;10:868-76.
27. Bahl R, Baqui A, Bahn MK, editors. Effect of zinc supplementation on acute diarrhea. New Delhi, India, 2001;p:338-46.
28. Baqui AH, Black RE, El-Arifteen S, et al. Effect of zinc supplementation started during diarrhea on morbidity and mortality in Bangladeshi children: community randomised trial. *Br Med J* 2002;325:1059.
29. Gupta DN, Mondal SK, Ghosh S, et al. Impact of zinc supplementation on diarrheal morbidity in rural children of West Bengal, India. *Acta Paediatr* 2003;92:531-36.
30. Haidar J, Tsegaye D, Mariam DH, et al. Vitamin A supplementation on child morbidity. *East Afr Med J* 2003;80:17-21.
31. Rahman MM, Vermund SH, Wahed MA, et al. Simultaneous zinc and vitamin A supplementation in Bangladeshi children: randomised double blind controlled trial. *Br Med J* 2001;323:314-18.
32. Hunter PR, Hughes S, Woodhouse S, et al. Sporadic cryptosporidiosis case-control study with genotyping. *Emerg Infect Dis* 2004;10:1241-49.

**128 Diarrhea and micronutrient deficiency**

33. Dore K, Buxton J, Henry B, et al. Risk factors for Salmonella typhimurium DT104 and non-DT104 infection: a Canadian multi-provincial case-control study. *Epidemiol Infect* 2004;132:485-93.
34. Nimri LF, Hijazi SS. Cryptosporidium; A cause of gastroenteritis in preschool children in Jordan. *J Clin Gastroenterol* 2003;19:288-91.
35. Shield J, Baumer JH, Dawson JA, et al. Cryptosporidiosis-an educational experience. *J Infect* 1990; 21:297-301.
36. Shukla R, Slack R, George A, et al. Escherichia coli O157 infection associated with a farm visitor centre. *Commun Dis Rep CDR Rev* 1995;5:R86-90.
37. Valentiner-Branth P, Steinsland H, Fischer TK, et al. Cohort study of Guinean children: incidence, pathogenicity, conferred protection, and attributable risk for enteropathogens during the first 2 years of life. *J Clin Microbiol* 2003;41:4238-45.