

## Frequent use of stun gun may be associated with vibration-induced Raynaud's phenomenon: A case series

**Hafez Ghaheri, Mehrdad Karimi, Shervin Assari**

### ABSTRACT

**Introduction:** Stun guns are energy-conducting weapons that are becoming more frequently used by law enforcement officials to subdue combative individuals, or civilians for the case of security. Although complications related to stun guns are well known for the victim, less is known about stun guns complications among their users.

**Case Series:** In this case series, we reported vibration-induced Raynaud's phenomenon in four individuals who were recurrent users of stun guns. Two patients were male and two were female. Symptoms varied from moderate to severe. Although both hands were involved, symptoms were more severe in the right hand. Nifedipine was prescribed for all cases. Two patients received Losartan, as well. All patients reported a decrease in their symptoms in follow-up visits.

**Conclusion:** While further research is needed, clinicians should remain vigilant to stun gun usage as a possible cause of Raynaud's phenomenon.

 All Articles Open Access

## International Journal of Case Reports and Images (IJCRI)



International Journal of Case Reports and Images (IJCRI) is an international, peer reviewed, monthly, open access, online journal, publishing high-quality, articles in all areas of basic medical sciences and clinical specialties.

Aim of IJCRI is to encourage the publication of new information by providing a platform for reporting of unique, unusual and rare cases which enhance understanding of disease process, its diagnosis, management and clinico-pathologic correlations.

IJCRI publishes Review Articles, Case Series, Case Reports, Case in Images, Clinical Images and Letters to Editor.

**Website: [www.ijcasereportsandimages.com](http://www.ijcasereportsandimages.com)**

# Frequent use of stun gun may be associated with vibration-induced Raynaud's phenomenon: A case series

Hafez Ghaheri, Mehrdad Karimi, Shervin Assari

## ABSTRACT

**Introduction:** Stun guns are energy-conducting weapons that are becoming more frequently used by law enforcement officials to subdue combative individuals, or civilians for the case of security. Although complications related to stun guns are well known for the victim, less is known about stun guns complications among their users. **Case Series:** In this case series, we reported vibration-induced Raynaud's phenomenon in four individuals who were recurrent users of stun guns. Two patients were male and two were female. Symptoms varied from moderate to severe. Although both hands were involved, symptoms were more severe in the right hand. Nifedipine was prescribed for all cases. Two patients received Losartan, as well. All patients reported a decrease in their symptoms in follow-up visits. **Conclusion:** While further research is needed, clinicians should remain vigilant to stun gun usage as a possible cause of Raynaud's phenomenon.

**Keywords:** Raynaud's phenomenon, Vibration, Stun gun

### How to cite this article

Ghaheri H, Karimi M, Assari S. Frequent use of stun gun may be associated with vibration-induced Raynaud's phenomenon: A case series. *International Journal of Case Reports and Images* 2014;5(2):95–98.

doi:10.5348/ijcri-2014-02-449-CS-1

## INTRODUCTION

Stun guns have been developed as non-lethal devices for law enforcement officers to control potentially violent subjects. Thus, stun guns are used as alternatives for firearms. Stun guns apply high voltage, pulsatile electric shocks to the subject, which result in involuntary skeletal muscle contraction. As a result, the subject is unable to further resist [1].

The use of stun guns is showing an increase among law enforcement officials because of the increasing demand for non-lethal weapons to subdue combative individuals. In the United States, in the year 2005, stun guns were being used by about 40% of law enforcement agencies, with about 150,000 devices in use by police officers. However, the use of stun guns is not limited to law enforcement officers. Worldwide, 100,000 civilians own a stun gun [2, 3]. In Canada, in only one province, more than 10,000 stun gun uses were recorded during six years [4].

Literature has provided conflicting results regarding the safety profile of stun guns for the victims. Although some studies among healthy volunteers have shown safety of these weapons [5–8], severe injuries have also been reported with its daily usage [9–14]. Main

Hafez Ghaheri<sup>1</sup>, Mehrdad Karimi<sup>2</sup>, Shervin Assari<sup>3,4</sup>  
**Affiliations:** <sup>1</sup>Isfahan University of Medical Sciences, Isfahan, Iran; <sup>2</sup>Shahrekord University of Medical Sciences, Shahrekord, Iran; <sup>3</sup>Department of Health Behavior and Health Education, School of Public Health, University of Michigan, Ann Arbor, MI, USA; <sup>4</sup>Center for Research on Ethnicity, Culture, and Health (CRECH), University of Michigan School of Public Health, Ann Arbor, MI, USA.  
**Corresponding Author:** Mehrdad Karimi, MD Department of Surgery, Shahrekord University of Medical Sciences, Shahrekord, Iran, Address: Ayatollah Kashani Hospital Shariati Ave, Shahrekord, Iran; Ph: +98 983812224445; Email: mehrdad\_karimi@sums.ac.ir

Received: 09 August 2013  
Accepted: 05 September 2013  
Published: 01 February 2014

injuries reported in victims include basilar skull fracture, comminuted nasal fracture, and orbital floor fracture; concussion, facial laceration, and subarachnoid and epidural hemorrhage necessitate craniotomy. Penetration of the outer table and cortex of the cranium by stun guns are also reported [8, 15]. One study in the United States reported an incidence of 0.25% for significant injuries following the use of electrical weapons. Main complications in this study included intracranial injuries and rhabdomyolysis. Mortality has been also reported [16].

Published literature about the safety of stun guns is limited to the victims, not to the users. Herein, we reported four cases of vibration-induced Raynaud’s phenomenon in the persons who fire the weapon, recurrently. In all cases, diagnosis was based on the user’s history of firing the weapon, in addition to a clinical and laboratory examination. Standard clinical tests included cold stimulation test and nail-fold capillaroscopy. Antinuclear antibody (ANA), erythrocyte sedimentation rate (ESR), and C-reactive protein (CRP) were also measured for some cases [17]. In all cases, a thorough investigation ruled out diagnosis of vasculitis.

## CASE REPORT

Herein, we reported four cases of Raynaud’s phenomenon that may be related to stun gun use. Out of four patients, two individuals were users and the other two were sellers of stun guns. Two patients were men, and other two were women. Age ranged from 22 to 34 years.

Symptoms of Raynaud’s phenomenon had developed between 1 and 48 months after the initial use of stun guns. Stress or cold was the stimulant in three patients. Symptoms were severe in only one patient, other patients reported mild to moderate problems. Although both hands were involved in some cases, symptoms were always more prominent in one hand.

In all cases, treatment included avoiding use of stun guns in addition to medications. Nifedipine (for 18 to 90 days) was prescribed for all cases and two patients received Losartan (for 21 and 38 days), as well. All our patients reported decrease in symptoms in follow-up visits (Table 1).

## DISCUSSION

We believe that Raynaud’s phenomenon in our patients may be related to vibration. Thus, we believe the diagnosis in our patients is vibration-induced Raynaud’s phenomenon. This diagnosis should be considered as a new possible complication among the users of stun guns.

As explained before, most previously reported stun gun injuries of eyes, genitalia, and large blood vessels are related to the victims, not the weapon users [18, 19]. The mechanism of vibration-induced Raynaud’s phenomenon is as follows:

Raynaud’s phenomenon has been explained by pathology in the central nervous system, autonomic dysfunction, and vasodysregulation. Symptoms are known to be a result of receptor and nerve ending dysfunction [20]. Stoyneva et al. reported in 2003, that vibration induces endothelial damage, as a result of an increase in shear stresses and elevated plasma level of thrombomodulin and of von Willebrand factor and reduced endothelium-dependent vasodilator responses. In patients with vibration induced Raynaud’s phenomenon, high concentrations of endothelin-1 especially in most advanced stages, and a decreased plasma thiol level have shown an increased production and activity of free radicals, which contributes to vasospastic paroxysms [21].

We used Nifedipine for all our cases in combination with Losartan for two of our patients. Calcium channel blockers such as Nifedipine and angiotensin II receptor antagonist alone or in combination have shown therapeutic effects in Raynaud’s phenomenon [22, 23].

Table 1: Summaries of data of four patients with vibration-induced Raynaud’s phenomenon due to frequent use of stun gun.

Code	demographic data		History	Exposure to stun guns			Presentation			Treatment		Outcome
	Gender	Age (years)	Smoking	Exposure nature	Exposure (months)	Stress or cold stimulant	Right hand symptoms	left hand symptoms	Dry gangrene	Nifedipine (days)	Losartan (days)	Improved
#1	male	34	Yes	Seller	48	Yes	Moderate	Mild	No	28	-	Yes
#2	female	22	No	User	3	No	Moderate	-	No	60	21	Yes
#3	male	28	Yes	Seller	1	Yes	Moderate	Mild	No	90	-	Yes
#4	female	28	No	User	2	Yes	Severe	Moderate	Yes	18	38	Yes

Although vibration-induced Raynaud's phenomenon is a well-defined clinical entity, it is not very clear how stun gun users may develop this condition. Future research is needed on epidemiology, etiology, and natural history of Raynaud's phenomenon among individuals who use stun guns.

## CONCLUSION

Here we reported vibration-induced Raynaud's phenomenon in users of stun guns. Stun gun users, including both law enforcement officials and civilians, should be informed of the potential serious complications of recurrent firing. However, any further conclusion needs further studies.

\*\*\*\*\*

## Author Contributions

Hafez Ghaheri – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Mehrdad Karimi – Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Shervin Assari – Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

## Guarantor

The corresponding author is the guarantor of submission.

## Conflict of Interest

Authors declare no conflict of interest.

## Copyright

© Hafez Ghaheri et al. 2014; This article is distributed under the terms of Creative Commons attribution 3.0 License which permits unrestricted use, distribution and reproduction in any means provided the original authors and original publisher are properly credited. (Please see [www.ijcasereportsandimages.com/copyright-policy.php](http://www.ijcasereportsandimages.com/copyright-policy.php) for more information.)

## REFERENCES

1. McDaniel WC, Benwell A, Kovaleski S. Electrical parameters of projectile stun guns. *Conf Proc IEEE Eng Med Biol Soc* 2009;2009:3184–7.
2. United States Government Accountability Office. *TASER weapons: use of TASERs by selected law enforcement agencies*. Washington (DC): The Office
3. Hamilton A. Stun guns for everyone. *Time* 2002 Feb 4;159(5):50.
4. Review of the policing standards advisory committee 2009 available at: <http://www.mcscs.jus.gov.on.ca/stellent/groups/public/@mcscs/@www/@com/documents/webasset/eco81155.pdf>
5. Ho JD, Dawes DM, Bultman LL, et al. Respiratory effect of prolonged electrical weapon application on human volunteers. *Acad Emerg Med* 2007 Mar;14(3):197–201.
6. Ho JD, Miner JR, Lakireddy DR, Bultman LL, Heegaard WG. Cardiovascular and physiologic effects of conducted electrical weapon discharge in resting adults. *Acad Emerg Med* 2006 Jun;13(6):589–5.
7. Vilke GM, Sloane CM, Bouton KD, et al. Physiological effects of a conducted electrical weapon on human subjects. *Ann Emerg Med* 2007 Nov;50(5):569–75.
8. Mangus BE, Shen LY, Helmer SD, Maher J, Smith RS. Taser and Taser associated injuries: A case series. *Am Surg* 2008 Sep;74(9):862–5.
9. Winslow JE, Bozeman WP, Fortner MC, Alson RL. Thoracic compression fractures as a result of shock from a conducted energy weapon: A case report. *Ann Emerg Med* 2007 Nov;50(5):584–6.
10. Mehl LE. Electrical injury from Taser and miscarriage. *Acta Obstet Gynecol Scand* 1992 Feb;71(2):118–23.
11. Strote J, Range Hutson H. Taser use in restraint-related deaths. *Prehosp Emerg Care* 2006 Oct-Dec;10(4):447–50.
12. Kornblum RN, Reddy SK. Effects of the Taser in fatalities involving police confrontation. *J Forensic Sci* 1991 Mar;36(2):434–8.
13. Lee BK, Vittinghoff E, Whiteman D, Park M, Lau LL, Tseng ZH. Relation of Taser (electrical stun gun) deployment to increase in in-custody sudden deaths. *Am J Cardiol* 2009 Mar 15;103(6):877–80.
14. Reed LD. Deaths of people who received an electrical shock from conducted energy devices (CEDs) or “stun guns”. *Public Health Rep* 2009 Mar-Apr;124(2):187.
15. Injury Profile weapons <http://test.president.ufl.edu/committees/civil/documents/TaserArticles.pdf>
16. Bozeman WP, Hauda WE 2nd, Heck JJ, Graham DD Jr, Martin BP, Winslow JE. Safety and injury profile of conducted electrical weapons used by law enforcement officers against criminal suspects. *Ann Emerg Med* 2009 Apr;53(4):480–9.
17. National Heart, Lung, and Blood Institute (NHLBI); How Is Raynaud's Diagnosed? Available at: <http://www.nhlbi.nih.gov/health/health-topics/topics/raynaud/diagnosis.html>
18. Bleetman A, Steyn R, Lee C. Introduction of the TASER into British policing. Implications for UK emergency departments: An overview of electronic weaponry. *Emerg Med J* 2004 Mar;21(2):136–40.
19. Koscove EM. The taser weapon: A new emergency medicine problem. *Ann Emerg Med* 1985 Dec;14(12):1205–8.
20. Herrick AL. Pathogenesis of Raynaud's phenomenon. *Rheumatology (Oxford)* 2005 May;44(5):587–96.

21. Stoyneva Z, Lyapina M, Tzvetkov D, Vodenicharov E. Current pathophysiological views on vibration-induced Raynaud's phenomenon. *Cardiovasc Res* 2003 Mar;57(3):615–24.
22. Thompson AE, Pope JE. Calcium channel blockers for primary Raynaud's phenomenon: A meta-analysis. *Rheumatology (Oxford)* 2005 Feb;44(2):145–50.
23. Dziadzio M, Denton CP, Smith R, et al. Losartan therapy for Raynaud's phenomenon and scleroderma: clinical and biochemical findings in a fifteen-week, randomized, parallel-group, controlled trial. *Arthritis Rheum* 1999 Dec;42(12):2646–55.

## ABOUT THE AUTHORS

**Article citation:** Ghaheri H, Karimi M, Assari S. Frequent use of stun gun may be associated with vibration-induced Raynaud's phenomenon: A case series. *International Journal of Case Reports and Images* 2014;5(2):95–98.



**Hafez Ghaheri** is a vascular surgeon and an assistant professor at the Department of Surgery, Isfahan University of Medical Sciences, Isfahan, Iran. His research is focused on patients outcomes following surgical interventions.



**Mehrdad Karimi** is an assistant professor at Shahrekord University of Medical Sciences, Shahrekord, Iran. His research focus is on outcomes of different treatment modalities.



**Shervin Assari** is a postdoctoral research fellow at University of Michigan School of Public Health, Ann Arbor, Michigan. His research interests include health behaviors, mental health, and well-being. His research uses advanced quantitative techniques to explore determinants of health across different groups. He has published more than 100 research papers in peer review journals and 4 book chapters.

Access full text article on  
other devices



Access PDF of article on  
other devices





## Edorium Journals: An introduction

Edorium Journals Team

### About Edorium Journals

Edorium Journals is a publisher of high-quality, open access, international scholarly journals covering subjects in basic sciences and clinical specialties and subspecialties.

#### Invitation for article submission

We sincerely invite you to submit your valuable research for publication to Edorium Journals.

### But why should you publish with Edorium Journals?

In less than 10 words - we give you what no one does.

#### Vision of being the best

We have the vision of making our journals the best and the most authoritative journals in their respective specialties. We are working towards this goal every day of every week of every month of every year.

#### Exceptional services

We care for you, your work and your time. Our efficient, personalized and courteous services are a testimony to this.

#### Editorial Review

All manuscripts submitted to Edorium Journals undergo pre-processing review, first editorial review, peer review, second editorial review and finally third editorial review.

#### Peer Review

All manuscripts submitted to Edorium Journals undergo anonymous, double-blind, external peer review.

#### Early View version

Early View version of your manuscript will be published in the journal within 72 hours of final acceptance.

#### Manuscript status

From submission to publication of your article you will get regular updates (minimum six times) about status of your manuscripts directly in your email.

### Our Commitment

#### Six weeks

You will get first decision on your manuscript within six weeks (42 days) of submission. If we fail to honor this by even one day, we will publish your manuscript free of charge.

#### Four weeks

After we receive page proofs, your manuscript will be published in the journal within four weeks (31 days). If we fail to honor this by even one day, we will publish your manuscript free of charge and refund you the full article publication charges you paid for your manuscript.

#### Mentored Review Articles (MRA)

Our academic program "Mentored Review Article" (MRA) gives you a unique opportunity to publish papers under mentorship of international faculty. These articles are published free of charges.

#### Most Favored Author program

Join this program and publish any number of articles free of charge for one to five years.

#### Favored Author program

One email is all it takes to become our favored author. You will not only get fee waivers but also get information and insights about scholarly publishing.

#### Institutional Membership program

Join our Institutional Memberships program and help scholars from your institute make their research accessible to all and save thousands of dollars in fees make their research accessible to all.

#### Our presence

We have some of the best designed publication formats. Our websites are very user friendly and enable you to do your work very easily with no hassle.

#### Something more...

We request you to have a look at our website to know more about us and our services.

We welcome you to interact with us, share with us, join us and of course publish with us.



Edorium Journals: On Web



Browse Journals

CONNECT WITH US

