

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Iontophoretic skin permeation of peptides: an investigation into the influence of molecular properties, iontophoretic conditions and formulation parameters

¹, Michael S. Roberts^{2, 3}, Jeffrey Grice², Yuri G. Anissimov⁴, Hamid R. Moghimi^{3, 5}, Heather A. E. Benson^{1,*}

¹School of Pharmacy, Curtin Health Innovation Research Institute, Curtin University, Perth WA, ²Therapeutics Research Centre, School of Medicine, University of Queensland, QLD, ³School of Pharmacy and Medical Science, University of South Australia, Adelaide, South Australia, ⁴School of Biomolecular and Physical Sciences, Griffith University, QLD, Australia, and ⁵School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

* Corresponding author: Dr Heather Benson, School of Pharmacy, Curtin University, GPO Box U1987, Perth WA 6845, Australia. Tel: (+61)-8-92662338; Email: h.benson@curtin.edu.au

ABSTRACT:

The transdermal route offers advantages for delivery of peptides and proteins. However, these polar and large molecules do not permeate the skin barrier well. Various enhancement methods have been employed to address this problem. Iontophoresis is one of the methods that showing promise but its application to peptide delivery has yet to be fully explored. This study investigates the effects of different molecular properties and iontophoretic conditions on the skin permeation of peptides.

In this study, the permeation of alanine-tryptophan dipeptide (Mw = 276), alanine-alanine-proline-valine tetrapeptide (MW = 355), Argireline[®] (Acetyl hexapeptide-3, MW = 889) and Triptorelin acetate (decapeptide, MW = 1311) through excised human skin under passive or iontophoretic current of 0.4mA was studied using Franz diffusion cells. The effects of pH (3.0-7.4, to provide different net negative, neutral and positive charges), donor concentration (1-10 mg/ml), background electrolyte (34-137 mM NaCl and/or 5-20 mM HEPES) and current direction (anodal vs cathodal) were also investigated. Peptides were analysed by HPLC or liquid scintillation counting.

Iontophoresis led to a 30 times increase in peptide permeation relative to the passive flux for the peptides. Electroosmosis was an important determinant of the total flux for and even high molecular weight charged peptides. Electrorepulsion was found to be considerable for low molecular weight charged moieties. Permeation flux was decreased at lower pH, possibly due to decreased electroosmosis. Results also showed that 10 times increase in donor peptide concentration increases flux of peptides by about 2-4 times and decreases permeability coefficients by about 2.5-5 times. The addition of extra background electrolyte decreased the permeation coefficient of peptides by 2-60 times.

This study shows that iontophoretic permeation of peptides is affected by a number of parameters that can be optimized for effective transdermal peptide delivery.

Key words: Percutaneous absorption; Peptide and proteins delivery; Iontophoresis; Molecular weight and charge, Electro-osmosis; Skin penetration

INTRODUCTION

1
2 Many proteins and peptides have been identified that have therapeutic potential for skin
3 conditions [1], cardiovascular disease, Parkinson's disease, Alzheimer's disease, depression,
4 anxiety, attention deficit hyperactivity disorder, and many other disorders. Poor oral
5 bioavailability has led to ongoing research for a convenient and noninvasive delivery method.
6
7 The transdermal route offers different advantages and potential for protein and peptide
8 delivery if the considerable barrier to permeation presented by the stratum corneum can be
9 overcome [2]. Proteins and peptides present a significant challenge to successful transdermal
10 delivery due to the size and complexity of their molecular structure. In addition, most are
11 hydrophilic and often exist in a charged state, thus passive skin diffusion across the highly
12 structured lipid domains of the stratum corneum is unlikely to be effective. To overcome the
13 skin's barrier to peptides and proteins delivery, different enhancement methods have been
14 employed over the last decades, amongst which iontophoresis looks promising.
15
16

17
18 Iontophoresis involves the use of a mild electrical current (up to $0.5\text{mA}/\text{cm}^2$) to drive charged
19 and uncharged molecules across the skin. This method is considered non-invasive and
20 delivers consistent quantities of drug through skin. The positive or negative drug ions act as
21 charge carriers across the high impedance stratum corneum to enhance delivery of the drug
22 into the skin. There have been a number of detailed analyses of the mechanistic aspects of
23 iontophoresis and iontophoretic enhancement [3, 4]. Electrorepulsion (electromigration) is
24 known to be the main mechanism by which iontophoresis exerts its enhancement effect on
25 ionised solutes, but other factors including increased stratum corneum permeability in the
26 presence of an electric current flow, and electroosmosis (movement due to convection flow)
27 are also important, especially for uncharged molecules. Proteins and peptides are usually
28 charged at physiological pH or can be rendered charged by altering pH and are therefore
29 considered ideal candidates for iontophoretic delivery. Besides charge, other factors like
30 molecular weight and mobility of molecules also affect the permeation of peptides. However,
31 the iontophoresis of peptides and proteins requires further investigation, especially in the
32 areas of the effects of molecular weight on the mechanism of iontophoretic permeation
33 through the skin, that is the subject of the present investigation.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 In this study the influence of molecular size of peptides, the direction of applied current and
55 the effect of formulation parameters such as concentration, pH and presence of competing
56 ions on the iontophoretic delivery of five peptides (Figure 1) across human epidermal
57
58
59
60

1 membrane in vitro were investigated. The peptides investigated were 5-Aminolevulinic acid
2 (small peptide-like drug), a model dipeptide (L-Ala-L-Trp), a therapeutic tetrapeptide L-Ala-
3 L-Ala-L-Pro-L-Val, a cosmetic hexapeptide (Acetyl-hexapeptide-3 or Argireline[®]) and a
4 therapeutic decapeptide (Triptorelin acetate). These molecules provide a wide molecular
5 weight range of 130-1300 and different net charges of negative, neutral and positive. Where
6 possible peptides with potential therapeutic relevance have been chosen, as discussed below.
7
8
9

10
11
12
13 <Insert Figure 1>
14
15
16
17

18 5-Aminolevulinic acid (ALA) is a small molecular weight (131.1 g/mol), highly polar,
19 hydrophilic (pKa 4.0 and 8.4) molecule. It is the intermediate in heme biosynthesis and is the
20 precursor of protoporphyrin IX (PpIX), which is strongly fluorescent and undergoes
21 photobleaching rapidly producing singlet oxygen. Consequently, ALA is used as an
22 endogenous photosensitizer in photodynamic therapy for a range of non-melanoma skin
23 cancers like basal cell carcinoma, squamous cell carcinoma, actinic keratoses and Bowen's
24 disease, and in photo diagnosis [5]. ALA-PDT requires a long administration period of
25 approximately 4h at the skin surface to provide homogenous delivery at the target site.
26 Chemical and physical penetration enhancement techniques have been investigated to
27 improve the skin penetration of ALA [6-9].
28
29
30

31 L-Ala-L-Trp (Alanine-Tryptophan, called Ala-Trp in this manuscript) is a model dipeptide
32 molecule (molecular weight 275.9 g/mol), hydrophilic (alanine pKa 2.34 and 9.69,
33 tryptophan pKa 2.28 and 9.39), log P 0.61 ± 0.52.
34
35
36

37 L-Ala-L-Ala-L-Pro-L-Val (Alanine-Alanine-Proline-Valine; AAPV) is a small molecular
38 weight (355.4 g/mol), hydrophilic (logP -0.45 ± 0.65) tetrapeptide. It fits the P-P₁ subsites of
39 elastase and inhibits human neutrophil elastase (HNE). HNE acts as a pro-inflammatory
40 agent that is implicated in a number of conditions including rheumatoid arthritis, psoriasis,
41 atopic dermatitis and allergic contact dermatitis. Peptidic HNE inhibitors have a common
42 hydrophobic peptide sequence which partially mimics certain amino acid sequences found in
43 elastin. [10].
44
45
46
47
48
49
50
51
52
53
54

55 AC-Glu-Glu-Met-Gln-Arg-Arg-NH₂ (EEMQRR; Acetyl hexapeptide-3; Argireline[®]) is a
56 synthetic peptide (molecular weight 888.6 g/mol and log P -4.5 ± 0.95) that has been
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

incorporated into cosmeceutical products for anti-ageing properties [11]. It is designed from the N-terminal end of the protein SNAP-25 that inhibits Ca^{2+} -dependent catecholamine release from chromaffin cells [12]. It acts by down-regulation of muscle action and is applied to reduce wrinkles around the eyes.

p-Glu-His-Trp-Ser-Tyr-[D-Trp]-Leu-Arg-Pro-Gly-NH₂ (EHWSYWLRPG; Triptorelin available as acetate and pamoate) is a synthetic gonadotropin-releasing hormone (GnRH) agonist and a long-acting LHRH analogue (molecular weight: 1311.5 g/mol, pKa: 7.2, 9.5, 12 and log P 0.3 ± 1.41). It is marketed as a sustained release injection for treatment of infertility, endometriosis and hormone-responsive cancers.

MATERIALS AND METHODS

Materials

All chemicals and reagents listed below were used as supplied: 5-aminolevulinic acid hydrochloride (ALA) (A3785 ~98%), L-Alanyl-L-Trptophan (05400 ≥ 99%), N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid HEPES (H3375 ≥99.5% titration) and fluorescamine (F9015 ≥98%) were purchased from Sigma-Aldrich (Sydney, NSW, Australia). AAPV (AV-4-NH₂, P100401-SY121097, ≥96% purity), Acetyl-hexapeptide-3 (Ac-ER-6, 54947, ≥95% purity) and Triptorelin acetate (HG-9, 5773-63-4, ≥98% purity) were purchased from GL Biochem Ltd (Shanghai, China), Phosphate buffer saline was prepared according to United States Pharmacopeia. Sodium chloride (NaCl) from Asia Pacific Speciality Chemicals Ltd (Seven Hills, NSW, Australia), analytical grade potassium chloride (KCl) and sodium hydroxide (NaOH) were purchased from Merck Pty Ltd. (Darmstadt, Germany) and boric acid was purchased from Ajax Finechem (Taren Point, NSW, Australia). Analytical grade acetonitrile was purchased from JT Baker (Philipsburg, NJ, USA) and trifluoroacetic acid (TFA) from Sigma-Aldrich (Sydney, NSW Australia). Triethylamine (TEA) from Fluka Chemika (Switzerland) and orthophosphoric acid from Ajax Finechem (Australia). Silver (Ag)-wire (99.99%, 1 mm diameter) was obtained from Sigma-Aldrich and purified deionised Milli-Q water from Millipore (North Ryde, NSW Australia).

Quantitative Analysis

1 The quantitative analysis of all peptides was by reverse phase high performance liquid
2 chromatography (HPLC) and liquid scintillation counting for ALA. The chromatographic
3 system consisted of an Agilent 1100 series equipped with a quaternary pump (G1311A),
4 autosampler (G1313A), solvent-degasser (G1312A) with a photo diode array
5 detector/fluorescence detector (G1321A). Separation was achieved on a Phenomenex Jupiter
6 C18 300A column (5 μ m, 150mm \times 4.6 mm) with a guard column (wide pore C18). Peak
7 integration was undertaken on a personal computer using Chemstation A08.01 software. All
8 standards and samples were prepared at room temperature. The radiolabelled samples were
9 analysed on a liquid scintillation counter (Packard LSC: Canberra, CT, AUS).

16 The HPLC mobile phase consisted of water and acetonitrile buffered with trifluoroacetic acid
17 (TFA), triethylamine or phosphoric acid. The ratio of aqueous phase to organic phase was
18 dependent on the compound: 30:70 (0.1% TFA) isocratic system for ALA. ALA belongs to
19 the class of alpha amino ketones with a weak chromophoric carbonyl group and is therefore
20 unsuitable for quantification by ultraviolet (UV) absorption spectroscopy. Precolumn
21 derivatisation step with fluorescamine generated a fluorescent compound that was detected
22 using an excitation wavelength of 395nm and emission wavelength of 480nm [8]. For Ala-
23 Trp a linear gradient system, aqueous (0.045% TFA): acetonitrile (0.036% TFA) from 10% to
24 100% (UV detection at 210nm), for AAPV a gradient system, aqueous (0.1% TFA):
25 acetonitrile (0.1% TFA) from 5% to 20% (UV detection at 220nm), for acetyl hexapeptide
26 90:10 (0.05% TFA) isocratic system (UV detection at 210nm) and for triptorelin 75:25
27 (20mM triethylamine and 50mM phosphoric acid) isocratic system (UV detection at 280nm).
28 The R.S.D of repeatability was less than 1% and the lowest limit of quantification was in the
29 range of 12 to 320 *ng* for all the peptides assayed.

43 Radiolabelled ALA (0.5 μ Ci/diffusion cell) were mixed with scintillation cocktail (Emulsifier
44 safe, Perkin Elmer, Waltham, MA, USA) and analysed for radioactivity.

50 **Skin Preparation**

51 Ethical approval for using human skin was obtained from the Human Research Ethics
52 Committee of Curtin University prior to the study. Full thickness human skin (abdominal
53 section) excised from female patients following abdominoplasty surgery at Perth hospitals
54 was used. Skin was prepared by removing the subcutaneous tissue by dissection then
55
56
57
58
59

1 isolating the epidermal membranes from human skin by the heat separation method [13]
2 where the full thickness human skin was immersed in water at 60°C for 1 min. The epidermal
3 membrane was then teased off the dermis; placed onto aluminium foil with the stratum
4 corneum layer facing upward, air dried and then stored at -20°C (for not more than 2 months)
5 until required.
6
7
8
9

10 11 **Iontophoretic in-vitro Permeation Studies**

12 All iontophoresis experiments were conducted across human epidermal membrane using
13 Pyrex glass side-by-side diffusion cells, enabling permeation across skin sections of cross
14 sectional area 0.95 cm²; donor and the receptor volume of 1.8 ± 0.1 mL. The membrane was
15 allowed to equilibrate for 1 h in a water bath maintained at 35 ± 0.5°C. Membrane integrity
16 was then determined by visual inspection over a bright light and by electrical resistance (kΩ)
17 measurement using a digital portable LCR multimeter (TH2821/A/B, Changzhou Tonghui
18 Electronic Co Ltd, Jiangsu Province, China). The measurements were taken by immersing the
19 silver-silver chloride electrode attached to the probe lead tips, one each in the donor and
20 receptor compartments. Membranes exhibiting an electrical resistance less than 20 kΩ were
21 rejected.
22
23
24
25
26
27
28
29
30
31

32 The receptor compartment was always filled with electrolyte solution (20mM HEPES +
33 137mM NaCl). The donor compartment consisted of ALA, Ala-Trp, AAPV,
34 acetylhexapeptide-3 or triptorelin acetate prepared with water or receptor electrolyte solution
35 at various concentrations.
36
37
38
39

40 A constant iontophoretic current of 0.4mA was supplied by a custom-built power supply unit
41 (Art Electro NSW, Australia) for 2 h unless otherwise stated. Passive diffusion cells were
42 treated in the same way but without the electrodes. Samples were collected from the receptor
43 compartment and replaced with fresh pre-warmed electrolyte solution throughout the
44 experimental period. The data was plotted as mean cumulative amount permeating to the
45 receptor compartment versus time. Transdermal permeation flux and permeability coefficient
46 were calculated over the linear portion of the plot using Fick's first law.
47
48
49
50
51
52

53 ***Influence of concentration:*** The donor compartment consisted of Ala-Trp at 3.6 and 36 mM
54 (1 and 10 mg/ml), AAPV at 2.84 and 28.4 mM (1 and 10 mg/ml), acetylhexapeptide-3 at 1.13
55 and 11.3mM (1 and 10 mg/ml) and triptorelin at 0.9 and 9mM. The solutions were prepared
56
57
58
59
60
61
62
63
64
65

1
2 in water or 20 mM HEPES + 137 mM NaCl. The later electrolyte was also used as the
3 receptor electrolyte solution.

4 ***Influence of pH:*** The donor solution pH was maintained at physiological pH 7.4 or at pHs
5 6.0, 5.5, 5.0, and 3 using 0.1M or 0.01M sodium hydroxide (NaOH) and/or 0.1 or 0.01M
6 hydrochloric acid (HCl) solutions. The net electrical charges of peptides were calculated at all
7 pHs using Calculator Plugins for structure property prediction and calculation, Marvin 6.0.0,
8 2013, ChemAxon (<http://www.chemaxon.com>) [14].
9

10
11
12 ***Influence of background electrolyte:*** The donor solutions of Ala-Trp, AAPV and triptorelin
13 were prepared in water or receptor electrolyte solutions (20 mM HEPES + 137 mM NaCl).
14 The donor solution for acetylhexapeptide-3 (EEMQRR) was prepared at six different
15 combinations (20, 10, 5 mM HEPES + 137, 68.5, 34.2 mM NaCl respectively and 137, 68.5
16 and 34.2 mM NaCl alone).
17

18
19 ***Influence of direction of current:*** For anodal iontophoresis, active cells received the silver
20 electrode in the donor compartment and silver chloride electrode in the receptor compartment,
21 and vice versa for cathodal iontophoresis.
22

23 24 25 26 27 28 29 30 31 **Statistical Analysis**

32 The permeation data for ALA was analyzed using SPSS 16.0 for Windows (SPSS Inc.,
33 Chicago IL). The data for other peptides were analyzed using The Mixed Procedure in SAS
34 version 9.1 for Windows (SAS Institute Inc., Cary, NC, USA). The Mann-Whitney U Test (1-
35 tailed) was used to determine significant differences between all parameters.
36
37
38
39
40
41
42
43

44 **RESULTS**

45 Table 1 provides the passive and iontophoretic flux and permeability coefficients of peptides
46 with neutral or positive net charges under anodal iontophoresis. The results show that
47 iontophoresis has been able to increase permeability coefficient of these peptides by about 7-
48 27 times. Table 1 shows that, except for ALA, irrespective of the molecular weight (300-
49 1300 Da) and charge (0 to 1.1) of the corresponding peptides, the permeability coefficients of
50 di, tetra, hexa and deca-peptides are very close ($12-17 \times 10^{-3}$ cm/h). This means that, at
51 similar concentrations, these molecules permeate the skin at almost identical fluxes,
52
53
54
55
56
57
58
59

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

regardless of their differences in physicochemical properties. Such behaviour might indicate that electroosmosis is the main driving force for iontophoresis of these molecules. Consistent with this suggestion, the estimate electroosmotic volume flux, based on the peptide fluxes and concentrations, ranges from 10 to 17 $\mu\text{l}/\text{cm}^2/\text{h}$ for these four peptides and the conditions used. However, as these peptides are zwitterionic and the skin is cation permselective, electromigration as a determinant of peptide permeation is a potential confounder.

<Insert Table 1>

Influence of charge and pH: Table 2 provides the effects of pH on charge and iontophoretic permeation of peptides. The iontophoretic permeability coefficient of ALA increased from $1.3 \times 10^{-3} \text{ cm/h}$ to $11 \times 10^{-3} \text{ cm/h}$ when the pH was increased from 5.5 to 7.4 [15]. In contrast, Ala-Trp showed a significant increase ($P < 0.0001$) in the permeability coefficient from $0.4 \times 10^{-3} \text{ cm/h}$ to $13 \times 10^{-3} \text{ cm/h}$ when the pH was decreased from 7.4 to pH 5.5. The permeability coefficient of AAPV reduced significantly from $18 \times 10^{-3} \text{ cm/h}$ at pH 7.4 to $0.4 \times 10^{-3} \text{ cm/h}$ ($P < 0.0001$) at pH 3.0 and no change at pH 5.5 ($18 \times 10^{-3} \text{ cm/h}$). Acetylhexapeptide-3 also demonstrated a significantly reduced permeability coefficient from $17 \times 10^{-3} \text{ cm/h}$ at pH 7.4 to $4.8 \times 10^{-3} \text{ cm/h}$ at pH 5.0 ($P < 0.0001$).

<Insert Table 2>

Influence of concentration: The effect of donor peptide concentration is presented in Table 3. A 10x increase in donor concentration of Ala-Trp, from 3.6 mM to 36 mM (from 1mg/ml to 10 mg/ml) at pH 5.5 resulted in a 4.2x increase in iontophoretic flux but reduced the permeability coefficient from $13.0 \times 10^{-3} \text{ cm/h}$ to $5.5 \times 10^{-3} \text{ cm/h}$. A 10x increase in donor concentration of acetylhexapeptide-3 from 1mg/ml to 10 mg/ml at physiological pH, resulted in a 2.5x increase in the iontophoretic epidermal flux but decreased the permeability coefficient by about 4 times. The same trend was observed for triptorelin where a 2x increase in flux and 5x decrease in permeability coefficient was observed after a 10x increase in the peptide concentration (Table 3).

<Insert Table 3>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

Influence of background electrolyte:

The presence of HEPES and Na⁺ ions significantly reduced the iontophoretic delivery of all peptides (Table 4 and Table 5). A 14x reduction in permeability coefficient of Ala-Trp resulted from the donor solution of 20mM HEPES and 137mM NaCl at pH 5.5 compared to water. A less pronounced reduction of less than 2x in the permeability coefficient of AAPV, EEMQRR and Triptorelin occurred with 20 mM HEPES and 137 mM NaCl (Table 4).

To investigate the effects of electrolytes type and concentration, the influence of different HEPES (10 and 5 mM) + NaCl (68.5 and 34.2 mM) systems and different NaCl solutions alone (137, 68.5 and 34.2 mM) were studied for EEMQRR iontophoretic permeation. The permeability coefficient data for all donor solutions is presented in Table 5 in comparison to base system. The results clearly indicate that a high concentration of ions in the delivery vehicle reduces the iontophoretic permeability coefficient of peptides. The extent of reduction was more for NaCl alone (11-57x) in comparison to HEPES-containing systems (less than 2x reduction). The permeability reduction effects of background electrolytes were also concentration-dependent and the effect increased with increase in electrolyte concentration (Table 5).

<Insert Table 4 and Table 5>

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

Influence of direction of current: anodal v cathodal: The direction of the current can influence the iontophoretic skin permeation depending on the predominant charge carrier. The permeability coefficient of AAPV and acetylhexapeptide-3 that permeated human epidermis under the influence of anodal versus cathodal current is presented in Table 6. The direction of current did not influence iontophoretic delivery of AAPV. In contrast the iontophoretic permeability coefficient of acetylhexapeptide-3 under cathodal current was similar to passive delivery $[(1.2 \pm 0.3) \times 10^{-3} \text{ cm/h}]$ compared to $(17 \pm 2) \times 10^{-3} \text{ cm/h}$ with anodal current.

<Insert Table 6>

55
56
57
58
59
60
61
62
63
64
65

DISCUSSION:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 shows that, regardless of the molecular weight (300-1300 Da) and charge (0 to 1.1) of the peptides, the permeability coefficients are very similar ($12-17 \times 10^{-3}$ cm/h). This suggests that, irrespective of differences in their physicochemical properties, these molecules permeate the skin with effectively the same fluxes. Such behaviour is consistent with electroosmosis being the main determinant of the anodal iontophoretic flux for each of these molecules. The other mechanisms involved in the iontophoretic enhancement of skin permeation of drugs are: electro-repulsion (electromigration) and changes in skin permeability (skin damage) due to electrical current [16]. In contrast to passive skin permeation, where the unionised form of a drug molecule permeates better than its ionised form, it is the ionised solute that is transported across the skin in iontophoresis. Here, the iontophoretic electrical potential acts as an external force to push ionized drug molecules across the skin (electromigration). Cathodal iontophoresis is applied to repel a negatively charged drug from the negative electrode across an epidermal membrane, whilst anodal iontophoresis is used for positively charged drug molecules [17]. Electroosmosis produces bulk fluid flow in the membrane and occurs when a voltage difference is imposed across a charged membrane. This fluid stream carries all cargos (ions or neutral species) dissolved within it and always flows in the same direction as the flow of counterions (i.e. ions of opposite charge to the membrane charge) [16]. Human skin is negatively charged above about pH 4 and counterions are positive ions at this condition, thus electroosmotic flow occurs from anode to cathode [16]. Electroosmosis applies for both charged and uncharged drugs and its rate depends on solvent volume flux and concentration of the cargo. On the other hand, dependency of electroosmosis on molecular weight is expected to be lower than other mechanisms, as discussed later in this paper. Therefore, the similarity of permeability coefficients for different molecules (charged and uncharged with different molecular weight, Table 1) might show the important role of electroosmosis in peptides penetration in the present investigation. Considering this argument, the electroosmotic volume flux in the conditions used in the present study was calculated to be about $10-17 \mu\text{l}/\text{cm}^2/\text{h}$. An electroosmotic flow of up to $37 \mu\text{l}/\text{cm}^2/\text{h}$ in hairless mouse skin, was reported by Pikal and Shah [18, 19]. However, as discussed later, the presence of other mechanisms, including electrorepulsion, cannot be dismissed specially for low molecular weight peptides.

In contrast to the above-mentioned argument, ALA showed a low permeability coefficient. This molecule is slightly positively charged and its low permeation cannot be justified based on electroattraction. Its low permeation might be due to its high concentration of 60 mM. As

1 ALA is a small molecule with a positive charge, the high concentration might cause
2 neutralization of skin negative charge and therefore decrease the electroosmosis. Our studies
3 show that the same drug at pH 7.4 [15], where it contains a net charge of -0.3, shows an
4 anodal permeability coefficient of about 11×10^{-3} cm/h, while negative charges are expected
5 to show lower permeability in anodal iontophoresis in comparison to neutral moities.
6
7 Considering the above-mentioned volume flow and the fact that this compound is negative
8 and small, these data might show that both electromigration and electroosmosis are playing
9 important roles in permeation of this low molecular weight compound.
10

11
12
13
14
15 As shown in Table 1, triptorelin has a positive net charge of about +1 at pH = 7.4. Therefore,
16 at least theoretically, a good contribution from electromigration is expected for this molecule
17 for anodal iontophoresis. Thus a higher permeability is expected in comparison to those
18 obtainable by electroosmosis only. As discussed by Pikal [16], electroosmotic flow increases
19 in importance as the size of the drug ion increases. The 'ionic' or Nernst-Planck effect
20 (electromigration) is the largest contributor to flux enhancement for small ions. Increased
21 skin permeability or the skin 'damage effect', is a significant factor for both large and small
22 ions, particularly for experiments at high current density. For monovalent ions with Stokes
23 radii larger than about 1 nm, electroosmotic flow is the dominant flow mechanism [16]. This
24 effect is in a way that transdermal delivery of a large anion (or negatively charged protein)
25 from the anode compartment can be more effective than delivery from the cathode
26 compartment [16]. The present data might show that permeation of triptorelin under the
27 effects of iontophoresis is mainly ruled by electroosmosis.
28
29

30
31
32
33
34
35
36
37
38
39 Table 2 shows the effects of pH on iontophoretic permeation of peptides. pH affects
40 permeation of drugs in iontophoresis in different ways. The pH of the donor solution plays an
41 important role in ionisation of molecules (their electromigration) and the charge of the skin
42 (skin permselectivity) and therefore electroosmosis [20]. The pH of the solution can also
43 affect the chemical stability of peptides and proteins, as was previously reported for ALA [7,
44 15]. Isoelectric point (pI) of skin is about 4-5 [21] and skin shows negative charges at pHs
45 above this value and positive charges at lower pHs. Bath et al. [22] showed that
46 electroosmotic flow is highly pH dependent and is from anode to cathode at pHs > 3.5 and
47 from cathode to anode at pHs < 3.5. They also showed that anodal electroosmosis increases
48 with pH increase above 3.5 and cathodal electroosmosis increases with pH decrease below
49 3.5.
50
51
52
53
54
55
56
57
58
59

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

As shown in Table 2, both Ala-Trp and AAPV are neutral/slightly positive at pH 5.5 and show similar permeability coefficients that are explainable by electroosmosis as discussed above. When the pH is increased to 7.4 both peptides become negative. In this condition, an electromigration in an opposite direction to that of electroosmosis is expected. This might explain the sharp decrease in permeability of Ala-Trp from 13 to 0.4, but cannot explain the similarity of permeation of AAPV at pHs 5.5 and 7.4. This might show the importance of electroosmosis for permeation of AAPV and electromigration for permeation of Ala-Trp. Such behaviour might be due to the difference in their molecular weight as discussed above, indicating that electroosmosis is important for higher molecular weights and electromigration for lower molecular weights. However, we cannot dismiss the importance and the role of electromigration at these conditions.

As discussed above, decreased pH will decrease electroosmotic flow from anode to cathode and can even reverse it when the pH becomes lower than 3.5. This might explain the decreased K_p of EEMQRR from 17×10^{-3} cm/h at pH = 7.4 to 4.8×10^{-3} cm/h (close to its passive permeation) when pH is decreased from 7.4 to 5. For AAPV, permeability coefficient decreases from 18×10^{-3} cm/h at 7.4 to 0.4×10^{-3} cm/h at pH 3.0. These reductions happen in spite of the positive charges of these molecules (0.2 and 0.9 for EEMQRR and AAPV respectively) at lower pHs (Table 2) that is expected to increase the flux through electrorepulsion. This again shows that electroosmosis is very important in the permeation of these higher molecular weight peptides. A similar finding was reported for thyrotropin-releasing hormone (TRH) [23]. Greater iontophoretic transport was achieved in its unionised form at pH 8 by electroosmosis than at pH 4 where it is 99% cationic; this was attributed to the reverse in skin permselectivity from cationic to anionic at acidic pH, and electroosmosis occurs from cathode to anode.

Increasing the donor peptide concentration increased the iontophoretic flux of the peptides, but the flux increase was not equivalent to increase in concentration. A 10x increase in the peptide concentration resulted in only 2-4 times increase in flux, indicating 2.5-5x decrease in the permeability coefficients (Table 3). The effect of concentration on the iontophoretic permeation of drugs is complicated. Delgado-Charro et al. [24] showed that when drugs neutralize the skin charge, increased drug concentration decreases the permeability through decreased and even reversal of electroosmosis. This might explain in part the present data. However, the same group suggested that when electrorepulsion is the major mechanism, the

1 permeability coefficient is expected to stay constant (linear relationship between
2 concentration and flux). Thus another explanation is the possible iontophoretic charge
3 saturation, as was reported for LHRH and two analogues [25].
4

5
6 Marro et al showed that in the absence of background electrolyte the iontophoretic fluxes of
7 the three drugs lidocaine, quinine and propranolol are essentially independent of donor
8 concentration over a 100-fold range, while in the presence of background electrolyte
9 lidocaine delivery increased linearly with concentration as it competed more and more
10 effectively with Na⁺ to carry the charge across the skin. However, iontophoretic delivery of
11 quinine and propranolol increased non-linearly with concentration [26].
12
13
14
15
16

17 The sum of the flux of individual ions must equal the current applied by the power source; in
18 other words, there is competition among all the ions present to carry the charge.
19 Physicochemical characteristics such as electrical mobility and concentration of the ion
20 concerned, and the properties of the media through which the ion is moving, determine the
21 probability of the ion being the major carrier, and eventually being efficiently transported
22 across the skin [27, 28]. The concentration of an ionised species is said to influence the
23 electro-migration component of the iontophoretic flux, when it is applied along with
24 counterions and co-ions (buffers, electrolytes and preservatives in the formulation) [29]. In
25 such cases, molar fraction of the drug molecule that plays a major role in iontophoretic
26 delivery, as this is the concentration relative to that of the competing ions [26, 28]. The
27 electro-diffusion model developed by Kasting and Keister based on single ion transport
28 suggests that, in the absence of competing cations, iontophoretic flux is independent of
29 concentration and dependent only on the ratio of mobilities of the cation and the main
30 counterion (usually Cl⁻) arriving from beneath the skin [30].
31
32
33
34
35
36
37
38
39
40
41
42

43 The present results show that the presence of 20 mM HEPES + 137 mM NaCl reduced the
44 permeability coefficient of peptides (Table 4). The highest reduction was for Ala-Trp (about
45 14x). Other peptides showed about 2x difference regardless of their charges and molecular
46 weight. To further investigate the effects of addition of background electrolytes, the effects of
47 5-20 mM HEPES and 34-137mM NaCl was investigated for the hexapeptide (EEMQRR:
48 Table 5). The presence of co-ions and counterions revealed a linear trend in the permeation of
49 acetyl hexapeptide-3, with the linear reduction in concentration of co-ions and counterions
50 present (20, 10, 5 mM HEPES with 137, 68.5, 34.2 mM NaCl respectively). Significantly
51 lower amounts of acetyl hexapeptide-3 permeated when the peptide donor solution was
52
53
54
55
56
57
58
59

1 prepared in NaCl (137, 68.5, 34.2 mM) alone. Results show that the reduction effects of
2 added electrolyte is both electrolyte type and concentration dependent. In agreement with the
3 present data, Bellantone et al [31] demonstrated a reduction in benzoate flux by more than
4 half in the presence of equimolar amounts of sodium chloride. Cázares-Delgado et al [32]
5 presented a similar observation with cytochrome C-A, a 12.4 kDa protein. The iontophoretic
6 delivery of the macromolecule was reduced 3.9-fold when the sodium ion concentration
7 increased to 170 mM. These effects can be related to the effects of added electrolytes on both
8 electrorepulsion and also electroosmosis, as described below.
9

10
11
12
13
14
15 Addition of a background electrolyte, buffer or substance that could increase the ionic
16 strength of the formulation increases the competition between the charge carriers and can
17 affect iontophoretic drug flux. In practice, multiple ions exist as co-ions and counterions.
18 Hence knowledge of the transport number of a drug molecule becomes essential. Transport
19 numbers are complex functions of the concentration and mobility of all the ions present in a
20 system. They are always less than unity due to the presence of relatively small and mobile
21 endogenous ions such as sodium and chloride ions that have transport numbers of $t_{Na^+} = 0.6$
22 and $t_{Cl^-} = 0.4$ [29]. So, for electromigration, drug should compete with such ions. Smaller and
23 more mobile ions (higher transport number) can carry a significant fraction of charge and
24 reduce permeation of the drug.
25

26
27
28
29
30
31
32
33
34 Electrolytes can also affect electroosmotic flow. It has been demonstrated that increasing the
35 electrolyte ionic strength in the electrode chambers (relative to the physiological level present
36 beneath the tissue) decreased mannitol extraction (i.e., decreased electroosmosis) [33]. Santi
37 and Guy [33] investigated the electro-osmotic extraction of mannitol (a non-ionizable, non-
38 metabolizable model) in vitro using current density of 0.6 mA/cm² via Ag/AgCl electrodes
39 and pH 7.4 HEPES-buffered saline (0.14 M). They found that iontophoresis of divalent ions
40 from the anode chamber (i.e. an anode formulation with CaCl₂ or MgCl₂ instead of NaCl)
41 increased electro-osmotic flow from beneath the skin surface towards the anode (i.e.
42 modulation of the normal situation in which the skin's permselectivity dictates net electro-
43 osmosis in the opposite direction). Shielding of the net negative charge on the skin was
44 considered as a possible mechanism for this phenomenon. In the cathode chamber, to which
45 electro-osmosis predominates, ~3-fold levels of enhancement were achieved by formulating
46 the electrode bathing solution with either 2 mM calcein, 50–300 USP U/ml heparin, or 10
47 mM EDTA. It was postulated that this involves either binding of endogenous Ca²⁺ (and hence
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 decreased shielding of the negative charge on the skin), or simply the presence of more
2 negativity in the membrane (or a combination of the two) [33].
3

4 The above mentioned data and discussion show that regardless of the mechanism,
5 (electrorepulsion or electroosmosis), the concentration of the electrolyte should be at its
6 optimum level to allow charge transfer whilst at the same time not neutralize the skin charge
7 and not prevent the drug carrying the charge. A minimum amount of electrolyte, however,
8 must be present in order to provide the required electrochemistry of the system.
9

10
11
12
13 Table 6 presents the effects of direction of the current on the permeation of two peptides.
14 AAPV shows a similar flux from both anodal and cathodal concentration, in spite of its
15 negative charge. This might show the importance of both electrorepulsion and electroosmosis
16 for permeation of AAPV. However, acetyl hexapeptide-3 is benefiting from electroosmosis
17 due to lack of charge and high molecular weight, therefore showing a high anodal flux and a
18 low cathodal permeation (Table 6).
19
20
21
22
23

24
25 The peptides were chosen here to provide a series with increasing molecular weight and
26 different charges. Figure 2 show the relationship between iontophoretic permeability
27 coefficient and molecular weight and molecular charge for the peptides used in the present
28 investigation. As is seen, the relationship between these variables looks complex. As
29 discussed above, different mechanisms are involved in iontophoretic drug delivery including
30 passive diffusion, electrorepulsion and electroosmosis. Whatever the mechanism, it has been
31 shown that iontophoretic permeation of drugs decreases with increased molecular size [34-
32 37]. However, the size-dependency for iontophoretic permeation under these mechanisms is
33 different and the size-dependency of electroosmosis is expected to be lower than that of
34 passive diffusion or electrorepulsion [3]. Also pH-dependency and charge-dependency of
35 iontophoretic permeation under different mechanisms are also expected to be different. Using
36 the ionic mobility-pore model, Roberts et al [35] correlated the molecular weight of the
37 peptides with their iontophoretic permeability coefficient. The same has been reported with
38 experimental evidence using small molecular weight local anaesthetics. The ionic mobility
39 pore model identified ionic mobility and molecular size to be the major predictors of
40 iontophoretic permeability. Ionic mobility of a molecule depends on its pK_a , molecular
41 weight and conductivity.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 <Insert Figure 2>
58
59

1
2 Here and on the basis of the mechanisms described above, iontophoretic permeation of higher
3 weight molecular weight peptides, AAPV (tetrapeptide), Argirelin (hexapeptide) and
4 Triptoreline (decapeptide), were analysed against their molecular weight using linear
5 regression and a line with a negative slope of 0.0002 was observed (Figure 3). Such
6 relationship has been reported in the literature, with Green et al [34, 38] reporting a negative
7 slope of 0.002 for cathodal iontophoresis of a series of amino acids. Similarly DelTerzo et al
8 [39] reported a negative slope of 0.006 for alkanolic acids and Phipps et al [40] reported a
9 negative slope of 0.003 for inorganic cationic drugs. Based on the free volume model by
10 Yoshida et al [41, 42] a mean slope of 0.0032 was interpreted to give an average free volume
11 equivalent to the molecular volume of an ionised solute with molecular weight of 135.
12
13
14
15
16
17
18
19
20
21
22

23 <Insert Figure 3>
24
25
26
27

28 **CONCLUSION:**

29 This investigation showed that transdermal permeation of peptides is a complex process and
30 is affected by different factors including molecular weight, charge, donor pH, donor drug
31 concentration, background electrolytes and direction of the iontophoresis. Results show that
32 although both electroosmosis and electrorepulsion play important roles in iontophoretic
33 permeation of peptides, the importance of each depends on the molecular weight of the
34 peptide, pH of the environment and molecular charge. Electroosmosis plays a very important
35 role in permeation of higher molecular weight peptides, even when they are charged, while
36 lower weight peptides are affected more by electromigration. The present data clearly shows
37 that these different parameters need to be adjusted for optimized peptide delivery.
38
39
40
41
42
43
44
45
46
47
48

49 **Acknowledgement**

50 The authors would like to thank Dr Tom A Robertson, University of South Australia, for his
51 calculation of peptides net charges and related discussions. GK acknowledges a Curtin
52 University Postgraduate Scholarship. This work was financially supported by grants from an
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Australian Research Council Linkage Grant and from the National Health & Medical
2 Research Council of Australia.
3
4
5

6 **Conflict of Interest Statement**
7

8
9
10 All authors declare that have no conflict of interest related to this research.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES:

- [1] S. Namjoshi, R. Cacetta, H.A. Benson, *Skin peptides: biological activity and therapeutic opportunities*, *J Pharm Sci*, 97 (2008) 2524-2542.
- [2] H.A. Benson, S. Namjoshi, *Proteins and peptides: strategies for delivery to and across the skin*, *J Pharm Sci*, 97 (2008) 3591-3610.
- [3] R.H. Guy, Y.N. Kalia, M.B. Delgado-Charro, V. Merino, A. Lopez, D. Marro, *Iontophoresis: electrorepulsion and electroosmosis*, *J Control Rel*, 64 (2000) 129-132.
- [4] N. Abla, A. Naik, R.H. Guy, Y.N. Kalia, *Contributions of electromigration and electroosmosis to peptide iontophoresis across intact and impaired skin*, *J Control Rel*, 108 (2005) 319-330.
- [5] S. Gerscher, J.P. Connelly, J. Griffiths, S.B. Brown, A.J. MacRobert, G. Wong, L.E. Rhodes, *Comparison of the pharmacokinetics and phototoxicity of protoporphyrin IX metabolized from 5-aminolevulinic acid and two derivatives in human skin in vivo*, *Photochem Photobiol*, 72 (2000) 569-574.
- [6] D.M. Ferreira, Y.Y. Saga, E. Aluicio-Sarduy, A.C. Tedesco, *Chitosan nanoparticles for melanoma cancer treatment by photodynamic therapy and electrochemotherapy using aminolevulinic acid derivatives*, *Curr Med Chem*, (2013).
- [7] G. Krishnan, J.E. Grice, M.S. Roberts, H.A. Benson, T.W. Prow, *Enhanced sonophoretic delivery of 5-aminolevulinic acid: preliminary human ex vivo permeation data*, *Skin Res Technol*, 19 (2013) e283-289.
- [8] S. Namjoshi, R. Cacetta, J. Edwards, H.A. Benson, *Liquid chromatography assay for 5-aminolevulinic acid: application to in vitro assessment of skin penetration via Dermaportation*, *J Chromatogr B*, 852 (2007) 49-55.
- [9] P. Mikolajewska, R.F. Donnelly, M.J. Garland, D.I. Morrow, T.R. Singh, V. Iani, J. Moan, A. Juzeniene, *Microneedle pre-treatment of human skin improves 5-aminolevulinic acid (ALA)- and 5-aminolevulinic acid methyl ester (MAL)-induced PpIX production for topical photodynamic therapy without increase in pain or erythema*, *Pharm Res*, 27 (2010) 2213-2220.
- [10] W. Hornebeck, E. Moczar, J. Szecsi, L. Robert, *Fatty acid peptide derivatives as model compounds to protect elastin against degradation by elastases*, *Biochem Pharmacol*, 34 (1985) 3315-3321.
- [11] C. Blanes-Mira, J. Clemente, G. Jodas, A. Gil, G. Fernandez-Ballester, B. Ponsati, L. Gutierrez, E. Perez-Paya, A. Ferrer-Montiel, *A synthetic hexapeptide (Argireline) with antiwrinkle activity*, *Int J Cosmet Sci*, 24 (2002) 303-310.
- [12] C. Blanes-Mira, J.M. Merino, E. Valera, G. Fernandez-Ballester, L.M. Gutierrez, S. Viniestra, E. Perez-Paya, A. Ferrer-Montiel, *Small peptides patterned after the N-terminus domain of SNAP25 inhibit SNARE complex assembly and regulated exocytosis*, *J Neurochem*, 88 (2004) 124-135.
- [13] A. Kligman, E. Christophers, *Preparation of isolated sheets of human stratum corneum*, *Arch Dermatol*, 88 (1963) 70-73.
- [14] <http://www.chemaxon.com/>
- [15] G. Krishnan, M.S. Roberts, J. Grice, Y.G. Anissimov, H.A. Benson, *Enhanced transdermal delivery of 5-aminolevulinic acid and a dipeptide by iontophoresis*, *Biopolymers*, 96 (2011) 166-171.
- [16] M.J. Pikal, *The role of electroosmotic flow in transdermal iontophoresis*, *Adv Drug Deliv Rev*, 46 (2001) 281-305.
- [17] M.B. Delgado-Charro, R.H. Guy, *Iontophoresis: Application in drug delivery and noninvasive monitoring*, in: J.H. R.H. Guy (Ed.) *Transdermal drug delivery*, Marcel Dekker, New York, 2003, pp. 199-225.
- [18] M.J. Pikal, S. Shah, *Transport mechanisms in iontophoresis. III. An experimental study of the contributions of electroosmotic flow and permeability change in transport of low and high molecular weight solutes*, *Pharm Res*, 7 (1990) 222-229.
- [19] M.J. Pikal, S. Shah, *Transport mechanisms in iontophoresis. II. Electroosmotic flow and transference number measurements for hairless mouse skin*, *Pharm Res*, 7 (1990) 213-221.

- 1 [20] B. Mudry, R.H. Guy, M. Begona Delgado-Charro, Prediction of iontophoretic transport across
2 the skin, *J Control Rel*, 111 (2006) 362-367.
- 3 [21] P.G. Green, Iontophoretic delivery of peptide drugs, *J Control Rel*, 41 (1996) 33-48.
- 4 [22] B.D. Bath, E.R. Scott, J.B. Phipps, H.S. White, Scanning electrochemical microscopy of
5 iontophoretic transport in hairless mouse skin. Analysis of the relative contributions of diffusion,
6 migration, and electroosmosis to transport in hair follicles, *J Pharm Sci*, 89 (2000) 1537-1549.
- 7 [23] Y.Y. Huang, S.M. Wu, C.Y. Wang, Response surface method: a novel strategy to optimize
8 iontophoretic transdermal delivery of thyrotropin-releasing hormone, *Pharm Res*, 13 (1996) 547-
9 552.
- 10 [24] M.B. Delgado-Charro, A.M. Rodrguez-Bayon, R.H. Guy, Iontophoresis of nafarelin: effects of
11 current density and concentration on electrotransport in vitro, *J Control Rel*, 35 (1995) 35-40.
- 12 [25] L.L. Miller, C.J. Kolaskie, G.A. Smith, J. Rivier, Transdermal iontophoresis of gonadotropin
13 releasing hormone (LHRH) and two analogues, *J Pharm Sci*, 79 (1990) 490-493.
- 14 [26] D. Marro, Y.N. Kalia, M.B. Delgado-Charro, R.H. Guy, Contributions of electromigration and
15 electroosmosis to iontophoretic drug delivery, *Pharm Res*, 18 (2001) 1701-1708.
- 16 [27] B.H. Sage, J.E. Riviere, Model systems in iontophoresis transport, *Adv Drug Deliv Rev*, 9 (1992)
17 265-287.
- 18 [28] J.B. Phipps, J.R. Gyory, Transdermal ion migration, *Adv Drug Deliv Rev*, 9 (1992) 137-176.
- 19 [29] B. Mudry, R.H. Guy, M.B. Delgado-Charro, Iontophoresis, in: E. Touitou, B.A. Barry (Eds.)
20 *Transdermal Delivery*, CRC Press, 2006.
- 21 [30] G.B. Kasting, J.C. Keister, Application of electrodiffusion theory for a homogeneous membrane
22 to iontophoretic transport through skin., *J Control Rel*, 8 (1989) 195-210.
- 23 [31] N.H. Bellatone, S. Rim, M.L. Francoeur, B. Rasadi, Enhanced percutaneous absorption via
24 iontophoresis I. Evaluation of an in vitro system and transport of model compounds., *Int J Pharm*,
25 30 (1986) 63-72.
- 26 [32] J. Cazares-Delgadillo, A. Naik, A. Ganem-Rondero, D. Quintanar-Guerrero, Y.N. Kalia,
27 Transdermal delivery of cytochrome C--A 12.4 kDa protein--across intact skin by constant-current
28 iontophoresis, *Pharm Res*, 24 (2007) 1360-1368.
- 29 [33] P. Santi, R.H. Guy, Reverse iontophoresis — Parameters determining electroosmotic flow: I. pH
30 and ionic strength, *J Control Rel*, 38 (1996) 159-165.
- 31 [34] P.G. Green, R.S. Hinz, C. Cullander, G. Yamane, R.H. Guy, Iontophoretic delivery of amino acids
32 and amino acid derivatives across the skin in vitro, *Pharm Res*, 8 (1991) 1113-1120.
- 33 [35] M.S. Roberts, P.M. Lai, Y.G. Anissimov, Epidermal iontophoresis: I. Development of the ionic
34 mobility-pore model, *Pharm Res*, 15 (1998) 1569-1578.
- 35 [36] P.M. Lai, M.S. Roberts, Epidermal iontophoresis: II. Application of the ionic mobility-pore
36 model to the transport of local anesthetics, *Pharm Res*, 15 (1998) 1579-1588.
- 37 [37] P.M. Lai, M.S. Roberts, An analysis of solute structure-human epidermal transport
38 relationships in epidermal iontophoresis using the ionic mobility: pore model, *J Control Rel*, 58
39 (1999) 323-333.
- 40 [38] P.G. Green, R.S. Hinz, A. Kim, F.C. Szoka, Jr., R.H. Guy, Iontophoretic delivery of a series of
41 tripeptides across the skin in vitro, *Pharm Res*, 8 (1991) 1121-1127.
- 42 [39] S. DelTerzo, C.R. Behl, R.A. Nash, Iontophoretic transport of a homologous series of ionized
43 and nonionized compounds: influence of hydrophobicity and mechanistic interpretation, *Pharm*
44 *Res*, 6 (1989) 89-90.
- 45 [40] J.B. Phipps, R.V. Padmanabhan, G.A. Lattin, Iontophoretic delivery of model inorganic and
46 drug ions, *J Pharm Sci*, 78 (1989) 365-369.
- 47 [41] N.H. Yoshida, M.S. Roberts, Structure-transport relationships in transdermal iontophoresis.,
48 *Adv Drug Deliv Rev*, 9 (1992) 239-264.
- 49 [42] N.H. Yoshida, M.S. Roberts, Solute molecular size and transdermal iontophoresis across exised
50 human skin., *J Control Rel*, 25 (1993) 177-195.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends:

Figure 1. Molecular structures, molecular weight and some other properties of peptides used in the present investigation

Figure 2. Relationship between molecular weight, molecular charge and iontophoretic permeability coefficient of peptides through human skin.

Figure 3. Linear relationship between molecular weight and iontophoretic permeability coefficient of AAPV (tetrapeptide), Argirelin (hexapeptide) and Triptorelin (decapeptide) at pH 7.4 through human skin

Tables:**Table 1.** Permeation of peptides across human skin under 0.4 mA/cm² anodal iontophoresis in relation to their molecular properties. Data are mean \pm SEM.

Peptide	MW	Conc (mM)	pH	Net charge	Iontophoretic flux ($\mu\text{g}/\text{cm}^2/\text{h}$)	Permeability coefficient ($\text{cm}/\text{h} \times 10^3$)			
						Iontophoresis	Passive	Pure (Ionto-Passive)	ER
ALA	131	60	5.5	0.06	12.5 \pm 1	1.3 \pm 1.1	0.08 \pm 4.8	1.22	16
Ala-Trp	276	3.6	5.5	0.04	13 \pm 2.8	13 \pm 3.0	ND	~13	>13
AAPV	355	28.4	5.5	0.03	175 \pm 7.1	18 \pm 0.1	ND	~18	>18
EEMQRR	889	1.13	7.4	0.0	15.7 \pm 2.3	17 \pm 2.0	2.5 \pm 0.05	14.5	6.8
Triptorelin	1311	0.9	7.4	1.1	12.1 \pm 3.6	12 \pm 3	0.4 \pm 0.0	11.6	27.3

ND: Not detectable
ER: Enhancement ratio; Iontophoresis/passive

Table 2. Effect of pH on permeability coefficient ($\text{cm/h} \times 10^3$) of peptides across human skin under 0.4 mA/cm^2 anodal iontophoresis. Data are mean \pm SEM.

Peptide	MW	pH = 7.4		pH = 5.5		pH = 5		pH = 3	
		Charge	Kp	Charge	Kp	Charge	Kp	Charge	Kp
ALA	131	-0.27	11 ± 10	0.06	1.3 ± 1.1	-	-	-	-
Ala-Trp	276	-0.11	0.4 ± 0.1	0.04	13 ± 3.0	-	-	-	-
AAPV	355	-0.2	18 ± 0.7	0.03	18 ± 0.1	-	-	0.9	0.4 ± 0.1
EEMQRR	889	0.0	17 ± 2	-	-	0.18	4.8 ± 1.4	-	-

Table 3. Effect of concentration on flux and permeability coefficient (Kp) of peptides across human skin under 0.4 mA/cm² anodal iontophoresis. Data are mean ± SEM.

Peptide	MW	Donor pH	Charge	Flux (µg/cm ² /h)		Kp (cm/h × 10 ³)	
				1 mg/ml	10 mg/ml	1 mg/ml	10 mg/ml
Ala-Trp	276	5.5	0.04	13 ± 2.8	52 ± 11	13 ± 2.8	5.5 ± 1.2
EEMQRR	889	7.4	0	16 ± 2.3	41 ± 6.1	17 ± 2	4.3 ± 0.6
Triptorelin*	1311	7.4	1.1	12 ± 3.6	21 ± 6.5	12 ± 3	2.3 ± 0.6

Table 4. Effect of added background electrolyte on permeability coefficient (Kp, cm/h) of peptides across human skin under 0.4 mA/cm² anodal iontophoresis. Data are mean ± SEM.


Peptide	No added electrolyte (Control)			Added 20/137mM HEPES/NaCl			Kp reduction ratio
	pH	Charge	Kp ($\times 10^3$)	pH	Charge	Kp ($\times 10^3$)	
Ala-Trp	5.5	0.04	13 ± 3.0	5.5	0.04	0.9 ± 0.06	14.4
AAPV	7.4	-0.2	18 ± 0.7	6.0	0	10 ± 1	1.8
EEMQRR	7.4	0.0	17 ± 2	6.0	0.02	9.3 ± 2.4	1.8
Triptorelin	7.4	1.1	12 ± 3	7.4	1.1	6.8 ± 2.6	1.8

Table 5. Effect of added background electrolyte on permeability coefficient (Kp) of EEMQRR across human skin under 0.4 mA/cm² anodal iontophoresis. Data are mean ± SEM.


Added electrolyte	pH	Kp (cm/h × 10 ³)	Kp reduction ratio
No added electrolyte (Control)	7.4	17 ± 2	1
137 mM NaCl + 20 mM HEPES	6.0	9.3 ± 2.4	1.8
68.5 mM NaCl+ 10 mM HEPES	6.0	12 ± 0.2	1.4
34.2 mM NaCl + 5 mM HEPES	6.0	14 ± 1	1.2
137 mM NaCl	6.0	0.5 ± 0.1	34
68.5 mM NaCl	6.0	1.6 ± 1.2	11
34.2 mM NaCl	6.0	0.3 ± 0.0	57

Table 6. Effect of current direction on permeability coefficient (Kp) of tetra and deca-peptides across human skin under 0.4 mA/cm² iontophoresis at pH 7.4. Data are mean ± SEM.


Peptide	MW	Charge	Kp ($\times 10^3$) and [Charge]	
			Anodal	Cathodal
AAPV	355	-0.2	18 ± 0.7	17 ± 1
EEMQRR	889	0	17 ± 2	1.2 ± 0.3




L-Ala-L-Trp MW
275.9; logP 0.6




L-Ala-L-Ala-L-Pro-L-Val
MW 355.4; logP -0.45



**p-GLU-His-Trp-Ser-Tyr-[D-Trp]-
Leu-Arg-Pro-Gly-NH₂ (Triptorelin)**
MW 1131.5; logP -0.3



Ac-Glu-Glu-Met-Gln-Arg-Arg-NH₂
(Acetyl hexapeptide-3) MW 888.6; logP -4.5



5-Aminolevulinic acid
MW 131.1; logP -1.3




Figure 2

Figure 3

