

1 **Climate variability over the last 35,000 years recorded in marine and terrestrial**  
2 **archives in the Australian region: an OZ-INTIMATE compilation.**

3 **Jessica M. Reeves<sup>a</sup>, Timothy T. Barrows<sup>b</sup>, Timothy J. Cohen<sup>c</sup>, Anthony S. Kiem<sup>d</sup>, Helen**  
4 **C. Bostock<sup>e</sup>, Kathryn E. Fitzsimmons<sup>f</sup>, John D. Jansen<sup>g</sup>, Justine Kemp<sup>h</sup>, Claire Krause<sup>i</sup>,**  
5 **Lynda Petherick<sup>j</sup>, Steven Phipps<sup>k,l</sup> and OZ-INTIMATE members.**

6

7 <sup>a</sup>Centre for Environmental Management, University of Ballarat, Australia

8 <sup>b</sup>Geography, College of Life and Environmental Sciences, University of Exeter, United  
9 Kingdom

10 <sup>c</sup>GeoQuEST Research Centre, School of Earth & Environmental Sciences, The University of  
11 Wollongong, New South Wales, Australia

12 <sup>d</sup>School of Environmental and Life Sciences, University of Newcastle, New South Wales,  
13 Australia

14 <sup>e</sup> National Institute of Water and Atmospheric Research, Wellington, New Zealand

15 <sup>f</sup>Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology,  
16 Leipzig, Germany

17 <sup>g</sup>Department of Physical Geography and Quaternary Geology, Stockholm University, Sweden

18 <sup>h</sup>Australian Rivers Institute, Griffith University, Queensland, Australia

19 <sup>i</sup>Research School of Earth Sciences, The Australian National University, Canberra, Australia

20 <sup>j</sup>School of Earth, Environment and Biological Sciences, Queensland University of  
21 Technology, Gardens Point, Queensland, Australia

22 <sup>k</sup>Climate Change Research Centre, University of New South Wales, Sydney, Australia

23 <sup>l</sup>ARC Centre of Excellence for Climate System Science, University of New South Wales,  
24 Sydney, Australia

25 **Abstract**

26 The Australian region spans some 60 degrees of latitude and 50 degrees of longitude and  
27 displays considerable regional climate variability both today and during the Late Quaternary.  
28 A synthesis of marine and terrestrial climate records, combining findings from the Southern  
29 Ocean, temperate, tropical and arid zones, identifies a complex response of climate proxies to  
30 a background of changing boundary conditions over the last 35,000 years. Climate drivers  
31 include the seasonal timing of insolation, greenhouse gas content of the atmosphere, sea level  
32 rise and ocean and atmospheric circulation changes. Our compilation finds few climatic  
33 events that could be used to construct a climate event stratigraphy for the entire region,  
34 limiting the usefulness of this approach. Instead we have taken a spatial approach, looking to  
35 discern the patterns of change across the continent.

36

37 The data identify the clearest and most synchronous climatic response at the time of the last  
38 glacial maximum (LGM) ( $21 \pm 3$  ka), with unambiguous cooling recorded in the ocean, and  
39 evidence of glaciation in the highlands of tropical New Guinea, southeast Australia and  
40 Tasmania. Many terrestrial records suggest drier conditions, but with the timing of inferred  
41 snowmelt, and changes to the rainfall/runoff relationships, driving higher river discharge at  
42 the LGM. In contrast, the deglaciation is a time of considerable south-east to north-west  
43 variation across the region. Warming was underway in all regions by 17 ka. Post-glacial sea  
44 level rise and its associated regional impacts have played an important role in determining the  
45 magnitude and timing of climate response in the north-west of the continent in contrast to the  
46 southern latitudes. No evidence for cooling during the Younger Dryas chronozone is evident  
47 in the region, but the Antarctic cold reversal clearly occurs south of Australia. The Holocene  
48 period is a time of considerable climate variability associated with an intense monsoon in the

49 tropics early in the Holocene, giving way to a weakened monsoon and an increasingly El  
50 Niño-dominated ENSO to the present. The influence of ENSO is evident throughout the  
51 southeast of Australia, but not the southwest. This climate history provides a template from  
52 which to assess the regionality of climate events across Australia and make comparisons  
53 beyond our region

#### 54 **Keywords**

55 Australia, tropics, temperate, arid zone, Southern Ocean, last glacial maximum, deglacial  
56 period, Holocene, INTIMATE

#### 57 **1. Introduction**

58 Australia, the “Island Continent”, spans the latitudinal range of 10°S to 43°S, and is bordered  
59 by New Guinea and Indonesia to the north, the Pacific Ocean to the east, the Indian Ocean to  
60 the west and the Southern Ocean to the south (Fig. 1). This location means that the continent  
61 and the modern climate are juxtaposed between the heat of the equatorial tropics from the  
62 Indo-Pacific Warm Pool (IPWP) and the cool waters of the Southern Ocean. Australia’s  
63 latitudinal position results in several synoptic-scale controls on the climate including dry  
64 descending air associated with the Hadley Cell and the resulting Sub-tropical High Pressure  
65 Belt, easterly zonal flow associated with the South East Trade winds and the summer  
66 monsoon in the north and prevailing westerlies in the south (Fig. 2). The size of the mainland  
67 produces strong continentality with precipitation and temperature gradients from the coast to  
68 the dry centre.

69

70 The mean climatic conditions across Australia primarily respond to the seasonal zonal  
71 circulation by continental heating and cooling and the land-sea temperature contrast (Gimeno

72 et al., 2010), and are driven by multiple atmospheric and oceanic influences and the  
73 interactions between them (e.g. Gallant et al., 2012). The non-stationarity, intensity and  
74 spatial variability of these climatic drivers, and the degree to which they have interacted  
75 through time, are the key criteria for addressing the reconstruction of Australian  
76 palaeoclimate.

77

78 The last 35 kyr encompasses the end of Marine Isotope Stage (MIS) 3, the transition into full  
79 glacial conditions, subsequent deglacial warming and establishment of the Holocene, and  
80 includes millennial-scale reversals (e.g. the Antarctic Cold Reversal; ACR) and centennial-  
81 scale variability of climate (e.g. the Holocene). This period, into and out of the last glacial, is  
82 of key interest to the INTegration of Ice core, Marine and Terrestrial (INTIMATE) project  
83 (Barrows et al., editorial). In this paper we bring together the records discussed in each of the  
84 four regional synthesis papers in this volume (Bostock et al., Fitzsimmons et al., Petherick et  
85 al., Reeves et al., this volume) as the ‘broader Australian region’. This broader region extends  
86 from 10°N encompassing New Guinea and much of Indonesia, to 65°S, to include the  
87 Australian segment of the Southern Ocean, and from 100-165°E (Fig. 3).

88

89 This paper provides a synthesis of major climatic events over the last 35 kyr and attempts to  
90 determine their regional distribution. We find that changing boundary conditions through  
91 time exert variable influence across the region, with regional controls such as the land:sea  
92 ratio, the monsoon and the influence of the El Niño Southern Oscillation (ENSO), and  
93 changes in the westerly winds and mean state of the Southern Ocean, important for  
94 determining the pattern of climate change.

## 96 **Climate Event Stratigraphy**

97 The original remit of the AUS-INTIMATE project was to develop a climate event  
98 stratigraphy for the Australian region (Barrows et al., editorial). From a global perspective,  
99 the phenomenon of the bi-polar see-saw (Broecker, 1998) and the continuing debate as to the  
100 relative timing of climate change between the hemispheres prompted the question of how  
101 global scale climate events are expressed in the Australian region. Clear differences are  
102 apparent between the high latitudes of both hemispheres (Blunier and Brook, 2001)  
103 indicating that a climate event stratigraphy based on a Greenland ice core record has limited  
104 meaning in the Australian region. There is potential to develop a Southern Hemisphere high-  
105 latitude stratigraphy based on Antarctic records (e.g. Pedro et al., 2011; EPICA, 2006), but it  
106 remains unclear as to how applicable this would be for the Australian region, given its  
107 distance from Antarctica. Clearly, we first need to understand how the Australian region  
108 responds to the competing influences of changing boundary conditions over such a large  
109 region.

110

111 Here we document changes across the region to better describe the driving mechanisms of  
112 climate change. We take a time-slice approach to synthesize terrestrial and marine records  
113 during the late MIS 3 period, the early glacial period, the LGM, the deglacial period  
114 including during the ACR and the early-to-mid and late Holocene. The regional compilations  
115 (e.g. Bostock et al. Fitzsimmons et al., Petherick et al., Reeves et al., this volume) examine  
116 various proxy archives from the equator to the Southern Ocean which together allow us to  
117 investigate the relative changes of the tropics versus high latitudes through time, and spatial  
118 coherence in response to global climate change across the vast Australian region. This

119 temporal and spatial synthesis then allows for the better characterisation of the drivers of the  
120 climatic changes over the last 35 kyr and the regionality of the climate response.

121

## 122 **1.1 Modern ocean-atmospheric climate drivers**

123 The following section provides a brief overview of the modern ocean-atmospheric drivers of  
124 the Australian climate. We interpret climatic changes over the last 35 kyr as modifications to  
125 the major elements of the modern climate system. The oceanographic setting of the region  
126 has been discussed elsewhere in this volume (see Bostock et al., Petherick et al., Reeves et  
127 al.) and the dominant features are shown in Figure 1.

128

129 The modern climate patterns of the Australian region may be considered as including the  
130 tropics, encompassing both the perennially wet equatorial and seasonally wet monsoonal  
131 regions; the temperate zone of the east, south and south-west coasts, including Tasmania; and  
132 the arid interior, which receives little rainfall (<250 mm per annum) (see Fig. 3 for a  
133 schematic representation of this subdivision). Australia receives predominantly winter rainfall  
134 in the south, summer rainfall in the north and sporadic, winter and summer rainfall in the  
135 central region of the east coast (Gentilli, 1971). The ocean circulation patterns and the heat  
136 distribution around the continent also play a role in determining the nature and distribution of  
137 rainfall patterns across Australia. With few exceptions beyond the eastern Great Dividing  
138 Range/Eastern Escarpment, the relatively low altitude of the Australian continent results in  
139 minimal orographic influences. The interannual variation in precipitation can exceed that of  
140 annual variation, particularly in the arid zone.

141

142 The climate of Australia (Fig. 3) is determined by continental-scale atmospheric circulation  
143 patterns. In the Southern Hemisphere, warm, moist air is lifted by the Hadley Cell at the  
144 equatorial low pressure or Inter-tropical Convergence Zone (ITCZ) and descends around  
145 30°S resulting in the Sub-Tropical High Pressure Belt (STHPB). The sinking air of this  
146 STHPB dominates the modern climate of the arid interior (Sturman and Tapper, 1996). The  
147 ITCZ moves into the north of Australia during the austral summer, bringing with it the  
148 summer monsoon. To the north of Australia lies the Indo-Pacific Warm Pool (IPWP), a  
149 region where mean annual sea surface temperatures (SSTs) exceed 28°C and provide a major  
150 global source of latent heat release (Gagan et al., 2004). Some of the descending air from the  
151 STHPB travels back to the equator along the surface of the Earth, creating the south easterly  
152 trade winds, which flow from the east to west across the equatorial Pacific. These trade winds  
153 in turn typically bring about deep atmospheric convection (Walker Circulation) over the  
154 tropics which result in heavy rainfall over the northeast coast of Australia during summer  
155 (November to April).

156

157 The climate patterns of the eastern half of Australia are partly modulated on an inter-annual  
158 scale by ENSO (e.g. Verdon et al., 2004). Variations in SST across the equatorial Pacific are  
159 associated with weakening (El Niño) or strengthening (La Niña) of this Walker Circulation  
160 over cycles of several years (see Diaz and Markgraf (2000) for further details). The Inter-  
161 decadal Pacific Oscillation (IPO) has similar characteristics to ENSO, but on longer  
162 timescales and affecting the wider Pacific Basin (Power et al., 1999). The IPO influences  
163 Australia via a modulation of both the magnitude and frequency of ENSO impacts leading to  
164 multi-decadal epochs that are significantly wetter or drier than others (e.g. Kiem et al., 2003;  
165 Kiem and Franks, 2004). This effect has been identified for at least the past 400 years  
166 (Verdon and Franks, 2006). Whilst other inter-annual drivers are relevant in the Australian

167 region (e.g. Indian Ocean Dipole or Southern Annular Mode) most palaeo-proxy data are of  
168 lower resolution preventing the identification of such drivers.

169

170 South of the Hadley Cell, descending air moves southward in the Ferrel Cell, before  
171 ascending at the margins of the polar front (~60°S). The surface wind flow of the Ferrel Cell  
172 is responsible for the mid-latitude westerly wind belt or “westerlies” which are largely  
173 responsible for winter rainfall in southern Australia (Pitman et al., 2004) and play a dominant  
174 role in circulation in the Southern Ocean (Varma et al., 2010). In addition, the cold fronts  
175 embedded in the STHPB and associated with the westerlies, influence the landscape through  
176 aeolian transport of surface sediments, particularly in the arid zone (Hesse and McTainsh  
177 1999; Hesse, 2010).

178

## 179 **1.2 Drivers of past changes**

180 The dominant external forcing on the global climate system over the past 35 kyr has been the  
181 changing seasonal and meridional distribution of insolation arising from cyclic changes in the  
182 Earth's orbital geometry (Berger, 1978). On glacial-interglacial timescales, feedbacks  
183 involving the global carbon cycle and atmospheric greenhouse gases have amplified the  
184 response of the climate system to this signal (e.g. Petit et al., 1999). In the Northern  
185 Hemisphere, this period has been characterised by the expansion and retreat of ice sheets in  
186 response to changes in summer insolation at high latitudes. This resulted in a long build-up of  
187 ice sheets during glacial phases and relatively rapid deglaciation, into interglacial phases. By  
188 contrast, the continents of the Southern Hemisphere did not possess ice sheets, excepting  
189 Antarctica, but the associated changes in sea level altered land:sea ratios and continental  
190 connections, and therefore altered ocean currents and regional climates. Changing  
191 temperature differentials between land and sea altered wind strengths and precipitation. The



192 variable seasonal cycle of insolation also drove changes in the background state of the  
193 tropical Pacific Ocean, with direct consequences for the climate of the Australian region.

194

195 The Southern Ocean has played a significant role on the balance of CO<sub>2</sub> between the oceans  
196 and atmosphere on centennial-millennial timescales (e.g. Toggweiler, 1996; Sigman et al.,  
197 2010). The expansion of Antarctic sea ice during the glacial acted to trap CO<sub>2</sub> in the deep  
198 ocean, lowering atmospheric CO<sub>2</sub> levels. Conversely, during the deglacial when sea ice  
199 retreated, outgassing of CO<sub>2</sub> occurred on a large scale (e.g. Laurantou et al., 2010; Bostock et  
200 al., this volume).

201

202 On decadal to centennial timescales, changes in solar irradiance and explosive volcanism are  
203 also significant. Existing reconstructions of solar forcing only extend back as far as the early  
204 Holocene (e.g. Steinhilber et al., 2009), while reconstructions of volcanic forcing (e.g.  
205 Plummer et al., 2012) only cover the last 2,000 years. Thus the potential role of these forcings  
206 in driving climatic changes over the past 35 kyr is extremely poorly understood, and changes  
207 seen within the Australian climate over this period may therefore have been driven by  
208 unknown external events. In particular, there is some evidence that volcanic eruptions,  
209 despite being only short in duration, can be sufficient to push the climate system past tipping  
210 points into alternative states that can persist for multiple centuries (Miller et al., 2012).

211

## 212 **2. Rationale for record inclusion**

213 The synthesis presented here builds on the four review papers in this issue, which deal with  
214 the Australian tropical (Reeves et al.), terrestrial (Petherick et al.) and arid interior

215 (Fitzsimmons et al.) zones and the Southern Ocean (Bostock et al.). Our synthesis builds on  
216 an earlier effort as part of the AUS-INTIMATE project by Turney et al. (2006a). Previous  
217 regional reviews of climate have been summarised by Williams et al. (2009), who also  
218 compiled key long records with quantifications of climate change. In this review we build on  
219 this earlier works by incorporating a larger contribution from deep-sea records, covering the  
220 Holocene and incorporating new records from all regions.

221

222

223 The criteria for inclusion of records into this synthesis would ideally be: continuity through  
224 the last 35 kyr, sound chronology, centennial-scale or better resolution and unambiguous and  
225 quantifiable palaeoclimate estimates. Although this has been achieved in many of the marine  
226 and speleothem records, their spatial coverage does not represent the greater Australian  
227 region. We have therefore also chosen here to also include: high-resolution short-term  
228 records (e.g. corals), discontinuous geomorphic records (e.g. fluvial, lake shore, dune,  
229 glacier) where the interpretation is robust (see Fitzsimmons et al., this volume for further  
230 discussion), as well as qualitative records (e.g. pollen), although noting the context of the site  
231 and the limitations of each record (Fig. 2).

232

233 The chronologies of the records were determined using a variety of methods. Radiocarbon is  
234 the most common method used, frequently on bulk sediment, charcoal or foraminifera for  
235 marine cores and calibrated here using INTCAL09 or MARINE09 (Reimer et al., 2009).  
236 Other methods include optically-stimulated luminescence (OSL) and thermoluminescence  
237 (TL) dating of sediments, predominantly applied to aeolian and fluvial records, exposure  
238 dating (especially  $^{10}\text{Be}$  and  $^{36}\text{Cl}$ ) of moraines; and U/Th dating of speleothems and corals. All  
239 ages are presented as calendar years ka (thousand years) for consistency.

240

### 241 **3. Synthesis**

242 During the first time slice (late MIS 3), we compare climatic conditions to modern mean  
243 annual temperature and precipitation. Other time slices are described with reference to the  
244 preceding time slice. The age boundaries of these time slices are chosen to coincide with the  
245 largest changes recorded in the climate proxy records. We refer here to effective  
246 precipitation, which is influenced not only by total precipitation, but also evaporation, wind  
247 and vegetation cover. The simplified patterns of temperature and effective precipitation  
248 between regions and through time are shown in Figure 4 a-h.

249

#### 250 **35-32 ka – late Marine Isotope Stage 3**

251 Figure 4ai,ii shows climatic conditions during the period 35-32 ka in comparison to the  
252 present day, with cooler climate in the NE tropics, arid interior and lower latitudes of the  
253 Southern Ocean, and transient or variable conditions in the other regions. Wet conditions are  
254 evident throughout most of the region due to decreased evaporation under cooler than present  
255 conditions (Galloway, 1965; Bowler and Wasson, 1984).

256

257 Leading up to the last glacial period, much of inland Australia had large permanent lakes in  
258 catchments which are presently dry or ephemeral (Bowler et al., 1976; 2012; Cohen et al.,  
259 2011, 2012; Fitzsimmons et al., this volume). Relatively high lake stands prevailed in both  
260 the temperate (Coventry, 1976; Bowler and Hamada, 1971) and tropical zones (Veth et al.,  
261 2009). This period witnessed a peak in fluvial activity across the continent, both in the north  
262 (Nanson et al., 2008; Veth et al., 2009) and in the temperate zone, particularly in the

263 tributaries of the Murray-Darling Basin (MDB) (Bowler and Wasson, 1984; Page et al., 1996;  
264 Page and Nanson, 1996; Kemp and Rhodes, 2010). Cool and humid conditions are inferred,  
265 with increased effective precipitation (Kemp and Rhodes, 2010). However, it is important to  
266 note that although this time was wet in comparison to present, it was dwarfed by the wet  
267 intervals of Marine Isotope Stage 5 and 4 (Nanson et al., 1992, Cohen et al., 2011, 2012).  
268 Cooler conditions in the arid zone (Miller et al., 1997) and wetter conditions in the north of  
269 the continent around this time are also suggested (Wallis, 2001; van der Kaars and De  
270 Deckker, 2002). Ice core records from Antarctica show millennial-scale variability through  
271 this period, corresponding to temperature shifts of up to 2°C (EPICA, 2006). These  
272 perturbations are also recorded in the Murray Canyons offshore South Australia (De Deckker  
273 et al., 2012) and in deep sea cores from the Southern Ocean (Barrows et al., 2007a; Armand  
274 and Leventer, 2010).

275

### 276 **32-22 ka - early glacial period**

277 Cooling began around 32 ka, with glacier advance in the Snowy Mountains (Barrows et al.,  
278 2001). Glacial conditions may have commenced in New Guinea by ~28 ka (Prentice et al.,  
279 2005). A peak in dune activity in both the central arid region (Fitzsimmons et al., this issue),  
280 and increased dust transport, most likely from the increasingly arid MDB, to the subtropics  
281 along the east coast (Petherick et al., 2008; 2009) is interpreted as indicating cooler, drier  
282 conditions. Significant vegetation change commenced, with the expansion of herbs and  
283 grasses at the expense of arboreal taxa, in southeastern Australia (Dodson, 1975; Colhoun et  
284 al., 1982; Colhoun, 2000; Kershaw et al., 2007). A drying trend was also present in the NE  
285 Indian Ocean region from 32 ka, noted in the speleothems of Flores (Lewis et al., 2011) and  
286 pollen records off Java (van der Kaars et al., 2010) and also off Cape Range, Western

287 Australia (van der Kaars and De Deckker, 2002). However relatively wet conditions persisted  
288 in Sumatra and West Papua (Newsome and Flenley, 1988; Hope and Tulip, 1994; van der  
289 Kaars et al., 2012).

290

291 After slightly warmer conditions in Antarctica centred around 30 ka, cooling commenced in  
292 earnest and near full glacial temperatures were achieved around 27 ka (EPICA, 2006). Strong  
293 cooling had also occurred in the Southern Ocean by ~26 ka, accompanied by a northward  
294 movement of the subtropical front by some 3-5°, bringing cool, sub-Antarctic waters to the  
295 south of Australia (De Deckker et al., 2012; Bostock et al., this volume). The wet conditions  
296 of the temperate region persisted until ~25 ka, followed by drying, bringing about a  
297 dominance of grass and herb vegetation (Petherick et al., this volume). Dune activity  
298 increased within the central arid zone (e.g. Hesse et al., 2004; Fitzsimmons et al., 2007) and  
299 expanded at the desert margins during the early glacial period (Lomax et al., 2011;  
300 Fitzsimmons et al., this volume). In the tropics, dry, stable conditions are evident in the  
301 vegetation and speleothem records of Indonesia (Reeves et al., this volume). Both the lake  
302 level and the speleothem records from the tropics show a close correlation between effective  
303 precipitation and regional insolation through this period. There is no evidence for a strong  
304 monsoon penetrating the Australian mainland at this time (Devriendt, 2011; Lewis et al.,  
305 2011; Reeves et al., this volume).

306

307 There is some evidence for a short-lived expansion of rainforest taxa in north Queensland  
308 ~26-24 ka (Moss and Kershaw, 2007). This coincides with lower, oscillating lake levels in  
309 the Willandra system (Bowler et al., 2012), an increase in dust transport to the subtropics  
310 (Petherick et al., 2008), abrupt drying and increased variability in the Borneo speleothem  
311 record 26.5-25 ka (Partin et al., 2007) and warming offshore South Australia (Calvo et al.,

312 2007) and the Southern Ocean (Armand and Leventer, 2010). These combined factors point  
313 to a warming and drying trend in the earlier glacial period (Fig. 4bi,ii). Although there are  
314 few records from this time period that have strong enough chronological control to resolve  
315 how widespread this climatic anomaly was, in most places conditions prior to 26 ka were  
316 substantially different to those prevailing after 24 ka, when the accelerated descent into the  
317 last glacial maximum (LGM) commenced. This period is consistent with the first interstadial  
318 of the Last Glacial Cold Period in New Zealand, coincident with the Kawakawa tephra  
319 (Barrell et al, this volume; Vandergoes et al., this volume) but earlier than the brief warming  
320 in Antarctica at ~24-23 ka (EPICA, 2006).

321

### 322 **22-18 ka – Last Glacial Maximum**

323 The LGM, globally the interval of greatest ice sheet extent over the last full glacial cycle,  
324 encompasses both maximum glacial extent and SST minima across the region (Fig. 4ci,ii).  
325 Sea surface temperature was typically in the order of 3-6°C cooler than present in the  
326 Southern Ocean (Barrows et al., 2007a). This was combined with an expansion of winter sea-  
327 ice as far north as 55°S (Gersonde et al., 2005; Armand and Levanter, 2010). Cooling in the  
328 tropical oceans was less pronounced, with SST 1-3°C cooler than present in the NW and 1-  
329 2.5°C cooler in the NE (e.g. Stott et al., 2002; Visser et al., 2003; Barrows and Juggins, 2005;  
330 Dunbar and Dickens, 2003; Jorry et al., 2008. These cooler ocean temperatures coincided  
331 with a constriction of both the ITF and the IPWP (Reeves et al., this volume). Some of the  
332 greatest differences in SST were felt along the northwest and south coast of Australia,  
333 attributed to the weakening of the Leeuwin Current and northward movement of the  
334 subtropical front (Petherick et al., this volume).

335

336 The most significant glacier advance is centred on ~19 ka, both in Tasmania and the Snowy  
337 Mountains of southeastern Australia (Barrows et al., 2001; 2002; Kiernan et al., 2004;  
338 Mackintosh et al., 2006). Peak glacial extent occurred at the same time (20.3-19.4 ka) at Mt  
339 Giluwe in New Guinea (Barrows et al., 2011). Estimates of the maximum expanse of ice are  
340 15 km<sup>2</sup> in the Snowy Mountains (Barrows et al., 2001), 1085 km<sup>2</sup> in Tasmania (Colhoun,  
341 1996) and 3400 km<sup>2</sup> in New Guinea (Prentice et al., 2011). Cooling in the upland areas is  
342 postulated to be as much as 11°C below present in New Guinea (Hope, 2009), 9°C in south-  
343 eastern Australia (Galloway, 1965) and between ~6.5- 4.2°C in Tasmania (Colhoun, 1985;  
344 Fletcher and Thomas, 2010) with a peak in periglacial activity at ~22 ka (Barrows et al.,  
345 2004). These estimates are based on pollen records, snowlines and lower limits of periglacial  
346 solifluction. The extent of cooling is generally at odds with the moderate cooling observed in  
347 the oceans (Barrows et al., 2000) and highlights their buffering capacity, except where  
348 movement of the oceanic fronts brought about significant cooling (e.g. Calvo et al., 2007;  
349 Bostock et al., this issue). Some of the differences may also be due to the accuracy and  
350 precision of the temperature estimates. For example, the snowline is affected by both  
351 temperature and precipitation, solifluction can occur below the treeline, and vegetation is  
352 affected by many variables including precipitation and carbon dioxide levels.

353

354 Both cooler and drier conditions are evident in the vegetation records from the Indonesian  
355 region and the east coast of Australia, including Tasmania, with a reduction in woody taxa, in  
356 particular rainforest species, and expansion of grasses and herbs (Petherick et al., this  
357 volume, Reeves et al., this volume). Estimates of temperature from vegetation changes in the  
358 lowlands of Tasmania are in the order of 4.2°C below present (Fletcher and Thomas, 2010).  
359 However, some minor rainforest persisted in refugia in the Australian subtropics (Donders et  
360 al., 2006; Petherick et al., 2008) and woodland and heath in the southwest (Dodson, 2001).

361

362 Average air temperature in the interior decreased significantly to as low as 9°C below  
363 present, determined from amino-acid racemisation of emu eggshells (Miller et al., 1997).  
364 Dune activity intensified in the arid core, but was most noticeable in the semi-arid desert  
365 margins (Fitzsimmons et al., 2007) where it was better preserved (Lomax et al., 2011;  
366 Fitzsimmons et al., this volume) and into the present-day temperate zone (Hesse et al., 2003;  
367 Duller and Augustinus, 2006; Gardner et al., 2006), indicating either less vegetation, windier  
368 conditions and/or increased sediment availability with the potential expansion of the arid  
369 zone beyond its modern extent. Lakes and rivers in the northern, monsoon-influenced arid  
370 zone generally experienced lower levels and reduced flow respectively, due to an absence of  
371 the monsoon in northern Australia during the LGM (Fitzsimmons et al., this volume).

372

373 By contrast, rainfall-fed lakes in the southeastern Australian highlands, record high or  
374 oscillating levels during the LGM (Coventry, 1976; Page et al., 1994; Bowler et al., 2012) .  
375 This is coeval with significant fluvial activity in the rivers of the Riverine Plain (Page et al.,  
376 1996; Kemp and Rhodes, 2010). These conditions are believed to reflect increased runoff  
377 from seasonal snow melt and reduced vegetation within the catchments, and do not preclude  
378 a drier climatic phase (Dosseto et al., 2010; Kemp and Rhodes, 2010; Bowler et al. 2012;  
379 Fitzsimmons et al., this volume). Evidence for at least periodically wet conditions exists also  
380 in the tropical region (Nott and Price, 1999; Reeves et al., 2007; Croke et al., 2010). In  
381 addition, there is evidence for increased precipitation in South Australia, from speleothem  
382 records from Naracoorte (Ayliffe et al., 1998) as well as recurrent large floods in silt-rich  
383 floodplains accumulating in the southern Flinders Ranges (Haberlah et al., 2010), attributed  
384 to a more northerly penetration of the westerlies.

385



386 **18-15 ka early deglacial period**

387 The first evidence of warming following the LGM comes from the Coral Sea, commencing  
388 around 20 ka (Tachikawa et al., 2009). Deglacial warming in Antarctica commenced around  
389 19 ka (Pedro et al., 2011), followed by the retreat of sea ice and increase in SST in the  
390 Southern Ocean, which occurred rapidly between 18 and 15 ka (Barrows et al., 2007a),  
391 accompanied by a dramatic increase in opal flux and atmospheric CO<sub>2</sub> (Armand and  
392 Levanter, 2010; Bostock et al., this volume). This warming resulted in the STF moving back  
393 to a more southerly position (Sikes et al., 2009). SST also increased off the east coast of  
394 Australia although lagged the warming in the south (Weaver et al., 2003; Petherick et al., this  
395 volume). Other than a brief decrease around 18 ka (Yokoyama et al., 2001), warming in the  
396 Indian Ocean lagged behind the Pacific Ocean, not commencing until ~15 ka (Martinez et al.,  
397 1999;). This time interval also saw glacial retreat in the Snowy Mountains around 16.8 ka  
398 (Barrows et al., 2001) and in Tasmania and New Guinea from ~18 ka, the latter becoming  
399 becoming more rapid after 15.4 ka (Barrows et al., 2011).

400

401 The rate of response in vegetation change during the deglacial period varied greatly across  
402 the region (Fig. 4di,ii). Wetter and warmer conditions are first noted in Indonesia around 17  
403 ka and NW Australia ~15 ka with more prolonged recovery in New Guinea and NE Australia  
404 (Reeves et al., this volume). Dry conditions persisted in Borneo, as recorded in speleothems,  
405 until ~15 ka, after which time there was a rapid shift to wetter conditions (Partin et al., 2007).  
406 Vegetation records from the temperate region showed a gradual response to warmer and  
407 wetter conditions, with an increase in arboreal taxa from ~15 ka (Petherick et al., this  
408 volume).

409

410 Although there was an expansion in Lake Carpentaria from 18 ka, the northern plunge pools  
411 were considered inactive and Lake Gregory may have been dry (Reeves et al., this volume).  
412 Other lakes in the arid interior and temperate regions were also low. High lake levels are  
413 recorded in Lake Frome from 18-16 ka, although dry conditions at Lake Eyre persisted (De  
414 Deckker et al., 2010; Cohen et al., 2011, 2012). However, fluvial discharge increased in the  
415 lower MDB and in coastal rivers of NSW, possibly in response to snow-melt (Page et al.,  
416 1996; 2009; Nanson et al., 2003). Although limited dune activity persisted throughout this  
417 period within the semi/arid zone (Fitzsimmons et al., 2007; Lomax et al., 2011), conditions in  
418 the southern part of the continent are likely to have become relatively more stable and humid.

419

#### 420 **15-12 ka late deglacial period**

421 Climate for the period from 15-12 ka was highly varied, with conflicting climatic responses  
422 across the Australian region (Fig 4ei,ii). Whilst glacier retreat was complete on the continent  
423 (Barrows et al., 2001, 2002), the re-advance of sea-ice and decrease in SST and atmospheric  
424 CO<sub>2</sub> in the Southern Ocean occurred between 14.5 and 13 ka during the Antarctic Cold  
425 Reversal (Pedro et al., 2011; Bostock et al., this volume). This resulted in a northward  
426 movement of the STF, again close to the southern Australian coast and suggests a lowering of  
427 SST offshore South Australia (Calvo et al., 2007). Wetter conditions are evident in western  
428 Tasmania around this time (14-11.7 ka; Fletcher and Moreno, 2011) and active river  
429 migration in the temperate zone continued throughout this period, perhaps due to snow melt,  
430 with evidence of increased precipitation in the lower MDB from 13.5 ka (Gingele et al.,  
431 2007; Petherick et al., this volume).

432

433 In contrast, SST warming commenced at 15 ka in the tropical north-west of Australia,  
434 coincident with the initial flooding of the Sunda Shelf (Bard, 1990). Warmer and wetter

435 conditions prevailed throughout the north, with the re-invigoration of the monsoon, bringing  
436 freshwater to the Gulf of Carpentaria and recovery of woodland and forest in PNG and NE  
437 Australia (Reeves et al., this volume). Increased precipitation associated with the monsoon  
438 from ~14 ka is evident in lake level transgression in the Gregory Lakes basin (Wyrwoll and  
439 Miller 2001) and by ~12 ka at Lake Eyre (Magee et al., 2004). The speleothem record of  
440 Borneo shows a rapid decrease in  $\delta^{18}\text{O}$ , reflecting an increase in precipitation after 15 ka,  
441 with a reversal to drier conditions between 13.3-12.3 ka (Partin et al., 2007). Desert dune  
442 records across both the arid zone and semi-arid desert margins of the MDB show aeolian  
443 activity persisted or possibly stabilised, during this period (Fitzsimmons et al. 2007; this  
444 volume; Lomax et al., 2011).

445

#### 446 **12-8 ka early Holocene**

447 Maximum warming in Antarctica occurred at 11.8 ka (Pedro et al., 2011). The peak in SSTs  
448 of the Southern Ocean (e.g. Barrows et al., 2007a) and ambient air temperature are coincident  
449 with the most poleward migration of the STF (Bostock et al., this volume). SST throughout  
450 the temperate and tropical regions approached modern values around 11-9 ka (Petherick et  
451 al.; Reeves et al, this volume), and saw a re-activation of the Leeuwin current extending to  
452 South Australia (Calvo et al., 2007; De Deckker et al., 2012). By 9.8 ka glaciers in New  
453 Guinea had disappeared (Barrows et al., 2011), most of the shallow shelf seas were re-  
454 established, and the land:sea ratio, particularly in the tropics, approached modern conditions.  
455 The mangrove swamps of the north and estuaries of the eastern seaboard had formed and  
456 growth of the Great Barrier Reef was re-initiated (Reeves et al., this volume).

457

458 Increasingly wet conditions in Borneo and Flores from 12 and 11 ka coincided with warming  
459 of the South China Sea (Partin et al., 2007; Griffiths et al, this volume) (Fig 4fi.ii). Wet

460 conditions are also evidenced by the active plunge pools in the north of Australia and  
461 vegetation similar to modern in Indonesia by 11 ka and NE Australia by 9 ka, with the  
462 establishment of rainforest taxa (Reeves et al., this volume). Interestingly, although the early  
463 Holocene fluvial records of the Lake Eyre Basin see a peak in discharge compared to present,  
464 those of the Riverine Plain of the lower MDB show a significant decline compared with the  
465 deglacial, with a shift from bedload to suspended-load sedimentation, caused primarily by a  
466 reduction in discharge (Fitzsimmons et al., this volume; Petherick et al., this volume).

467

468 Vegetation records from across the temperate region east of the Great Dividing Range show a  
469 shift at ~12-11 ka to include a greater representation of rainforest taxa, indicating both  
470 warmer and wetter conditions than previous (Petherick et al., this volume). Lake levels at  
471 Keilambete and George also increased after 11 ka (Wilkins et al., in press; Fitzsimmons and  
472 Barrows, 2010) and peaked after 9.6 ka in the Wimmera (Kemp et al., 2012).

473

#### 474 **8-5 ka mid-Holocene**

475 The mid-Holocene period represents maximum temperature in terrestrial records throughout  
476 the Australasian region, although it was expressed in different places at different times (Fig  
477 4gi,ii). By 8 ka both sea level and SST had reached essentially modern conditions (Lewis et  
478 al., this volume) and after a decline from the high temperatures of the early Holocene,  
479 Antarctic temperatures stabilised (EPICA, 2004). In the north, the thermal maximum of the  
480 IPWP was achieved by 6.8-5.5 ka (Abram et al., 2009). Peak wet conditions occurred on  
481 Flores by 8-6 ka, but only by ~5 ka in Borneo (Reeves et al., this volume). There is an  
482 increase in composite charcoal from the north at 8 ka, associated with a switch from grass-  
483 dominant to rainforest taxa from the northeast (Mooney et al., 2011; Moss and Kershaw,  
484 2007).

485

486 Offshore southern Australia experienced a peak in SST at ~6 ka of 18-21°C (Calvo et al.,  
487 2007). An increased representation of rainforest taxa, peak in representation in the macro-  
488 charcoal record especially in the north (Mooney et al., 2011) and a peak in discharge in the  
489 fluvial records of the east coast support both warmer and wetter conditions centred around 6  
490 to 5.5 ka (Petherick et al., this volume). Dune activity persisted in the desert margins of the  
491 MDB (Lomax et al., 2011). Increasingly humid conditions from 7 to 5 ka are evident around  
492 the Flinders Ranges and Lake Frome (Fitzsimmons et al., this volume). Both Lake George  
493 (Fitzsimmons and Barrows, 2010) and the lakes of western Victoria (Wilkins et al., in press)  
494 record high lake conditions, indicating peak effective precipitation, in the early-mid-  
495 Holocene. Although the peak warm and relatively wet conditions throughout the Australian  
496 region all occur sometime during this mid-Holocene period, the east coast appears to lag the  
497 more southerly sites.

498

#### 499 **5-0 ka late-Holocene**

500 Increasing variability with enhanced drier conditions was the overall characteristic of much  
501 of the Australasian region through the Late Holocene, indicative of ENSO in El Niño mode  
502 (Fig 4hi,ii). This is seen in the coral (e.g. Tudhope et al., 2001; Gagan et al., 2004; McGregor  
503 and Gagan, 2004) and speleothem (Griffiths et al., 2009, 2010 a,b) records in the north, and  
504 reactivation of dunes and dust deposits in the northern, interior and temperate zones (e.g.  
505 Shulmeister and Lees, 1995; Fitzsimmons et al., 2007; Marx et al., 2009, 2011; Fitzsimmons  
506 and Barrows, 2010). Fluvial activity decreased (e.g. Cohen and Nanson, 2007), with lower  
507 lake levels across much of the continent (e.g. De Deckker, 1982; Wilkins et al. in press). In  
508 contrast, the southwest was relatively consistently warm and moist (e.g. Gouramanis et al.,  
509 2012). An increasing number of records across the continent are showing variability in the

510 order of 1000-2000 years, attributed to El Niño in the north (e.g. McGregor and Gagan, 2004;  
511 Turney et al., 2004) and changes in the westerly wind belt in the south (e.g. Moros et al.,  
512 2009; Kemp et al., 2012; Wilkins et al., in press). Although there have been recent records  
513 which focus particularly on decadal and finer resolution of the last 2000 years in the  
514 Australasian region, these are covered in other reviews (e.g. Neukom and Gergis, 2011).

515

#### 516 **4. Discussion**

517 The changes in climate we have documented over the last 35 kyr highlight a complex  
518 response across a continent that spans a vast array of climatic zones. The latitudinal range,  
519 together with its extensive interior and fringing mountainous areas, results in the interplay of  
520 all the major circulation systems of the low and mid latitudes over the continent. The  
521 consequences of this are often diachronous, and sometimes contradictory, responses of the  
522 landscape, vegetation and sea surface when sites are compared across this range. This  
523 patchwork response is in strong contrast to the Northern Atlantic Ocean, where huge changes  
524 in heat flux from the ocean act to synchronise the climate of Europe through time. A  
525 consequence of this is that the climate signal preserved in the ice cores of Greenland can act  
526 as a template for the stratigraphy of a large proportion of the European region (e.g. Lowe et  
527 al., 2001). No such obvious simple approach exists for Australia that can adequately describe  
528 or characterise the changes that take place. Both the magnitude and pattern of climate change  
529 vary meridionally across the Australian continent. At the sea surface, the highest magnitude  
530 temperature changes occur in the high latitudes (cf. Barrows et al., 2007b) and on land the  
531 highest changes tend to be in the highest altitudes.

532

533 A major limitation on the characterisation of Australian climate change is the lack of  
534 quantitative proxies for both temperature and precipitation across a large proportion of the

535 continent, with exceptions being in the far north (e.g. Lewis et al., 2011; Griffiths et al., this  
536 volume) and far south (e.g. Rees et al., 2008; Fletcher and Thomas, 2010), confounded by  
537 limitations in the ability to date events precisely. Although chronologies are constantly  
538 improving, aided by increased luminescence, exposure age and radiocarbon data sets, the  
539 interpretation of many records remains difficult. Landscape responses such as lake level  
540 change, river discharge and dune activity can be produced under a range of different climate  
541 scenarios, from variable combinations of temperature and precipitation. Knowledge of  
542 temperature change across the continent is consequently mostly guided by sea-surface  
543 temperature records from around the periphery of the continent.

544

545 The changes in temperature and effective precipitation recorded across Australia during the  
546 last 35 kyr suggest large scale reorganisation of the circulation systems and major shifts in  
547 climate zones. However, current evidence indicates that the majority of these changes can be  
548 explained and accommodated using existing climate systems. Most of the changes in  
549 precipitation can be explained by the presence or absence of the monsoon in the northern half  
550 of the continent, and the meridional shifts in the westerlies and their intensity in the south.  
551 Temperature changes have not necessarily occurred in tandem with these moisture delivery  
552 changes, leading to effective precipitation at times being out of phase with major global  
553 temperature changes. This is illustrated for example during the deglaciation when both wet  
554 and arid periods are superimposed upon a dominantly warming trend.

555

556 The onset of glacial-magnitude cooling is recorded first in the high latitudes of the Australian  
557 region in the Southern Ocean, as early as 40 ka (Barrows et al., 2007a). Vandergoes et al.  
558 (2005) suggested that the early onset of cooling in the high southern latitudes relates to a  
559 minimum in Southern Hemisphere summer insolation around 30-35 ka. However, this

560 suggestion contrasts with the lock-step relationship of temperature between the hemispheres  
561 during full glacial conditions and the Holocene. The lag in cooling of the low latitudes  
562 resulted in a stronger meridional temperature gradient than at present, which is likely to have  
563 had a response in stronger westerly and trade wind systems, delivering more water to inland  
564 lake and river systems. However, the Asian monsoon probably migrated northward between  
565 ~40-35 ka, removing the major source of inflow for the monsoon dominated north and Lake  
566 Eyre basin (e.g. Magee et al., 2004).

567

568 High resolution SST records indicate one or more interstadial periods around 30 ka (e.g.  
569 Barrows et al., 2007a), which also occur in ice core records from Antarctica (e.g., EPICA,  
570 2006), but temperature remained well below Holocene levels. Most records indicate steady  
571 cooling from 30-20 ka. However, there is a brief reversal in the cooling trend during the  
572 period 26-24 ka recorded in Antarctic ice cores (Fig. 4bi,ii; EPICA, 2006), which produced  
573 warm and wetter conditions in the NE of Australia, and a minor interstadial in the southern  
574 temperate zone and the higher latitudes of the Southern Ocean (Petherick et al., this volume;  
575 Bostock et al., this volume). An interstadial is also recorded in Antarctic ice core records at  
576 the same time (EPICA, 2006). The cause of this event is unknown and it slightly leads  
577 Greenland Interstadial 2 in the North Atlantic (Lowe et al., 2001), indicating a relationship  
578 through the 'bipolar seesaw' (Broecker, 1998).

579

580 Rapid growth of the Northern Hemisphere ice sheets from 30-20 ka (Lambeck and Chappell,  
581 2001) resulted in lower sea levels, which exposed the continental shelves around the  
582 Australian margin. Land bridges were formed between New Guinea and the north of  
583 Australia, forming the Sahul Shelf. The Indonesian archipelago was also largely connected,  
584 forming the Sunda Shelf. This not only altered ocean circulation, but also reduced the area of



585 warm shallow seas. This had a major influence on the tropical climates of northern Australia  
586 because of a greater restriction on sources of tropical rainfall (Reeves et al., this volume). The  
587 exposure of the continental shelves also has an acute regional affect on moisture delivery in  
588 Bass Strait and Torres Strait.

589

590 The LGM heralded cooler, windier and drier conditions across most of the continent (Fig.  
591 4ci,ii). The synchronicity of this response both at the sea surface (Barrows and Juggins, 2005)  
592 and on land, across a range of climatic zones, indicates a common regional control through  
593 greenhouse gas forcing of temperature. Carbon dioxide levels fell below 200 ppm before 25  
594 ka, and remained between 180-200 ppm until 17 ka (Schmitt et al., 2012). This interval  
595 corresponds to maximum temperature depression and decline of woody taxa, despite a peak  
596 in summer insolation in the Southern Hemisphere. However, the period of maximum  
597 glaciation and periglacial activity in Australia and Papua New Guinea was only a brief  
598 interval within this period, centred at 19-22 ka, indicating that carbon dioxide cannot be the  
599 only control on surface temperature. Despite lower precipitation and presumably more  
600 northerly penetration of the westerlies, lower evaporation and increased seasonality of  
601 discharge (because of the timing of snowmelt), resulted in apparently wetter conditions in  
602 parts of inland Australia (Kemp and Rhodes, 2010; Fitzsimmons et al., this volume).

603

604 Warming in the deglacial period occurred much earlier than in the Northern Hemisphere,  
605 commencing as early as 19 ka in the south (Pedro et al., 2011; Bostock et al., this volume)  
606 and established by ~18-17 ka through much of the region (Turney et al., 2006b; Petherick et  
607 al., this volume; Reeves et al., this volume) (Fig. 4di, ii). Warming preceded increases in  
608 precipitation in most areas, particularly the northwest of Australia. The lag in this region may  
609 be attributed in part to the restricted influence of warmer tropical waters, and a less

610 pronounced Leeuwin Current (Spooner et al., 2005, 2011; De Deckker et al., 2012). The  
611 presence of relatively high levels at Lake Frome, but dry Lake Eyre during the early deglacial  
612 period, in the absence of a monsoon may be attributed to a southerly moisture supply in the  
613 westerlies (Cohen et al., 2011, 2012).

614

615 Temperature reached near Holocene levels by 15 ka in many records. An increase in dune  
616 activity occurs at this time, perhaps as a result of temperature outpacing increases in  
617 precipitation. Because this is the last major period of dune activity, it may also be an  
618 indication of stabilisation and preservation of dunes, as much as an increase in aridity. The  
619 warming also coincides with the resumption of the monsoon increasing moisture delivery to  
620 the north (Fig. 4ei,ii) as low pressure systems developed over the continent.

621

622 With the exception of the marine records offshore South Australia (Calvo et al., 2007) the  
623 ACR (14 – 13.5 ka) is only weakly recorded in the southern part of the continent, mostly as a  
624 pause in the warming rather than as a distinct cooling event, producing wetter conditions in  
625 the temperate zone. No cooling is recorded during the Younger Dryas interval, as is expected  
626 under the 'bipolar seesaw' of interhemispheric heat exchange, nor is there a mechanism for  
627 causing a cooling at this time in the Australian region (Barrows et al., 2007b; De Deckker et  
628 al., 2012; Tibby, 2012).

629

630

631 Although the timing of response is varied, the early Holocene is a time of warmer and wetter  
632 conditions across the region. Conditions are wetter in the north associated with high sea-  
633 surface temperatures and moisture delivery enhanced because of flooding of the shallow shelf  
634 seas. The monsoon delivered more precipitation than at present, as a function of the pressure

635 gradient established between northern Australia and Asia (Fig. 4fi,ii). The exception to this  
636 would be the Riverine Plain in the lower Murray-Darling Basin, where the trend is towards a  
637 decrease in river discharge. This is associated with a southerly migration of the westerly wind  
638 belt, restricting moisture supply west of the Great Dividing Range, coupled with increasing  
639 temperatures, decreasing snow cover and subsequent melt in the upland areas which fed the  
640 rivers.

641

642 The establishment of maximum sea level (~8-7.5 ka; Lewis et al, this volume) is coincident  
643 with many regions recording maximum temperature and precipitation. This period sees the  
644 reinitiation of the coral reefs on the Great Barrier Reef and the flooding and infilling of the  
645 tropical estuaries with sediment trapped by mangroves (Reeves et al., this volume). In some  
646 areas, particularly the temperate zone, vegetation change and establishment of woody taxa  
647 lags behind changes in climate and rise in CO<sub>2</sub>, presumably owing to slow migration and re-  
648 establishment processes in forest ecosystems. Compared with the early Holocene, the trend at  
649 ~5 ka is for cooling conditions in the NE and Southern Ocean and drying in the interior (Fig.  
650 4 gi,ii). The flooding of the broader continental shelf along the northeast coast, may have  
651 buffered the climatic response during this interval; however the driving mechanism remains  
652 unresolved. During the late Holocene,. a weakening of the Walker Circulation, and thus a  
653 weakening of the easterly trade winds over the tropical Pacific Ocean, has been attributed to a  
654 weaker Asian summer monsoon system due to decreasing insolation at northern mid-latitudes  
655 during the boreal summer (Zheng et al., 2008). This created conditions more favourable for  
656 the development of El Niño events, resulting in a progressive increase in El Niño frequency  
657 (Phipps and Brown, 2010). The El Niño mode starts to dominate from 7-5 ka (Rodbell et al.,  
658 1999; Tudhope et al., 2001; Moy et al., 2002; Gagan et al., 2004) and this has the effect of  
659 producing drier conditions in northern and eastern Australia.

660

661 The Late Holocene sees an increase in climatic variability, particularly with regards to  
662 moisture availability (Fig. 4 hi,ii) largely attributed to a stronger expression of El Niño  
663 (Moros et al., 2009). This is over-printed by the interaction with the Indian Ocean Dipole and  
664 a weakening of the monsoon (Shulmeister and Lees, 1995; Shulmeister, 1999). Thus, the late  
665 Holocene records warm conditions, with greater El Niño frequency, increased fire and  
666 extended droughts, alternating with shorter, wet La Niña events (e.g. Moy et al., 2002; Moros  
667 et al., 2009; Mooney et al., 2011), a pattern which persists today .

668

## 669 **5. Further work**

670 There are clear gaps in the types and distribution of data across the greater Australian region  
671 and areas where the chronology could be significantly improved. As a palaeoclimate  
672 community there is still a lack of consensus with regards to the hydrological balance and role  
673 of groundwater in key time intervals such as the LGM. Quantitative estimates are required on  
674 wind strength, evaporation and seasonality; parameters that are currently largely  
675 unconstrained. The Australian region still lacks sufficient sub-millennial-scale resolution  
676 terrestrial records to identify climatic leads and lags and there are many locations across the  
677 continent, such as the majority of Western Australia and the Northern Territory, coastal  
678 southeastern Australia and terrestrial southwest Queensland, where data is non-existent.  
679 Lastly, the role of humans in modifying the biophysical environment, which influences the  
680 interpretation of climate, is one area that is poorly known for much of the last 35 kyr.

681

682 The OZ-INTIMATE project aimed to consolidate the climate records from the broader  
683 Australian region over the past ~35 kyr and compare the response of each of the four major  
684 climatic regions to global climatic drivers. Future work will need to engage more actively

685 with climate modelling to test the longevity of dynamical mechanisms of the climate system  
686 of the modern day and how these have manifested in the past. In addition, targeted palaeo-  
687 proxy data gathering must continue, with a focus on filling the spatial gaps outlined above,  
688 developing well-resolved chronologies and quantitative reconstructions. This combined  
689 approach will enhance our ability to predict the evolution of the climate system and the  
690 impacts of future changes on the climate of the Australian region.

691

## 692 **Acknowledgements**

693 The OZ-INTIMATE working group, comprising >50 researchers, is part of Australasian  
694 INTIMATE project #0806 of PALCOMM and we acknowledge INQUA's support. We are  
695 grateful also for the support of AINSE in hosting and sponsoring 3 workshops which enabled  
696 the group to meet to develop both this and the four Australian regional compilations  
697 presented in this volume. This paper has been enhanced by the comments of Patrick De  
698 Deckker and two anonymous reviewers and we are grateful for their contribution. SJP was  
699 supported under the Australian Research Council's Discovery Projects funding scheme  
700 (project number DP1092945) and TTB was supported by DP1095053. JMR and TJC received  
701 funding (via PALCOMM and the CcASH project of ANSTO) and encouragement from NZ-  
702 INTIMATE to re-engage the Australian palaeo-community in this project.

703 **References**

704 Abram, N.J., McGregor, H.V., Gagan, M.K., Hantoro, W.S., Suwargadi, B.W., 2009.

705 Oscillations in the southern extent of the Indo-Pacific Warm Pool during the mid- Holocene.

706 Quaternary Science Reviews 28, 2794-2803.

707

708 Armand, L.K., Leventer, A., 2010. Palaeo sea ice distribution and reconstruction derived from

709 the geological record. In: Sea Ice, Eds Thomas, D.N., Dieckmann, G.S. Wiley-Blackwell,

710 Oxford, pp. 469-530.

711

712 Ayliffe, L. K., Marianelli, P.C., Moriarty, K.C., Wells, R.T., McCulloch, M.T., Mortimer,

713 G.E., Hellstrom, J.C., 1998. 500 ka precipitation record from southeastern Australia:

714 Evidence for interglacial relative aridity, Geology 26, 147-150.

715

716 Bard, E., Hamelin, B. and Fairbanks, R.G., 1990. U-Th ages obtained by mass spectrometry

717 in corals from Barbados: sea level during the past 130,000 years. Nature, 346, 456-458.

718

719 Barrell, D.J.A, Almond, P., Vandergoes, M.J., Lowe, D.J., Newnham, R.M. and INTIMATE

720 members, accepted. A composite terrestrial stratotype for regional comparison of climatic

721 events in New Zealand over the past 30,000years (NZ-INTIMATE project). Quaternary

722 Science Reviews.

723

724 Barrows, T.T. and Juggins, S., 2005. Sea-surface temperatures around the Australian margin

725 and the Indian Ocean during the Last Glacial Maximum. Quaternary Science Reviews, 24,

726 1017-1047.

727

728 Barrows, T.T., Hope, G.S., Prentice, M.L., Fifield, L.K., Tims, S.G., 2011. Late Pleistocene  
729 glaciation of the Mt Giluwe volcano, Papua New Guinea. *Quaternary Science Reviews*, 30,  
730 2676-2689.

731

732 Barrows, T.T., Juggins, S., De Deckker, P., Calvo, E., and Pelejero, C., 2007a, Long-term  
733 sea-surface temperature and climate change in the Australian-New Zealand region:  
734 *Paleoceanography*, 22, PA 2215, doi:10.1029/2006PA001328.

735

736 Barrows, T.T., Juggins, S., De Deckker, P., Thiede, J., Martinez, J.I., 2000. Sea-surface  
737 temperatures of the southwest Pacific Ocean during the Last Glacial Maximum.  
738 *Paleoceanography*, 15, 95–109.

739

740 Barrows, T.T., Lehman, S.J., Fifield, L.K., De Deckker, P., 2007b. Absence of Cooling in  
741 New Zealand and the Adjacent Ocean During the Younger Dryas Chronozone. *Science*, 318,  
742 86-89.

743

744 Barrows, T.T., Reeves, J.M., Alloway, B.V., in press. The AUS-INTIMATE project special  
745 volume – editorial. *Quaternary Science Reviews*, this volume.

746

747 Barrows, T.T., Stone, J.O., Fifield, L.K., 2004. Exposure ages for Pleistocene periglacial  
748 deposits in Australia. *Quaternary Science Reviews*, 23(5-6), 697-708.

749

750 Barrows, T.T., Stone, J.O., Fifield, L.K., Cresswell, R.G., 2001. Late Pleistocene glaciation  
751 of the Kosciuzsko Massif, Snowy Mountains, Australia. *Quaternary Research*, 55, 179-189.

752

753 Barrows, T.T., Stone, J.O., Fifield, L.K., Cresswell, R.G., 2002. The timing of the Last  
754 Glacial Maximum in Australia. *Quaternary Science Reviews*, 21, 159-173.  
755  
756  
757  
758  
759 Berger, A.L., 1978. Long Term Variations of Daily Insolation and Quaternary Climatic  
760 Changes. *Journal of the Atmospheric Sciences*, 35, 2362-2367.  
761  
762 Blunier, T., Brook, E.J., 2001. Timing of Millennial-Scale Climate Change in Antarctica and  
763 Greenland During the Last Glacial Period. *Science*, 291, 109-112.  
764  
765 Bostock, H.C., Armand, L.K., Barrows, T.T., Carter, L., Chase, Z., Cortese, G., Dunbar, G.,  
766 Ellwood, M., Hayward, B., Howard, W., Neil, H.L., Noble, T.L., Mackintosh, A., Moss, P.T.,  
767 Moy, A.D., White, D., Williams, M.J.M., this volume. A review of the Australian-New  
768 Zealand sector of the Southern Ocean over the last 30 ka (Aus-INTIMATE project).  
769 *Quaternary Science Reviews*, this volume.  
770  
771 Bowler, J. M., Hope, G.S., Jennings, J.N., Singh, G., Walker, D., 1976. Late Quaternary  
772 Climates of Australia and New Guinea. *Quaternary Research*, 6, 359-394.  
773  
774 Bowler, J. M., Gillespie, R., Bolkjovac, K., Johnson. H., 2012. Wind v Water: Glacial  
775 Maximum records from the Willandra Lakes. In: S.G. Haberle & B. David (eds.) *Peopled*  
776 *Landscapes: Archaeological and Biogeographic Approaches to Landscapes*. Terra  
777 *Australis* 34. Canberra: ANU E-Press. pp. 271-296.



778

779 Bowler, J.M., Hamada, T., 1971. Late Quaternary Stratigraphy and Radiocarbon Chronology  
780 of Water Level Fluctuations in Lake Keilambete, Victoria. *Nature*, 232(5309), 330-332.

781

782 Bowler, J.M. and Wasson, R.J., 1984. Glacial age environments of inland Australia. In: J.C.  
783 Vogel (Ed.) *Late Cainozoic Palaeoclimates of the Southern Hemisphere*. Balkema, Rotterdam.

784

785 Broecker, W., 1998. Paleocean circulation during the last deglaciation: A bipolar seesaw?  
786 *Paleoceanography*, 13, 119.

787

788 Calvo, E., Pelejero, C., De Deckker, P., Logan, G.A., 2007. Antarctic deglacial pattern in a  
789 30 kyr record of sea surface temperature offshore South Australia. *Geophysical Research*  
790 *Letters*, 34, L13707, doi:10.1029/2007GL029937.

791

792 Cohen, T. J., Nanson, G.C., 2007. Mind the gap: an absence of valley fill deposits identifying  
793 the Holocene hypsithermal period of enhanced flow regime in southeastern Australia. The  
794 Holocene 17, 1-8.

795

796 Cohen, T.J., Nanson, G.C., Jansen, J.D., Jones, B.G., Jacobs, Z., Treble, P., Price, D.M.,  
797 May, J.-H., Smith, A.M., Ayliffe, L.K., Hellstrom, J.C., 2011. Continental aridification and  
798 the vanishing of Australia's megalakes. *Geology*, 39, 167-170.

799

800 Cohen, T.J., Nanson, G.C., Jansen, J.D., Jones, B.G., Jacobs, Z., Larsen, J.R., May, J.-H.,  
801 Treble, P., Price, D.M., Smith, A.M., 2012. Late Quaternary mega-lakes fed by the northern

802 and southern river systems of central Australia: varying moisture sources and increased  
803 continental aridity. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 356-357, 89-108.  
804

805 Colhoun, E.A., van de Geer, G., Mook, W.G., 1982. Stratigraphy, pollen analysis, and  
806 palaeoclimatic interpretation of Pulbeena Swamp, northwestern Tasmania. *Quaternary*  
807 *Research*, 18, 108-126.  
808

809 Colhoun, E. A., Hannah, D., Kiernan, K., 1996. Late Wisconsin glaciation of Tasmania.  
810 *Papers and Proceedings - Royal Society of Tasmania* 130 (2):33-45.  
811

812 Colhoun, E., 2000. Vegetation and climate change during the Last Interglacial-Glacial cycle  
813 in western Tasmania, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 155,  
814 195-209.  
815

816 Coventry, R.J., 1976. Abandoned shorelines and the late Quaternary history of Lake George,  
817 New South Wales. *Journal of the Geological Society of Australia*, 23, 249-273.  
818

819 Croke, J., Jansen, J.D., Amos, K., Pietsch, T.J., 2010. A 100 ka record of fluvial activity in  
820 the Fitzroy River Basin, tropical northeastern Australia. *Quaternary Science Reviews*, 30,  
821 1681-1695.  
822

823 De Deckker, P., 1982. Holocene ostracods, other invertebrates and fish remains from cores of  
824 four maar lakes in southeastern Australia. *Proceedings of the Royal Society of Victoria* 94:  
825 183-220.

826

827 De Deckker, P., Magee, J.W., Shelley, J.M.G., 2010. Late Quaternary palaeohydrological  
828 changes in the large playa Lake Frome in central Australia, recorded from the Mg/Ca and  
829 Sr/Ca in ostracod valves and biotic remains. *Journal of Arid Environments* 75: 38-50.

830

831 De Deckker, P., Moros, M., Perner, K., Jansen, J., 2012. Influence of the tropics and southern  
832 westerlies on glacial interhemispheric asymmetry. *Nature Geoscience*, 5, 266-269.

833

834 Devriendt, L.S.J, 2011. Late Quaternary environment of Palaeolake Carpentaria inferred from  
835 the chemistry of ostracod valves. Unpublished MSc Thesis, University of Wollongong.  
836 <http://ro.uow.edu.au/theses/3319>

837 Diaz H.F., and V. Markgraf 2000. *El Niño and the Southern Oscillation: Multiscale*  
838 *Variability and Global and Regional Impacts*, Cambridge University Press, 496 pp.

839

840 Dodson, J.R., 1975. Vegetation history and water fluctuations at Lake Leake, south-eastern  
841 South Australia. 2. 50,000 B.P. to 10,000 B.P. *Australian Journal of Botany*, 23(5), 815-831.

842

843 Dodson, J.R., 2001. Holocene vegetation change in the Mediterranean-type climate regions  
844 of Australia. *The Holocene*, 11, 673-680.

845

846 Donders, T. H., Wagner, F., Visscher, H., 2006. Late Pleistocene and Holocene subtropical  
847 vegetation dynamics recorded in perched lake deposits on Fraser Island, Queensland,  
848 Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 241(3-4), 417-439.

849

850 Dosseto, A., Hesse, P.P., Maher, K., Fryirs, K., Turner, S., 2010. Climatic and vegetation  
851 control on sediment dynamics during the last glacial cycle. *Geology*, 38(5), 395-398.  
852

853 Duller, G. A. T., Augustinus, P. C. 2006. Reassessment of the record of linear dune activity  
854 in Tasmania using optical dating. *Quaternary Science Reviews*, 25, 2608-2618.  
855

856 Dunbar, G.B., Dickens, G.R., 2003. Late Quaternary shedding of shallow marine carbonate  
857 along a tropical mixed siliciclastic-carbonate shelf: Great Barrier Reef, Australia.  
858 *Sedimentology*, 50, doi: 10.1046/j.1365-3091.2003.00593  
859

860 EPICA community members. 2004. Eight glacial cycles from an Antarctic ice core. *Nature*,  
861 429, 623-628.  
862

863 EPICA, 2006. One-to-one coupling of glacial climate variability in Greenland and Antarctica.  
864 *Nature*, 444, 195-198.  
865

866 Fitzsimmons, K.E., Rhodes, E.J., Magee, J.W., Barrows, T.T. 2007. The timing of linear  
867 dune activity in the Strzelecki and Tirari Deserts, Australia. *Quaternary Science Reviews*, 26,  
868 2598-2616.  
869

870 Fitzsimmons, K.E., Cohen, T., Hesse, P.P., Jansen, J., Nanson, G.C., May, J.-H., Barrows,  
871 T.T., Haberlah, D., Hilgers, A., Kelly, T., Larsen, J., Lomax, J., Treble, P., in review. Late  
872 Quaternary palaeoenvironmental change in the Australian drylands. *Quaternary Science*  
873 *Reviews*, this volume.  
874

875 Fitzsimmons, K.E., Barrows, T.T., 2010. Holocene hydrologic variability in temperate  
876 southeastern Australia: An example from Lake George, New South Wales. *The Holocene*,  
877 20(4), 585–597.

878

879 Fletcher, M.-S., Thomas, I., 2010. A quantitative Late Quaternary temperature reconstruction  
880 from western Tasmania, Australia. *Quaternary Science Reviews*, 29, 2351-2361.

881

882 Fletcher, M.-S., Moreno, P.I. 2011. Zonally symmetric changes in the strength and position  
883 of the Southern Westerlies drove atmospheric CO<sub>2</sub> variations over the past 14 k.y. *Geology*,  
884 doi: 10.1130/G31807.1

885

886 Gagan, M.K., Hendy, E.J., Haberle, S.G. and Hantoro, W.S., 2004. Post-glacial evolution of  
887 the Indo-Pacific warm pool and El Nino-Southern Oscillation. *Quaternary International*, 118,  
888 127–143.

889

890 Gallant, A.J.E., Kiem, A.S., Verdon-Kidd, D.C., Stone, R.C. and Karoly, D.J. 2012.  
891 Understanding hydroclimate processes in the Murray-Darling Basin for natural resources  
892 management. *Hydrology and Earth System Sciences*, 16, 2049-2068, doi:10.5194/hess-16-  
893 2049-2012.

894

895 Galloway, R.W., 1965. Late Quaternary climates in Australia. *Journal of Geology*, 73, 603-  
896 18.

897

898 Gardner, T. W., Webb, J., Davis, A. G., Cassel, E. J., Pezzia, C., Merritts, D. J., Smith, B.,  
899 2006. Late Pleistocene landscape response to climate change: eolian and alluvial fan

900 deposition, Cape Liptrap, southeastern Australia. *Quaternary Science Reviews*, 25, 1552-  
901 1569.

902

903 Gentilli, J., 1971. *Climates of Australia and New Zealand. World Survey of*  
904 *Climatology*, Volume 13. Elsevier, Amsterdam, 405 pp.

905

906 Gersonde, R., Crosta, X., Abelmann, A., Armand, L., 2005. Sea-surface temperature and sea  
907 ice distribution of the Southern Ocean at the EPILOG Last Glacial Maximum – a circum-  
908 Antarctic view based on siliceous microfossil records. *Quaternary Science Reviews*, 24, 869-  
909 896.

910

911 Gimeno, L., Drumond, A., Nieto, R., Trigo, R. M., Stohl, A., 2010. On the origin of  
912 continental precipitation. *Geophysical Research Letters*, 37, L13804.

913

914

915 Gingele, F.X., De Deckker, P., Norman, M., 2007. Late Pleistocene and Holocene climate of  
916 SE Australia reconstructed from dust and river loads deposited offshore the River Murray  
917 mouth. *Earth Planetary Science Letters*, 255, 257-272.

918

919 Gouramanis, C., Dodson, J., Wilkins, D., De Deckker, P., Chase, B.M., 2012. Holocene  
920 palaeoclimate and sea level fluctuation recorded from the coastal Barker Swamp, Rottne  
921 Island, south-western Western Australia. *Quaternary Science Reviews*  
922 doi10.1016/j.quascirev.2012.05.007.

923

924 Griffiths, M.L., Drysdale, R.N., Gagan, M.K., Zhao, J.-x., Ayliffe, L.K., Hellstrom, J.C.,

925 Hantoro, W.S., Frisia, S., Feng, Y.-x., Cartwright, I., St Pierre, E., Fischer, M.J., Suwargadi,  
926 B.W., 2009. Increasing Australian-Indonesian monsoon rainfall linked to early Holocene sea  
927 level rise. *Nature Geoscience* 2, 636-639.

928

929 Griffiths, M.L., Drysdale, R.N., Gagan, M.K., Frisia, S., Zhao, J.-x., Ayliffe, L.K., Hantoro,  
930 W.S., Hellstrom, J.C., Fischer, M.J., Feng, Y.-x., Suwargadi, B.W., 2010a. Evidence for  
931 Holocene changes in Australian- Indonesian monsoon rainfall from stalagmite trace element  
932 and stable isotope ratios. *Earth and Planetary Science Letters* 292, 27-38.

933

934 Griffiths, M.L., Drysdale, R.N., Gagan, M.K., Zhao, J., Hellstrom, J.C., Ayliffe, L. And  
935 Hanton, W.S., this volume. Abrupt increase in east Indonesian rainfall from flooding of the  
936 Sunda Shelf ~9,500 years ago. *Quaternary Science Reviews*.

937

938 Griffiths, M.L., Drysdale, R.N., Vonhof, H.B., Gagan, M.K., Zhao, J.-x., Ayliffe, L.K.,  
939 Hantoro, W.S., Hellstrom, J.C., Cartwright, I., Frisia, S., Suwargadi, B.W., 2010b. Younger  
940 Dryas–Holocene temperature and rainfall history of southern Indonesia from  $\delta^{18}\text{O}$  in  
941 speleothem calcite and fluid inclusions. *Earth and Planetary Science Letters* 295, 30-36.

942 Haberlah, D., Williams, M.A.J., Halverson, G., Mctainsh, G.H., Hill, S.M., Hrstka, T., Jaime,  
943 P., Butcher, A.R., Glasby, P., 2010. Loess and floods: High –resolution multi-proxy data of  
944 Last Glacial Maximum (LGM) slackwater deposition in the Flinders Ranges, semi-arid Sout  
945 Australia. *Quaternary Science Reviews* 29: 2673-2693.

946

947 Hesse, P.P., Magee, J.W., van der Kaars, S., 2004. Late Quaternary climates of the Australian  
948 arid zone: A review. *Quaternary International*, 118-119, 87-102.

949

950 Hesse, P.P., 2010. The Australian desert dunefields: formation and evolution in an old, flat,  
951 dry continent. In *Australian Landscapes*, eds. P. Bishop & B. Pillans, 141-163. London:  
952 Geological Society.

953

954 Hesse, P.P., McTainsh, G.H., 1999. Last Glacial Maximum to Early Holocene wind strength  
955 in the Mid-latitudes of the Southern Hemisphere from aeolian dust in the Tasman Sea.  
956 *Quaternary Research*, 52, 343-349.

957

958 Hesse, P.P., Humphreys, G.S., Selkirk, P.M., Adamson, D.A., Gore, D.B., Nobes, D.C.,  
959 Price, D.M., Schwenninger, J.-L., Smith, B., Tulau, M., Hemmings, F., 2003. Late  
960 Quaternary aeolian dunes on the presently humid Blue Mountains, Eastern Australia.  
961 *Quaternary International* 108, 13-32.

962

963 Hope, G. and Tulip, J., 1994. A long vegetation history from lowland Irian Jaya, Indonesia.  
964 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 109, 385-398.

965

966 Hope, G., 2009. Environmental change and fire in the Owen Stanley Ranges, Papua New  
967 Guinea. *Quaternary Science Reviews*, 28, 2261-2276.

968

969 Jorry, S., Droxler, A.W., Mallarino, G., Dickens, G.R., Bentley, S.J., Beaufort, L., Peterson,  
970 L.C., Opdyke, B.N., 2008. Bundled turbidite deposition in the central Pandora Trough (Gulf  
971 of Papua) since Last Glacial Maximum: Linking sediment nature and accumulation to sea  
972 level fluctuations at millennial timescale. *Journal of Geophysical Research*, 113, F01S19,  
973 doi:10.1029/2006JF000649.

974



975 Kemp, J., Rhodes, E., 2010. Episodic fluvial activity of inland rivers in southeastern  
976 Australia: Palaeochannel systems and terraces of the Lachlan River Quaternary Science  
977 Reviews, 29, 732-752.

978

979 Kemp, J., Radke, L.C., Olley, J., Juggins, S., De Deckker, P., 2012. Holocene lake salinity  
980 changes in the Wimmera, southeastern Australia, provide evidence for millennial-scale  
981 climate variability. Quaternary Research, 77(1), 65-76.

982

983 Kershaw, A.P., McKenzie, G.M., Porch, N., Roberts, R.G., Brown, J., Heijnis, H., Orr, M.L.,  
984 Jacobsen, G., Newall, P.R., 2007. A high-resolution record of vegetation and climate through  
985 the last glacial cycles from Caledonia Fen, southeastern highlands of Australia. Journal of  
986 Quaternary Science, 22(5), 481-500.

987

988 Kiem, A.S., Franks, S.W., Kuczera, G., 2003. Multi-decadal variability of flood risk.  
989 Geophysical Research Letters, 30(2), 1035, 10.1029/2002GL015992.

990

991 Kiem, A.S., Franks, S.W., 2004. Multi-decadal variability of drought risk - Eastern  
992 Australia. Hydrological Processes, 18(11), 2039-2050.

993

994 Kiernan, K., Fifield, L.K., Chappell, J., 2004. Cosmogenic ages for Last Glacial Maximum  
995 moraine at Schnells Ridge, Southwest Tasmania. Quaternary Research, 61, 335-338.

996

997 Lambeck, K. and Chappell, J., 2001. Sea level change through the last glacial cycle. Science,  
998 295, 679-686.

999

1000 Lewis, S.C., Gagan, M.K., Ayliffe, L.K., Zhao, J.-x., Hantoro, W.S., Treble, P.C., Hellstrom,  
1001 J.C., LeGrande, A.N., Kelley, M., Schmidt, G.A., 2011. High-resolution stalagmite  
1002 reconstructions of Australian-Indonesian monsoon rainfall variability during Heinrich stadial  
1003 3 and Greenland interstadial 4. *Earth and Planetary Science Letters*, 303, 133-142.  
1004  
1005 Lewis, S.E., Sloss, C.R., Murray-Wallace, C.V., Woodroffe, C.D., Smithers, S.G., in review.  
1006 Postglacial sea-level change around the Australian mainland. *Quaternary Science Reviews*,  
1007 this volume.  
1008  
1009 Lomax, J., Hilgers, A., Radtke, U., 2011. Palaeoenvironmental change recorded in the  
1010 palaeodunefields of the western Murray Basin, South Australia - New data from single grain  
1011 OSL-dating. *Quaternary Science Reviews*, 30, 723-736.  
1012  
1013 Lourantou, A., Lavric, J.V., Kohler, P., Barnola, J.-M., Paillard, D., Michel, E., Raynaud, D.,  
1014 Chappellaz, J., 2010. Constraint on the CO<sub>2</sub> rise by new atmospheric carbon isotopic  
1015 measurements during the last deglaciation. *Global Biogeochemical Cycles* 24, GB2015,  
1016 doi:10.1029/2009GB003545.  
1017  
1018  
1019 Mackintosh, A.N., Barrows, T.T., Colhoun E.A., Fifield, L.K., 2006. Exposure dating and  
1020 glacial reconstruction at Mt. Field, Tasmania, Australia, identifies MIS 3 and MIS 2 glacial  
1021 advances and climatic variability. *Journal of Quaternary Science*, 21, 363-376.  
1022

1023 Magee, J.W., Miller, G.H., Spooner, N.A., Questiaux, D., 2004. Continuous 150 k.y.  
1024 monsoon record from Lake Eyre, Australia: Insolation-forcing implications and unexpected  
1025 Holocene failure. *Geology*, 32, 885-888.  
1026  
1027 Marx, S. K., McGowan, H. A., Kamber, B. S., 2009. Long-range dust transport from eastern  
1028 Australia: A proxy for Holocene aridity and ENSO-type climate variability. *Earth and*  
1029 *Planetary Science Letters* 282, 167-177.  
1030  
1031 Marx, S. K., Kamber, B. S., McGowan, H. A., Denholm, J., 2011. Holocene dust deposition  
1032 rates in Australia's Murray-Darling Basin record the interplay between aridity and the  
1033 position of the mid-latitude westerlies. *Quaternary Science Reviews* 30, 3290-3305.  
1034  
1035 Martinez, J.I., De Deckker, P., Barrows, T., 1999. Paleoceanography of the Last glacial  
1036 maximum in the eastern Indian Ocean: planktonic foraminifera evidence. *Palaeogeography,*  
1037 *Palaeoclimatology, Palaeoecology*, 147, 73-99.  
1038  
1039 McGregor, H.V., Gagan, M.K., 2004. Western Pacific coral  $\delta^{18}\text{O}$  record of anomalous  
1040 Holocene variability in the El Niño-Southern Oscillation. *Geophysical Research Letters* 31,  
1041 L11204, doi:10.1029/2004GL019972.  
1042 Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M.,  
1043 Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H.,  
1044 Thordarson, T., 2012. Abrupt onset of the Little Ice Age triggered by volcanism and  
1045 sustained by sea-ice/ocean feedbacks, *Geophysical Research Letters*, 39, L02708,  
1046 doi:10.1029/2011GL050168.  
1047

1048 Miller, G.H., Magee, J.W., Jull, A.J.T., 1997. Low-latitude glacial cooling in the southern  
1049 hemisphere from amino-acid racemisation in emu eggshells. *Nature*, 385, 241-244.  
1050

1051 Mooney, S.D., Harrison, S.P., Bartlein, P.J., Daniau, A.-L., Stevenson, J., Brownlie, K.C.,  
1052 Buckman, S., Cupper, M., Luly, J., Black, M., Colhoun, E., D'Costa, D., Dodson, J., Haberle,  
1053 S., Hope, G.S., Kershaw, P., Kenyon, C., McKenzie, M., Williams, N., 2011. Late Quaternary  
1054 Fire regimes in Australia. *Quaternary Science Reviews*, 30, 28-46.  
1055

1056 Moros, M., De Deckker, P., Jansen, E., Perner, K., Telford, R.J., 2009. Holocene climate  
1057 variability in the Southern Ocean recorded in a deep-sea sediment core off South Australia.  
1058 *Quaternary Science Reviews*, 28, 1932-1940.  
1059

1060 Moss, P.T., Kershaw, A.P., 2007. A late Quaternary marine palynological record (oxygen  
1061 isotope stages 1 to 7) for the humid tropics of northeastern Australia based on ODP Site 820.  
1062 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 251, 4-22.  
1063

1064 Moy, C.M., Seltzer, G.O., Rodbell, D.T., Anderson, D.M., 2002. Variability of El  
1065 Niño/SouthernOscillation activity at millennial timescales during the Holocene epoch.  
1066 *Nature*, 402, 162-165.  
1067

1068 Nanson, G. C., Price, D. M., Short, S. A., 1992. Wetting and drying of Australia over the  
1069 past. *Geology*, 20, 791-794.  
1070

1071 Nanson, G.C., Cohen, T.J., Doyle, J., Price, D.M., 2003. Alluvial evidence of late-Quaternary  
1072 climate and flow-regime changes on the coastal rivers of New South Wales, Australia.

1073 Palaeohydrology: Understanding Global Change. K. J. Gregory and G. Benito, Eds.  
1074 Chichester, Wiley, 233-258.  
1075  
1076 Nanson, G.C., Price, D.M., Jones, B.G., Maroulis, J.C., Coleman, M., Bowman, H., Cohen,  
1077 T.J., Pietsch, T.J., Larsen J.R., 2008. Alluvial evidence for major climate and flow regime  
1078 changes during the middle and late Quaternary in eastern central Australia. *Geomorphology*,  
1079 101, 109-129.  
1080  
1081 Neukom, R., Gergis, J., 2011: Southern Hemisphere high-resolution palaeoclimate records of  
1082 the last 2000 years. *The Holocene*, doi:10.1177/0959683611427335.  
1083  
1084 Newsome, J., Flenley, J.R., 1988. Late Quaternary vegetational history of the Central  
1085 Highlands of Sumatra. II. Palaeopalynology and vegetational history. *Journal of*  
1086 *Biogeography*, 15, 555-578.  
1087  
1088 Nott, J.F., Price, D.M., 1999. Waterfalls, floods and climate change: evidence from tropical  
1089 Australia. *Earth and Planetary Science Letters*, 171, 267-276.  
1090  
1091 Page, K.J., Nanson, G.C. Price, D., 1996. Chronology of Murrumbidgee river  
1092 palaeochannels on the Riverine Plain, southeastern Australia. *Journal of Quaternary Science*,  
1093 11, 311-326.  
1094  
1095  
1096 Page, K., Kemp, J., G.C. Nanson, G.C., 2009. Late Quaternary evolution of Riverine Plain  
1097 paleochannels, southeastern Australia. *Australian Journal of Earth Sciences*, 56, 19-33.

1098

1099 Page, K., Nanson, G., Price, D., 1996. Chronology of Murrumbidgee River palaeochannels  
1100 on the Riverine Plain, southeastern Australia. *Journal of Quaternary Science*, 11(4), 311-326.

1101

1102 Page, K. J. and Nanson, G.C., 1996. Stratigraphic architecture resulting from late Quaternary  
1103 evolution of the Riverine Plain, southeastern Australia. *Sedimentology*, 43, 927-945.

1104

1105 Partin, J.W., Cobb, K.M., Adkins, J.F., Clark, B., Fernandez, D.P., 2007. Millennial-scale  
1106 trends in west Pacific warm pool hydrology since the Last Glacial Maximum. *Nature*, 449,  
1107 452-455.

1108

1109 Pedro, J.B., van Ommen, T.D., Rasmussen, S.O., Morgan, V.I., Chappellaz, J., Moy, A.D.,  
1110 Masson-Delmotte, V., Delmotte, M., 2011. The last deglaciation: timing the bipolar see-saw.  
1111 *Climate of the Past*, 7, 671-683.

1112

1113 Petherick, L., McGowan, H., Moss P., 2008. Climate variability during the Last Glacial  
1114 Maximum in eastern Australia: evidence of two stadials? *Journal of Quaternary Science*, 23,  
1115 787-802.

1116

1117 Petherick, L., McGowan, H.A., Kamber, B.S., 2009. Reconstructing transport pathways for  
1118 late Quaternary dust from eastern Australia. *Geomorphology* 105, 67-79.

1119

1120 Petherick, L., Bostock, H.C., Cohen T., Fitzsimmons, K., Tibby, J., Moss, P., Mooney, S.,  
1121 Barrow, T., Reeves, J., Kemp, J., Jansen, J., Nanson, G., De Deckker, P., Fletcher, M. and  
1122 OZ-INTMATE members, this volume. *Quaternary Science Reviews*.

1123

1124 Petit, J.R., Jouzel, J., Raynaud, D., Barkov, N.I., Barnola, J.-M., Basile, I., Bender, M.,  
1125 Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V.M., Legrand, M.,  
1126 Lipenkov, V.Y., Lorius, C., Pépin, L., Ritz, C., Saltzman E., Stievenard M., 1999. Climate  
1127 and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica,  
1128 *Nature*, 399, 429-436.

1129

1130 Phipps, S.J. and Brown, J.N., 2010. Understanding ENSO dynamics through the exploration  
1131 of past climates. *Earth and Environmental Science*, 9, doi:10.1088/1755-1315/9/1/012010.

1132

1133 Pitman, A.J., Narisma, G.T., Pielke Sr., R.A., Holbrook, N.J., 2004. Impact of land cover  
1134 change on the climate of southwest Western Australia. *Journal of Geophysical Research-*  
1135 *Atmospheres*, 109, D18109.

1136

1137 Plummer, C.T., Curran, M.A.J., van Ommen, T.D., Rasmussen, S.O., Moy, A.D., Vance,  
1138 T.R., Clausen, H.B., Vinther, B.M., Mayewski, P.A., 2012. An independently dated 2000-yr  
1139 volcanic record from Law Dome, East Antarctica, including a new perspective on the dating  
1140 of the 1450s CE eruption of Kuwae, Vanuatu, *Climate of the Past*, 8, 1929-1940,  
1141 doi:10.5194/cp-8-1929-2012.

1142

1143 Power, S., Casey, T., Folland, C., Colman, A. Mehta, V., 1999. Inter-decadal modulation of  
1144 the impact of ENSO on Australia. *Climate Dynamics*, 15, 319-324.

1145

1146 Prentice, M.L., Hope, G.S., Maryunani, K., Peterson, J.A., 2005. An evaluation of snowline  
1147 data across New Guinea during the last major glaciation, and area-based glacier snowlines in

1148 the Mt. Jaya region of Papua, Indonesia, during the Last Glacial Maximum. *Quaternary*  
1149 *International*, 138-139, 93-117.

1150

1151 Prentice, M.L., Hope, G.S., Peterson, J.A., Barrows, T.T., 2011. Chapter 73 - The Glaciation  
1152 of the South-East Asian Equatorial Region. In: Ehlers, J., Gibbard, P.L., Hughes, P.D. (Eds.),  
1153 *Developments in Quaternary Sciences*. Elsevier, pp. 1023-1036.

1154

1155 Rees, A.B.H, Cwynar, L.C., Cranston, P.S., 2008. Midges (Chironomidae, Ceratopogonidae,  
1156 Chaoboridae) as a temperature proxy: a training set from Tasmania, Australia. *Journal of*  
1157 *Paleolimnology* 40: 1159-1178.

1158

1159 Reeves J.M., Bostock H.C., Ayliffe L.K., Barrows T.T., De Deckker P., Devriendt L.S.,  
1160 Dunbar G.B., Drysdale R.N., Fitzsimmons K.E., Gagan M.K., Griffiths M.L., Haberle S.G.,  
1161 Jansen, J.D., Krause C.E., Lewis S., McGregor H.V., Mooney S.D., Moss P., Nanson G.,  
1162 Purcell A. and van der Kaars, S., this volume. *Palaeoenvironmental change in tropical*  
1163 *Australasia over the last 30 000 years - A synthesis by the OZ-INTIMATE group*. *Quaternary*  
1164 *Science Reviews*, this volume.

1165

1166 Reeves, J.M., Chivas, A.R., Garcia, A., De Deckker, P., 2007. *Palaeoenvironmental change*  
1167 *in the Gulf of Carpentaria (Australia) since the last interglacial based on Ostracoda*.  
1168 *Palaeogeography, Palaeoclimatology, Palaeoecology* 246, 163-187.

1169

1170 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey,  
1171 C.B., Buck, C.E., Burr, G.S., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P.,  
1172 Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac,



1173 F.G., Manning, S.W., Reimer, R.W., Richards, D.A., Southon, J.R., Talamo, S., Turney,  
1174 C.S.M., van der Plicht, J., Weyhenmeyer, C.E., 2009. INTCAL09 and MARINE09  
1175 radiocarbon calibration curves, 0-50,000 years cal BP. *Radiocarbon*, 51, 1111-1150.  
1176  
1177 Rodbell, D.T., Selzer, G.O., Anderson, D.M., Abbott, M.B., Enfield, D.B., Newman, J.H.,  
1178 1999. An ~15,000-year record of El Niño-driven alleviation in southwestern Ecuador.  
1179 *Science*, 283, 516-519.  
1180  
1181 Schmitt, J., Schneider, R., Elsig, J., Leuenberger, D., Laurantou, A., Chappellaz, J.r.m.,  
1182 Kahler, P., Joos, F., Stocker, T.F., Leuenberger, M., and Fischer, H., 2012. Carbon isotope  
1183 constraints on the deglacial CO<sub>2</sub> rise from ice cores. *Science*, 336, 711-714.  
1184  
1185 Shulmeister, J. and Lees, B.G., 1995. Pollen evidence from tropical Australia for the onset of  
1186 an ENSO-dominated climate at c. 4000 BP. *The Holocene*, 5, 10-18.  
1187  
1188 Shulmeister, J., 1999. Australasian evidence for mid-Holocene climate change implies  
1189 precessional control of Walker Circulation in the Pacific. *Quaternary International*, 57-58, 81-  
1190 91.  
1191  
1192 Sigman, D.M., Hain, M.P., Haug, G.H., 2010. The polar ocean and glacial cycles in  
1193 atmospheric CO<sub>2</sub> concentration. *Nature*, 466, 47-55.  
1194  
1195 Sikes, E.L., Howard, W.R., Samson, C.R., Mahan, T.S., Robertson, L.G., Volkman, J.K.,  
1196 2009. Southern Ocean seasonal temperature and Subtropical Front movement on the South

1197 Tasman Rise in the late Quaternary. *Paleoceanography*, 24, PA2201,  
1198 doi:10.1029/2008PA001659.

1199

1200 Spooner, M.I., Barrows, T.T., De Deckker, P., Paterne, M., 2005. Palaeoceanography of the  
1201 Banda Sea, and Late Pleistocene initiation of the Northwest Monsoon. *Global and Planetary*  
1202 *Change*, 49, 28-46.

1203

1204 Spooner, M.I., De Deckker, P., Barrows, T.T., Liffeld, L.K., 2011. The behaviour of the  
1205 Leeuwin Current offshore NW Australia during the last five glacial-interglacial cycles.  
1206 *Global and Planetary Change*, 75, 119-132.

1207

1208 Steinhilber, F., Beer, J., Fröhlich, C., 2009. Total solar irradiance during the Holocene,  
1209 *Geophysical Research Letters*, 36, L19704, doi:10.1029/2009GL040142.

1210

1211 Stott, L., Poulsen, C., Lund, S., Thunell, R., 2002. Super ENSO and global climate oscillation  
1212 at millennial time scales. *Science*, 297, 222-226.

1213

1214 Sturman, A., Tapper, N., 1996. *The Weather and Climate of Australia and New Zealand*.  
1215 Oxford University Press, Melbourne.

1216

1217 Tachikawa, K., Vidal, L., Sonzogni, Bard, E., 2009. Glacial/interglacial sea surface  
1218 temperature changes in the southwest Pacific Ocean over the past 360 ka. *Quaternary Science*  
1219 *Reviews*, 28, 1160-1170.

1220

1221 Tibby, J., 2012. The Younger Dryas: Relevant in the Australian region? *Quaternary*  
1222 *International*, 253, 47-54.

1223

1224 Toggweiler, J. R., 1999. Variation of atmospheric CO<sub>2</sub> by ventilation of the ocean's deepest  
1225 water. *Paleoceanography*, 145, 571-588.

1226

1227 Tudhope, A.W., Chilcott, C.P., McCulloch, M.T., Cook, E.R., Chappell, J., Ellam, R.M., Lea,  
1228 D.W., Lough, J.M. and Shimmield, G.B., 2001. Variability in the El Niño-Southern  
1229 Oscillation through the glacial-interglacial cycle. *Science* 291, 1511-1517.

1230

1231 Turney, C.S.M., Kershaw, A.P., Clemens, S.C., Branch, N., Moss, P.T. and Fifield, L.K.  
1232 2004. Millennial and orbital variations of El Niño/Southern Oscillation and high-latitude  
1233 climate in the last glacial period. *Nature*, 428, 306-310.

1234

1235 Turney, C.S.M., Kershaw, A.P., Lowe, J.J., van der Kaars, S., Johnston, R., Rule, S., Moss,  
1236 P., Radke, L., Tibby, J., McGlone, M.S., Wilmshurst, J.M., Vandergoes, M.J., Fitzsimons,  
1237 S.J., Bryant, C., James, S., Branch, N.P., Cowley, J., Kalin, R.M., Ogle, N., Jacobsen, G.,  
1238 Fifield, K.L., 2006a. Climatic variability in the southwest Pacific during the Last Termination  
1239 (20–10ka BP). *Quaternary Science Reviews*, 25, 886-903.

1240

1241 Turney, C.S.M., Haberle, S., Fink, D., Kershaw, A.P., Barbetti, M., Barrows, T.T., Black, M.,  
1242 Cohen, T.J., Corrège, T., Hesse, P.P., Hua, Q., Johnston, R., Morgan, V., Moss, P., Nanson,  
1243 G., van Ommen, T., Rule, S., Williams, N.J., Zhao, J.X., D'Costa, D., Feng, Y.X., Gagan, M.,  
1244 Mooney, S., Xia, Q., 2006b. Integration of ice-core, marine and terrestrial records for the

1245 Australian Last Glacial Maximum and Termination: a contribution from the OZ INTIMATE  
1246 group. *Journal of Quaternary Science* 21, 751-761.

1247

1248 Vandergoes, M.J., Newnham, R.M., Preusser, F., Hendy, C.H., Lovll, T.V., Fitzsimons, S.J.,  
1249 Hogg, A.G., Kasper, H.U., Schlüter, C., 2005. Regional insolation forcing of late Quaternary  
1250 climate change in the Southern Hemisphere. *Nature*, 436, 242-245.

1251

1252 Vandergoes, M.J., Hogg, A.G., Lowe, D.J., Newnham, R.M., Denton, G.H., Southon, J.,  
1253 Barrell, D.J.A, Wilson, C.J.N., McGlone, M.S., Allan, A.S.R., Almond, P.C., Petchey, F.,  
1254 Dabell, K., Dieffenbacher-Krall, A.C., Blaauw, M., this volume. A revised age for the  
1255 Kawakawa/Oruanui tephra, a key marker for the Last Glacial Maximum in New Zealand.  
1256 *Quaternary Science Reviews*, this volume.

1257

1258 van der Kaars, S., De Deckker, P., 2002. A Late Quaternary pollen record from deepsea core  
1259 Fr10/95, GC17 offshore Cape Range Peninsula, northwestern Western Australia. Review of  
1260 *Palaeobotany and Palynology*, 120, 17-39.

1261

1262 van der Kaars, S, Bassinot, F., De Deckker, P. and Guichard, F., 2010. Changes in monsoon  
1263 and ocean circulation and the vegetation cover of southwest Sumatra through the last 83,000  
1264 years: The record from marine core BAR94-42. *Palaeogeography, Palaeoclimatology,*  
1265 *Palaeoecology*, 296, 52-78.

1266

1267 van der Kaars, S., Williams, M.A.J., Bassinot, F., Guichard, F., Moreno, E., Dewilde, F.,  
1268 Cook, E.K., 2012. The influence of the ~73 ka Toba super-eruption on the ecosystems of

1269 northern Sumatra as recorded in marine core BAR94-25. *Quaternary International* 258, 45-  
1270 53.  
1271  
1272 Varma, V., Prange, M., Lamy, F., Merkel, U., Schulz, M., 2010. Solar-forced shifts of the  
1273 Southern Hemisphere Westerlies during the late Holocene. *Climate of the Past*, 6, 369-384.  
1274  
1275 Verdon, D. C. and S. W. Franks (2006), Long-term behaviour of ENSO:  
1276 Interactions with the PDO over the past 400 years inferred from paleoclimate  
1277 records. *Geophysical Research Letters*, 33, L06712, doi:10.1029/2005GL025052.  
1278  
1279 Verdon, D.C., Wyatt, A.M., Kiem, A.S., Franks, S.W., 2004. Multi-decadal variability of  
1280 rainfall and streamflow - Eastern Australia. *Water Resources Research*, 40(10), W10201.  
1281  
1282 Veth, P., M. Smith, Bowler, J., Fitzsimmons, K.E., Williams, A., Hiscock, P., 2009.  
1283 Excavations at Parnkupirti, Lake Gregory, Great Sandy Desert: OSL ages for occupation  
1284 before the Last Glacial Maximum. *Australian Archaeology*, 69, 1-10.  
1285  
1286 Visser, K, Thunell, R., Stott, L., 2003. Magnitude and timing of temperature change in the  
1287 Indo- Pacific warm pool during the deglaciation. *Nature*, 421, 152-155.  
1288  
1289 Wallis, L., 2001. Environmental history of northwest Australia based on phytolith analysis at  
1290 Carpenter's Gap 1. *Quaternary International*, 83-85, 103-117.  
1291  
1292 Weaver, A.J., Saenko, O.A., Clark, P.U., Mitrovica, J.X., 2003. Meltwater pulse 1A from  
1293 Antarctica as a trigger of the Bølling-Allerød warm interval. *Science*, 299, 1709-1713.

1294

1295 Wilkins, D., Gouramanis, C., De Deckker, P., Fifield, L.K., Olley, J, in press. Revised  
1296 Holocene lake levels from Lake Keilambete and Lake Gnotuk, south-western Victoria,  
1297 Australia. *The Holocene*.

1298

1299 Williams, M., Cook, E., van der Kaars, S., Barrows, T., Shulmeister, J., Kershaw, P., 2009.  
1300 Glacial and deglacial climatic patterns in Australia and surrounding regions from 35 000 to  
1301 10 000 years age reconstructed from terrestrial and near-shore proxy data. *Quaternary*  
1302 *Science Reviews*, 28, 2398-2419.

1303

1304 Wyrwoll, K.-H., Miller, G.H., 2001. Initiation of the Australian summer monsoon 14,000  
1305 years ago. *Quaternary International*, 83-85, 119-128.

1306

1307 Yokoyama, Y., Purcell, A., Lambeck, K., Johnston, P., 2001. Shore-line reconstruction  
1308 around Australia during the Last Glacial Maximum and Late Glacial Stage. *Quaternary*  
1309 *International*, 83-85, 9-18.

1310

1311 Zheng, W., Braconnot, P., Guilyardi, E., Merkel, U., Yu, Y., 2008. ENSO at 6ka and 21ka  
1312 from ocean-atmospheric coupled model simulations. *Climate Dynamics*, 30, 745-762.

1313

1314 **Figures**

1315 Figure 1. Map of average modern sea surface temperature and the major oceanographic  
1316 features of the Australian region. The major currents are shown in solid lines: IPWP = Indo-  
1317 Pacific Warm Pool, ITF = Indonesian Throughflow, SEC = South Equatorial Current, LC =  
1318 Leeuwin Current (with dashed line showing extent during La Niña), EAC = East Australian  
1319 Current, ACC = Antarctic Circumpolar Current. The mean positions of the major fronts are  
1320 shown in dotted lines: TF = Tasman Front, STF = Subtropical Front, SAF = Subantarctic  
1321 Front, PF = Polar Front.

1322

1323 Figure 2. Map showing the major features of the modern climate system of the Australian  
1324 region. ITCZ = Inter-tropical Convergence Zone, ENSO = El Niño Southern Oscillation, IOD  
1325 = Indian Ocean Dipole, SAM = Southern Annular Mode.

1326

1327 Figure 3. Map of the greater Australian region, with site localities (white dots) considered  
1328 here in this study. The shadings are a stylistic representations of the four climate regions  
1329 referred to as tropics (yellow terrestrial, purple marine), arid interior (orange), temperate  
1330 (green terrestrial, royal blue offshore) and Southern Ocean (lower rectangle). The reader is  
1331 directed to the other papers within this volume, which deal with each of these regions  
1332 independently, for more information on the detailed boundaries (i.e. Reeves et al.,  
1333 Fitzsimmons et al., Petherick et al., Bostock et al.). The inset graphs show summer and winter  
1334 insolation at a) 20°S and b) 40°S, respectively.

1335

1336

1337 Figure 4. Schematic representation in the changes in temperature (i) and effective  
1338 precipitation (ii) through time. Change here is considered as a trend in relation to the previous  
1339 time slice, that is; are conditions hotter or wetter, for example, than the previous period. For  
1340 temperature, red = hot, blue = cold, purple = no change. For precipitation, green = wet,  
1341 orange = dry, yellow = no change. Grey indicates no data are available. Please refer to figure  
1342 2 for the division of the climate zones represented here by boxes, with reference to latitude  
1343 and longitude. The time slices are a = c. 32 ka, b = c. 25 ka, c = c. 20 ka, d = c. 17 ka, e = c.  
1344 14 ka, f = c. 9 ka, g = c. 5 ka, h = c. 2 ka  
1345



Figure  
[Click here to download high resolution image](#)




Figure 2  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)




Figure 4ai  
[Click here to download high resolution image](#)




Figure 4aii  
[Click here to download high resolution image](#)




Figure 4bi  
[Click here to download high resolution image](#)






Figure 4bii  
[Click here to download high resolution image](#)




Figure 4ci  
[Click here to download high resolution image](#)






Figure 4cii  
[Click here to download high resolution image](#)




Figure 4di  
[Click here to download high resolution image](#)




Figure 4dii  
[Click here to download high resolution image](#)




Figure 4ei  
[Click here to download high resolution image](#)




Figure 4eii  
[Click here to download high resolution image](#)




Figure 4fi  
[Click here to download high resolution image](#)




Figure 4fii  
[Click here to download high resolution image](#)




Figure 4gi  
[Click here to download high resolution image](#)






Figure 4gii  
[Click here to download high resolution image](#)




Figure 4hi  
[Click here to download high resolution image](#)




Figure 4hii  
[Click here to download high resolution image](#)







Figure 4 (layout)




a i




a ii




b i




b ii




c i




c ii




d i




d ii




e i




e ii




f i




f ii




g i



g ii



h i



h ii

Annotated manuscript

[Click here to download Supplementary Data: Reeves et al text\\_231212.docx](#)