

ORIGINAL PAPER

---

## QUALITY LOSSES IN TEMPORARY SUNFLOWER SEED STORES AND INFLUENCES OF STORAGE CONDITIONS ON QUALITY LOSSES DURING STORAGE

Can Burak SISMAN

Trakya Univ. Tekirdag Agricultural Faculty, Farm Structure and Irrigation Dept.59030 Tekirdag/TURKEY.  
Tel: +90 282 2931442, Fax: +90 282 2931454, e-mail: cbsisman@tu.tzf.edu.tr.

Manuscript received: January 18, 2005; Reviewed: February 18, 2005; Accepted for publication: April 27, 2005

### ABSTRACT

The aim of storage is to preserve properties of products and their freshness. If suitable storage conditions are not supplied according to product variety, quality and quantity losses increase. Decreasing these losses is possible with providing suitable storage conditions and storage management.

In this study were aimed at determining of storage losses in the temporary sunflower seed stores and investigating the influences of storage condition on quality losses of sunflower during storage.

**KEY WORDS:** Storage, sunflower, temporary storage, storage conditions, quality losses

## INTRODUCTION

Because of the nourishment necessity of population in the world, sources in the world must be used sustainably. Recently, increased farm production based on obtaining more product from a unit area. However processing and evaluating of these products are more important. Majority of farm products are consumed after the process. Hence, storage of products after the process until marketing is a necessity. The aim of the storage is to preserve properties of products and their freshness until marketing or consuming. If suitable storage conditions are not supplied according to product variety, quality and quantity losses increase.

Majority of oil, which is one of the most important food for human because of high energy, are vegetable origin (86%). Sunflower in the oil plants leads due to 25-50 % oil content. Approximately 12.6 % of vegetable oil production in the world is covered by sunflower [1]

Turkey is the 9<sup>th</sup> largest sunflower producer in the world with the average of 843 000 t from 578 000 ha area. An important part of sunflower production (60 %) in Turkey (500 000 t) is produced in the Thrace Region [2]. Majority of sunflower produced in the Thrace Region (70%) are stored in temporary stores. Because these stores are very sensitive to weather condition, losses during storage are economically very important. Using temporary stores widespreadly in this region leads to losses of approximately 10 million US \$ per year [3].

The objective of this study is to determine quality losses in temporary sunflower seed storage in Thrace Region, to identify the reason effective for the quality losses during storages.

## MATERIAL AND METHODS

### Material

The study was carried out in one of the stores of Thrace Union, which is the state oil production company in Tekirdag, Yagcı Village temporary store and a model store built specifically for this research in the Agricultural Faculty's area. Sunflower seeds were collected from these stores every month regularly during nine-month storage period started from September 2001 and all analysis were done on these samples.

Temporary sunflower store in Yagcı village was settled up in East-West direction and this store was 10 m in width and 3.6 m in height. During its construction, first soil was compressed, drainage ditches were opened at side of the store and a nylon canvas was spread over the soil. 1.3 m height store walls were formed by stacking

sacks full of sunflower one on the top of the other. Sunflower seeds were heaped between the walls and covered with canvas and then it was tied firmly. Sunflower seeds harvested in 2001 having 6 % moisture content and 3 % foreign material were placed in this store.

A model store having an aeration system and 2 m<sup>3</sup> capacity was built and the quality losses in this store were compared to the losses occurred in the temporary store in the region. The floor of the model store was concrete, walls were bricks and plaster. The aeration in the store was done using two duct systems on the floor having 1.5x 0.1 m cross sectional area for 2 m<sup>3</sup> store capacity as suggested by Hellevang [4] and Proctor [5]. Air flow was supplied with a fan having 0.3 m<sup>3</sup>/min capacity and 0.2 m in diameter for 2 m<sup>3</sup> store capacity as proposed by Hall [6], Hellevang [4], Cloud & Morey [7], Bloome et al. [8], Hofman & Hellevang [9], Harner et al. [10]. The size of the open area in the roof was 0.75 m<sup>2</sup> (cross sectional area) for 2 m<sup>3</sup> store capacity as recommended by Hellevang [4], Proctor [5] and Bloome et al. [8]. Sunflower seeds having 6 % moisture content and 3 % foreign material was placed in this store.

### Methods

Mass temperature and relative humidity into the stores as storage conditions and moisture content due to its effect on quality criteria, oil content and free fatty acidity as sunflower quality criteria were determined on the sample collected from each stores during the storage.

Measurements of mass temperature and relative humidity in the stores were recorded using a digital humidity/temperaturemeter recorder reweekly as recommended by Harrier [11], Thompson & Shelton [12] and Noyes et al. [13]. Temperature and humidity measurements were done at eight different points in each store.

Weather temperature and relative humidity were measured by a thermohydrograph and these records regulated when to operate the aeration system in the model store. Aeration was operated as soon as the air temperature was 8°C cooler than that of the mass and was continued until the temperatures became equal. Aeration system was also shut and covered when the relative humidity exceeded 75 % and during the rainy period [11] [4] [7].

Sunflower samples were taken every month regularly in the stores to determine the losses occurred in the stores during storage period. The samples were taken in eight different points of the mass according to Turkish Standards no 163 related to Taking Sample from Oil Seed (Turkish Standard Institute, 1980) [14]. Partitioned hand probe was used for taking samples.

Analyzes of moisture content, oil content and free fatty acidity in the laboratory were done on the samples. Moisture contents were determined on basis of dry weight [15]. To determine the oil contents Soxhlet method, which adjust oil content to a constant moisture, (IUPAC method no: 1.122) was used [16]. Free fatty acidity was determined using titratable acidity in IUPAC method no: 2.201 [16].

The results obtained were evaluated using SPSS computer programs for statistical analysis.

## RESULTS AND DISCUSSIONS

### Storage Conditions

Changes in the average temperatures and humidities during the storage period in the temporary and model stores were presented in Figure 1 and 2.

As seen in Figure 1, in general, mass temperature in the stores varied parallel to changing weather temperature. Mass temperature decreased in the autumn and increased in the spring. In the autumn, mass temperature in the model store was lower than temporary store because of aeration. Temperature in the stores decreased to 0 °C during December and January. In the spring, mass temperature in the stores began to increase apart from increasing the weather temperature and especially in the temporary store it quickly increased beginning from March because of increasing the outside temperature, crop respiration and insect's activity. In this period, increasing temperature in the model store was less than temporary store due to operate the aeration system. Generally mass temperature in the model store was maintained below 17°C as recommended by [4], Cloud & Morey [7], Brooker et al. [17] and Navarro [18] for optimum storage in terms of insect control as a result of aerating 220 h during autumn and 125 h during spring. Owing to aeration (total 345 h), temperature differences occurred between the zones in the store were prevented and homogenous temperature distribution in the model store was provided according to the measurements. Therefore moisture migration in the store in a great extent was prevented.

As seen in Figure 2, in general, while mass humidity in the model stores varied parallel to changing weather humidity, mass humidity in the temporary store increased especially during autumn. The reason of this increase in the humidity of the temporary store was the decreasing weather temperature and rains in this period. Mass humidity in the spring decreased slightly in windy days and low relative humidity days, but it rose again because of crop respiration.

The most suitable storage conditions were recorded in the model store with 16°C temperature and 78 % mass humidity and this value were around the recommended value for suitable sunflower seed storage conditions in terms of insect control [11; 19; 4, 7, 17; 9; 20; 21]. While temperature differences and moisture migration between the zones in the model store was prevented through aeration, important temperature differences between the zones in the temporary stores were determined and moisture migration occurred. This was because the temporary store was affected by the weather temperature changes more.

### Sunflower Quality Criteria

#### Moisture Content

Moisture content is one of the most important criteria effective on the losses during the storage. Therefore, moisture contents of sunflower samples taken regularly from the stores were determined and changes in the moisture contents with time during the storage were presented in Figure 3.

As shown in Figure 3, while the moisture content in the temporary store increased during initial storage months, it did not change in the model store because of aeration. This increase in the temporary store resulted from entering rain water in this months and the moisture migration which arose from temperature differences between the zones. Afterwards, moisture content in this store decreased with warming up the outside temperature, but moisture content increased again because of rainy periods, high relative humidity and increasing the crop respiration and mass temperature. Although moisture content decreased minimally in windy and cloudless days, increasing outside temperature, crop respiration and insect's activity caused to increase in the moisture content. In the model store, increase in the moisture content was prevented by means of aeration, but when the aeration stopped in the winter, it increased. When the moisture content in the temporary store investigated during storage period, it is seen that the moisture content in this store exceeded 8% suggested level for safely storage by Harrier [11], Patterson [19], Brooker et al. [17], Anonymous [22], Hofman & Hellevang [89], Hellevang [20] and Hellevang [21] in the beginning from second month of the storage and it remained over 8% until end of the storage.

#### Oil Content

Changes in the oil contents, which is the most important feature of sunflowers, was determined by analyzes that was regularly done on the samples taken from stores and changes in the oil contents with time were presented in Figure 4.




Figure 1. Changes in the average temperatures during the storage period in the stores




Figure 2. Changes in the average humidity during the storage period in the stores




Figure 3. Changes in the average moisture content during the storage period in the stores




Figure 4. Changes in the average oil content during the storage period in the stores




Figure 5. Changes in the free fatty acid during the storage period in the stores

As shown in Figure 4, oil contents of both temporary and model store decrease continuously however the magnitude of decrease in the temporary store was higher, from about 43.2 % to 36.5 % corresponding 0.3 to 0.7 % each month against only 2.9 % decrease, 0.1 to 0.3 % each month in the model store. A rapid decrease in the oil content of temporary store during the last three months when compared to the previous months and a relatively more decrease in the temporary store in comparison to the model store was because of keeping low temperature and humidity by means of aeration and moisture content which was lower than 8 %, suggested for a safe storage [11; 4; 17; 5; 21]. Decreasing moisture content in the model store continued to same level during storage and it decreased 0.1-0.3% per month.

According to these results, temporary stores must not be used for long time because of high losses as determined by Hall [6], Patterson [19], Hellevang [4], Jones & Shelton [23], Hellevang [24], Hofman & Hellevang [9], Hellevang [20], Hellevang [21].

### Free Fatty Acid

Free fatty acid which affects losses occurred during the refinement was determined by analysis that was regularly done on the sample taken from the stores and changes in the free fatty acid with time were presented in Figure 5.

Free fatty acid of sunflowers in both of the stores continuously increased during the storage. While free fatty acid in the model stores did not exceed 1.2-1.5 % suggested for suitable free fatty acid by Nas et al. [15], in the temporary store, beginning from February, acidity in this store quickly increased and it rose 7.683 % in June. In this period, high temperature and moisture content in the store caused to increase in the free fatty acid.

As seen in Figure 5, free fatty acid in the model store increased 0.05 to 0.06 % per month during the first five months of storage and then the monthly increase in the free fatty acid was recorded as 0.2% because of the increase in the mass temperature and moisture content in

this period. The decreases in the mass temperature and moisture content with aeration limited the increase the acidity in this period in some extends.

### 3.3. Effects of Storage Conditions on the Quality Losses

Statistical evaluation done to determine the relationship between the sunflower quality losses and storage conditions and correlation coefficients for temporary store and model store were given Table 1 and Table 2 respectively.

As seen in Table 3.1, influences of storage conditions on sunflower quality losses were significant at the 0.01 confidence level. While temperature and humidity of the mass affected adversely, moisture content and oil content of the sunflower affected free fatty acidity positively. A strong negative correlation between oil content and free fatty acidity was determined.

When correlation coefficients for model store were investigated, in this store a negative correlation between the oil content and only humidity whereas a positive relation between free fatty acidity and humidity were determined.

When the stores were compared, negative influences of mass temperature and moisture content on oil content and free fatty acidity disappeared, influence of humidity on these quality criteria was reduced approximately 50 % due to aeration. This proves the importance of aeration.

## CONCLUSION AND RECOMMENDATION

In this study conducted in the Thrace Region, influences of the storage conditions of temporary stores widely used for storing sunflower on the quality losses occurred during storage was investigated. For this aim, in the selected temporary store and a model store built specifically for this research, mass temperature and humidity as storage conditions and moisture content, oil content and free fatty acidity as sunflower quality criteria were determined and evaluated during the storage.

In the temporary store, mass temperature decreased and mass humidity increased during autumn when the outside temperature decreased, because the store was quickly affected by the outside conditions. In this period, while the side of the store on the blowing direction of wind got quickly cold, the other side and the middle of the store got colder by time. Therefore, significant temperature differences in the store occurred and lead to increase the magnitude of moisture migration. In this period in which mass was generally cold and the mass temperature of 35°C at the beginning decreased to 2.29°C in January and mass humidity of 74 % increased 98 % in January. Mass temperature fell below the suitable storage temperature

of 17°C given by Cloud & Morey [7], Thompson & Shelton [12], Jones & Shelton [23], Hellevang [24] and Hellevang [21] after November. From the early January, mass temperature increased until end of storage and it exceeded suitable storage temperature after March. In the same period, although mass humidity decreased slightly with increasing the temperature sometimes, generally it remained at the level of 97 % because of crop respiration and raining.

The criteria for sunflower in the temporary store were negatively affected by weather condition more and rain water entrance into the store. Moisture content of the crop in this store exceeded 8 % which was recommended level for a safe storage by Harrier [11], Patterson [19], Brooker et al. [17], Anonymous [22], Hofman & Hellevang [9], Hellevang [20] and Hellevang [21] and the highest moisture content occurred in January and April. This was because of warming weather conditions and rain. In April, sunflower in the upper part of the mass germinated owing to increase in the moisture content and high temperature. The result of unfavorable storage conditions in the store, oil content and free fatty acidity of crop were negatively affected. While oil content and free fatty acidity of crop did not change significantly during the Autumn, in the Spring, oil content quickly decreased unlike acidity. At the end of the storage, oil content of the crop decreased by 6.60 %, acidity of crop increased 7.33 %.

Temporary storage widely used in the Thrace Region caused losses in significant amount. The raw oil losses of crops stored in the temporary stores in the Thrace region are around 19 500 tone per year and this corresponds 16 million US \$ according to the price of 2002. Therefore temporary stores should not be used for sunflower storage, but if used, the durations should be less than five months in Autumn when the weather temperature decreases continuously. This kind of stores should not be also used in the districts having much rain.

Mass temperature of 32°C in the model store at the beginning decreased below 17°C that is recommended for safely storage when the aeration began from the end of October and remained below this level until June [7; 12; 23; 24; 21]. Mass humidity in this store increased 81 % from December to March because of not operating the aeration in this period, but it again decreased with aeration. Moisture content of crop during the storage period except for February kept under 8 % suggested for safely storage by Harrier [11], Patterson [19], Brooker et al. [17], Hofman & Hellevang [9], Hellevang [21]. Supplying suitable storage conditions and recommended moisture content in this store lowered the losses. At the

Table 1. Corelation coefficients of covariance analiysis for temporary store

|  | Oil Content | Moisture Content | Free Faty Acidity | Temperature | Humidity |
|--|-------------|------------------|-------------------|-------------|----------|
| Oil Content<br>(Pearson Correlation)       | 1,000       |                  |                   |             |          |
| Moisture Content<br>(Pearson Correlation)  | -0,411**    | 1,000            |                   |             |          |
| Free Faty Acidity<br>(Pearson Correlation) | -0,871**    | 0,218*           | 1,000             |             |          |
| Temperature<br>(Pearson Correlation)       | -0,343**    | -0,417**         | 0,578**           | 1,000       |          |
| Humidity<br>(Pearson Correlation)          | -0,583**    | 0,620**          | 0,421**           | -0,384**    | 1,000    |

\*\* Corelation is significant at the 0,01 confidance level

\* Corelation is significant at the 0,05 confidance level

Table 2. Corelation coefficients of covariance analiysis for model store

|  | Oil Content | Moisture Content | Free Faty Acidity | Temperature | Humidity |
|--|-------------|------------------|-------------------|-------------|----------|
| Oil Content<br>(Pearson Correlation)       | 1,000       |                  |                   |             |          |
| Moisture Content<br>(Pearson Correlation)  | -0,168      | 1,000            |                   |             |          |
| Free Faty Acidity<br>(Pearson Correlation) | -0,921**    | 0,149            | 1,000             |             |          |
| Temperature<br>(Pearson Correlation)       | 0,059       | -0,611**         | -0,001            | 1,000       |          |
| Humidity<br>(Pearson Correlation)          | -0,278**    | 0,630**          | 0,281**           | -0,675**    | 1,000    |

\*\* Corelation is significant at the 0,01 confidance level

end of the storage, oil losses occurred 2.8 % and free fatty acidity increased 1.44 %. Losses in the model store appeared quite low level compairing with temporary store and losses reduced approximately 70 %. Therefore using this store for sunflower storage will provide important economical benefit.

## REFERENCE

- [1] Anoglu H. Cultivation of Oil Seed and Improvement. Cukurova Üniv. Agricultural Fac. Pub. Number 220, 190 p. Adana. 1999.
- [2] Anonymous. Agricultural Structure (Production, Price, Value) 2000. Republic Of Turkey, Government Statistical Enstitute. Publication Nuber 2547, Ankara. (2002)
- [3] Gaytancıoğlu, O. Farming Policies on Sunflower and Economical Analıysis of Storage Losses. T.U. Tekirdag Agricultural Fac, Unpublished, Tekirdag. (1999)
- [4] Hellevang K.J. Crop Storage Management. NDSU. Extension Service, ND 58105-AE-791, North Dakota, USA. (1990).
- [5] Proctor D.L. Grain Storage Techniques Evolution and Trends in Developing Countries. FAO Agricultural Service Bulletin No 109, ISBN 92-5-103456-7, Roma. (1994).
- [6] Hall C.W. Drying and Storage of Agricultural Crops. The AVI Publishing Company Inc. ISBN 0-87055-364-X, USA. 1980.
- [7] Cloud H.A., Morey R.V. Management of Stored Grain with Aeration, University of Minnesota. Collage of Agricultural Food and Enviromental Science, FO 1327-GO, USA. (1991).
- [8] Bloome P., Brusewitz G., Harp S. Aeration and Cooling of Stored Grain. Oklahoma State Univ. Cooperative Extension Service, Division of Agricultural

**QUALITY LOSSES IN TEMPORARY SUNFLOWER SEED STORES AND INFLUENCES OF STORAGE CONDITIONS ON  
QUALITY LOSSES DURING STORAGE**

Science and Natural Resource, F-1101, Oklahoma. (1995).

[9] Hofman V.L., Hellewang K.J. Harvesting Drying and Storage of Sunflower. Sunflower Technology and Production, American Society of Agronomy, Soil Science Society of American, Agronomy Monograph No 35, Madison, USA. (1997).

[10] Harner, J.P., Herrman T.J., Reed C. Temporary Grain Storage Considerations. Kansas State Univ. Agricultural Experiment Station and Cooperative Extension Service, MF 2362, Kansas. (1998).

[11] Harrier J.P. Drying and Storing Sunflowers. Kansas State Univ. Cooperative Extension Service, Ag. Facts 158, Kansas. (1987).

[12] Thompson T.L., Shelton D.P. Aeration of Stored Grain. Nort Dakota State Univ. Extension Service, G84-692, USA. (1993).

[13] Noyes R.T., Clary B.L., Cuperus G.W. Maintaining Quality of Stored Grain by Aeration. Oklahoma State Univ. Extension Service Division of Agricultural Science and Natural Resources, USA. (1998).

[14] Ekmeklier H.Y., Geçit H.H. Standardization and Storage of Field. Ankara Üniv. Agricultural Fac Pub. No: 989, Ankara. 1986

[15] Nas S., Gökalp H.Y., Ünsal M. Bitkisel Yağ Teknolojisi. Pamukkale Üniv. Mühendislik Fak. Ders Kitapları Yayın No 005, 329s, Denizli. 1998.

[16] Anonymous. International Union for Pure and Applied Chemistry, Standard Methods of Analysis for the

Analysis of Oil. Fats and Derivatives, 7<sup>th</sup>. End., Blacwell, Jevent Publishers, Oxford. 1987

[17] Brooker D.B., Arkema F.B., Hall C.W. Drying and Storage of Grains and Oilseeds. An AVI Book, Published by Van Nostrand Reinhold, ISBN 0-442-20515-5, New York. 1992.

[18] Navarro S. Aeration and Cooling for Control of Stored Grain Insect. International Course on Agricultural Engineering Technology of Grain Storage, The Volcani Center, Israel. (1996).

[19] Patterson H.B.W. Handling and Storage of Oilseed, Oils, Fats and Meal. Elsevier Applied Science, 294 p., New York. 1989.

[20] Hellevang K. Temporary Storage of Sunflower Seed. The Sunflower Magazine, September, National Sunflower Association. (1998).

[21] Hellevang K. Storing Wet Sunflower. Irrigator's Workshops, Bismarck Radisson Inn. 4 December 2000. (2000).

[22] Anonymous. Drying and Storage Sunflower Production. NDSU Extension Service, Agricultural Experiment Station, EB 25, USA. 1995.

[23] Jones D., Shelton P. Management to Maintain Stored Grain Quality. Nebraska State Univ. Cooperative Extension Service, Institute of Agriculture and Natural Researces, G 94-1199-A, USA. (1994).

[24] Hellevang K.J. Grain Moisture Content Effects and Management. NDSU. Extension Service AE-905, North Dakota. (1995).

