



original scientific paper / izvorni znanstveni rad

FAUNA OF THE ADRIATIC DECAPOD CRUSTACEANS (CRUSTACEA: DECAPODA) – STATUS AND OUTLOOK

MARIN KIRINČIĆ¹ & ZDRAVKO ŠTEVČIĆ²

¹Natural History Museum Rijeka, Lorenzov prolaz 1, HR-51000 Rijeka, Croatia
(marin@prirodoslovni.com)

²Lacosercio 19, HR-52210 Rovinj, Croatia (stevcic@cim.irb.hr)

Kirinčić, M. & Števcic, Z.: Fauna of the Adriatic decapod crustaceans (Crustacea: Decapoda) – status and outlook. Nat. Croat., Vol. 17, No. 2., 131–139, 2008, Zagreb.

The checklist of the Adriatic decapod fauna is re-examined and supplemented. Several new species for the area are included in the list, new immigrants are noted and some species excluded. The species names are updated and their status and prospects commented on.

Key words: decapod crustaceans, Adriatic Sea, new records, revision

Kirinčić, M. & Števcic, Z.: Fauna jadranskih desetonožnih rakova (Crustacea: Decapoda) – stanje i perspektive. Nat. Croat., Vol. 17, No. 2., 131–139, 2008, Zagreb.

Rad donosi pregled i nadopunu popisa vrsta jadranskih desetonožnih rakova. U popis je dodano nekoliko novih vrsta za ovo područje, zabilježene su imigrantske vrste, a neke vrste su izbačene. Revidirana su imena vrsta te se raspravlja o njihovom stanju i perspektivama.

Ključne riječi: desetonožni raci, Jadran, novi nalazi, revizija

Adriatic decapod crustaceans (Crustacea: Decapoda) have been the subject of numerous investigations, starting from the 16th century (ŠTEVČIĆ, 1993). Papers dealing with this issue are numerous and new ones appear nearly every year. Such intensive surveys of the Adriatic Sea result in a constantly increasing number of decapod species that have been noted for this region. The last complete checklist of the Adriatic decapods appeared in 1990 (ŠTEVČIĆ, 1990) and has thereafter been updated twice (ŠTEVČIĆ, 1995; 2002). In the meantime, FROGLIA (1995) published a checklist of Italian Malacostraca including the Adriatic decapods. In the last several years decapodologists from the Northern and Southern Adriatic Sea have been very active and with other researchers on the Adriatic Sea contributed to the increasing number of known Adriatic decapod species (MIZZAN, 1993, 1995; MIZZAN & ZANELLA,

1996; UNGARO, 2000; UNGARO & MARANO, 2002; VUKANIĆ, 2003; UNGARO *et al.*, 2005; KASALICA & JOKSIMOVIĆ, 2005; KASALICA, 2006; KASALICA *et al.*, 2006). New records following the latest supplements to the Adriatic decapod fauna (ŠTEVČIĆ, 2002) are presented in Tab. 1.

All species mentioned in Tab. 1 are relatively well known from the Mediterranean Sea and Eastern Atlantic Ocean. *Munida perarmata* A. Milne-Edwards & Bouvier, 1894 mentioned by UNGARO *et al.* (2005), however needs further confirmation in order to be considered an Adriatic species because its presence in the area is rather questionable (D'UDEKEM D'ACCOZ, 1999: 164).

Several species have been reported for the Adriatic Sea the occurrence of which is highly questionable or even impossible: *Platymaia wyvillethomsoni* Miers, 1886, *Portunus sanguinolentus* (HERBST, 1793), *Thenus orientalis* (LUND, 1793) and *Uca coarctata* (H. MILNE EDWARDS, 1852). These species were either sampled only once, probably transported by ship ballast waters or mistakenly identified (PESTA, 1918). Two more species should also be mentioned here, *Portunus hastatus* (LINNAEUS, 1767) and *Atelecyclus undecimdentatus* (HERBST, 1783) which were noted once long time ago and whose presence has not been confirmed by subsequent explorations. *Scyllarus*

Tab. 1. New records for the Adriatic Sea

PANDALIDAE
<i>Plesionika gigliolii</i> (Senna, 1902) (cf. UNGARO <i>et al.</i> , 2005)
HIPPOLYTIDAE
<i>Hippolyte prideauxiana</i> Leach, 1817 (cf. KIRINČIĆ, 2006)
<i>Hippolyte varians</i> Leach 1814 (cf. D'UDEKEM D'ACCOZ, 1996)
PALAEONIDAE
<i>Brachycarpus biunguiculatus</i> (Lucas, 1846) (cf. KIRINČIĆ, 2003)
SCYLLARIDAE
<i>Scyllarus caparti</i> Holthuis, 1952 (cf. FROGLIA, 1995)
CALLIANASSIDAE
<i>Callianassa truncata</i> Giard & Bonier, 1890 (cf. ABED-NAVANDI & DWORSCHAK, 1997)
PAGURIDAE
<i>Pagurus chevreuxi</i> (Bouvier, 1896) (cf. ARKO <i>et al.</i> , 2001)
GALATHEIDAE
<i>Munida perarmata</i> A. Milne-Edwards & Bouvier, 1894 (cf. UNGARO <i>et al.</i> , 2005)
<i>Munida ruillanti</i> Zariquiey Alvarez, 1952 (cf. UNGARO <i>et al.</i> , 2005)
CALAPPIDAE
<i>Calappa tuerkayana</i> Pastore, 1995 (cf. UNGARO <i>et al.</i> , 2005)
PINNOTHERIDAE
<i>Pinnotheres marioni</i> Gourret, 1887 (cf. FROGLIA, 1995)
GRAPSIDAE
<i>Hemigrapsus sanguineus</i> (de Haan, 1835) (cf. SCHUBART, 2003)
<i>Eriocheir sinensis</i> H. Milne Edwards, 1854 (cf. MIZZAN, 2005)

caparti Holthuis, 1952, which was sampled just once nearly 30 years ago (FROGLIA, 1979) has to be mentioned as yet another rather problematic species. The same can be said for *Pinnotheres marioni* Gourret, 1887, which is also a doubtful species (D'UDEKEM D'ACCOZ, 1999: 244).

It is worth noting that the presence of *Maja goitziana* d'Oliveira, 1888 has been confirmed just recently. ŠTEVČIĆ (1990: 255) expressed some doubt about its occurrence in the Adriatic Sea due to their rarity and the very few records so far (PALLAORO & DULČIĆ, 2004). Furthermore, *Paragalene longicrura* (Nardo, 1868) has also been found recently, a long time after its description [PESTA (1918: 431–433) reported this species for the first time], in its »locus typicus« – i.e. Middle Dalmatia (PALLAORO, 2005).

During the last fifteen years several species from Asian and North American waters have been reported from the Mediterranean (GALIL *et al.*, 2002). Eight such species were reported for the Adriatic Sea (Tab. 2). The mode of introduction for these immigrant species is probably by ship ballast waters, except for *Marsupenaeus japonicus* (BATE, 1888), which most likely escaped from mariculture facilities, and *Scyllarus caparti* Holthuis, 1952, probably aquarium-released. Among exotic species two types have to be distinguished: alien species and established ones. Established species have self maintaining populations as opposed to alien species who do not have such populations and are recorded only once. All species in Tab. 2 except *Dyspanopeus sayi* (SMITH, 1869), *Rhithropanopeus harrisi* (GOULD, 1841) and *Callinectes sapidus* Rathbun, 1869 can be considered alien because they are just occasionally found in the area. Up to now, no Lessepsian migrant has been reported from the area.

The Adriatic decapod fauna shows a high diversity. The current information regarding families and their pertaining numbers of genera and species is presented in Tab. 3.

So far, 241 decapod species have been noted for the Adriatic Sea (KIRINČIĆ & ŠTEVČIĆ, in prep). If this is compared to PESTA (1918) who noted only 143 species for the same area or D'UDEKEM D'ACCOZ (1999) who reported 340 species for the entire Mediterranean Sea, we can conclude that the Adriatic Sea seems to be qualitatively very well explored so far. The level of diversity in the Adriatic Sea according to the data in Tab. 3 is as follows: 4.55 species per family, 2.01 species per genus and 2.26 genera per family.

Tab. 2. Exotic species recorded at least once in the Adriatic Sea

<i>Marsupenaeus japonicus</i> (Bate, 1888)
<i>Scyllarus caparti</i> Holthuis, 1952
<i>Dyspanopeus sayi</i> (Smith, 1869)
<i>Rhithropanopeus harrisi</i> (Gould, 1841)
<i>Callinectes danae</i> Smith, 1869
<i>Callinectes sapidus</i> Rathbun, 1869
<i>Hemigrapsus sanguineus</i> (de Haan, 1835)
<i>Eriocheir sinensis</i> H. Milne Edwards, 1854

Tab. 3. Adriatic decapod fauna: number of genera and species

FAMILIES	NO. OF GENERA	NO. OF SPECIES
ARISTEIDAE	3	3
PENAEIDAE	3	3
SOLENOCERIDAE	1	1
SICYONIDAE	1	1
SERGESTIDAE	2	5
LUCIFERIDAE	1	1
PASIPHAEIDAE	1	2
OPLOPHORIDAE	1	1
PANDALIDAE	3	10
ALPHEIDAE	5	9
HIPPOLYTIDAE	5	14
PALAEMONIDAE	6	13
PROCESSIDAE	1	7
GNATHOPHYLLIDAE	1	1
CRANGONIDAE	4	11
STENOPODIDAE	1	1
POLYCHELIDAE	1	1
PALINURIDAE	1	1
SCYLLARIDAE	2	4
NEPHROPIDAE	2	2
AXIIDAE	1	1
CALOCARIDIDAE	1	1
LAOMEDIIDAE	1	1
UPOGEBIIDAE	1	4
CALLIANASSIDAE	3	7
CTENOCHELIDAE	1	1
DIOGENIDAE	5	7
PAGURIDAE	3	15
GALATHEIDAE	2	12
PORCELLANIDAE	2	4
HOMOLIDAE	2	2
LATREILLIIDAE	1	1
DROMIIDAE	1	1
CANCRIDAE	1	1
ATELECYCLIDAE	1	2
CORYSTIDAE	1	1
THIIDAE	1	1
PIRIMELIDAE	2	2
ERIPHIIDAE	1	1
XANTHIDAE	3	6

PANOPEIDAE	2	2
CALAPPIDAE	1	2
PILUMNIDAE	1	4
GERYONIDAE	2	2
GONEPLACIDAE	1	1
PARTHENOPIIDAE	2	4
PORTUNIDAE	7	16
MAJIDAE	12	28
DORIPPIDAE	2	2
PALICIDAE	1	1
LEUCOSIIDAE	3	7
PINNOTHERIDAE	2	3
GRAPSIDAE	5	7
Σ 53	120	241

Because of permanent revision, species names have to be changed and adjusted to the current state of nomenclature of decapod Crustacea. According to that the names of some decapod species listed by ŠTEVČIĆ (1990, 2002) are updated and replaced by the valid names given in Tab. 4.




Fig. 1. Number of references and number of decapod species

It is worthy mentioning that since the mid 18th century carcinologists have been reporting nearly one decapod crustacean species per year (in the Adriatic Sea) and have published a similar number of papers dealing with Adriatic decapods (Fig. 1). In recent years the average number of species and papers has been 3.6 and 2.3, respectively per year from 2001 on. It is very indicative that the number of references and new species for this small area is not decreasing, due to the significant number of carcinologists in the area (Italy, Croatia, Slovenia, Montenegro, Albania, Austria and even Germany) and therefore we expect that prospects for this group of crustaceans are favorable.

All presented facts indicate that the Adriatic decapod fauna is qualitatively very well explored. Therefore, the increasing amount of data on decapoda fauna also provides information on the frequency and distribution of species in the area. The increase in the number of species recorded in the recent past is the result of intensified classical oceanographic research and monitoring studies based on samples utilizing grabs, dredges, and bottom trawls. In addition, it is also the result of an increased interest of carcinologists in *in situ* research and collection while SCUBA

Tab. 4. Updated names of the Adriatic decapod species

PREVIOUSLY USED NAME (Števcic, 1990; 2002)	CURRENT NAMES
<i>Penaeus japonicus</i> (Bate, 1888)	<i>Marsupenaeus japonicus</i> (Bate, 1888)
<i>Penaeus kerathurus</i> Forskål, 1775	<i>Melicertus kerathurus</i> (Forskål, 1775)
<i>Sergestes corniculum</i> Krøyer, 1855	<i>Sergestes arachnipodus</i> (Cocco, 1832)
<i>Sergestes robustus</i> (S.I. Smith, 1882)	<i>Sergia robusta</i> (S. I. Smith, 1882)
<i>Parapandalus narval</i> (Fabricius, 1787)	<i>Plesionika narval</i> (Fabricius, 1787)
<i>Hippolyte longirostris</i> (Czerniavsky, 1868)	<i>Hippolyte varians</i> , Leach 1814
<i>Pontocaris cataphractus</i> (Olivi, 1792)	<i>Aegaeon cataphractus</i> (Olivi, 1792)
<i>Pontocaris lacazei</i> (Gourett, 1887)	<i>Aegaeon lacazei</i> (Gourett, 1887)
<i>Callianassa candida</i> (Olivi, 1792)	<i>Pestarella candida</i> (Olivi, 1792)
<i>Callianassa tyrrhena</i> (Petagna, 1792)	<i>Pestarella tyrrhena</i> (Petagna, 1792)
<i>Callianassa whitei</i> Sakai, 1999	<i>Pestarella whitei</i> (Sakai, 1999)
<i>Anapagurus brevicarpus</i> A. Milne-Edwards & Bouvier, 1892	<i>Anapagurus chiroacanthus</i> (Liljeborg, 1856)
<i>Pagurus sculptimanus</i> Lucas, 1846	<i>Pagurus forbesi</i> Bell, 1845
<i>Xantho incisus</i> Leach, 1814	<i>Xantho hydrophilus</i> (Herbst, 1790)
<i>Liocarcinus arcuatus</i> (Leach, 1814)	<i>Liocarcinus navigator</i> (Herbst, 1794)
<i>Portumnus pestai</i> Forest, 1967	<i>Portumnus lysianassa</i> (Herbst, 1796)
<i>Macropodia longipes</i> (A. Milne-Edwards & Bouvier, 1899)	<i>Macropodia tenuirostris</i> (Leach, 1814)
<i>Pisa corallina</i> (Risso, 1816)	<i>Pisa hirticornis</i> (Herbst, 1804)
<i>Anamathia rissoana</i> (Roux, 1828)	<i>Rochinia rissoana</i> (Roux, 1828)
<i>Pinnotheres pinnotheres</i> (Linnaeus, 1758)	<i>Nepinnotheres pinnotheres</i> (Linnaeus, 1758)

diving. This allows the investigation of previously unexplored microhabitats of the infra-littoral zone such as marine caves and holes, as well as pebbly interstia, etc. We also expect the addition of Lessepsian immigrants, which may expand their distribution due to the increase of maritime transportation and the phenomenon of global warming. A lack of quantitative, zoogeographic and ecological studies forces us to undergo new explorations and to compile already existing together with newly obtained data. The increasing amount of data on decapod fauna will also affect our knowledge on the frequency and distribution of species in the area. There have been insufficient investigations dealing with decapods carried out in the deep sea, river estuaries and underwater springs.

Therefore research into Adriatic decapods will be continue and intensify in a way that gives good prospects of future knowledge about this group.

ACKNOWLEDGMENTS

We wish to express our sincere gratitude for helpful suggestions on the MS to Dr. M. Kovačić, Dr. García Raso and Dr. C. D'Udekem D'Acoz. Many thanks also to Dr. P. C. Dworschak, for sending us papers on the topic, which were most helpful.

Received August 2, 2007

REFERENCES

- ABED-NAVANDI, D. & P. DWORSCHAK, 1997: First record of the thalassinid *Callianassa truncata* Giard & Bonnier, 1890 in the Adriatic Sea (Crustacea: Decapoda: Callianassidae). *Annalen des Naturhistorischen Museums in Wien*, 99B, 565–570
- ARKO-PIJEVAC, M., Č. BENAC, M. KOVAČIĆ & M. KIRINČIĆ, 2001: A submarine cave at the Island of Krk (North Adriatic Sea). *Natura Croatica* 10 (3), 163–184.
- FROGLIA, C., 1979: Segnalazione di alcuni Crostacei Decapodi nuovi per la fauna adriatica. *Quaderni del Laboratorio di Tecnologia della Pesca, Ancona*, 2(4), 191–196.
- FROGLIA, C., 1995: Crustacea Malacostraca III (Decapoda). In: A. MINELLI, S. RUFFO & S. LA POSTA (Eds.). *Checklist delle Specie della Fauna Italiana*, Fascicolo 31, Edizioni Calderini. Bologna. 1–17.
- GALIL, B., C. FROGLIA & P. NOËL, 2002: CIESM atlas of exotic species in the Mediterranean. Vol.2, CIESM Publishers, Monaco. 192.
- KASALICA, O. & A. JOKSIMOVIĆ, 2005: Prostorna distribucija i relativna biomasa kozice, *Parapenaeus longirostris* (Lucas, 1846) na šelfu Crnogorskog primorja. II Međunarodna Konferencija »Ribarstvo«, 10–12. februar 2005. Zemun-Beograd. Zbornik predavanja, 219–224.
- KASALICA, O., 2006: Distribucija Penaeidae u vodama Jadrana sa osvrtom na *Parapenaeus longirostris* (Lucas, 1846). Konferencija »Vode, vodovodi, sanitarne tehnologije« 8–11. maj 2006. Budva. Zbornik radova, 129–132.
- KASALICA, O., S. REGNER & A. JOKSIMOVIĆ, 2006: Estimate of Length-Weight relations of deep-water Pink shrimp in trawler fishing at Montenegrin coast. II International Symposium of ecologist of the Republic of Montenegro, 20–24th September 2006, Kotor. Proceedings of the Symposium, 143–147.

- KIRINČIĆ, M., 2003: *Brachycarpus biunguiculatus* (Lucas, 1846): a new species of shrimp for the Adriatic Sea (Decapoda, Caridea). *Crustaceana*, **76** (7), 891–894.
- KIRINČIĆ, M., 2006: First record of *Hippolyte prideauxiana* Leach, 1817 (Crustacea, Decapoda, Caridea) in the Adriatic Sea. *Acta Adriat.*, **47** (1), 85–88.
- MIZZAN, L., 1993: Presence of swimming crabs of the genus *Callinectes* (Stimpson) (Decapoda, Portunidae) in the Venice lagoon (North Adriatic Sea – Italy). First record of *Callinectes danae* Smith in European waters. *Bull. Mus. civ. nat. Venezia*, **42**: 31–44. (1991).
- MIZZAN, L., 1995: Notes on the presence and diffusion of *Dyspanopeus sayi* (Smith, 1869) (Crustacea, Decapoda, Xanthidae) in the Venetian lagoon. *Bull. Mus. civ. nat. Venezia*, **44**, 121–129. (1993).
- MIZZAN, L. & L. ZANELLA, 1996: First record of *Rhitropanopeus harrisi* (Gould, 1841) (Crustacea, Decapoda, Panopeidae) e *Eriocheir sinensis* H. Milne Edwards, 1854. (Crustacea, Decapoda, Grapsidae): due nuovi granchi esotici in laguna di Venezia. *Bull. Mus. civ. St. nat. Venezia*, **56**, 89–96.
- MIZZAN, L., 2005: *Rhitropanopeus harrisi* (Gould, 1841) (Crustacea, Decapoda, Xanthidae) in the Venetian lagoon. *Bull. Mus. civ. nat. Venezia*, **44**, 121–129. (1993).0
- MÜLLER C. H. G. & C. D. SCHUBART, 2007: Insight into the Crustacea Decapoda of the Adriatic Sea. Observations from four sampling locations along the Croatian coast. *Rostocker Meeresbiologische Beiträge*, **18**, 112–130.
- PALLAORO, A., 2005: The rediscovery of the crab, *Paragalene longicrura* (Nardo, 1868) (Decapoda, Brachyura, Xanthidae) in the Adriatic Sea. *Crustaceana* **78** (6), 749–753.
- PALLAORO, A. & J. DULČIĆ, 2004: On the occurrence of the spider crab *Maja goletziana* (Crustacea: Majidae) an alien species in the Adriatic Sea. *J. Mar. biol. Ass. U.K.*, **84**: 1007–1008.
- PESTA, O., 1918: Die Decapodenfauna der Adria. Versuch einer Monographie. Deuticke. Leipzig and Vienna. X+ 500.
- SCHUBART, C. D., 2003: The East Asian shore crab *Hemigrapsus sanguineus* (Brachyura: Varunidae) in the Mediterranean Sea: an independent human-mediated introduction. *Sci. Mar.*, **67** (2), 195–200.
- ŠTEVČIĆ, Z., 1990: Check list of the Adriatic decapod Crustacea. *Acta Adriat.*, **31**, 183–274.
- ŠTEVČIĆ, Z., 1993: History of investigations of the Adriatic decapod fauna. *Bios*, **1** (1), 151–161.
- ŠTEVČIĆ, Z., 1995: Contribution to the faunistic list of Adriatic decapod Crustacea. *Nat. Croat.*, **4** (2), 113–115.
- ŠTEVČIĆ, Z., 2002: New observation on the Adriatic decapod fauna (Years 1990–2000). *Crustaceana*, **75** (3–4), 643–647.
- UDEKEM D'ACQZ, C. D', 1996: The genus *Hippolyte* Leach, 1814 (Crustacea: Decapoda: Caridea: Hippolytidae) in the East Atlantic Ocean and Mediterranean sea, with the checklist of all species in the genus. *Zool. Verh. Leiden*, **303**, 1–133.
- UDEKEM D'ACQZ, C. D', 1999: Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au nord de 25° N (Inventory and distribution of the Crustacea Decapoda in Northeastern Atlantic Ocean, Mediterranean Sea and adjacent continental waters, north of 25° N). *Patrimoines naturels (M.N.H.N./S.P.N.)*, **40**, 1–383.
- UNGARO, N., 2000: Ritrovamenti di *Paromola cuvieri* (Risso, 1816) (Dromiacea, Homolidae) nel bacino Adriatico meridionale. *Biol. Mar. Medit.*, **7**(2), 744–747.
- UNGARO, N. & G. MARANO, 2002: On the distribution and demography of the deep-water pink shrimp and Norway lobster fishery stocks in the south Adriatic Sea. *Studia Marina*, **23** (1), 15–24.

- UNGARO, N., G. A. MARANO, L. CERIOLO & M. MARTINO, 2005: Distribution of demersal crustaceans in the southern Adriatic Sea. *Acta Adriat.*, **46** (1), 27–40.
- VUKANIĆ, V., 2003: Contribution to diversity of crustaceans in the Adriatic. *Natura Montenegrina*, Podgorica. **2**, 139–154.

SAŽETAK

Fauna jadranskih deseteronožnih rakova (Crustacea: Decapoda) – stanje i perspektive

M. Kirinčić & Z. Števcić

Jadranski deseteronožni raci (Crustacea: Decapoda) predmet su istraživanja mnogih znanstvenika počevši još od 16. stoljeća. Objavljeno je mnogo radova o ovoj skupini, a i novi se stalno pojavljuju. Od izlaska zadnje nadopune jadranske liste dekapoda (ŠTEVČIĆ, 2002) zabilježeno je 13 novih vrsta. Identificirano je i 8 imigrantskih vrsta koje su u Jadran dospjele balastnim vodama iz brodova ili bijegom iz akvakulturnih uzgajališta. Jadranska dekapodna fauna sadrži 241 vrsta, koje su klasificirane u 120 rodova, te 53 porodice. Uspoređujući te podatke s PESTOM (1918) koji navodi 143 vrste za isto područje, te s D'UDEKEM D'ACOSOM (1999) koji bilježi 340 vrsta za cijeli Mediteran, Jadran možemo smatrati do sada kvalitativno vrlo dobro istraženim morem. Povećano nalaženje vrsta zadnjih godina možemo pripisati intenzivnijim oceanografskim istraživanjima temeljenima na uporabi grabila, dredža, koča, te povećanom interesu znanstvenika-karcinologa za *in situ* istraživanje i sakupljanje materijala tehnikom autonomnog ronjenja koja dozvoljava istraživanja do tada nedostupnih mikrostaništa infralitoralne zone kao što su podmorske špilje i rupe, te valutićavi intersticij. Nedostaju još sustavna istraživanja dubokih voda (otvorenog mora) i riječnih ušća. U nadolazećim godinama možemo očekivati pojavljivanja egzotičnih vrsta iz Crvenog mora (Lessepsijske vrste) i iz ostalih dijelova svijeta u Jadransko more, zbog najava povećanog brodskeg prometa našim morem, kao i opće pojave globalnog zatopljenja mora. Iz svega rečenog proizlazi da je fauna jadranskih deseteronožaca dobro istražena, te spada među najistraženije uopće, a zbog stalnih revizija svojiti i useljavanja novih potrebno je stalno revidirati listu pa stoga možemo očekivati da će se znanstveni rezultati na ovom polju još dugo nastaviti.