

VETERINARSKI ARHIV 75 (4), 317-327, 2005

## Mucosubstances of the digestive tract mucosa in northern pike (*Esox lucius* L.) and european catfish (*Silurus glanis* L.)

Zdravko Petrincec<sup>1</sup>\*, Srebrenka Nejedli<sup>2</sup>, Snježana Kužir<sup>2</sup>,  
and Anđelko Opačak<sup>3</sup>

<sup>1</sup>Department for Biology and Pathology of Fish and Bees, Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

<sup>2</sup>Department of Anatomy, Histology and Embryology, Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

<sup>3</sup>Department of Zootechnics, Faculty of Agriculture, University of J. J. Strossmayer, Osijek

---

**PETRINEC, Z., S. NEJEDLI, S. KUŽIR, A. OPAČAK: Mucosubstances of the digestive tract mucosa in northern pike (*Esox lucius* L.) and european catfish (*Silurus glanis* L.). Vet. arhiv 75, 317-327, 2005.**

### ABSTRACT

The distribution and quality of mucosubstances in the digestive tract of pike (*Esox lucius*) and catfish (*Silurus glanis*) is described. The digestive tract of pike and catfish is short, in accordance with their predatory nature, and consists of oesophagus, stomach and intestine. The oesophagus is short, with longitudinal folds. The mucosa of pike oesophagus is stratified and contains two types of mucous cells: in contrast to catfish oesophagus there is only one type of mucous cell. The stomach is an elongated sac-like structure with glands in lamina propria. In pike the gastric glands are simply tubular, but in catfish they are branched tubular and surrounded by connective tissue. The intestines in both fishes are short with many intestinal villi. Many goblet cells are situated between superficial epithelial cells.

**Key words:** pike, catfish, digestive tract, mucosubstances, histochemistry

---

### Introduction

The digestive tract of fishes show marked diversity in its morphology and function. This is in correlation with taxonomy and the different feeding habits, as well as with body shape (VERMA and TYAGI, 1974; VERMA et al., 1974; KAPOOR et al., 1975; RAY and MOITRA, 1982; BUDDINGTON et al., 1997). The presence of mucosubstances in mucosa

---

\* Contact address:

Prof. Dr. Zdravko Petrincec, Department of Biology and Pathology of Fish and Bees, Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, 10000 Zagreb, Croatia, Phone: +385 1 2390 151; Fax: +385 1 2441 390; E-mail: [zdravko.petrincec@vef.hr](mailto:zdravko.petrincec@vef.hr)

of the digestive tract has been observed in most teleosts (OSTOS GARRIDO et al., 1993; NACHI et al., 1998; DOMENEGHINI et al., 1998; DOMENEGHINI et al., 1999; DOMENEGHINI et al., 2002). The gut mucins lubricate and protect the tunica mucosa against chemicals, parasites and acidity and form a diffusion barrier for various ions (PABST, 1987; GUPTA, 1989; VEGETTI et al., 1999). MURRAY et al. (1994) suggests that mucins in the posterior oesophagus of the same fish may play a role in pregastric digestion. FIERTAK and KILARSKI (2002) concluded that the glycoconjugate composition of cyprinid digestive tract depends on the species as well as on the intestinal region, and that there is no correlation between its composition and type of ingested food.

Both investigated fishes are active predators. Pike feed exclusively on small fish, while catfish have a broad range of prey, such as fish, frogs and crayfish and sometime small mammals (BEKBERGENOV and SAGITOV, 1984; OMAROV and POPOVA, 1985; STEPHENSON and MOMOT, 1991; ELVIRA et al., 1996; KANGUR and KANGUR, 1998).

The purpose of the present study was to describe histochemically the gut mucosubstances of two carnivorous fish species which display different feeding habits.

### Materials and methods

Adult northern pike (*Esox lucius*) and catfish (*Silurus glanis*) specimens collected from the Lonja River (NW Croatia) were used for investigation. Body weight ranged between 1-2 kg. Samples of digestive tract (oesophagus, stomach and middle part of intestine) were fixed in buffered 10% formalin and embedded in paraffin. Dewaxed sections (8 µm) were stained for general morphological purposes with haematoxylin and eosin (HE). Periodic acid-Shiff (PAS) (McMANUS, 1948) was used for demonstration of neutral glycoconjugates. Alcian blue (AB) at pH 2.5, (MOWRY, 1956) and metachromasia with toluidine blue (TB) were used for demonstration of various kinds of acid glycoconjugates (PEARSE, 1968).

### Results

*Pike*. Large numbers of distinct longitudinal folds extend along the entire length of the oesophagus. A differentiation exists between major and minor folds. The mucosa is lined with a stratified epithelium and two basic mucous cells are present. One type is round, the other elongated. Round cells are typically mucous and they are numerous on the basal part of the epithelium. Elongated cells, on the other hand, have a fine granular cytoplasm and are dominant in the superficial part of the epithelium. Both cell types were stained purple with periodic acid-Shiff (Fig. 1). But with AB and TB, only round mucous cells were stained (Fig. 2).

The mucosa of the stomach is raised into thick longitudinal folds. The epithelium of mucosa consists of very high, simple columnar cells with large nuclei located on the basal



Fig. 1. Pike, oesophagus, PAS-positive stained cells (arrows), scale bar = 200  $\mu$ m




Fig. 2. Pike, oesophagus, AB-positive stained cells (arrows), note unstained cells in apical part of mucosa, scale bar = 200  $\mu$ m



Fig. 3. Pike, stomach, PAS, epithelial cells show strong PAS reaction (arrows), scale bar = 200  $\mu$ m




Fig. 4. Pike, stomach AB-positive mucosubstances in apical part of epithelial cells (arrows), scale bar = 200  $\mu$ m




Fig. 5. Pike, intestine, AB, goblet cells show acid mucosubstances stained AB positive (arrow), scale bar = 100  $\mu$ m




Fig. 6. Catfish, oesophagus, PAS-positive mucous cells in oesophageal mucosa (arrows), scale bar = 200  $\mu$ m




Fig. 7. Catfish, stomach, PAS-positive mucosubstance present in apical part of epithelial cells (arrows), scale bar = 200  $\mu$ m




Fig. 8. Catfish, intestine, AB, numerous goblet cells stained AB-positive (arrows), scale bar = 200  $\mu$ m

side of the cells. The apical cytoplasm of epithelial cells shows presence of substances stained strongly with PAS and AB, and slightly with TB (Fig. 3 and 4).

The glandular cells are arranged as straight tubular glands and occupy the entire mucosal layer beneath the superficial epithelium. Tubular gastric glands are composed of one type of polygonal cell, with fine granular cytoplasm. There is no presence of any kinds of mucous substance in these glands.

The intestine is short and shows a uniform histological structure throughout its entire length. The mucosa is folded into long and thin intestinal villi consisting of a single layer of columnar epithelial cells. Mucous goblet cells are interspersed among them and connective lamina propria. Mucosubstances are present only in goblet cells and all were stained very strongly with PAS and AB (Fig. 5), but only of them is also stained with TB.

*Catfish.* The mucosa of the oesophagus is very thick and lined with stratified epithelium with only one type of intraepithelial mucous cell. All mucous cells show a moderate to high content of PAS-positive stained mucosubstances (Fig. 6). But only some of them were positively stained by AB, and metachromatic by TB.

The mucosa of the stomach is raised into fine longitudinal folds. The epithelium of the mucosa consists of simple columnar cells. The apical cytoplasm of the epithelial cells shows the presence of mucous substances stained strongly with PAS (Fig. 7), and slightly with AB and metachromatic with TB. The gastric glands were branched tubular and surrounded by a layer of connective tissue and opened into the bottom of gastric pits. The glands consisted of a single layer of cells of uniform appearance. The high content of PAS-positive, as well as TB-metachromatic substances, are present in the surface epithelial cell. A small amount of PAS-positively stained substances is observed in lamina propria. Glandular cells show no presence of mucosubstances.

The intestine is short and its mucosa has an almost uniform structure throughout its entire length, with some minor difference in the shape of the villi. The villi are long, branched and lined by columnar cells with rare goblet cells, which are found at all levels of the intestinal villi. The goblet cell contents stain intensively with AB (Fig. 8) and they are also PAS-positive. The epithelial cells show presence of a small amount of TB metachromatic substances, which are also present in some goblet cells.

## Discussion

Mucous is an important layer protecting the epithelia either from mechanical injuries or bacterial invasion (HUMBERT et al., 1984).

Mucous cells in fish oesophagus differentiated early during embryonic development and their appearance corresponds to the opening of the mouth. Mucous cells were found

early in the developing oesophagus of *Sparus aurata* (DOMENEGHINI et al., 1998), in *Chanos chanos* (FERRARIS et al., 1987) and *Solea solea* (BOULHIC and GABAUDAN, 1992). In this period these cells are PAS negative but stained by alcian blue, and contained acid mucosubstances only.

The mucosa of pike oesophagus is lined by the epithelium with two types of mucous cell. These two mucous cells appear to have different secretions, since the round mucous cells are PAS and AB positive, and TB-metachromatic, but elongated cells stained with AB only. Mucous finds in round mucous cells exhibit properties of neutral and acid sialosulfoglycoproteins as well as carboxylated and weak sulfated mucosubstances. Periodic acid-Schiff and alcian blue positive mucous cells in oesophagus of fishes were described by AL-HUSSAINI and KHOLY (1953), YAMADA and YOKOTE (1975), GARGIULIO et al. (1996), SCOCCO et al. (1998), and DOMENEGHINI et al. (1999). In elongated mucous cells there are neutral mucosubstances only. Two types of mucous cell described here were found in the oesophagus of tilapia (PASHA, 1964; MORRISON and WRIGTH, 1999). Mucous cells in catfish oesophagus were mainly PAS positive, although some contain AB positive and TB metachromatic mucosubstances. These histochemical staining properties of mucosubstances in catfish mucous cells point to the presence of a high content of neutral mucosubstances. Acid mucosubstances stained with AB and TB metachromatic suggest the presence of sialic acid-rich and weak sulfated mucosubstances. As reported by TIBBETTS (1997), mucous cells containing sialo- and sulfoglycoproteins lead to an increase in the viscosity of secretion, which is likely to be related to a higher protection role (SUPRASERT et al., 1987).

The stomach of pike displays uniform histological features. Superficial epithelial cells and simple tubular gastric glands in lamina propria form the mucosa. In catfish, gastric glands are arranged as tubular branched glands surrounded by connective tissue. The apical borders of the superficial epithelia in both fish show strong PAS reaction and mild AB and TB reaction. These reactions mostly indicated the presence of neutral mucosubstances and a small amount of acid mucosubstances. The acid mucosubstances are shown to be chiefly of the carboxylate and sialylated type. In fish, neutral mucosubstances secreted by the stomach epithelium have been related to the absorption of easily digested molecules, such as disaccharide and short-chain fatty acids (GRAU et al., 1992). The mucosubstances facilitate the movement of large-sized food particles, as well as protecting the gastric mucosa from mechanical injury. The presence of both neutral and acid mucosubstances was observed in the superficial gastric epithelium of the white sturgeon (DOMENEGHINI et al., 1999) and in developing larvae and adults of *Sparus aurata* (DOMENEGHINI et al., 1998).

The tubular gastric glands of pike and catfish show the presence of only one type of secretor cells, but they show no presence of any kind of mucosubstances. Many authors agree that the gastric glands of teleosts contain only one type of secretory cell (WESTERN and JENNINGS, 1970; VERMA and TYAGI, 1974; RAY and MOITRA, 1982). ARELLANO

et al. (2001) reported the presence of light and dark cells in the gastric glands of *Solea senegalensis*. It is generally thought that fish gastric glands synthesise both pepsinogen and acid (SMITH, 1989) and that neutral mucosubstances have a buffering effect on the high acidity of the stomach content (SCOCCO et al., 1996).

The intestine of both investigated fish shows uniform histological structure throughout its entire length. The mucosa of pike intestine consists of a single layer of columnar cells with many mucous goblet cells. The catfish intestinal mucosa is similar to the pike, but there are more mucous cells than in pike. The intestinal goblet cells of both investigated fishes synthesise neutral and acidic mucosubstances. The acid mucosubstances are made by carboxylated and sulfated mucoconjugates. The presence of TB metachromatic substances, in particular goblet cells, shows in the presence of a small amount of sialic-rich nonsulfated mucoconjugates.

The first goblet cells can be observed early during the differentiation of the intestinal mucosa of fish (BOULHIC and GABAUDAN, 1992; MIYASHITA et al., 1998; GARCIA HERNANDEZ et al., 2001; MORRISON et al., 2001). The presence of mucous-producing goblet cells in the intestinal mucosa has been reported in many fish studied earlier (NARASIMHAM and PARVATHESWARAO, 1974; CHATURVEDI and GUPTA, 1976; RAY and MOITRA, 1982; DOMENEGHINI et al., 1998; NACHI et al., 1998; DOMENEGHINI et al., 1999; MORRISON and WRIGHT, 1999). The mucous secreted by goblet cells in the intestine has many functions. For example, it lubricates undigested materials for onward progression into the rectum. NARASIMHAM and PARVATHESWARAO (1974) suggested a possible role of intestinal mucins in osmoregulation. The studies of RIBELLES et al. (1995) have shown that the quality of gut mucosubstances is directly related to environmental conditions, which in turn may directly affect the function of the alimentary tract. The presence of mucosubstances, especially those sulfated in the intestine, possibly regulate the transfer of proteins, or a fragment of them, as well as of ions and fluids (STROMBAND and VAN DER VEEN, 1981; GUPTA, 1989; SIRE and VERNIER, 1992; SEGNER et al., 1994; DOMENEGHINI et al., 1998). The possible role of mucosubstances in intestinal absorption processes is supported by the findings of BOŽIĆ et al. (2001) who observed that starvation induced an increase in the number of intestinal goblet cells in carp.

This investigation, as well as previous investigations on other fishes, demonstrates that the quality of gut mucosubstances varies in different regions of pike and catfish alimentary canals and that it sustains functional harmony of the digestive tract. It appears that the quality of mucoconjugates in particular parts of the digestive tract is similar in both investigated carnivorous fish, and that there is no correlation between mucous composition and food habits.

## References

- AL-HUSSAINI, A. H., A. A. KHOLY (1953): On the functional morphology of the alimentary tract of some omnivorous fish. Proceedings of the Egyptian Academy of Sciences 4, 17-39.
- ARELLANO, J. M., V. STORCH, C. SARASQUETE (2001): Histological and histochemical observations in the stomach of the Senegal sole, *Solea senegalensis*. Histol. Histopathol. 16, 511-521.
- BEKBERGENOV, ZH., N. I. SAGITOV (1984): Feeding habits of juveniles of some comercial fishes in the Amu Dar'ya River. J. Ichthyol. 124, 18-22.
- BOULHIC, M., J. GABAUDAN (1992): Histological study of the organogenesis of the digestive system and swim bladder in the Dover sole, *Solea solea* (Linnaetus 1758). Aquaculture 102, 373-396.
- BUDDINGTON, R. K., Å. KROGDAHL, A. M. BAKKE-MC KELLEP (1997): The intestines of carnivorous fish: structure and function and the relations with diet. Acta Physiol. Scand. 161, suppl. 638, 67-80.
- BOŽIĆ, F., E. SREBOČAN, Z. KOZARIĆ (2001): Starvation- induced pathobiology in the gut of carp (*Cyprinus carpio* L.). Berl. Münch. Tierärztl. Wschr. 114, 134-138.
- CHATURVEDI, L., R. K. D. GUPTA (1976): Studies on the morphology and histology of the digestive tract of *Bagarius bagarius* (Ham). Gegenbaurs. morph. Jahrb. 122, 636-645.
- DOMENEGHINI, C., R. P. STRANINI, A. VEGGETTI (1998): Gut glycoconjugates in *Sparus aurata* L. (Pisces, Teleostei), a comparative histochemical study in larval and adult ages. Histol. Histopathol. 13, 359-372.
- DOMENEGHINI, C., S. ARRIGHI, G. RADAELLI, G. BOSI, G. S. MASCARELLO (1999): Morphological and histochemical peculiarities of the gut in white sturgeon, *Acipenser transmontanus*. European J. Histochemistry 43, 135-145.
- DOMENEGHINI, C., G. RADAELLI, G. BOSI, S. ARRIGHI, A. DI GIANCAMILLO, M. PAZZAGIA, F. MASCARELLO (2002): Morphological and histochemical differences in the structure of alimentary canal in feeding and runt (feed deprived) white sturgeon (*Acipenser transmontanus*). J. Appl. Ichthyol. 18, 341-346.
- ELVIRA, B., G. G. NICOLA, A. ALMODOVAR (1996): Pike and red swamp crayfish: a new case on predator-prey relationship between aliens in central Spain. J. Fish Biol. 48, 437-446.
- FERRARIS, R. P., J. D. TAN, M. C. DE LA CRUZ (1987): Development of the digestive tract of milkfish, *Chanos chanos* (Forsskal): histology and histochemistry. Aquaculture 61, 241-257.
- FIERTAK, A., W. M. KILARSKI (2002): Glycoconjugates of the intestinal goblet cells of four cyprinids. Cell. Mol. Life. Sci. 59, 1724-1733.
- GARCÍA HERNÁNDEZ, M. P., M. T. LOZANO, M. T. ELBAL, B. AGULLEIRO (2001): Development of the digestive tract of sea bass (*Dicentrarchus labrax* L.). Light and electron microscopic studies. Anatomy Embryol. 204, 39-57.
- GARGIULO, A. M., C. DALL'AGRILO, Z. TSOKU, P. CACCARELLI, V. PEDINI (1996): Morphology and histology of the oesophagus of a warmwater tilapine fish (Teleostei). J. Appl. Ichthyol. 12, 121-124.



- Z. Petrinc et al.: Mucosubstances of the digestive tract mucosa in northern pike (*Esox lucius* L.) and european catfish (*Silurus glanis* L.)
- GRAU, A., S. CRESPO, M. C. SARASQUETE, M. L. GONZALES de CANALES (1992): The digestive tract of the amberjack *Seriola dumerii*, Risso: a light and scanning electron microscope study. *J. Fish Biol.* 41, 287-303.
- GUPTA, B. J. (1989): The relationship of mucoid substances and ion and water transport, with new data on intestinal goblet cells and a model for gastric secretion. *Symp. Soc. Exp. Biol.* 43, 81-110.
- HUMBERT, W., R. KIRSCH, M.F. MEISTER (1984): Scanning electron microscopic study of the oesophageal mucous layer in the eel *Anguilla anguilla* L. *J. Fish Biol.* 25, 117-122.
- KANGUR, A., P. KANGUR (1998): Diet composition and size-related changes in the feeding of pikeperch, *Stizostedion lucioperca* (Percidae) and pike, *Esox lucius* (Esocidae) in the Lake Peipsi (Estonia). *Ital. J. Zool.* 65, 255-259.
- KAPOOR, B. G., H. SMITH, I. A. VERIGHINA (1975): The alimentary canal and digestion in teleosts. In: *Advances in Marine Biology* (Russel, F. S., M. Yonge, Eds.), Vol. 13. Acad. Press. London pp. 109-239.
- McMANUS, J. F. A. (1948): Histological and histochemical uses of periodic acid. *Stain Technol.* 23, 99-108.
- MIYASHITA, S., K. KATO, Y. SAWADA, O. MURATA, Y. ISHITANI, K. SHIMIZU, S. YAMAMOTO, H. KUMAI (1998): Development of digestive system and digestive enzyme activities of larval and juvenile Bluefin tuna, *Thunnus thynnus*, reared in the laboratory. *Suisanzoshoku* 46, 111-120.
- MORRISON, C. M., J. R. WRIGHT JR (1999): A study of the histology of the digestive tract of the Nile tilapia. *J. Fish Biol.* 54, 597-606.
- MORRISON, C. M., T. MIYAKE, J. R. WRIGHT (2001): Histological study of the development of the embryo and early larva of *Oreochromis niloticus* (Pisces: Cichilidae). *J. Morphol.* 247, 172-195.
- MOWRY, R. W. (1956): Alcian blue technics for the histochemical study of acidic carbohydrates. *J. Histochem. Cytochem.* 4, 407-408.
- MURRAY, H. M., G. M. WRIGHT, G. P. GOFF (1994): A study of the posterior oesophagus in the winter flounder, *Pleuronectes americanus*, and the yellowtail flounder, *Pleuronectes ferrugineus*: morphological evidence of pregastric digestion? *Can. J. Zool.* 72, 1191-1198.
- NACHI, A. M., F. J. HERNANDETBLASZQUEZ, R. L. BARBIERI, R. G. LEITE, S. FERRI, M. T. PHAN (1998): Intestinal histology of a detritivorous (iliophagous) fish *Prochilodus scrofa* (Characiformes, Prochilodontidae). *Annales des Sciences Naturelles-Zoologie et Biologie Animale* 19, 81-88.
- NARASIMHAM, C., V. PARATHESWARAO (1974): Adaptation to osmotic stress in a fresh-water euryhaline teleost, *Tilapia mossambica*. X. Role of mucopolysaccharides. *Acta histochemica* 51, 37-49.
- OMAROV, O. P., O. A. POPOVA (1985): Feeding behavior of pike, *Esox lucius* and catfish, *Silurus glanis*, in the Arakum Reservoirs of Dagestan. *J. Ichthyol.* 25, 25-36.

Z. Petrínek et al.: Mucosubstances of the digestive tract mucosa in northern pike (*Esox lucius* L.) and european catfish (*Silurus glanis* L.)

- OSTOS GARRIDO, M. V., M. I. NUNEZ TORRES, M. A. ABAURREA EQUIAOAIN (1993): Histological, histochemical and ultrastructural analysis of the gastric musosa in *Oncorhynchus mykiss*. *Aquaculture* 115, 121-132.
- PABST, R. (1987): The anatomical basis for the immune function of the gut. *Anat. Embriol.* 176, 135-144.
- PASHA, K. (1964): The anatomy and histology of alimentary canal of herbivorous fish *Tilapia mossambica* (Peters). *Proceedings Indian Acad. Sci.* B54, 340-349.
- PEARSE, A. E. G. (1968): *Histochemistry. Theoretical and Applied.* J. & A. Churchill Ltd. London.
- RAY, A. K., S. K. MOITRA (1982): On the morpho-histology of the alimentary tract in the Indian Climbing Perch, *Anabas tastudineus* (Bloch), in relation to food and feeding habits. *Gegenbaurs. morph. Jahrb.* 128, 778-798.
- RIBELLES, A., M. C. CARRASCO, M. ROSETY, M. ALDANA (1995): A histochemical study of the biological effects of sodium dodecyl sulfate on the intestine of the gilthead seabream, *Sparus aurata* L. *Ecotox. Environ. Safe.* 32, 131-138.
- SCOCCO, P., D. ACCILI, G. MENGHI, P. CECCARELLI (1998): Unusual glycoconjugates in the oesophagus tilapine polyhybrid. *J. Fish Biol.* 53, 39-48.
- SCOCCO, P., P. CECCARELLI, G. MENGHI (1996): Glycohistochemistry of the *Tilapia* sp. stomach. *J. Fish Biol.* 49, 584-593.
- SENGER, H., V. STORCH, M. REINECKER, W. KLOAS, W. HANKE (1994): The development of functional digestive and metabolic organs in turbot, *Scophthalmus maximus*. *Marine Biol.* 119, 471-486.
- SIRE, M. F., J. M. VERNIER (1992): Intestinal absorption of protein in teleost fish. *Comp. Biochem. Physiol.* 103A, 771-781.
- SMITH, L. S. (1989): Digestive functions in teleost fishes. In: *Fish nutrition*, 2<sup>nd</sup> ed. (Halver J. E., ed.). Academic Press. London. pp. 331-421.
- STEPHENSON, S. A., W. T. MOMOT (1991): Food habits and growth of walleye *Stizostedion vitreum*, smallmouth bass, *Micropterus dolomieu*, and northern pike, *Esox lucius*, in the Kaministiquia River, Ontario. *Can. Field Nat.* 105, 517-521.
- STROMBAND, H. W. J., H. VAN DER VEEN (1981): Localization of protein absorption during transport of food in the intestine of the grasscarp, *Ctenopharyngodon idella*. *J. Exp. Zool.* 218, 149-156.
- SUPRASERT, A., T. FUJIOKA, K. YAMADA (1987): The histochemistry of glycoconjugates in the colonic epithelium of the chicken. *Histochemistry* 86, 491-497.
- TIBBETTS, I. R. (1997): The distribution and function of mucous cells and their secretions in the alimentary tract of *Arrhamphus sclerolepis krefftii*. *J. Fish Biol.* 50, 809-820.
- VEGGETTI, A., A. ROWLERSON, G. RADAELLI, S. ARRIGHI, C. DOMENEGHINI (1999): Post-hatching development of the gut and lateral muscle in the sole, *Solea solea* (L). *J. Fish Biol.* 55 (Suppl. A), 44-65.

Z. Petrinc et al.: Mucosubstances of the digestive tract mucosa in northern pike (*Esox lucius* L.) and european catfish (*Silurus glanis* L.)

VERMA, S. R., M. P. TYAGI (1974): A comparative histological study of the stomach of a few teleost fishes (Part I). Gegenbaurs. morph. Jahrb.120, 244-253.

VERMA, S. R., M. P. TYAGI, R. C. DALELA (1974): Morphological variation in the stomach of few teleost fishes in relation to food and feeding habits. Gegenbaurs. Morph. Jahrb.120, 367-380.

WESTERN, J. R. H., J. B. JENNINGS (1970): Histochemical demonstration of hydrochloric acid in the gastric tubules of teleosts using an *in vivo* Prussian blue technique. Comp. Biochem. Physiol. 35, 879-884.

YAMADA, K., M. YOKOTE (1975): Morphochemical analysis of mucosubstances in same epithelial tissues of the eel (*Angulia japonica*). Histochemistry 43, 161-172.

Received: 16 June 2004

Accepted: 28 June 2005

---

**PETRINEC, Z., S. NEJEDLI, S. KUŽIR, A. OPAČAK: Mukopolisaharidi u sluznici probavnog trakta štuke (*Esox lucius* L. ) i soma (*Silurus glanis* L.). Vet. arhiv 75, 317-327, 2005.**

**SAŽETAK**

Istraživana su svojstva i raspored mukosubstancija u probavnom traktu štuke i soma. Probavni trakt istraživanih riba je jednostavne građe i sastoji se od jednjaka, želuca i relativno kratkog crijeva. U sluznici štuke opažaju se dva tipa stanica koje pokazuju prisutnost neutralnih mukopolisaharida, a kiselih mukopolisaharidi nađeni su samo u mukoznim stanicama bazalnog dijela epitela. U jednjaku soma opisan je samo jedan tip okruglih mukoznih stanica u kojima su nađeni neutralni mukopolisaharidi, a samo neke od njih sadrže i kisele mukopolisaharide. Želudac je vrečastog oblika sa žlijezdama u lamini propriji. U žlijezdama nisu dokazani mukopolisaharidi, no nađeni su u površinskom epitelu. U štuke su neutralnih i kiselih osobina, a u soma su uglavnom neutralni s manjom količinom kiselih. Mješavina neutralnih i kiselih mukopolisaharida prisutna je u vrčastim stanicama crijeva. Broj vrčastih stanica veći je u crijevu soma nego štuke.

**Ključne riječi:** štuca, som, probavni trakt, mukopolisaharidi, histokemija

---

