

VETERINARSKI ARHIV 75 (4), 359-368, 2005

## **Determination of levels of some essential (iron, copper, zinc) and toxic (lead, cadmium) metals in the blood of sheep and in samples of water, plants and soil in Northwest Turkey**

**Mehmet Erman Or<sup>1\*</sup>, Abdullah Kayar<sup>1</sup>, Ali Rıza Kıziler<sup>2</sup>, Çağla Parkan<sup>1</sup>,  
Remzi Gönül<sup>1</sup>, Bora Barutçu<sup>2</sup>, and Hazım Tamer Dodurka<sup>1</sup>**

<sup>1</sup>*Istanbul University, Veterinary Faculty, Department of Internal Medicine, Avcılar-Istanbul, Turkey*

<sup>2</sup>*Istanbul University, Cerrahpaşa Medicine Faculty, Department of Biophysics, Cerrahpaşa-Istanbul, Turkey*

---

**OR, M. E., A. KAYAR, A. R. KIZILER, Ç. PARKAN, R. GÖNÜL, B. BARUTÇU, H. T. DODURKA: Determination of levels of some essential (iron, copper, zinc) and toxic (lead, cadmium) metals in the blood of sheep and in samples of water, plants and soil in Northwest Turkey. Vet. arhiv 75, 359-368, 2005.**

### **ABSTRACT**

The aim of this investigation was to determine the levels of trace elements (zinc, iron, copper), as well as of lead and cadmium, in both the soil and in organism in order to obtain more efficient economical results and healthier sheep breeds in Northwest Turkey. Based on the results of these analyses, we plan to make suggestions to veterinarians and sheep breeders in that region on how to improve their performance. The study was carried out on 400 sheep from different sites in Northwest Turkey and included the collection of water (n = 5), plant (n = 5) and soil (n = 5) samples from every district. When serum parameters were examined an important decrease in iron levels was found in the 8<sup>th</sup>, 9<sup>th</sup> and 10<sup>th</sup> districts, while the lowest zinc levels were found in the 1<sup>st</sup> and 7<sup>th</sup> districts, the lowest copper levels in the 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and 6<sup>th</sup> districts, the highest lead levels in the 9<sup>th</sup> and 10<sup>th</sup> districts, and the highest cadmium levels in the 2<sup>nd</sup>, 5<sup>th</sup> and 6<sup>th</sup> districts. Analyses of water, plant and soil samples showed significant differences in zinc and copper levels between districts. High levels of lead were detected in water and plant samples. We are of the opinion that important improvements will be obtained in the treatment and control of diseases due to deficiency of trace elements and of parasitic, bacterial and viral diseases due to related systems affected by such deficiencies if these results are appropriately communicated to animal breeders and veterinary surgeons operating in the regions in question.

**Key words:** sheep, trace element, zinc, iron, copper, lead, cadmium

---

\* Contact address:

Assoc. Prof. Dr. Mehmet Erman Or, Department of Internal Medicine, Faculty of Veterinary Medicine, Istanbul University, Avcılar 34320, Istanbul-Turkey, Phone: +90 212 473 70 70 (15 Line)-17279; Fax: +90 212 473 72 41; E-mail: [ermanor@istanbul.edu.tr](mailto:ermanor@istanbul.edu.tr)

## **Introduction**

The importance of many inorganic elements classified as macro- and micro-elements in the diet and on the growth of humans and animals has meanwhile been accepted and recognized beyond doubt (ALTINTAŞ et al., 1990). Diseases due to deficiencies or an excess of macro- and micro-elements in animals are of unquestionable importance. While the deficiency or excess of one or a few elements damages normal bodily functions, imbalance among elements impairs the regular function of organism (ÇAMAŞ et al., 1994).

While deficiencies or excess of trace elements in animals result in serious clinical disorders, the important economic losses they cause have been highlighted only in recent years. Clinical signs of trace element deficiencies in animals are diarrhoea, anemia, alopecia, depigmentation, development disorders in the osseous tissues, impaired movement, seborrhea, hyperkeratosis, parakeratosis, anorexia, fertility problems, development problems in young animals, decreased production, tetany, decrease in protein synthesis, insufficiencies in the immune system, abortions not related to infections and pica (ERGÜN, 1983; MARTIN et al., 1983; KUMARESAN and KAPIOH 1984; AĞAOĞLU, 1991; ÇAMAŞ et al., 1994; YILDIZ et al., 1995; UNDERWOOD, 1997; ZANTOPOULOS et al., 1999).

It has been reported that trace elements are of equal importance as hormone and vitamins and have an impact in all climatic zones (OZAN, 1985; ALTINTAŞ et al., 1990; AĞAOĞLU, 1991; ÇAMAŞ et al., 1994). It is also been pointed out that losses caused by the effects of trace elements are as important as the losses incurred as a result of infectious and parasitary diseases (ÇAMAŞ et al., 1994), because trace elements have a significant impact on resistance against diseases in living organisms.

Various systems are affected by a deficiency or excess of trace elements such as zinc, copper and iron, or by the negative effects of elements that contribute to metal pollution poisoning such as lead and cadmium. As a consequence, any of the specific diseases caused by any of the elements in question may occur, or they may contribute as important factors in the occurrence of various diseases (MARTIN et al., 1983; ALTINTAŞ et al., 1990; AĞAOĞLU, 1991; ÇAMAŞ et al., 1994).

The aim of this study was to determine the levels of trace elements in sheep and water, soil and plants in order to define the influence of trace elements in the natural environment on sheep in Northwest Turkey and consequently to provide recommendations to veterinary surgeons and sheep breeders in this region.

## **Materials and methods**

The research involved a total of 400 sheep of the Kıvırcık breed, all between 2-3 years old, which were selected from different flocks in Northwest Turkey. The randomly chosen sheep were neither pregnant in a late state nor had they recently given birth. Sheep were

regularly administered antiparasits after faeces examination. The study was carried out from April to June while the sheep were grazing in the pasture. As a first step, the northwestern region of Turkey was divided into 12 districts, as shown in Table 1, and water (n = 5), plant (n = 5) and soil (n = 5) samples were collected from every region.

Table 1. Sampling districts in Northwest Turkey

| District Number | Name of the District                                      | Number of Collected Blood Samples |
|-----------------|---|-----------------------------------|
| 1               | Babaeski, Sofuhali, Pehlivan köy and surrounding villages | 25                                |
| 2               | Kırklareli, Kofçaz and surrounding villages               | 35                                |
| 3               | Istanbul University Veterinary Faculty Farm               | 35                                |
| 4               | Çorlu, Çerkezköy and surrounding villages                 | 35                                |
| 5               | Kırklareli, Dereköy and surrounding villages              | 30                                |
| 6               | Çatalca and surrounding villages                          | 30                                |
| 7               | Babaeski, Lüleburgaz and surrounding villages             | 25                                |
| 8               | Tekirdağ, Malkara and surrounding villages                | 35                                |
| 9               | Tekirdağ, Keşan and surrounding villages                  | 40                                |
| 10              | Saray, Vize, Pınarhisar and surrounding villages          | 40                                |
| 11              | Kırcaali, Uzunköprü and surrounding villages              | 35                                |
| 12              | Edirne and surrounding villages                           | 35                                |

Samples were collected in sterile plastic containers in order to prevent any contamination. Water, plant and soil samples were evaluated by the Department of Biophysics at the Cerrahpaşa Faculty of Medicine of Istanbul University. 10 cc blood were taken from the v. jugularis of sheep selected from each region, both with and without anticoagulant. Seven cc anticoagulated blood was centrifuged at 5000 rpm for 3-5 minutes for zinc, iron and copper measurements. These sera were stored at -18 °C until analysis. For the lead and cadmium measurements an equal amount of 20% TCA (Trichloroacetic acid) was added to the tubes with 3 cc anticoagulated blood and this mixture was then centrifuged. Trace elements (Zn, Fe, Cu) as well as lead and cadmium measurements were carried out with a Shimadzu Atomic Absorption Spectrometry model AA-680 in accordance with the technique described in the references (BROWN et al., 1986; BROWN and TAYLOR, 1995).

Mean values, standard deviations and statistical differences were calculated using the Duncan F Test (DUNCAN, 1955).

## Results

The mean values and standard deviations of the levels of non-toxic (zinc, iron, copper) and toxic (lead and cadmium) elements in the serum of 400 sheep, as well as the examined water, plant and soil samples taken in Northwest Turkey are shown in Tables 2, 3, 4 and 5.

Results summarized in Table 2 show the lowest serum zinc levels in the 1<sup>st</sup> and 7<sup>th</sup> districts, a slight decrease in serum iron levels in the 8<sup>th</sup>, 9<sup>th</sup> and 10<sup>th</sup> districts, the lowest copper levels in the 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup> and 6<sup>th</sup> districts, the highest lead levels in the 9<sup>th</sup>, 10<sup>th</sup> and 11<sup>st</sup> districts, and the highest cadmium levels in the 2<sup>nd</sup>, 5<sup>th</sup> and 6<sup>th</sup> districts.

Results presented in Table 3 reveal the lowest zinc levels in the 1<sup>st</sup> and 2<sup>nd</sup> districts, a significant decrease in the iron level in the 2<sup>nd</sup> district, the lowest copper levels in the 1<sup>st</sup> and 7<sup>th</sup> districts, the highest lead levels in the 11<sup>th</sup> and 12<sup>th</sup> districts, and the highest cadmium level in the 7<sup>th</sup> district.

When plant samples were examined, the lowest zinc level was found in the 3<sup>rd</sup> district, the lowest copper levels in the 3<sup>rd</sup>, 8<sup>th</sup> and 9<sup>th</sup> districts, the highest lead levels in the 4<sup>th</sup> and 7<sup>th</sup> districts, and the highest cadmium level in the 1<sup>st</sup> district.

Table 2. Mean values and standard deviations of essential elements (Zn, Fe, Cu) in serum and toxic metal (Pb, Cd) levels in full blood of 400 sheep from different districts of Northwest Turkey

|          |    | Zn (µg/dL)         |     | Fe (µg/dL)          |      | Cu (µg/dL)         |      | Pb (µg/dL)           |     | Cd (µg/dL)         |     |
|----------|----|--------------------|-----|---------------------|------|--------------------|------|----------------------|-----|--------------------|-----|
| District | N  | X                  | Sx  | X                   | Sx   | X                  | Sx   | X                    | Sx  | X                  | Sx  |
| 1        | 25 | 22.5 <sup>e</sup>  | 4.6 | 88.1 <sup>cb</sup>  | 11.8 | 29.9 <sup>ed</sup> | 4.6  | 4.7 <sup>cha</sup>   | 2.9 | 0.2 <sup>f</sup>   | 0.1 |
| 2        | 35 | 31.1 <sup>c</sup>  | 5.3 | 92.2 <sup>b</sup>   | 14.6 | 30.1 <sup>ed</sup> | 12.2 | 4.2 <sup>fc</sup>    | 1.4 | 1.8 <sup>ba</sup>  | 0.5 |
| 3        | 35 | 28.8 <sup>d</sup>  | 4.8 | 88.6 <sup>cb</sup>  | 12.7 | 29.6 <sup>ed</sup> | 5.6  | 3.9 <sup>f</sup>     | 0.5 | 1.2 <sup>c</sup>   | 0.7 |
| 4        | 35 | 29.1 <sup>d</sup>  | 4.1 | 87.1 <sup>deb</sup> | 13.3 | 30.4 <sup>ed</sup> | 5.7  | 4.4 <sup>fedcb</sup> | 0.5 | 0.9 <sup>dc</sup>  | 0.4 |
| 5        | 30 | 32.1 <sup>cb</sup> | 3.5 | 89.9 <sup>cb</sup>  | 7.3  | 29.1 <sup>e</sup>  | 3.1  | 4.2 <sup>fed</sup>   | 0.3 | 2.1 <sup>a</sup>   | 0.4 |
| 6        | 30 | 28.6 <sup>d</sup>  | 3.8 | 89.6 <sup>cb</sup>  | 10.4 | 27.7 <sup>e</sup>  | 4.7  | 4.3 <sup>fedcb</sup> | 0.4 | 1.8 <sup>ba</sup>  | 0.4 |
| 7        | 25 | 20.9 <sup>e</sup>  | 1.7 | 88.3 <sup>cb</sup>  | 11.7 | 32.6 <sup>dc</sup> | 8.1  | 4.4 <sup>edcb</sup>  | 0.3 | 0.3 <sup>f</sup>   | 0.1 |
| 8        | 35 | 27.7 <sup>d</sup>  | 3.4 | 85.9 <sup>cdc</sup> | 2.5  | 35.2 <sup>c</sup>  | 2.2  | 4.8 <sup>cha</sup>   | 0.2 | 0.9 <sup>cdc</sup> | 0.5 |
| 9        | 40 | 34.1 <sup>b</sup>  | 2.5 | 81.4 <sup>e</sup>   | 2.2  | 35.7 <sup>c</sup>  | 2.7  | 5.1 <sup>a</sup>     | 0.2 | 1.1 <sup>c</sup>   | 0.5 |
| 10       | 40 | 36.4 <sup>a</sup>  | 3.7 | 82.9 <sup>ed</sup>  | 3.8  | 35.6 <sup>c</sup>  | 3.3  | 5.0 <sup>a</sup>     | 2.6 | 0.9 <sup>ed</sup>  | 0.4 |
| 11       | 35 | 28.9 <sup>d</sup>  | 3.9 | 89.2 <sup>cb</sup>  | 4.4  | 41.2 <sup>b</sup>  | 3.1  | 4.9 <sup>ba</sup>    | 0.3 | 0.9 <sup>cdc</sup> | 0.5 |
| 12       | 35 | 32.8 <sup>cb</sup> | 4.9 | 97.2 <sup>a</sup>   | 6.7  | 44.9 <sup>a</sup>  | 5.7  | 4.3 <sup>fedc</sup>  | 0.6 | 0.7 <sup>e</sup>   | 0.6 |

There is a statistically significant difference ( $P < 0.001$ ) between the mean values (X), which are indicated with different letters in each column. N = Number of animals, X = Mean values, Sx = Standard deviation.

Table 3. Mean values and standard deviations of essential (Zn, Fe, Cu) and toxic (Pb, Cd) metal levels in water samples from different districts of Northwest Turkey

| District | N | Zn (µg/L)        |     | Fe (µg/L)       |     | Cu (µg/L)        |     | Pb (µg/L)         |     | Cd (µg/L)           |     |
|----------|---|------------------|-----|-----------------|-----|------------------|-----|-------------------|-----|---------------------|-----|
|          |   | X                | Sx  | X               | Sx  | X                | Sx  | X                 | Sx  | X                   | Sx  |
| 1        | 5 | 19 <sup>g</sup>  | 3   | 29 <sup>d</sup> | 5   | 7 <sup>h</sup>   | 2   | 1.45 <sup>i</sup> | 3   | 0.05 <sup>f</sup>   | 1   |
| 2        | 5 | 7 <sup>h</sup>   | 3   | 4 <sup>g</sup>  | 1   | 39 <sup>e</sup>  | 5   | 1.24 <sup>h</sup> | 0.6 | 0.78 <sup>deb</sup> | 3   |
| 3        | 5 | 140 <sup>c</sup> | 5.3 | 32 <sup>c</sup> | 0.9 | 40 <sup>ed</sup> | 2.9 | 2.45 <sup>e</sup> | 0.2 | 0.54 <sup>ed</sup>  | 0.2 |
| 4        | 5 | 125 <sup>e</sup> | 5.8 | 21 <sup>f</sup> | 0.5 | 64 <sup>a</sup>  | 4.2 | 1.78 <sup>g</sup> | 0.1 | 0.56 <sup>edc</sup> | 4   |
| 5        | 5 | 135 <sup>d</sup> | 5.2 | 36 <sup>b</sup> | 0.5 | 47 <sup>c</sup>  | 0.3 | 2.65 <sup>d</sup> | 0.2 | 0.85 <sup>cb</sup>  | 0.2 |
| 6        | 5 | 120 <sup>f</sup> | 5.4 | 35 <sup>b</sup> | 0.8 | 32 <sup>f</sup>  | 2.6 | 2.1 <sup>f</sup>  | 0.1 | 0.45 <sup>e</sup>   | 0.3 |
| 7        | 5 | 148 <sup>b</sup> | 5.3 | 39 <sup>a</sup> | 0.8 | 27 <sup>g</sup>  | 1.1 | 2.03 <sup>f</sup> | 0.2 | 1.23 <sup>a</sup>   | 0.6 |
| 8        | 5 | 135 <sup>d</sup> | 4.9 | 32 <sup>c</sup> | 1.2 | 41 <sup>d</sup>  | 2.3 | 2.8 <sup>c</sup>  | 0.1 | 0.89 <sup>b</sup>   | 0.3 |
| 9        | 5 | 120 <sup>f</sup> | 5.2 | 30 <sup>d</sup> | 0.6 | 59 <sup>b</sup>  | 3.3 | 2.48 <sup>e</sup> | 0.1 | 0.56 <sup>edc</sup> | 0.6 |
| 10       | 5 | 170 <sup>a</sup> | 4.6 | 33 <sup>c</sup> | 3.2 | 63 <sup>a</sup>  | 1.6 | 2.05 <sup>f</sup> | 0.1 | 0.74 <sup>deb</sup> | 0.3 |
| 11       | 5 | 150 <sup>b</sup> | 3.2 | 27 <sup>e</sup> | 0.8 | 48 <sup>c</sup>  | 1.3 | 5.2 <sup>a</sup>  | 0.1 | 0.85 <sup>cb</sup>  | 0.5 |
| 12       | 5 | 140 <sup>c</sup> | 5.6 | 32 <sup>c</sup> | 1.2 | 40 <sup>d</sup>  | 1.8 | 4.5 <sup>b</sup>  | 0.1 | 0.12 <sup>f</sup>   | 0.6 |

There is a statistically significant difference ( $P < 0.001$ ) between the mean values (X), which are indicated with different letters in each column.

Table 4. Mean values and standard deviations of essential (Zn, Fe, Cu) and toxic (Pb, Cd) metal levels in plant samples from different districts of Northwest Turkey

| District | N | Zn (µg/g)          |      | Fe (µg/g)           |      | Cu (µg/g)         |     | Pb (µg/g)          |      | Cd (µg/g)          |     |
|----------|---|--------------------|------|---------------------|------|-------------------|-----|--------------------|------|--------------------|-----|
|          |   | X                  | Sx   | X                   | Sx   | X                 | Sx  | X                  | Sx   | X                  | Sx  |
| 1        | 5 | 167.4 <sup>a</sup> | 9.8  | 399.3 <sup>a</sup>  | 26.4 | 47.3 <sup>a</sup> | 3.8 | 1.99 <sup>b</sup>  | 0.29 | 1.04 <sup>a</sup>  | 0.9 |
| 2        | 5 | 95.2 <sup>c</sup>  | 3.7  | 250 <sup>c</sup>    | 21.4 | 8.1 <sup>c</sup>  | 2.4 | 1.26 <sup>d</sup>  | 0.21 | 0.68 <sup>e</sup>  | 0.2 |
| 3        | 5 | 39.6 <sup>hj</sup> | 4.6  | 115.3 <sup>k</sup>  | 11.7 | 5.5 <sup>d</sup>  | 1.6 | 0.85 <sup>f</sup>  | 0.04 | 0.76 <sup>d</sup>  | 0.6 |
| 4        | 5 | 85.4 <sup>f</sup>  | 4.7  | 185.5 <sup>h</sup>  | 14.7 | 7.6 <sup>c</sup>  | 0.8 | 2.14 <sup>a</sup>  | 0.26 | 0.56 <sup>g</sup>  | 0.5 |
| 5        | 5 | 90.2 <sup>d</sup>  | 2.6  | 245.5 <sup>cd</sup> | 26.7 | 9.6 <sup>b</sup>  | 1.4 | 1.49 <sup>c</sup>  | 0.16 | 0.6 <sup>f</sup>   | 0.1 |
| 6        | 5 | 87.6 <sup>e</sup>  | 4.8  | 178 <sup>i</sup>    | 13.7 | 7.2 <sup>c</sup>  | 0.8 | 1.05 <sup>e</sup>  | 0.32 | 0.75 <sup>d</sup>  | 0.7 |
| 7        | 5 | 85.5 <sup>f</sup>  | 5.4  | 135 <sup>j</sup>    | 8.9  | 7.8 <sup>c</sup>  | 0.8 | 2.15 <sup>a</sup>  | 0.24 | 0.95 <sup>b</sup>  | 0.2 |
| 8        | 5 | 80.2 <sup>g</sup>  | 3.2  | 225 <sup>e</sup>    | 15.5 | 5 <sup>d</sup>    | 0.8 | 1.35 <sup>cd</sup> | 0.06 | 0.53 <sup>h</sup>  | 0.3 |
| 9        | 5 | 96.4 <sup>c</sup>  | 5.4  | 265 <sup>b</sup>    | 11.3 | 5.9 <sup>d</sup>  | 0.5 | 1.12 <sup>e</sup>  | 0.21 | 0.45 <sup>i</sup>  | 0.2 |
| 10       | 5 | 78.2 <sup>h</sup>  | 2.1  | 200 <sup>f</sup>    | 12.6 | 7.7 <sup>c</sup>  | 0.8 | 1.07 <sup>e</sup>  | 0.07 | 0.82 <sup>c</sup>  | 0.9 |
| 11       | 5 | 74 <sup>i</sup>    | 1.8  | 195 <sup>g</sup>    | 10.8 | 7.4 <sup>c</sup>  | 0.5 | 1.48 <sup>c</sup>  | 0.11 | 0.46 <sup>i</sup>  | 0.2 |
| 12       | 5 | 112.5 <sup>b</sup> | 17.8 | 175 <sup>i</sup>    | 8.7  | 7.5 <sup>c</sup>  | 0.4 | 1.35 <sup>cd</sup> | 0.14 | 0.55 <sup>gh</sup> | 0.5 |

There is a statistically significant difference ( $P < 0.001$ ) between the mean values (X), which are indicated with different letters in each column.

Table 5. Mean values and standard deviations of essential (Zn, Fe, Cu) and toxic (Pb, Cd) metal levels in soil samples from different districts of Northwest Turkey

| District | N | Zn (µg/g)          |     | Fe (µg/g)           |      | Cu (µg/g)         |     | Pb (µg/g)          |      | Cd (µg/g)          |     |
|----------|---|--------------------|-----|---------------------|------|-------------------|-----|--------------------|------|--------------------|-----|
|          |   | X                  | Sx  | X                   | Sx   | X                 | Sx  | X                  | Sx   | X                  | Sx  |
| 1        | 5 | 58.7 <sup>b</sup>  | 2.1 | 162.5 <sup>fg</sup> | 10.8 | 13.7 <sup>j</sup> | 1.6 | 1.16 <sup>g</sup>  | 0.4  | 0.66 <sup>e</sup>  | 0.2 |
| 2        | 5 | 61.1 <sup>a</sup>  | 4.2 | 159.4 <sup>g</sup>  | 9.6  | 39.7 <sup>b</sup> | 2.7 | 2.46 <sup>a</sup>  | 0.2  | 0.88 <sup>c</sup>  | 0.4 |
| 3        | 5 | 41.5 <sup>cd</sup> | 5.8 | 195.7 <sup>c</sup>  | 12.8 | 20.9 <sup>h</sup> | 1.3 | 2.39 <sup>ab</sup> | 0.3  | 0.21 <sup>j</sup>  | 0.4 |
| 4        | 5 | 38.8 <sup>ef</sup> | 3.6 | 125.4 <sup>k</sup>  | 11.3 | 26.8 <sup>e</sup> | 1.6 | 1.76 <sup>de</sup> | 0.1  | 1.23 <sup>b</sup>  | 0.8 |
| 5        | 5 | 35.5 <sup>g</sup>  | 5.1 | 225.5 <sup>b</sup>  | 21.4 | 35.4 <sup>c</sup> | 2.1 | 0.98 <sup>h</sup>  | 0.2  | 0.52 <sup>gh</sup> | 0.3 |
| 6        | 5 | 41.4 <sup>cd</sup> | 4.2 | 149.5 <sup>h</sup>  | 11.6 | 27.8 <sup>e</sup> | 1.3 | 1.45 <sup>f</sup>  | 0.1  | 0.48 <sup>h</sup>  | 0.4 |
| 7        | 5 | 40.5 <sup>de</sup> | 5.4 | 135.8 <sup>j</sup>  | 14.3 | 32.5 <sup>d</sup> | 2.4 | 1.87 <sup>d</sup>  | 0.1  | 1.25 <sup>b</sup>  | 0.5 |
| 8        | 5 | 36.9 <sup>fg</sup> | 4.2 | 165.3 <sup>f</sup>  | 18.7 | 39.5 <sup>b</sup> | 2.1 | 2.15 <sup>c</sup>  | 0.1  | 0.35 <sup>i</sup>  | 0.4 |
| 9        | 5 | 38.5 <sup>ef</sup> | 3.1 | 145.4 <sup>i</sup>  | 14.8 | 17.8 <sup>i</sup> | 1.1 | 1.74 <sup>e</sup>  | 0.09 | 0.78 <sup>d</sup>  | 0.5 |
| 10       | 5 | 42.7 <sup>cd</sup> | 3.2 | 178.3 <sup>d</sup>  | 10.4 | 25.4 <sup>f</sup> | 1.3 | 2.28 <sup>b</sup>  | 0.1  | 0.54 <sup>fg</sup> | 0.7 |
| 11       | 5 | 40.5 <sup>de</sup> | 4.2 | 256.2 <sup>a</sup>  | 11.4 | 41.6 <sup>a</sup> | 2.6 | 2.05 <sup>c</sup>  | 0.1  | 1.45 <sup>a</sup>  | 0.6 |
| 12       | 5 | 43.4 <sup>c</sup>  | 4.2 | 174.6 <sup>e</sup>  | 9.8  | 22.5 <sup>g</sup> | 1.6 | 0.94 <sup>h</sup>  | 0.07 | 0.58 <sup>f</sup>  | 0.4 |

There is a statistically significant difference ( $P < 0.001$ ) between the mean values (X), which are indicated with different letters in each column.

Table 6. Correlation between essential (Zn, Fe, Cu) and toxic (Pb, Cd) metal levels in water, soil and plant samples collected from different districts of Northwest Turkey

|       | Zn (n = 60)                 | Fe (n = 60)    | Cu (n = 60)                 | Pb (n = 60)             | Cd (n = 60)              |
|-------|-----------------------------|----------------|-----------------------------|-------------------------|--------------------------|
| Water | 117.41 ± 49.0**             | 29.16 ± 8.89   | 42.25 ± 15.56**             | 2.56 ± 1.13             | 0.63 ± 0.36              |
| Plant | 91.01 ± 28.59 <sup>++</sup> | 214.05 ± 71.06 | 10.55 ± 11.26 <sup>++</sup> | 1.44 ± 0.4 <sup>+</sup> | 0.67 ± 0.36              |
| Soil  | 43.27 ± 7.91 <sup>00</sup>  | 172.79 ± 36.37 | 28.63 ± 8.88                | 1.76 ± 0.5              | 0.74 ± 0.37 <sup>0</sup> |

\*Water-Plant, <sup>0</sup>Water-Soil, <sup>+</sup>Plant-Soil, \* $P < 0.05$ , \*\*  $P < 0.01$ . (<sup>0</sup>, <sup>+</sup> similar)

Soil analyses results produced the lowest zinc levels for the 5<sup>th</sup> and 8<sup>th</sup> districts. In the 4<sup>th</sup> district a dramatically low iron level was detected. The lowest copper levels were found in the 1<sup>st</sup> and 9<sup>th</sup> districts. Lead levels were higher in the 2<sup>nd</sup>, 3<sup>rd</sup>, 8<sup>th</sup> and 10<sup>th</sup> districts, while cadmium levels were somewhat elevated in the 4<sup>th</sup>, 7<sup>th</sup> and 11<sup>th</sup> districts.

## Discussion

When the number of animals with economical value is compared with those of other countries, Turkey is placed rather high up in the list. However, it would be closer the bottom of the list if products such as meat, milk, wool and leather obtained from these animals are considered (AĞAOĞLU, 1991; ÇAMAŞ et al., 1994). In this respect, deficiencies of trace elements play a role which is as important as infectious and parasitary diseases (UNDERWOOD, 1997). A similar condition became obvious in northwest Turkey, where the study was carried out. After interviews with animal breeders and veterinary surgeons it emerged that the level of diseases due to parasitary and trace element deficiencies was very high. In particular, animal health problems such as abortion, trouble in walking in new-born lambs, fertility problems, weight loss, lack of putting on weight despite proper rations, diarrhoea, growth disorders, susceptibility to infectious diseases and decrease in quality of wool were in conformity with data reported in the literature (ALTINTAŞ et al., 1990).

While NELSON et al. (1987) reported serum zinc levels of 0.32-0.40 µg/ml in a flock which suffered from cases of anorexia, wool eating, alopecia, hyperkeratosis and parakeratosis, ÇAMAŞ et al. (1994) have reported a level of 25-48 µg/dL for similar symptoms. Our zinc values were within the above levels, with the exception of the 1<sup>st</sup> and 7<sup>th</sup> districts. The low levels found only in the 1<sup>st</sup> and 7<sup>th</sup> districts, and especially in Babaeski, Lüleburgaz and Pehlivan köyü, are an indication of the problem. In particular, the zinc level in those districts was found to be lower in water and plant samples as compared to other districts. This indicates that sheep there do not benefit from water and plants as a source of zinc. Moreover, we are of the opinion that high cadmium levels in plant samples from those districts are associated with zinc deficiency, as has been suggested by researchers (MARTIN et al., 1983; AĞAOĞLU, 1991).

While normal serum copper levels for sheep as reported by various researchers (MARTIN et al., 1983; GHOSAL and MATHUR, 1992) are 59-101 µg/dL and 80-120 µg/dL, respectively, the lowest values determined in our studies prove how serious the problem is in northwestern Turkey. The correlation between blood Cu levels below 50 µg/dL level and the increased risk of enzootic ataxia risk constitutes an important threat to the region where this study was carried out (WILSON et al., 1991). To increase this risk, zinc levels in plant sheep grazing areas in the 1<sup>st</sup> and 12<sup>th</sup> districts were high, while copper levels were low in plants of some districts (3<sup>rd</sup>, 8<sup>th</sup> and 9<sup>th</sup>). Clinical signs such as depigmentation, weight loss, growth retardation, diarrhoea and ataxia due to low blood serum level of trace elements which correlated with the low levels found in soil, plant and water samples, conform to previously reported results (FERM and HANLON, 1974; ERGÜN, 1983; WILSON et al., 1991).

For normal sheep UNDERWOOD (1997) stated that serum iron levels should be in the order of 102-304 µg/dL, while NAZKI and RATTAN (1990) suggested a level of 115-234

$\mu\text{g/dL}$  for the same trace element. Although serum iron level below  $40 \mu\text{g/dL}$ , as reported by GHOSAL and MATHUR (1992), was not observed in any district of the region, in our opinion, the existence of a potential risk for the region should not be disregarded. In this connection, it is well known that lack of the trace element copper together with deficiency in the trace element iron leads to serious anaemia, which in consequence results in predisposition to parasitary diseases. It has been reported (PIRKLE et al., 1985; AĞAOĞLU, 1991) that high levels especially of cobalt, zinc, cadmium, manganese and copper were reducing the absorption of iron. In parallel with industrial development in the districts of Tekirdağ, Keşan and Kırklareli, increased levels of lead and cadmium in sheep have been observed in those districts and their rations correlated with their content in plants in the region. High lead levels in sheep, especially in the districts of Edirne, Kırcaşalih and their surroundings, and the level of the same element in plants from Çorlu, Çerkezköy and Lüleburgaz districts constitute the undesirable side effects of industrialisation such as environmental pollution, serious metallic toxication risks and negative effects on trace elements. As has been reported (MEREDITH, 1978; KUMARESAN and KAPIOH, 1984), the presence of excessive levels of lead and cadmium negatively affect primarily the hemopoietic system but also the urogenital system, the gastrointestinal system and the neurologic system, as well as the absorption of other trace elements.

We are of the opinion that by informing veterinary surgeons in the region about the results of soil, water and plant analyses, they would be in a position to prescribe supplements of those elements where deficient; the results would also contribute to the treatment of abortus cases which are regularly observed and which are not due to any infectious or parasitary diseases. And finally, the results could help to improve quality of wool.

Similar studies within the scope of the existing possibilities in the same region comprising other trace elements, as well as extended studies in other regions, would help to determine the levels of trace elements in soil, water and plants everywhere in Turkey and would thus provide an important tool to reduce incidences of disease, morbidity and mortality ratios.

This study, which found important differences in trace element levels in the investigated region, allows a certain generalisation as to the solution of problems regarding sheep breeding, which is an important contributor to the country's economy in the form of meat, milk, wool and leather, with respect to the effects of environmental factors.

If the results of this study are communicated to animal breeders and to local veterinary surgeons, they may, in our opinion, contribute to important developments both in treatment and control of diseases directly related to the lack of trace elements, as well as in the treatment and control of parasitary, bacterial and viral diseases caused as a consequence of trace element deficiency.



M. E. Or et al.: Determination of levels of some essential (iron, copper, zinc) and toxic (lead, cadmium) metals in the blood of sheep and in samples of water, plants and soil in Northwest Turkey

### Acknowledgement

This work was supported by the Research Fund of The University of Istanbul (Project Number: UDP-62/15082002) and TUBITAK (Project Number: VHAG- 1482).

### References

- AĞAOĞLU Z. T. (1991): Ülkemiz hayvancılığında bazı iz elementler ve önemleri. Veteriner Hekimler Vakfı Dergisi, 57-62.
- ALTINTAŞ, A., H. UYSAL, S. YILDIZ, T. GONCAGÜL (1990): Akkaraman ve melezlerinde serum ve yapağı örneklerinde karşılaştırmalı mineral durumu. Lalahan Hayv. Araş. Derg. 30, 40-57.
- BROWN, A., J. D. HALLS, A. TAYLOR (1986): Atomic spectrometry update-clinical materials, foods and beverages. J. Anal. Atom. Spect. 1, 21-35.
- BROWN, A. A., A. TAYLOR (1995): Applications of a slotted quartz tube and flame atomic absorption spectrophotometer to the analysis of biological samples. Analyst. 110, 579-582.
- ÇAMAŞ, H., A. BİLDİK, F. GÜLSER (1994): Toprak, bitki ve koyunların kanında bazı iz elementlerle (Cu, Mo, Zn, Co, Mn) Sülfat (SO<sub>4</sub>) miktarlarının araştırılması. Pro.no: VHAG-966. Van.
- DUNCAN, B. D. (1955): Multiple Range and Multiple F - tests. Biometrics 11, 1-42.
- ERGÜN, A.(1983): Zinc metabolism and deficiency in domestic animals. Vet. J. Ankara Univ. 30, 308-316.
- FERM, V. H., D. P. HANLON (1974): Placental transfer of zinc in the syriyan hamster during early embriyogenesis, J. Reprod. Fert. 39, 49-52.
- GHOSAL, A.K., G. N. MATHUR (1992): Zinc, copper and iron contents of blood serum of cattle, sheep in semiarid Tract of Rjasthan. Indian J. Anim. Sci. 62, 441-442.
- KUMARESAN, A., M. A. KAPIOH (1984): Hair as indicator of mineral status in yankassa sheep, Rew. Elev. Med. Vet. Pays Trop. 37, 61-64.
- MARTIN, D. W., P. A. MAYES, V. W. RODWELL (1983): Harper's Review of Biochemistry. 19<sup>th</sup> Edition Lance Medical Publications, Los Altos, California.
- MEREDITH, P. A. (1978): Delta aminolevulinic acid metabolism in normal and exposed Humans. Toxicology. 19, 1-9.
- NAZKI, A. R., J. S. RATTAN (1990): Status of blood micro-element during different seasons in sheep. Ind. Vet. J. 67, 274-276.
- NELSON, D. R., W. A. WOLFF, D. J. BLODGETT, B. LEUCKE, R. W. ELY, J. F. ZACHARY (1987): Zinc deficiency in sheep and goats: three field cases. J. Am. Vet. Med. Assoc. 184, 1480-1485.
- OZAN, S. (1985): The relation between fleece losses and zinc, copper levels in blood serum of Merine Sheep in Karacabey State Farm Flock. Vet. J. Selçuk Univ. 1, 133-142.
- PIRKLE J. L., J. SCHWARTZ, R. LANDIS, R. W. HARLAN (1985): The relationship between blood lead and blood pressure and its cardiovascular risk implications. Am. J. Epidemiol. 121, 246-258.

M. E. Or et al.: Determination of levels of some essential (iron, copper, zinc) and toxic (lead, cadmium) metals in the blood of sheep and in samples of water, plants and soil in Northwest Turkey

UNDERWOOD, E. J. (1997): Trace Elements in Human and Animal Nutrition. Academic Press, London.

WILSON J. D., E. BRAUNWALD, K. J. ISSELBACHER, R. G. PETERSDORF, J. B. MARTIN, A. S. FAUCI, R. K. ROOT (1991): Principles of Internal Medicine. Twelfth Edition 1, 443.

YILDIZ, G., K. KÜÇÜKERSAN, S. KÜÇÜKERSAN (1995): Yapağı dökmeye ve yapağı yemeye semptomları gösteren akkaraman koyunlarda kan serumu ve yapağıda meydana gelen mineral madde değişimi. Vet. J. Ankara Univ. 42, 251-256.

ZANTOPOULOS, N., V. ANTONIOU, E. NIKOLAIDIS (1999): Copper, zinc, cadmium, and lead in sheep grazing in North Greece. Bull. Environ. Contam. Toxicol. 62, 691-699.

Received: 31 October 2003

Accepted: 28 June 2005

---

**OR, M. E., A. KAYAR, A. R. KIZILER, Ç. PARKAN, R. GÖNÜL, B. BARUTÇU, H. T. DODURKA: Određivanje razine nekih esencijalnih (željezo, bakar, cink) i toksičnih (olovo, kadmij) metala u krvi ovaca te u uzorcima vode, biljaka i tla u sjeverozapadnoj Turskoj. Vet. arhiv 75, 359-368, 2005.**

**SAŽETAK**

Cilj istraživanja bio je odrediti razinu elemenata u tragovima (cink, željezo, bakar), te olova i kadmija u tlu i organizmu ovaca kako bi se polučili bolji gospodarski rezultati i uzgajale zdravije pasmine ovaca u sjeverozapadnoj Turskoj. Na temelju rezultata, za navedena područja planiraju se preporuke veterinarima i uzgajivačima na osnovi kojih bi se unaprijedio njihov rad. U istraživanje je bilo uključeno 400 ovaca iz različitih područja sjeverozapadne Turske. Uzimani su uzorci vode (n = 5), biljaka (n = 5) i tla (n = 5). Utvrđena je značajno niža razina željeza u 8., 9. i 10. području. Najniža razina cinka ustanovljena je u 1. i 7. području, najniža razina bakra u 1., 3., 5. i 6. području, dok su najviše razine olova utvrđene u 9. i 10. području, a kadmija u 2., 5. i 6. području. Analiza uzoraka vode, biljaka i tla pokazala je da između različitih područja postoje značajne razlike u razini cinka i bakra. Visoke razine olova ustanovljene su u uzorcima vode i biljaka. Prijenosom ovih rezultata do veterinarima i uzgajivača iz promatranih područja, ostvarit će se značajna poboljšanja u liječenju i kontroli bolesti uzrokovanih deficijencijom elemenata u tragovima. Isto se može očekivati i za bolesti uzrokovane parazitima, bakterijama i virusima u kojih navedene deficijencije imaju određenu ulogu.

**Glavne riječi:** ovaca, elementi u tragovima, cink, željezo, bakar, olovo, kadmij

---