

Revised Middle Miocene datum for initial marine flooding of North Croatian Basins (Pannonian Basin System, Central Paratethys)



Stjepan Ćorić¹, Davor Pavelić², Fred Rögl³, Oleg Mandić³, Sejfudin Vrabac⁴, Radovan Avanić⁵, Lazar Jerković⁶ and Alan Vranjković²

¹ Geological Survey of Austria, Neulingg. 38, A-1030 Wien, Austria; (stjepan.coric@geologic.ac.at)

² Faculty of Mining, Geology and Petroleum Engineering, University of Zagreb, Pierottijeva 6, HR-10000 Zagreb, Croatia; (dpavelic@rgn.hr, avranjko@rgn.hr)

³ Geological-Palaeontological Department, Natural History Museum Vienna, Burgring 7, A-1010 Wien, Austria; (fred.roegl@nhm-wien.ac.at, oleg.mandic@nhm-wien.ac.at)

⁴ Department of Geology, Faculty of Mining, Geology and Civil Engineering, University of Tuzla, Univerzitetska 2, BIH-75000 Tuzla, Bosnia and Herzegovina; (jvrabac@yahoo.com)

⁵ Croatian Geological Survey, Sachsova 2, P.O.Box 268, HR-10000 Zagreb, Croatia; (radovan.avanic@hgi-cgs.hr)

⁶ Faculty of Sciences, University of Podgorica, 81000 Podgorica, Montenegro; 202-444 rue Poudriere, Verdun, PQ, H4G 3L9, Canada; (dobj@estart.com)

Geologia Croatica

ABSTRACT

The Pannonian Basin System (PBS) originated during the Early Miocene as a result of extensional processes between the Alpine-Carpathian and the Dinaride Orogenic Belts. The Paratethys Sea flooded the new basins successively during the Karpatian (late Burdigalian, Early Miocene) and the Early Badenian (middle Langhian, Middle Miocene). The North Croatian Basins (NCB) occupied the south-western margin of the PBS and the Central Paratethys Sea. Their initial marine flooding has until now been dated as Karpatian in age. The transgression into the NCB invaded a lacustrine environment, representing the northern prolongation of the vast Dinaride Lake System extending southwards as far as the Adriatic Plate. We reinvestigate two sections from opposite margins of the NBS – from Mt. Medvednica in the west and from Mt. Požeška in the east, including the corresponding lowermost marine Miocene deposits, in order to critically examine the Karpatian datum. Our new biostratigraphic data, integrating calcareous nannoplankton, planktic and benthic foraminifera, diatom and mollusc records, have substantially revised the previous interpretation. The presence of a calcareous nannoplankton assemblage of the NN5 Zone and the planktic and benthic foraminifera of the regional Lower Lagenidae Zone now place the transgression into the main Early Badenian transgressive pulse of Central Paratethys. Consequently, the initial marine transgression correlates accurately with the middle part of the Early Badenian, which is more than 2 m.y. younger than the previously inferred datum, and at least 1 m.y. younger than the lower boundary of the Badenian and the Middle Miocene, respectively. Finally, the basal lacustrine infill of the NCB, previously dated as Otnangian (middle Burdigalian, Early Miocene) and continuously grading into marine deposits, has also to be reconsidered as Early Badenian.

Keywords: Pannonian Basin System, North Croatian Basins, Central Paratethys, Dinaride Lake System, Middle Miocene, biostratigraphy

1. INTRODUCTION

North Croatian Basins (NCB) occupies the south-western margin of the Pannonian Basin System (Fig. 1). Their basal infill comprises freshwater sediments with fluvial deposition, grading upwards into open lake deposits. The marine

transgression intruded into the basins during the open lake phase, as suggested by an unchanged lithology across the transition. The boundary can be traced exclusively by the first occurrence of marine fauna such as planktic and benthic foraminifera, indicating a salinity increase and an established connection to the open sea. The age inference of the bound-



Figure 1: Position of the North Croatian Basins in the Pannonian Basin System and the Central Paratethys Sea (after PAVELIĆ et al., 2003).

ary was determined solely on the interpretation of an impoverished, species-poor foraminiferal assemblage that has been originally regarded to correlate with Karpatian assemblages from the northern and western Central Paratethys (ŠIKIĆ, 1968) (Fig. 2).

The lake deposits of the basal infill bear freshwater molluscs and ostracods absent from Central Paratethys and endemic to the Dinaride Neogene basins adjoining the NCB to the south (KOCHANSKY-DEVIDÉ & SLIŠKOVIĆ, 1978; KOCHANSKY-DEVIDÉ, 1979; KRSTIĆ et al., 2003). Such a faunal composition of the basal parts of the NCB classifies it with the palaeobiogeographic entity termed the Dinaride Lake System (DLS) (KRSTIĆ et al., 2003; HARZHAUSER & MANDIĆ, 2008a, b). Consequently DLS includes the basal NCB together with the Dinaride Neogene basin fresh water deposits. The dating of the DLS deposits is still subject to debate, especially due to a lack of the possibility for marine biostratigraphy (HARZHAUSER & MANDIĆ, 2008a, b). The original correlation of the basal DLS deposits with the Otnangian, is rooted in the interpretation of a small characean flora combined with their stratigraphic position directly below the formerly Karpatian-dated deposits (ŠIKIĆ, 1968; KOCHANSKY-DEVIDÉ & SLIŠKOVIĆ, 1978). As will be shown by new data, the latter age inference turned out, however, to be incorrect.

This paper critically examines the datum for the initial marine flooding of the NCB by means of integrative bios-

stratigraphy, combining data on calcareous nannoplankton, foraminifera, molluscs and diatoms. This yields a more precise dating of the geodynamic history and rifting of the Pannonian Basin System (PBS). It also provides more insight into the palaeoenvironmental turnover in the south-western margin of the PBS resulting from the back-stepping of the DLS and the landward progradation of the Central Paratethys Sea.

2. HISTORY OF INVESTIGATION

PAUL (1872, 1874) was the first author to consider the basal freshwater deposits exposed at Mt. Medvednica N of Zagreb, as temporal equivalents of the Socka beds in NE Slovenia. Accordingly, he dated them as Late Oligocene. GORJANOVIĆ-KRAMBERGER (1908), setting the initial milestone in the development of stratigraphic nomenclature for the North Croatian Basin deposits, followed Pauls opinion. Furthermore, he referred the superposed sediments, representing the initial marine transgression into the basins, to “Early Mediterranean” based on correlation with the Eggenburgian strata of Lower Austria. Finally, the younger marine strata on top of the latter deposits were dated as “Younger Mediterranean” (Fig. 3).

POLIĆ (1935) advocated the earlier correlation of freshwater deposits. Based on the terrestrial macroflora he also suggested an Oligocene age. KOCHANSKY-DEVIDÉ (1944) provided the first extensive biostratigraphic investigation on the macro- and micro-fauna from marine deposits of the Mt. Medvednica environs. Following the advance in regional stratigraphic nomenclature, she referred the “Early Mediterranean beds” to the Burdigalian and the “Younger Mediterranean beds” to the Helvetian and Tortonian (note, however, the historical misuse of Helvetian and Tortonian in Central Paratethys as discussed by HARZHAUSER et al., 2003).

The intensive application of micropalaeontology for solving the stratigraphy of the NCB goes back to the 1960s. MULDINI-MAMUŽIĆ (1965) interpreted the ostracod record as support for the Late Oligocene age of the basal freshwater deposits. Furthermore, the foraminifera seemed to indicate Oligocene and Burdigalian age for the initial marine beds. This was discussed in detail and substantially revised by ŠIKIĆ (1966, 1968), ultimately establishing the currently applied stratigraphic concept for the marine part of the NCB infill (PAVELIĆ et al., 2003). Based on foraminiferal distributions, the lowermost marine beds were correlated with the Karpatian (therein also termed the Late Helvetian) beds from the Alpine Carpathian Foredeep and the western and north-western Pannonian Basin System. Furthermore, the underlying freshwater deposits bearing Charophyta related to the Upper Freshwater Molasse in Bavaria are referred to as being Helvetian s.str. and Otnangian in age (Fig. 3).

Subsequently, the Otnangian correlation was adopted by KOCHANSKY-DEVIDÉ & SLIŠKOVIĆ (1978). They regarded the freshwater deposits of the NCB as the initial deposits of the “Early lacustrine beds with congerians”, currently termed the Dinaride Lake System (KRSTIĆ et al., 2003; HARZHAUSER & MANDIĆ, 2008a, b). Additionally, at

M. A.	EPOCH	AGE	REGIONAL STAGES	CALCAREOUS NANNOPLANKTON	
15	MIOCENE Middle	SERRAVALLIAN 13,65	SARMATIAN	NN6	
		LANGHIAN 15,97 ?	BADENIAN	NN5	
		BURDIGALIAN	KARPATIAN	NN4	
	20	MIOCENE Early	EGGENBURGIAN 20,43	OTTNANGIAN	NN3
			AQUITANIAN 23,03	EGGERIAN	NN2
			CHATTIAN	EGGERIAN	NN1
25	OLIGOCENE Late	KISCELLIAN	EGGERIAN	NP25	
			KISCELLIAN	NP24	

Figure 2: Oligocene–Middle Miocene chronostratigraphy of the Central Paratethys correlated to the standard geochronologic scale and the calcareous nonnoplankton zone of MARTINI (1971) (simplified, after PILLER et al., 2007). Note the correlation of the Middle Miocene boundary after GRADSTEIN et al. (2004), defining it with the base of the palaeomagnetic chron C5Br. In contrast, biostratigraphically, the Middle Miocene boundary is defined by the First Appearance Datum for the planktonic foraminiferal genus *Praeorbulina* at 16.3 Ma, as indicated by RÖGL et al. (2007).

Deposition	GORJANOVIĆ-KRAMBERGER (1908)	KOCHANSKY (1944)	ŠIKIĆ (1968)	KOCHANSKY-DEVIDÉ & SLIŠKOVIĆ (1978)	KOCHANSKY-DEVIDÉ (1979)	This paper
Marine	II. Mediterranean	Tortonian	Late Tortonian	Ottngangian	Karpatian	Early Badenian
		Helvetian, "Schlier"	Early Tortonian (previous Helvetian "Schlier" and Čučerje beds)			
	Older Mediterranean	Burdigalian	Karpatian (Late Helvetian)			
Fresh-water	Late Oligocene		Helvetian s. str. (Early Helvetian)		Ottngangian	

Figure 3: Milestones in the history of stratigraphic correlation of the North Croatian Basins.

Mt. Požeška, besides an Ottngangian age a Karpatian age has also been presumed for the upper part of the basal, lacustrine NCB deposits. This was based on the more advanced evolutionary stage of the recorded dreissenid bivalves (KOCHANSKY-DEVIDÉ, 1979).

Given the above considerations, further palaeontological studies (as a part of a geological mapping effort), assigned the freshwater deposits to the Lower Helvetian, i.e. Ottngangian, and the first marine Miocene sediments to the Upper Helvetian, i.e. Karpatian (BASCH, 1983; JAMIČIĆ et al., 1987, 1989; ŠPARICA & BUZALJKO, 1984; ŠPARICA et al., 1980). Analyses of freshwater ostracods suggested a "Middle Miocene", "Helvetian" and Karpatian age (SOKAČ, 1987; SOKAČ & KRSTIĆ, 1987). More modern studies of terrestrial flora from the freshwater deposits indicated an Ottngangian age (JUNGWIRTH & ĐEREK, 2000), as did the mollusc associations (Unionidae) from some localities in North Croatia (ŽAGAR-SAKAČ, 2003, 2004).

Complex sedimentological studies, micropalaeontological and palynological analyses dated these freshwater deposits as Ottngangian, and the overlying marine sediments as Karpatian on Mounts Požeška, Mt. Medvednica and Mt. Kalnik (AVANIĆ et al., 1995a, 1995b, 1995c; AVANIĆ, 1997; PAVELIĆ et al., 1998, 2001, 2003) (Fig. 1). The same concept was applied in some regional works (LUČIĆ et al., 2001; PAVELIĆ, 2001, 2002, 2005; SAFTIĆ et al., 2003). A list of important foraminifera and mollusc species of these Croatian initial marine deposits, attributed to the Karpatian, has been compiled by BAJRAKTAREVIĆ & PAVELIĆ (2003). More recent research on ostracods at several localities in North Croatia indicates an Ottngangian age for the freshwater deposits, a latest Ottngangian to earliest Karpatian age for the brackish-water sediments, and a Karpatian age for the marine deposits (HAJEK-TADESSE, 2006).

Based on these former results, all currently available reconstructions of the general evolution of Central Paratethys presumed erroneously a vast marine cover over the NCB during the Karpatian (RÖGL & STEININGER, 1983; PAVELIĆ, 2001, 2005; BAJRAKTAREVIĆ & PAVELIĆ, 2003; RÖGL, 1996; KOVÁČ et al., 2003). Only very few studies suggested a younger age for those freshwater and early marine deposits. One such example is the palynological-based study by KRIZMANIĆ (1995). It related the bentonite clays contain-

ing remains of *Prodeinotherium bavaricum* KAUP on Mt. Moslavačka to the Miocene climatic optimum, specifically correlating them with the time-span between the Late Karpatian and the Early Badenian. The overlying marine beds bearing foraminiferal associations of the Upper Lagenidae Zone and the calcareous nannoplankton association of NN5 were previously dated as Early Badenian (BAJRAKTAREVIĆ, 1981; KRIZMANIĆ, 1995).

3. GEOLOGICAL SETTING

The Paratethys Sea evolved in the Early Oligocene at the northern margin of the pre-Mediterranean Tethys Ocean after Alpine and Dinaride orogenesis triggered the first disturbance of water mass circulation patterns. This was followed by faunal extinction and endemic events (RÖGL, 1996). The distribution of the Paratethys Sea was mainly controlled by regional geodynamic evolution, resulting in a separate chronostratigraphic scale (Fig. 2). The Western Paratethys of the Alpine Foredeep had already dried out in the Early Miocene. The Central Paratethys of the Alpine-Carpathian Foredeep and the Pannonian Basin System disintegrated at the end of the Middle Miocene, giving rise to the Late Miocene Lake Pannon. The disintegration of Eastern Paratethys, today divided into the Black Sea, Caspian Sea and Aral Lakes, is still ongoing.

The Dinaride Lake System (DLS) developed during the Early Miocene on the land block located between Central Paratethys and the Mediterranean Tethys (MANDIĆ et al., 2008). It was inhabited by highly diversified but almost completely endemic aquatic organisms such as molluscs or freshwater ostracods (KRSTIĆ et al., 2003; HARZHAUSER & MANDIĆ, 2008a, b). It represented, as shown by HARZHAUSER & MANDIĆ (2008b), a separate palaeobiogeographic entity, different from any marginal environment of Paratethys or of the Mediterranean Tethys. Extending originally into the Pannonian Basin System, the DLS retreated mainly due to Paratethys Sea transgression, becoming mainly restricted to the Dinaride Block.

North Croatia occupies the south-western margin of the Pannonian Basin System (PBS). The PBS, induced by extensional processes and rifting between the Carpathian, Alpine and Dinaride Chains, did not appear to exist prior to the

4. RESULTS

Two sections representing the initial marine NCB deposits have been studied in detail. The Čučerje Section is located on the south-eastern slope of Mt. Medvednica, about 10 km NE of the city of Zagreb. The Sokolovac Section is located on the southern slope of Mt. Papuk, about 1 km SSW of the village Gornji Vrhovci and 15 km NNW of the city of Požega (Fig. 1). The micropalaeontological samples comprised nanoplankton, diatoms, as well as planktonic and benthic foraminifera. Additionally, for Čučerje (Fig. 1), the mollusc macrofauna from the Coll. Kochansky collection (Croatian Natural History Museum) has been taxonomically revised and biostratigraphically reevaluated.

4.1. Čučerje Section (Mt. Medvednica)

The Čučerje Section (Fig. 5) shows a 42 m thick succession composed of 3 parasequences. The base of the section is at GPS N 45° 53' 45.5" E 16° 03' 17.3". For sample positions see Fig. 5. The section was initially described by GORJANOVIĆ-KRAMBERGER (1923) as the "Plaz" locality. KOCHAN-SKY-DEVIDÉ (1944) described it as a large outcrop above the brook SW of Podplaz and NW of the church of Čučerje. She attributed its deposits, based on micro- and macrofaunal content, to the Burdigalian, correlating them with the lower Ottnangian beds from Lower Austria. Although she originally noticed the conspicuous difference in faunal composition to the latter beds, in a subsequent publication she erroneously identified one Eggenburgian–Ottngian index fossil from the same locality (KOCHAN-SKY-DEVIDÉ, 1956). ŠIKIĆ (1968) investigated foraminifera samples from that locality, attributing them to the Karpatian. Finally, AVANIĆ et al. (1995b) (see also AVANIĆ, 1997) provided a detailed description and interpretation of the depositional history for the section. Although already correlating its middle and upper parts with the Lower Badenian, the authors still referred its lower parasequence to the Karpatian.

4.1.1. Sedimentology

The lower parasequence is 15 m thick and represents a coarsening upward succession with marls at the base followed by predominantly silt deposition on top, intercalated with thin sandy layers. The coarsening has been interpreted as an environmental shift and shallowing, grading from shelf conditions up to the transitional zone close to the palaeoshore. The middle parasequence comprises partly bedded sands and sandstones representing a high-energy, shallow-water shoreface to nearshore depositional environment. The upper parasequence superposes a tephra layer at 36 m up of the section. It reflects renewed deepening of the environment, comprising predominantly silt deposition.

4.1.2. Calcareous nanoplankton biostratigraphy

Ten samples (Fig. 5) have been investigated for the biostratigraphic evaluation of calcareous nanoplankton.

The assemblages from this section are dominated by high percentages of helicoliths and small reticulofenestrids (Table 1). Helicoliths are represented by: *Helicosphaera car-*

teri (WALLICH, 1877) KAMPTNER (1954), *H. walbersdorffensis* MÜLLER (1974), *H. wallichii* (LOHMANN, 1902) BOUDREAUX & HAY (1969), and *H. cf. waltrans* THEODORIDIS (1984), whereas the small reticulofenestrids are represented by *Reticulofenestra minuta* ROTH (1970) and *R. haqii* BACKMAN (1978). The accompanying forms are: *Braarudosphaera bigelowii* (GRAN & BRAARUD, 1935) DEFLANDRE (1947), *Coccolithus pelagicus* (WALLICH, 1877) SCHILLER (1930), *Coronosphaera mediterranea* (LOHMANN, 1902) GAARDER (1977), *Cyclicargolithus floridanus* (ROTH & HAY, 1967) BUKRY (1971), *Perforocalcinella fusiformis* BÓNA (1964), *Pontosphaera multipora* (KAMPTNER, 1948) ROTH (1970), *Reticulofenestra pseudo-umbilica* (GARTNER, 1967) GARTNER (1969), *Rhabdosphaera sicca* STRADNER (1963), *Sphenolithus heteromorphus* DEFLANDRE (1953), *S. moriformis* (BRÖNNIMANN & STRADNER, 1960) BRAMLETTE & WILCOXON (1967), *Triquetrorhabdulus milowii* BUKRY (1971), and *Umbilicosphaera jafari* MÜLLER (1974).

The regular occurrence of the zonal marker *Sphenolithus heteromorphus* DEFLANDRE (1953) and the absence of *Helicosphaera ampliaperta* BRAMLETTE & WILCOXON (1967) in all investigated samples allow an identification of nanoplankton Zone NN5 (MARTINI, 1971).

4.1.3. Foraminifera biostratigraphy

The foraminiferal associations were analyzed in samples HR 9, no. 1, no. 2, no. 3, no. 4, and HR 10 covering the lower parasequence and the lowermost 10 m of the section, respectively (Fig. 5). See Table 2 for the complete list of identified species.

The lowermost sample in section (HR 9) originated from grey silty micaceous shale. The rich assemblage is characterized by coarse-grained agglutinated foraminifera, typically Badenian calcareous benthic species, and stratigraphically indifferent planktonic forms. The occurrences of the following microfossils are interesting, *incertae sedis*: *Bachmayerella laqueata* RÖGL & FRANZ, and *Bolboforma moravica* REDINGER. Benthic foraminifera species such as *Uvigerina grilli* SCHMID, *U. macrocarinata* PAPP & TURNOVSKY, *U. uniseriata* JEDLITSCHKA and *Pappina parkeri* (KARRER) are of stratigraphic importance. The planktonic assemblage consists mainly of five-chambered globigerinids (e.g. *Globigerina ottmangiensis* RÖGL) together with several *Globigerinoides* species (e.g. *G. bisphericus* TODD). The uppermost studied sample HR 10, comprises gray-brown sediment residue, with cemented fine-grained quartz and silt and contains a mixed assemblage of shallow and deep-water species. Stratigraphically important taxa occur together in the sample HR9 (Table 2). The poor planktonic fauna is again dominated by small species such as *Globigerina prae-bulloides* BLOW or *Tenuitellinata selleyi* LI, RADFORD & BANNER. Microfossils *incertae sedis* *Bachmayerella tenuis* RÖGL & FRANZ and *Bolboforma* sp. are also present.

The lower part of the section represents a rather deep, probably outer shelf environment as indicated by the presence of agglutinated foraminiferal taxa (*Bathysiphon*, *Hy-*




Figure 5: Studied sections with indicated sample positions.

perammina, *Reophax*, *Reticulophragmium*, *Cyclammina*). A larger sediment input from shallow water is documented by the occurrence of miliolids, elphidiids, *Ammonia*, *Pararotalia*, and *Amphistegina*. *Asterigerinata planorbis* (D'ORBIGNY), *Ammonia viennensis* (D'ORBIGNY), *Cibicoides ungeri-*

anus ungerianus (D'ORBIGNY), and *Amphistegina mam-*
milla (FICHTEL & MOLL) and characterized by a large number of individuals.

Biostratigraphically, this part of the section belongs to the Early Badenian, corresponding with the Lower Lageni-

Table 3: Nannofossil distribution (following ODP terminology) in samples from the Sokolovac Section.

Samples	preservation	abundance	<i>Coccolithus pelagicus</i> (WALLICH, 1877) SCHILLER (1930)	<i>Helicosphaera carteri</i> (WALLICH, 1877) KAMPTNER (1954)	<i>Helicosphaera walbersdorffensis</i> MÜLLER (1974)	<i>Perforacina fusiformis</i> BONA (1964)	<i>Reticulofenestra gelida</i> (GETZENAUER, 1972) BACKMAN (1978)	<i>Reticulofenestra haqii</i> BACKMAN (1978)	<i>Reticulofenestra minuta</i> ROTH (1970)	<i>Reticulofenestra pseudumbilica</i> (GARTNER, 1967) GARTNER (1969)	<i>Umbilicosphaera jafari</i> MÜLLER (1974)	reworked	<i>Watznaueria barnesae</i> (BLACK, 1959) PERCH-NIELSEN (1968)
Sokolovac 7	G	C	f	r	r	r	r	r	c	r			
Sokolovac 6	barren												
Sokolovac 5	barren												
Sokolovac 4	barren												
Sokolovac 3	barren												
Sokolovac 2	barren												
Sokolovac 1	G	F	f	f				r	c		r		
Sokolovac 0	p	R	x	x					x				x
Sokolovac -1	barren												

fossil resting spore determined as *Truncatulus tortonicus* (HAJOS) SUTO. Although the zonal marker *Sphenolithus heteromorphus* DEFLANDRE (1953) was not identified in the samples containing nannofossils (Sokolovac 0, 1 and 7) the composition of the nannoplankton assemblage dominated by *Reticulofenestra minuta* indicates nannoplankton Zone NN5. This biostratigraphic contribution can be confirmed by the presence of *Reticulofenestra pseudumbilica* with diameters from 5 to 7 µm.

Truncatulus tortonicus (HAJOS) SUTO is related to the marine planktonic diatoms *Chaetoceros* EHRENBERG, described by HAJÓS (1986) from the Hungarian Middle Miocene diatomite (locality Szurdokpüspöki). It occurs from the early Middle Miocene diatom Zone NP4 4B to the upper late Pliocene Zone NP9. The genus *Chaetoceros* is an important marine producer in nearshore upwelling regions (SUTO, 2006).

4.2.3. Foraminifera biostratigraphy

Three samples from the lower part of the section have been studied. The samples no. 0 (54) and no. 1 (54) lacked microfossils. Sample no. 7 (55) contained a poorly preserved, but diverse assemblage of 45 species. With the exception of one specimen of *Cassigerinella globulosa* (EGGER), only benthic species are present, dominated by small agglutinated forms. Calcified casts of pteropods (*Spiralis*) are frequent.

The faunal assemblage is characteristic for the Karpatian/Badenian transition. *Uvigerina macrocarinata* PAPP & TURNOVSKY is a marker species for the Early Badenian (Lower Lagenidae Zone).

5. DISCUSSION AND CONCLUSION

This study revises the age for the lowermost marine beds of the North Croatian Basins. These originally Karpatian-dated sediments are now provided with a new age framework. Accordingly, our results based on integrative biostratigraphic studies prove a Middle Miocene, Early Badenian age.

All investigated samples from Čučerje and the uppermost sample from Sokolovac indicate nannoplankton Zone NN5, with the zonal marker species *Sphenolithus heteromorphus* DEFLANDRE (1953). Nannoplankton assemblages with blooms of helicoliths and small reticulofenestrids indicate a nearshore, nutrient-rich environment. This coincides with the results from the benthic and planktic foraminiferal assemblages. *Uvigerina macrocarinata* PAPP & TURNOVSKY, *Pseudogaudryina lapugyensis* (CUSHMAN), *P. sturi* (KARRER), *Amphistegina mammilla* (FICHTEL & MOLL), and *Orbulina suturalis* BRÖNNIMANN characterize the Lower Lagenidae Zone of the Badenian in the Vienna and Styrian Basins (RÖGL et al., 2007). Additionally, the presence of the microfossil *Bolboforma moravica* REDINGER, together with the mass occurrence of the planktonic gastropod "*Spiralis*", confirm the Early Badenian age. The diatom flora with *Truncatulus tortonicus* (HAJOS) SUTO determined from the transitional part of the Sokolovac Section suggests a Badenian age. The pectinid bivalve *Flabelliptecten solarium* (LAMARCK) from the Čučerje Section also indicates a Badenian age.

The assemblages point to a Badenian age correlated with the upper part of the Lower Lagenidae Zone (Fig. 6). The first occurrence of *Orbulina suturalis* BRÖNNIMANN, dated at 14.74 Ma and found at about 5 m up the Čučerje




Figure 6: Stratigraphic time scale and correlation of investigated sections (modified after Rögl et al., 2007).

Section, demonstrates a much higher stratigraphic level for the succession than previously considered. The depositional evolution in both sections suggests a position above the maximal flooding surface and within the High Stand System Tract, which can be biostratigraphically correlated with the 3rd order sequence TB 2.4 (Fig. 6). The age of the marine transgression lies within the lower part of nannoplankton Zone NN5, having its base at 14.91 Ma. This places the initial marine sedimentation in the NCB at least 1 m.y. above the Middle Miocene lower boundary. The position of the Early-Middle Miocene boundary is currently under discussion. Hence, GRADSTEIN et al. (2004) placed it on top of the palaeomagnetic Chron C5Cn, whereas its biostratigraphic definition relates to the First Appearance Datum of the planktic foraminiferal genus *Praeorbulina* (see LOURENS et al., 2004).

The transition from the lake deposits to the marine environment was observed in the Sokolovac Section. There is no sedimentological break between the lacustrine and marine deposits, suggesting continuous sedimentation throughout the Lower Badenian. Based on a time interval of 1 m.y., the resulting mean sedimentation rate for freshwater sedimentation would be 0.45 mm/y.

Surprisingly the present study suggests a distinctly younger age for the Neogene sedimentation in the North Croatian Basins than formerly considered. Probably the complete depositional cycle of its lower basinal infill, comprising the lacustrine and the early marine sediments, belongs to the Middle Miocene, Badenian stage. This completely changes the palaeogeographic picture for the south-western part of the

Pannonian Basin System. Instead of Ottnangian freshwater beds and a Karpatian marine transgression, aquatic environments existed solely during the Badenian.

ACKNOWLEDGEMENT

The study was funded by Ministry of Science, Education and Sports of the Republic of Croatia, Project: ‘Neogene terrestrial environments of the Pannonian Basin and karst region’ no. 195-1951293-2703, and by the Austrian FWF Project P18519-B17: ‘Mollusc Evolution of the Neogene Dinaride Lake System’. OM is highly indebted to Zlata JURISIĆ-POLŠAK (Croatian Natural History Museum, Zagreb) for facilitating access to Collection KOCHANSKY-DEVIDÉ. The authors are indebted to Mathias HARZHAUSER (NHM Vienna) and Zlatan BAJRAKTA-REVIĆ (PMF Zagreb) for critically reviewing the manuscript.

REFERENCES

AVANIĆ, R. (1997): Analiza facijesa srednjeg miocena na jugoistočnim padinama Medvednice [*Facies analysis of Middle Miocene on southern slopes of Mt. Medvednica – in Croatian*].– Unpubl. MSc Thesis, University of Zagreb, Zagreb, 54 p.

AVANIĆ, R., KOVAČIĆ, M., PAVELIĆ, D., ŠIMUNIĆ, AL., MIKNIĆ, M. & VRSALJKO, D. (1995a): Kongerijski vapnenci i klastiti ot-nanga – geološki stup Vidovec–2 [*Ottangian Congeria-limestone and clastics – geological column Vidovec–2 – in Croatian*].– In: ŠIKIĆ, K. (ed.): Geološki vodič Medvednice, Inst. za geol. istraž, INA – Industrija nafte d.d., Zagreb, 154.

AVANIĆ, R., PAVELIĆ, D., MIKNIĆ, M., BRKIĆ, M. & ŠIMUNIĆ, AL. (1995b): Karpatsko–donjobadenske naslage Čučerje [*Karpatian–Lower Badenian beds from Čučerje – in Croatian*].– In: ŠIKIĆ, K. (ed.): Geološki vodič Medvednice, Inst. za geol. istraž., INA – Industrija nafte d.d., Zagreb, 156–158.

- AVANIĆ, R., PAVELIĆ, D., VRSALJKO, D., ŠIMUNIĆ, AL., MIKNIĆ, M., HAJEK-TADESSE, V. & JERINIĆ, G. (1995c): Otnanški klastiti Planina [*Ottanian clastics from Planina* – in Croatian].– In: ŠIKIĆ, K. (ed.): Geološki vodič Medvednice. Inst. za geol. istraž., INA – Industrija nafte d.d., Zagreb, 164–168.
- BAJRAKTAREVIĆ, Z. (1981): Donji baden (donji “torton”) južnog dijela Moslavačke gore [*Early Badenian (Early “Tortonian”) of southern part of Mt. Moslavačka* – in Croatian].– Geol. vjesnik, 34, 9–13.
- BAJRAKTAREVIĆ, Z. & PAVELIĆ, D. (2003): The Karpatian in Croatia.– In: BRZOBOHATÝ, R., CICHA, I., KOVÁČ, M. & RÖGL, F. (eds.): The Karpatian. A Lower Miocene stage of the Central Paratethys. Masaryk University, Brno, 141–144.
- BASCH, O. (1983): Osnovna geološka karta SFRJ 1: 100000. Tumač za list Ivanić-Grad [*Basic Geological Map of SFRY 1:100000, Geology of the Ivanić-Grad sheet* – in Croatian].– Geol. zavod, Zagreb, Sav. geol. zavod, Beograd, 66 p.
- GORJANOVIĆ-KRAMBERGER, D. (1908): Geologijaska prijedlogna karta Kraljevine Hrvatske – Slavonije. Tumač geologijaskoj karti Zagreb [*Geological Descriptive Map of the Kingdom of Croatia – Slavonia, Geology for Geological Map of Zagreb* – in Croatian].– Nakl. Kralj. zemalj. vlade, Odjel za unutar. poslove, Zagreb, 75 p.
- GORJANOVIĆ-KRAMBERGER, D. (1923): Der Bruchrand des Zagreber Gebirges zwischen Podused und Zagreb und seine Bedeutung zur Heranbildung der Zagreber Terasse.– Glasn. Hrv. prirodosl. društva, Zagreb, 34, 3–15.
- GRADSTEIN, F.M., OGG, J.G. & SMITH, A.G. (2004): A geologic time scale 2004. Cambridge University Press, Cambridge, UK, 610 p.
- HAIJEK-TADESSE, V. (2006): Miocenski ostrakodi sjeverne Hrvatske [*Miocene ostracods of northern Croatia* – in Croatian].– Unpubl. PhD Thesis, University of Zagreb, Zagreb, 179 p.
- HAIJÓS, M. (1986): Stratigraphy of Hungary's Miocene diatomaceous earth deposits.– Geol. Hung., 49, 1–401.
- HAQ, B. (1991): Sequence stratigraphy, sea-level change, and significance for deep sea.– Spec. Publ. Int. Assoc. Sediment., Blackwell, 12, 3–39.
- HARZHAUSER, M. & MANDIĆ, O. (2008a): Neogene Dreissenids in Central Europe – exploring the limits.– In: VAN DER VELDE, G., RAJAGOPAL, S. & BIJ DE VAAT, A. (eds.): Zebra Mussels in Europe. (In press).
- HARZHAUSER, M. & MANDIĆ, O. (2008b): Neogene lake systems of Central and South-Eastern Europe: Faunal diversity, gradients and interrelations.– Palaeogeogr., Palaeoclimatol., Palaeoecol., 260, 417–434.
- HARZHAUSER, M., MANDIĆ, O. & ZUSCHIN, M. (2003): Changes in Paratethyan marine molluscs at the Early/Middle Miocene transition: diversity, palaeogeography and palaeoclimate.– Acta Geol. Pol. 53/4, 323–339.
- HARZHAUSER, M. & PILLER, W.E. (2007): Benchmark data of a changing sea – palaeogeography, palaeobiogeography and events in the Central Paratethys during the Miocene.– Palaeogeogr., Palaeoclimatol., Palaeoecol., 253, 8–31.
- JAMIĆIĆ, D., BRKIĆ, M., CRNKO, J. & VRAGOVIĆ, M. (1987): Osnovna geološka karta SFRJ 1:100000. Tumač za list Orahovica [*Basic Geological Map of SFRY 1:100000, Geology of the Orahovica sheet* – in Croatian].– Geol. zavod, Zagreb, Sav. geol. zavod, Beograd, 72 p.
- JAMIĆIĆ, D., VRAGOVIĆ, M. & MATIČEĆ, D. (1989): Osnovna geološka karta SFRJ 1:100000. Tumač za list Daruvar [*Basic Geological Map of SFRY 1:100000, Geology of the Daruvar sheet* – in Croatian].– Geol. zavod, Zagreb, Sav. geol. zavod, Beograd, 55 p.
- JELEN, B., MÁRTON, E., FODOR, L., BÁLDI, M., ČAR, J., RIFELJ, H., SKABERNE, D. & VRABEC, M. (1998): Palaeomagnetic, tectonic and stratigraphic correlation of Tertiary formations in Slovenia and Hungary along the Periadriatic and Mid-Hungarian tectonic zone (Preliminary communication).– Geologija, Ljubljana, 40, 325–331.
- JUNGWIRTH, E. & ĐEREK, T. (2000): Osobitosti paleoflore lokaliteta Planina [*Some characteristics of Palaeoflora of the Planina locality* – in Croatian].– In: VLAHOVIĆ, I. & BIONDIĆ, R. (eds.): Sec. Croat. Geol. Cong., Cavtat–Dubrovnik, 17–20.05.2000, Proceedings, Institute of Geology, Zagreb, Zagreb, 225–229.
- KOCHANSKY-DEVIDÉ, V. (1944): Fauna marinskog miocena južnog pobočja Medvednice (Zagrebačke gore) [*Fauna of marine Miocene of southern flanks of Mt. Medvednica (Mt. Zagrebačka)* – in Croatian].– Vjesnik Hrv. drž. geol. zav. i Hrv. drž. geol. muz., Zagreb, 2–3, 171–280.
- KOCHANSKY-DEVIDÉ, V. (1956): O fauni marinskog miocena i oortonskom “šliru” Medvednice (Zagrebačka Gora).– Geol. vjesnik, 10, 39–50.
- KOCHANSKY-DEVIDÉ, V. (1979): Srednjomiocenske kongerijske naslage Požeške gore [*Middle Miocene Congeria – beds of Mt. Požeška* – in Croatian].– Geol. vjesnik, 31, 69–72.
- KOCHANSKY-DEVIDÉ, V. & SLIŠKOVIĆ, T. (1978): Miocenske kongerije Hrvatske, Bosne i Hercegovine [*Miocene congerias of Croatia, Bosnia and Herzegovina* – in Croatian].– Palaeont. Jugosl., Zagreb, 19, 1–98.
- KOVÁČ, M., ANDREYEVA-GRIGOROVICH, A., BAJRAKTAREVIĆ, Z., BRZOBOHATÝ, R., FILIPESCU, S., FODOR, L., HARZHAUSER, M., NAGYMAROSY, A., OSZCZYPKO, N., PAVELIĆ, D., RÖGL, F., SAFTIĆ, B., SLIVA, E. & STUDENCKA, B. (2007): Badenian evolution of the Central Paratethys Sea: palaeogeography, climate and eustatic sea level changes.– Geol. Carpat., 58, 579–606.
- KOVÁČ, M., ANDREYEVA-GRIGOROVICH, A., BRZOBOHATÝ, R., FODOR, L., HARZHAUSER, M., OSZCZYPKO, N., PAVELIĆ, D., RÖGL, F., SAFTIĆ, B., SLIVA, E. & STRÁNIK, Z. (2003): Karpatian palaeogeography, tectonics and eustatic changes.– In: BRZOBOHATÝ, R., CICHA, I., KOVÁČ, M. & RÖGL, F. (eds.): The Karpatian. A Lower Miocene stage of the Central Paratethys, Masaryk University, Brno, 49–72.
- KRIZMANIĆ, K. (1995): Palynology of the Miocene bentonite from Gornja Jelenska (Mt. Moslavačka Gora, Croatia).– Geol. Croat., 48, 147–154.
- KRSTIĆ, N., SAVIĆ, LJ., JOVANOVIĆ, G. & BODOR, E. (2003): Lower Miocene lakes of the Balkan Land.– Acta Geol. Hung., 46, 291–299.
- LOURENS, L., HILGEN, F., SHACKLETON, N.J., LASKAR, J. & WILSON, J. (2004): The Neogene Period.– In: GRADSTEIN, F.M., OGG, J.G. & SMITH, A.G. (eds.): A Geologic Time Scale 2004., Cambridge University Press, Cambridge, UK, 409–440.
- LUČIĆ, D., SAFTIĆ, B., KRIZMANIĆ, K., PRELOGOVIĆ, E., BRITVIĆ, V., MESIĆ, I. & TADEJ, J. (2001): The Neogene evolution and hydrocarbon potential of the Pannonian Basin in Croatia.– Mar. Petrol. Geol., 18, 133–147.
- LUKENEDER, A. & HARZHAUSER, M. (2002): Shell Accumulations of the Nautilidae Aturia (Aturia) aturi in the Lower Miocene Paratethys.– In: SUMMESBERGER, H., HILTON, K. & DAURER, A. (eds.): Cephalopods – Present and Past. Abhandlungen der Geologischen Bundesanstalt, 57, 459–466.
- MANDIĆ, O., PAVELIĆ, D., HARZHAUSER, M., ZUPANIĆ, J., REISCHENBACHER, D., SACHSENHOFER, R.F., TADEJ, N. & VRANJKOVIĆ, A. (2008): Depositional history of the Miocene Lake Sinj (Dinaride Lake System, Croatia): a long-lived hard-water lake in a pull-apart tectonic setting.– J. Palaeolimnol., doi: 10.1007/s10933-008-9235-1
- MARTINI, E. (1971): Standard Tertiary and Quaternary calcareous nanoplankton zonation.– Proceedings of the II Planktonic Conference., Ed. Tecnoscienza, Roma, 739–785.
- MÁRTON, E., PAVELIĆ, D., TOMLJENOVIĆ, B., MÁRTON, P. & AVANIĆ, R. (2005): Palaeomagnetic investigations in the Croatian part of the Pannonian Basin: a review.– Acta Geol. Hung., 48, 225–233.

- MULDINI-MAMUŽIĆ, S. (1965): Rezultati mikrofaunističkog istraživanja oligocenskih i miocenskih naslaga Panonske kotline [*Results of microfaunistic study of the Oligocene and Miocene beds of the Požega Valley* – in Croatian]. – Acta geol., 5, (Prir. istraž. Jugosl. akad. znan. umjet., 35), 289–312.
- PAUL, M. (1872): Die Kohlenablagerungen bei Agram und Brod. – Verh. geol. Reichsanst., Wien, 6.
- PAUL, M. (1874): Die Braunkohlen – Ablagerungen von Croatien und Slavonien. – Jahrb. Geol. Reichsanst., 24/3.
- PAVELIĆ, D. (2001): Tectonostratigraphic model for the North Croatian and North Bosnian sector of the Miocene Pannonian Basin System. – Basin Res., 13, 359–376.
- PAVELIĆ, D. (2002): The south-western boundary of Central Paratethys. – Geol. Croat., 55, 83–92.
- PAVELIĆ, D. (2005): Cyclicity in the evolution of the Neogene North Croatian Basin (Pannonian Basin System). – In: MABESOONE, J.M. & NEUMANN, V.H. (eds.): Cyclic Development of Sedimentary Basins. Dev. Sedim., Elsevier, 57, 273–283.
- PAVELIĆ, D., AVANIĆ, R., BAKRAČ, K. & VRSALJKO, D. (2001): Early Miocene braided river and lacustrine sedimentation in the Kalnik Mountain Area (Pannonian Basin System, NW Croatia). – Geol. Carpat., 52, 375–386.
- PAVELIĆ, D., AVANIĆ, R., KOVAČIĆ, M., VRSALJKO, D. & MIKNIĆ, M. (2003): An Outline of the Evolution of the Croatian Part of the Pannonian Basin System. – In: VLAHOVIĆ, I. & TIŠLJAR, J. (eds.): Evolution of Depositional Environments from the Palaeozoic to the Quaternary in the Karst Dinarides and the Pannonian Basin. 22nd IAS Meeting of Sedimentology, Opatija – Sept. 17–19, 2003, Field Trip Guidebook, Zagreb, 155–161.
- PAVELIĆ, D., MIKNIĆ, M. & SARKOTIĆ ŠLAT, M. (1998): Early to Middle Miocene facies succession in lacustrine and marine environments on the southwestern margin of the Pannonian basin system. – Geol. Carpat., 49, 433–443.
- PILLER, W., HARZHAUSER, M. & MANDIĆ, O. (2007): Miocene Central Paratethys stratigraphy – current status and future directions. – Stratigraphy, 4, 151–168.
- POLIĆ, A. (1935): O oligocenu i njegovoj flori kod Planine u Zagrebačkoj gori [*About the Oligocene and its flora by Planina on Mt. Zagrebačka* – in Croatian]. – Rad Jugosl. akad. znan. umjetn., Zagreb, 251, 61–90.
- RÖGL, F. (1996): Stratigraphic correlation of the Paratethys Oligocene and Miocene. – Mitt. Ges. Bergbaustud. Österr., 41, 65–73.
- RÖGL, F., ČORIĆ, S., HOHENEGGER, J., PERVESLER, P., ROETZEL, R., SCHOLGER, R., SPEZZAFERRI, S. & STINGL, K. (2007): Cyclostratigraphy and transgressions at the Early/Middle Miocene (Karpatian/Badenian) boundary in the Austrian Neogene basins (Central Paratethys). – Scripta Fac. Sci. Natur. Univ. Masarykianae Brunensis, ser. Geology, Brno, 36, 7–13.
- RÖGL, F., SPEZZAFERRI, S. & ČORIĆ, S. (2002): Micropalaeontology and biostratigraphy of the Karpatian-Badenian transition (Early-Middle Miocene boundary) in Austria (Central Paratethys). – Courier Forsch. Inst. Senckenberg, Frankfurt a. M., 237, 47–67.
- RÖGL, F. & STEININGER, F.-F. (1983): Vom Zerfall der Tethys zu Mediterran und Paratethys. Die Neogene Palaeogeographie und Palinspastik des zirkum-mediterranen Raumes. – Ann. Naturhist. Mus. Wien, 85, 135–163.
- SAFTIĆ, B., VELIĆ, J., SZTANÓ, O., JUHÁSZ, G. & IVKOVIĆ, Ž. (2003): Tertiary subsurface facies, source rocks and hydrocarbon reservoirs in the SW part of the Pannonian Basin (Northern Croatia and South-Western Hungary). – Geol. Croat., 56, 101–122.
- SOKAČ, A. (1987): *Pannoninotus* n. gen. (Crustacea, Ostracoda) from the Middle Miocene of Bjelanovac (Psnj Mountain, North Croatia). – Geol. vjesnik, 40, 39–44.
- SOKAČ, A. & KRSTIĆ, N. (1987): Ostracod fauna of some non-marine Neogene basins in Yugoslavia. – Geol. vjesnik, 40, 45–52.
- SPEZZAFERRI, S., RÖGL, F., ČORIĆ, S. & HOHENEGGER, J. (2004): Palaeoenvironmental changes and agglutinated foraminifera across the Karpatian/Badenian (Early/Middle Miocene) boundary in the Styrian Basin. – In: BUBIK, M. & KAMINSKI, M.A. (eds.): Proceedings of the 6th International Workshop on agglutinated foraminifera (Prague, Czech Republic, Sept. 1–7, 2001). Grzybowski Foundation Spec. Publ., Krakow, 8, 423–459.
- SPIEGLER, D. & RÖGL, F. (1992): Bolboforma (Protophyta, incertae sedis) im Oligozän und Miozän des Mediterran und der Zentralen Paratethys. – Annalen Naturhist. Mus. Wien, 95A, 59–95.
- SUTO, I. (2006): *Truncatulus* gen. nov., a new fossil resting spore morphogenus related to the marine diatom genus *Chaetoceros* (Bacillariophyceae). – Phycologia, 45/5, 585–601.
- ŠIKIĆ, L. (1966): New concepts on the age of the hitherto existing Burdigalian and Upper Oligocene deposits in the Zagrebačka gora mountain. – Bull. Sci. Cons. Acad. Yougosl., Zagreb, (A), 11/1–12, 246–247.
- ŠIKIĆ, L. (1968): Stratigrafija miocena sjeveroistočnog dijela Medvednice na osnovu faune foraminifera [*Stratigraphy of the Miocene based on foraminifera fauna* – in Croatian]. – Geol. vjesnik, 21, 213–227.
- ŠIMUNIĆ, AL., AVANIĆ, R., & ŠIMUNIĆ, AN. (1990): “Macejski pješčenjaci” i vulkanizam zapadnog dijela Hrvatskog zagorja (Hrvatska, Jugoslavija) [*“Macej Sandstones” and volcanism in the western part of Hrvatsko Zagorje (Croatia, Yugoslavia)* – in Croatian]. – Rad JAZU, Zagreb, 449, 179–194.
- ŠPARICA, M. & BUZALJKO, R. (1984): Geološka karta SFRJ 1:100000. Tumač za list Nova Gradiška [*Basic Geological Map of SFRJ 1:100000, Geology of the Nova Gradiška sheet* – in Croatian]. – Geol. zavod, Zagreb, Inst. za geologiju, Sarajevo, Sav. geol. zavod, Beograd, 54 p.
- ŠPARICA, M., JURIŠA, M., CRNKO, J., ŠIMUNIĆ, AN., JOVANOVIĆ, Č. & ŽIVANOVIĆ, D. (1980): Osnovna geološka karta SFRJ 1:100000. Tumač za list Nova Kapela [*Basic Geological Map of SFRJ 1:100000 Geology of the Nova Kapela sheet* – in Croatian]. – Inst. za geol. istraž., Zagreb, Inst. za geol., Sarajevo, Sav. geol. zavod, Beograd, 55 p.
- ŽAGAR-SAKAČ, A. (2003): *Hyriopsis (Limnoscapha) slavonicus* n. sp. (Lamellibranchiata, Unionidae, Hyriopsinae) from Miocene strata in Slavonia, Croatia. – Nat. Croat., Zagreb, 2/1, 73–82.
- ŽAGAR-SAKAČ, A. (2004): The Otmangian Unionaceae from Jelov Dol. – Nat. Croat., 3/2, 225–252.

Manuscript received July, 7, 2008
 Revised manuscript accepted November 4, 2008