

**DRAFT
TOXICOLOGICAL PROFILE FOR
ACRYLAMIDE**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Agency for Toxic Substances and Disease Registry

September 2009

DISCLAIMER

The use of company or product name(s) is for identification only and does not imply endorsement by the Agency for Toxic Substances and Disease Registry.

This information is distributed solely for the purpose of pre dissemination public comment under applicable information quality guidelines. It has not been formally disseminated by the Agency for Toxic Substances and Disease Registry. It does not represent and should not be construed to represent any agency determination or policy.

UPDATE STATEMENT

Toxicological profiles are revised and republished as necessary. For information regarding the update status of previously released profiles, contact ATSDR at:

Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine/Applied Toxicology Branch
1600 Clifton Road NE
Mailstop F-62
Atlanta, Georgia 30333

This page is intentionally blank.

FOREWORD

This toxicological profile is prepared in accordance with guidelines developed by the Agency for Toxic Substances and Disease Registry (ATSDR) and the Environmental Protection Agency (EPA). The original guidelines were published in the *Federal Register* on April 17, 1987. Each profile will be revised and republished as necessary.

The ATSDR toxicological profile succinctly characterizes the toxicologic and adverse health effects information for these toxic substances described therein. Each peer-reviewed profile identifies and reviews the key literature that describes a substance's toxicologic properties. Other pertinent literature is also presented, but is described in less detail than the key studies. The profile is not intended to be an exhaustive document; more comprehensive sources of specialty information are referenced.

The focus of the profiles is on health and toxicologic information. Each toxicological profile begins with a public health statement that describes, in nontechnical language, a substance's relevant toxicological properties. Following the public health statement is information concerning levels of significant human exposure and, where known, significant health effects. The adequacy of information to determine a substance's health effects is described in a health effects summary. Data needs that are of significance to protection of public health are identified by ATSDR and EPA.

Each profile includes the following:

- (A) The examination, summary, and interpretation of available toxicologic information and epidemiologic evaluations on a toxic substance to ascertain the levels of significant human exposure for the substance and the associated acute, subacute, and chronic health effects;
- (B) A determination of whether adequate information on the health effects of each substance is available or in the process of development to determine levels of exposure that present a significant risk to human health of acute, subacute, and chronic health effects; and
- (C) Where appropriate, identification of toxicologic testing needed to identify the types or levels of exposure that may present significant risk of adverse health effects in humans.

The principal audiences for the toxicological profiles are health professionals at the Federal, State, and local levels; interested private sector organizations and groups; and members of the public. We plan to revise these documents in response to public comments and as additional data become available. Therefore, we encourage comments that will make the toxicological profile series of the greatest use.

Comments should be sent to:


Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine
1600 Clifton Road, N.E.
Mail Stop F-62
Atlanta, Georgia 30333

The toxicological profiles are developed under the Comprehensive Environmental Response, Compensation, and Liability Act of 1980, as amended (CERCLA or Superfund). CERCLA section 104(i)(1) directs the Administrator of ATSDR to "...effectuate and implement the health related authorities" of the statute. This includes the preparation of toxicological profiles for hazardous substances most commonly found at facilities on the CERCLA National Priorities List and that pose the most significant potential threat to human health, as determined by ATSDR and the EPA. Section 104(i)(3) of CERCLA, as amended, directs the Administrator of ATSDR to prepare a toxicological profile for each substance on the list. In addition, ATSDR has the authority to prepare toxicological profiles for substances not found at sites on the National Priorities List, in an effort to "...establish and maintain inventory of literature, research, and studies on the health effects of toxic substances" under CERCLA Section 104(i)(1)(B), to respond to requests for consultation under section 104(i)(4), and as otherwise necessary to support the site-specific response actions conducted by ATSDR.

This profile reflects ATSDR's assessment of all relevant toxicologic testing and information that has been peer-reviewed. Staffs of the Centers for Disease Control and Prevention and other Federal scientists have also reviewed the profile. In addition, this profile has been peer-reviewed by a nongovernmental panel and is being made available for public review. Final responsibility for the contents and views expressed in this toxicological profile resides with ATSDR.



Howard Frumkin, M.D., Dr. P.H.
Director, National Center for Environmental
Health/Agency for Toxic Substances and Disease
Registry



Thomas R. Frieden, M.D., M.P.H.
Administrator, Agency for Toxic Substances and
Disease Registry

Managing Hazardous Materials Incidents is a three-volume set of recommendations for on-scene (prehospital) and hospital medical management of patients exposed during a hazardous materials incident. Volumes I and II are planning guides to assist first responders and hospital emergency department personnel in planning for incidents that involve hazardous materials. Volume III—*Medical Management Guidelines for Acute Chemical Exposures*—is a guide for health care professionals treating patients exposed to hazardous materials.

Fact Sheets (ToxFAQs) provide answers to frequently asked questions about toxic substances.

Other Agencies and Organizations

The National Center for Environmental Health (NCEH) focuses on preventing or controlling disease, injury, and disability related to the interactions between people and their environment outside the workplace. Contact: NCEH, Mailstop F-29, 4770 Buford Highway, NE, Atlanta, GA 30341-3724 • Phone: 770-488-7000 • FAX: 770-488-7015.

The National Institute for Occupational Safety and Health (NIOSH) conducts research on occupational diseases and injuries, responds to requests for assistance by investigating problems of health and safety in the workplace, recommends standards to the Occupational Safety and Health Administration (OSHA) and the Mine Safety and Health Administration (MSHA), and trains professionals in occupational safety and health. Contact: NIOSH, 200 Independence Avenue, SW, Washington, DC 20201 • Phone: 800-356-4674 or NIOSH Technical Information Branch, Robert A. Taft Laboratory, Mailstop C-19, 4676 Columbia Parkway, Cincinnati, OH 45226-1998 • Phone: 800-35-NIOSH.

The National Institute of Environmental Health Sciences (NIEHS) is the principal federal agency for biomedical research on the effects of chemical, physical, and biologic environmental agents on human health and well-being. Contact: NIEHS, PO Box 12233, 104 T.W. Alexander Drive, Research Triangle Park, NC 27709 • Phone: 919-541-3212.

Referrals

The Association of Occupational and Environmental Clinics (AOEC) has developed a network of clinics in the United States to provide expertise in occupational and environmental issues. Contact: AOEC, 1010 Vermont Avenue, NW, #513, Washington, DC 20005 • Phone: 202-347-4976 • FAX: 202-347-4950 • e-mail: AOEC@AOEC.ORG • Web Page: <http://www.aoec.org/>.

The American College of Occupational and Environmental Medicine (ACOEM) is an association of physicians and other health care providers specializing in the field of occupational and environmental medicine. Contact: ACOEM, 25 Northwest Point Boulevard, Suite 700, Elk Grove Village, IL 60007-1030 • Phone: 847-818-1800 • FAX: 847-818-9266.

CONTRIBUTORS

CHEMICAL MANAGER(S)/AUTHOR(S):

Patricia Ruiz, Ph.D.
Obaid Faroon, Ph.D.
Moiz Mumtaz, Ph.D.
ATSDR, Division of Toxicology and Environmental Medicine, Atlanta, GA

David W. Wohlers, Ph.D.
Lynn E. Barber, M.S.
Peter R. McClure, Ph.D., DABT
Christine Carnegie, B.A.
SRC, Inc., North Syracuse, NY

THE PROFILE HAS UNDERGONE THE FOLLOWING ATSDR INTERNAL REVIEWS:

1. Health Effects Review. The Health Effects Review Committee examines the health effects chapter of each profile for consistency and accuracy in interpreting health effects and classifying end points.
2. Minimal Risk Level Review. The Minimal Risk Level Workgroup considers issues relevant to substance-specific Minimal Risk Levels (MRLs), reviews the health effects database of each profile, and makes recommendations for derivation of MRLs.
3. Data Needs Review. The Applied Toxicology Branch reviews data needs sections to assure consistency across profiles and adherence to instructions in the Guidance.
4. Green Border Review. Green Border review assures the consistency with ATSDR policy.

This page is intentionally blank.

PEER REVIEW

A peer review panel was assembled for acrylamide. The panel consisted of the following members:

1. Herman Bolt, M.D., Ph.D., Professor Emeritus, Leibniz Research Centre for Working Environment and Human Factors, Dortmund, Germany
2. Timothy Fennell, Ph.D., Senior Research Chemist, RTI International, Drug Metabolism and Pharmacokinetics Department, Research Triangle Park, NC
3. James Klaunig, Ph.D., Professor of Public Health, Director, Center for Environmental Health, Indiana University School of Medicine, Indianapolis, IN

These experts collectively have knowledge of acrylamide's physical and chemical properties, toxicokinetics, key health end points, mechanisms of action, human and animal exposure, and quantification of risk to humans. All reviewers were selected in conformity with the conditions for peer review specified in Section 104(I)(13) of the Comprehensive Environmental Response, Compensation, and Liability Act, as amended.

Scientists from the Agency for Toxic Substances and Disease Registry (ATSDR) have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.

This page is intentionally blank.

CONTENTS

DISCLAIMER	ii
UPDATE STATEMENT	iii
FOREWORD	v
QUICK REFERENCE FOR HEALTH CARE PROVIDERS	vii
CONTRIBUTORS	ix
PEER REVIEW	xi
CONTENTS	xiii
LIST OF FIGURES	xvii
LIST OF TABLES	xix
1. PUBLIC HEALTH STATEMENT	1
1.1 WHAT IS ACRYLAMIDE?	1
1.2 WHAT HAPPENS TO ACRYLAMIDE WHEN IT ENTERS THE ENVIRONMENT?	2
1.3 HOW MIGHT I BE EXPOSED TO ACRYLAMIDE?	2
1.4 HOW CAN ACRYLAMIDE ENTER AND LEAVE MY BODY?	2
1.5 HOW CAN ACRYLAMIDE AFFECT MY HEALTH?	3
1.6 HOW CAN ACRYLAMIDE AFFECT CHILDREN?	3
1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO ACRYLAMIDE?	3
1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO ACRYLAMIDE?	4
1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?	4
1.10 WHERE CAN I GET MORE INFORMATION?	5
2. RELEVANCE TO PUBLIC HEALTH	7
2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO ACRYLAMIDE IN THE UNITED STATES	7
2.2 SUMMARY OF HEALTH EFFECTS	8
2.3 MINIMAL RISK LEVELS (MRLs)	10
3. HEALTH EFFECTS	15
3.1 INTRODUCTION	15
3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE	15
3.2.1 Inhalation Exposure	16
3.2.1.1 Death	16
3.2.1.2 Systemic Effects	17
3.2.1.3 Immunological and Lymphoreticular Effects	21
3.2.1.4 Neurological Effects	21
3.2.1.5 Reproductive Effects	27
3.2.1.6 Developmental Effects	27
3.2.1.7 Cancer	27
3.2.2 Oral Exposure	30
3.2.2.1 Death	30
3.2.2.2 Systemic Effects	30
3.2.2.3 Immunological and Lymphoreticular Effects	60
3.2.2.4 Neurological Effects	60
3.2.2.5 Reproductive Effects	62
3.2.2.6 Developmental Effects	63

3.2.2.7	Cancer	65
3.2.3	Dermal Exposure.....	70
3.2.3.1	Death	70
3.2.3.2	Systemic Effects.....	71
3.2.3.3	Immunological and Lymphoreticular Effects	74
3.2.3.4	Neurological Effects	74
3.2.3.5	Reproductive Effects.....	74
3.2.3.6	Developmental Effects.....	75
3.2.3.7	Cancer	75
3.3	GENOTOXICITY	75
3.4	TOXICOKINETICS.....	88
3.4.1	Absorption.....	88
3.4.1.1	Inhalation Exposure	89
3.4.1.2	Oral Exposure	90
3.4.1.3	Dermal Exposure.....	93
3.4.2	Distribution	94
3.4.2.1	Inhalation Exposure	94
3.4.2.2	Oral Exposure	94
3.4.2.3	Dermal Exposure.....	95
3.4.3	Metabolism	95
3.4.4	Elimination and Excretion	102
3.4.4.1	Inhalation Exposure	102
3.4.4.2	Oral Exposure	102
3.4.4.3	Dermal Exposure.....	104
3.4.5	Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models.....	104
3.5	MECHANISMS OF ACTION	115
3.5.1	Pharmacokinetic Mechanisms.....	115
3.5.2	Mechanisms of Toxicity.....	115
3.5.3	Animal-to-Human Extrapolations.....	119
3.6	TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS	119
3.7	CHILDREN'S SUSCEPTIBILITY	120
3.8	BIOMARKERS OF EXPOSURE AND EFFECT	124
3.8.1	Biomarkers Used to Identify or Quantify Exposure to Acrylamide.....	125
3.8.2	Biomarkers Used to Characterize Effects Caused by Acrylamide.....	125
3.9	INTERACTIONS WITH OTHER CHEMICALS	125
3.10	POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE	126
3.11	METHODS FOR REDUCING TOXIC EFFECTS.....	127
3.11.1	Reducing Peak Absorption Following Exposure	128
3.11.2	Reducing Body Burden.....	128
3.11.3	Interfering with the Mechanism of Action for Toxic Effects.....	128
3.12	ADEQUACY OF THE DATABASE.....	129
3.12.1	Existing Information on Health Effects of Acrylamide	129
3.12.2	Identification of Data Needs	131
3.12.3	Ongoing Studies.....	139
4.	CHEMICAL AND PHYSICAL INFORMATION.....	141
4.1	CHEMICAL IDENTITY.....	141
4.2	PHYSICAL AND CHEMICAL PROPERTIES.....	141

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL	145
5.1 PRODUCTION	145
5.2 IMPORT/EXPORT	149
5.3 USE.....	149
5.4 DISPOSAL	150
6. POTENTIAL FOR HUMAN EXPOSURE	151
6.1 OVERVIEW	151
6.2 RELEASES TO THE ENVIRONMENT	153
6.2.1 Air	154
6.2.2 Water.....	157
6.2.3 Soil	158
6.3 ENVIRONMENTAL FATE.....	158
6.3.1 Transport and Partitioning	158
6.3.2 Transformation and Degradation	159
6.3.2.1 Air	159
6.3.2.2 Water.....	160
6.3.2.3 Sediment and Soil	160
6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT	160
6.4.1 Air	161
6.4.2 Water.....	161
6.4.3 Sediment and Soil	162
6.4.4 Other Environmental Media.....	162
6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE	168
6.6 EXPOSURES OF CHILDREN	170
6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES	171
6.8 ADEQUACY OF THE DATABASE.....	171
6.8.1 Identification of Data Needs	172
6.8.2 Ongoing Studies.....	174
7. ANALYTICAL METHODS.....	177
7.1 BIOLOGICAL MATERIALS	177
7.2 ENVIRONMENTAL SAMPLES	179
7.3 ADEQUACY OF THE DATABASE.....	182
7.3.1 Identification of Data Needs	182
7.3.2 Ongoing Studies.....	183
8. REGULATIONS, ADVISORIES, AND GUIDELINES.....	185
9. REFERENCES	191
10. GLOSSARY	241

APPENDICES

A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETS A-1

B. USER’S GUIDE..... B-1

C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS..... C-1

D. INDEX D-1

LIST OF FIGURES

3-1. Levels of Significant Exposure to Acrylamide - Inhalation	20
3-2. Levels of Significant Exposure to Acrylamide - Oral	52
3-3. Metabolic Scheme for Acrylamide and Its Metabolite Glycidamide	97
3-4. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance	106
3-5. Schematic of the Kirman et al. (2003) Physiologically Based Pharmacokinetic (PBPK) Model for Acrylamide.....	108
3-6. Schematic of the Young et al. (2007) Physiologically Based Pharmacokinetic/Toxicodynamic (PBPK/TD) Model For Acrylamide and Glycidamide	113
3-7. Existing Information on Health Effects of Acrylamide.....	130
6-1. Frequency of NPL Sites with Acrylamide Contamination	152

This page is intentionally blank.

LIST OF TABLES

3-1. Levels of Significant Exposure to Acrylamide - Inhalation	18
3-2. Self-Reported Neurological Symptoms and Observed Clinical Signs Among Acrylamide Workers and Nonexposed Workers	23
3-3. Prevalence of Symptoms and Signs of Adverse Health Effects in Acrylamide-Exposed Workers and Controls	25
3-4. Levels of Significant Exposure to Acrylamide - Oral	31
3-5. Incidence of Tumors with Statistically Significant Increases in 2-Year Bioassays with F344 Rats Exposed to Acrylamide in Drinking Water	69
3-6. Levels of Significant Exposure to Acrylamide - Dermal	72
3-7. Genotoxicity of Acrylamide In Vivo	76
3-8. Genotoxicity of Acrylamide In Vitro	84
3-9. Original Model Parameter Values for Rats in the Kirman et al. (2003) Physiologically Based Toxicokinetic (PBTK) Model	109
4-1. Chemical Identity of Acrylamide	142
4-2. Physical and Chemical Properties of Acrylamide	143
5-1. Facilities that Produce, Process, or Use Acrylamide	146
5-2. 2008 Acrylamide Production in the United States.....	148
6-1. Releases to the Environment from Facilities that Produce, Process, or Use Acrylamide.....	155
6-2. Acrylamide Levels in Different Food and Food Product Groups from Norway, Sweden, Switzerland, the United Kingdom, and the United States.....	164
6-3. Acrylamide in Potato Products	165
6-4. Acrylamide Content of Food	166
6-5. Acrylamide Levels in Food Products Sampled for the 2003–2006 Total Diet Studies (TDS) Summary	167
6-6. Ongoing Research Regarding the Environmental Fate and Exposure to Acrylamide	175
7-1. Analytical Methods for Determining Acrylamide in Biological Samples.....	178
7-2. Analytical Methods for Determining Acrylamide in Environmental Samples.....	180

8-1. Regulations and Guidelines Applicable to Acrylamide..... 187

1. PUBLIC HEALTH STATEMENT

This public health statement tells you about acrylamide and the effects of exposure to it.

The Environmental Protection Agency (EPA) identifies the most serious hazardous waste sites in the nation. These sites are then placed on the National Priorities List (NPL) and are targeted for long-term federal clean-up activities. Acrylamide has been found in at least 3 of the 1,699 current or former NPL sites. Although the total number of NPL sites evaluated for this substance is not known, the possibility exists that the number of sites at which acrylamide is found may increase in the future as more sites are evaluated. This information is important because these sites may be sources of exposure and exposure to this substance may be harmful.

When a substance is released either from a large area, such as an industrial plant, or from a container, such as a drum or bottle, it enters the environment. Such a release does not always lead to exposure. You can be exposed to a substance only when you come in contact with it. You may be exposed by breathing, eating, or drinking the substance, or by skin contact.

If you are exposed to acrylamide, many factors will determine whether you will be harmed. These factors include the dose (how much), the duration (how long), and how you come in contact with it. You must also consider any other chemicals you are exposed to and your age, sex, diet, family traits, lifestyle, and state of health.

1.1 WHAT IS ACRYLAMIDE?

White or colorless, odorless crystalline solid	Acrylamide can violently react when melting. When heated, acrid fumes may be released.
Used in industry	Acrylamide is used to make polyacrylamide, which is mainly used in treating effluent from water treatment plants and industrial processes.

1. PUBLIC HEALTH STATEMENT

1.2 WHAT HAPPENS TO ACRYLAMIDE WHEN IT ENTERS THE ENVIRONMENT?

Most commonly found in water	Acrylamide may enter drinking water if polyacrylamide is used in the treatment process. It can be found in soils, but is rarely found in air.
Rapidly broken down in soil and water	If acrylamide enters soil or water, it will be broken down quickly by bacteria.

1.3 HOW MIGHT I BE EXPOSED TO ACRYLAMIDE?

Water and soil	Drinking water can sometimes contain acrylamide. It can enter drinking water from the treatment process of municipal supplies as well as from substances used to construct dams and wells. Acrylamide breaks down quickly in water and soil, but there is still a chance of exposure if you live near a plastics or dye plants.
Inhalation and dermal contact	If you smoke, or breathe second-hand tobacco smoke, you might be exposed to acrylamide. Tobacco smoke is a major source of acrylamide exposure within the general population. People involved in the production or use of acrylamide and acrylamide-containing products are exposed if they breathe in air that contains acrylamide. They may also be exposed by coming into skin contact with acrylamide.
Food	Acrylamide is formed in foods that are rich in carbohydrates (particularly potatoes) when they are fried, grilled, or baked at normal cooking temperatures. Levels of acrylamide in these foods increase with higher temperatures and longer cooking times. Protein-based foods (such as meats) probably contain low amounts of acrylamide. Ingestion of foods that contain acrylamide is a primary source of exposure.

1.4 HOW CAN ACRYLAMIDE ENTER AND LEAVE MY BODY?

May enter your body through food, drinking water, breathing, and skin contact	Acrylamide can enter your body when you eat foods containing acrylamide. Drinking water that contains acrylamide will have a similar effect. Breathing tobacco smoke may cause some level of acrylamide to enter your lungs. Acrylamide can also enter your body if it comes in contact with your skin. Dermal contact with acrylamide can occur if you work in the manufacture of acrylamide or polyacrylamide gels.
Leaves through bodily fluids	Once in your body, acrylamide enters your body fluids. Acrylamide and its breakdown products leave your body mostly through urine; small amounts may leave through feces, exhaled air, and breast milk.

1. PUBLIC HEALTH STATEMENT

1.5 HOW CAN ACRYLAMIDE AFFECT MY HEALTH?

This section looks at studies concerning potential health effects in animal and human studies.

Nervous system effects	Nervous system effects such as muscle weakness, numbness in hands and feet, sweating, unsteadiness, and clumsiness were reported in some acrylamide workers. However, most people are not exposed to acrylamide levels high enough to cause these effects.
Reproductive effects	Acrylamide reduces the ability of male animals to produce offspring and could cause similar effects in humans, but not likely at exposure levels experienced by most people.
Cancer	Acrylamide has caused several types of cancer in animals. We do not know whether acrylamide causes cancer in humans.

1.6 HOW CAN ACRYLAMIDE AFFECT CHILDREN?

This section discusses potential health effects in humans from exposures during the period from conception to maturity at 18 years of age.

Effects in children	Acrylamide is expected to affect children in the same manner as adults. It is not known whether children are more susceptible than adults to the effects of acrylamide.
Developmental effects	Effects such as decreased body weight, decreased startle responses, and decreased levels of some chemicals involved in transmission of brain signals were seen in some animals exposed to acrylamide before or shortly following birth. There are no reports of acrylamide causing developmental effects in humans.

1.7 HOW CAN FAMILIES REDUCE THE RISK OF EXPOSURE TO ACRYLAMIDE?

Limit exposure to tobacco and second-hand smoke	Tobacco smoke contains acrylamide. Avoid smoking or breathing in second-hand smoke.
Reduce consumption of foods that contain acrylamide	Avoid eating a lot of carbohydrate-rich foods that are cooked at high temperatures (e.g., French fries). Foods with higher protein content appear to have lower amounts of acrylamide. Avoid overcooking foods.

1. PUBLIC HEALTH STATEMENT

1.8 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO ACRYLAMIDE?

Can be measured in blood and urine	Acrylamide and its breakdown products can be measured in blood and urine. These measurements may be useful in estimating how much acrylamide has entered the body.
---	--

1.9 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?

The federal government develops regulations and recommendations to protect public health. Regulations can be enforced by law. The EPA, the Occupational Safety and Health Administration (OSHA), and the Food and Drug Administration (FDA) are some federal agencies that develop regulations for toxic substances. Recommendations provide valuable guidelines to protect public health, but cannot be enforced by law. The Agency for Toxic Substances and Disease Registry (ATSDR) and the National Institute for Occupational Safety and Health (NIOSH) are two federal organizations that develop recommendations for toxic substances.

Regulations and recommendations can be expressed as “not-to-exceed” levels. These are levels of a toxic substance in air, water, soil, or food that do not exceed a critical value. This critical value is usually based on levels that affect animals; they are then adjusted to levels that will help protect humans. Sometimes these not-to-exceed levels differ among federal organizations because they used different exposure times (an 8-hour workday or a 24-hour day), different animal studies, or other factors.

Recommendations and regulations are also updated periodically as more information becomes available. For the most current information, check with the federal agency or organization that provides it.

Some regulations and recommendations for acrylamide include the following:

Levels in drinking water set by EPA	The EPA has determined that exposure to acrylamide in drinking water at concentrations of 1.5 mg/L for one day or 0.3 mg/L for 10 days is not expected to cause any adverse effects in a child.
Levels in workplace air set by OSHA	OSHA set a legal limit of 0.3 mg/m ³ for acrylamide in air averaged over an 8-hour work day.

1. PUBLIC HEALTH STATEMENT

1.10 WHERE CAN I GET MORE INFORMATION?

If you have any more questions or concerns, please contact your community or state health or environmental quality department, or contact ATSDR at the address and phone number below.

ATSDR can also tell you the location of occupational and environmental health clinics. These clinics specialize in recognizing, evaluating, and treating illnesses that result from exposure to hazardous substances.

Toxicological profiles are also available on-line at www.atsdr.cdc.gov and on CD-ROM. You may request a copy of the ATSDR ToxProfiles™ CD-ROM by calling the toll-free information and technical assistance number at 1-800-CDCINFO (1-800-232-4636), by e-mail at cdcinfo@cdc.gov, or by writing to:

Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine
1600 Clifton Road NE
Mailstop F-62
Atlanta, GA 30333
Fax: 1-770-488-4178

Organizations for-profit may request copies of final Toxicological Profiles from the following:

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161
Phone: 1-800-553-6847 or 1-703-605-6000
Web site: <http://www.ntis.gov/>

1. PUBLIC HEALTH STATEMENT

This page is intentionally blank.

2. RELEVANCE TO PUBLIC HEALTH

2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO ACRYLAMIDE IN THE UNITED STATES

Acrylamide is an industrial chemical used mainly in the production of polyacrylamides, which are used primarily as flocculants for clarifying drinking and treating municipal and industrial effluents. In the oil industry, acrylamide is used as a flow control agent to enhance oil production from wells. Acrylamide and polyacrylamides are also used in the production of dyes and organic chemicals, contact lenses, cosmetics and toiletries, permanent-press fabrics, textiles, pulp and paper products; also in ore processing, sugar refining, and as a chemical grouting agent and soil stabilizer for the construction of tunnels, sewers, wells, and reservoirs. In 2008, approximately 141,000 metric tons of acrylamide were produced in the United States.

Acrylamide may be released to the environment during production and use of polyacrylamides, which are used as clarifiers in water treatment. Residual monomer in the polyacrylamide is the main source of drinking water contamination by acrylamide. It can be released to soil and land from plastics and dye industries and from acrylamide-containing grouting agents used in reservoirs and wells. Acrylamide is rarely found in the atmosphere.

Acrylamide is not considered highly persistent in the environment. It is expected to be highly mobile in soil and water and is highly susceptible to biodegradation in both media. It can be removed from soils by hydrolysis, though it is typically not present in soil or the atmosphere. Acrylamide is not expected to significantly bioconcentrate.

Acrylamide is sometimes present in drinking water due to leaching of monomer during treatment processes. Acrylamide was identified in food samples cooked at high temperatures. Concentrations of acrylamide in food vary with the type of food, cooking method, temperature, water content, food thickness, and length of heating. Carbohydrate-rich foods typically contain the highest levels of acrylamide.

Exposure to acrylamide occurs mainly through ingestion, dermal contact, and inhalation. Ingestion of foods containing acrylamide appears to be one of the most common methods of exposure for the general public. Average estimated intake of acrylamide from food sources ranged from 0.3 to 0.8 $\mu\text{g}/\text{kg}$ bw/day. Children may be susceptible to food-borne exposure 2–3 times that of adults on a body weight basis.

2. RELEVANCE TO PUBLIC HEALTH

Ingestion of contaminated drinking water can result in exposure to acrylamide. Inhalation of tobacco smoke, including second-hand smoke, can result in inhalation exposure for both adults and children. In the body, acrylamide can cross the placenta and result in exposure to unborn children. Breast milk has been shown to contain acrylamide in mothers with diets high in acrylamide-containing foods. Occupational exposure to acrylamide is primarily due to dermal contact when handling bags and drums of the chemical, followed by inhalation of dust or aerosols.

2.2 SUMMARY OF HEALTH EFFECTS

Neurological Effects. Ataxia and skeletal muscle weakness are hallmark symptoms of acrylamide-induced central-peripheral neuropathy. Available information in humans includes case reports and cross-sectional studies in which reported symptoms of neuropathy were primarily associated with occupational exposure potential via both inhalation and dermal contact. Numerous animal studies confirm the neurological effects of acrylamide exposure. Available information in animals primarily involves oral exposure. Clinical signs can be elicited following single oral dosing in the range of 100–200 mg/kg; lower dose levels require repeated dosing to elicit clinical signs. Histopathological evidence of acrylamide-induced peripheral neuropathy has been observed in rats receiving oral doses as low as 1 mg/kg/day for 3 months; the observed degenerative effects in peripheral nerve fibers at such dose levels have been shown to be completely reversible within a few months following the cessation of exposure (Burek et al. 1980).

Reproductive Effects. Pre- and postimplantation losses and decreased numbers of live fetuses have been observed in rats and mice receiving repeated oral doses of acrylamide in the range of 3–60 mg/kg/day. Results of dominant lethality testing and crossover trials indicate that acrylamide-induced reproductive effects are male mediated. Other reported effects in acrylamide-treated laboratory animals include decreased sperm mobility, degenerative effects in spermatids, and testicular atrophy.

Cancer. Available cohort mortality studies of occupationally-exposed workers have not found associations between acrylamide and mortality from cancer. Available human data on cancer risk from acrylamide in food include case-control studies and reports from ongoing prospective studies. These studies provide little evidence for acrylamide-induced cancer in humans exposed to acrylamide in the diet.

2. RELEVANCE TO PUBLIC HEALTH

Results of two similarly-designed lifetime studies in orally-exposed rats provide clear evidence of acrylamide-induced carcinogenicity. Reported effects common to both studies include significantly increased incidences of thyroid (follicular cell) tumors, mesotheliomas of the tunica vaginalis testis, and mammary gland tumors. Findings of significantly increased incidences of adrenal pheochromocytomas in male rats and tumors of the oral cavity, central nervous system (glial origin), pituitary, and clitoris or uterus in female rats in the earlier bioassay were not replicated in the second bioassay.

Acrylamide was reported to initiate skin tumors in female mice administered repeated oral doses of acrylamide followed by dermal applications of 12-O-tetradecanoylphorbol-13-acetate (TPA, a tumor promoter) to the shaved back.

EPA assigned the descriptor B2 (probable human carcinogen) to acrylamide based on “inadequate human data and sufficient evidence of carcinogenicity in animals; significantly increased incidences of benign and/or malignant tumors at multiple sites in both sexes of rats, and carcinogenic effects in a series of one-year limited bioassays in mice by several routes of exposures.” EPA noted that positive genotoxicity data, adduct formation activity, and structure-activity relationships to vinyl carbamate and acrylonitrile provide support to the classification. The International Agency for Research on Cancer assigned acrylamide to Group 2A (probably carcinogenic to humans) based on similar assessment. The National Toxicology Program has determined that acrylamide is reasonably anticipated to be a human carcinogen.

Body Weight Effects. Depressed body weight, including actual body weight loss, and depressed body weight gain were consistently reported in laboratory animals following oral exposure to acrylamide. Significantly depressed mean body weight was observed in male prepubertal and adult ddy mice (range of 6–13% less than control mean body weight) was noted during 3–5 days following single oral dosing at 150 mg/kg (Sakamoto et al. 1988). Very slight to slight initial weight loss was reported following single oral administration of acrylamide at 126 mg/kg in rats and guinea pigs and 63 mg/kg in rabbits. Repeated-dose oral exposure of rats to doses as low as 15 mg/kg/day for 5 consecutive days resulted in significantly depressed body weight gain (approximately 40% less than controls); higher doses (30–60 mg/kg/day) caused actual weight loss. An 8-week dosing period to dogs at 7 mg/kg/day resulted in weight loss. Weight loss was reported in cats receiving acrylamide orally for 16 weeks at 15 mg/kg/day. A daily dose as low as 7.5 mg/kg/day during gestation days 6–20 resulted in approximately 12% depressed maternal weight gain in rats.

2. RELEVANCE TO PUBLIC HEALTH

2.3 MINIMAL RISK LEVELS (MRLs)

Estimates of exposure levels posing minimal risk to humans (MRLs) have been made for acrylamide. An MRL is defined as an estimate of daily human exposure to a substance that is likely to be without an appreciable risk of adverse effects (noncarcinogenic) over a specified duration of exposure. MRLs are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration within a given route of exposure. MRLs are based on noncancerous health effects only and do not consider carcinogenic effects. MRLs can be derived for acute, intermediate, and chronic duration exposures for inhalation and oral routes. Appropriate methodology does not exist to develop MRLs for dermal exposure.

Although methods have been established to derive these levels (Barnes and Dourson 1988; EPA 1990), uncertainties are associated with these techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of the procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

Inhalation MRLs

No inhalation MRLs were derived due to the lack of appropriate data on the effects of inhalation exposure to acrylamide.

Numerous accounts of associations between exposure to acrylamide and neurological impairment are available (Auld and Bedwell 1967; Bachmann et al. 1992; Calleman et al. 1994; Davenport et al. 1976; Donovan and Pearson 1987; Dumitru 1989; Fullerton 1969; Garland and Patterson 1967; Gjerloff et al. 2001; Hagmar et al. 2001; He et al. 1989; Igisu and Matsuoka 2002; Igisu et al. 1975; Kesson et al. 1977; Mapp et al. 1977; Mulloy 1996; Myers and Macun 1991; Takahashi et al. 1971). Most associations were made from occupational exposure scenarios that likely included inhalation and dermal (and possibly oral) exposure. Acrylamide levels in the workplace air were measured in some instances; however, reliable exposure-response data are not available.

2. RELEVANCE TO PUBLIC HEALTH

Acrylamide-induced health effects in laboratory animals following inhalation exposure to acrylamide were assessed in two studies that did not include multiple exposure levels. Mortalities and central nervous system effects were observed in dogs exposed to acrylamide dust at a concentration of 15.6 mg/m³ for 6 hours/day for up to 16 days (American Cyanamid Company 1953a). There were no signs of adverse effects in similarly-exposed guinea pigs. No adverse effects were seen in cats exposed to what was described as a “saturated” vapor for 6 hours/day, 5 days/week for 3 months at a mean analytical concentration of 1.65 ppm (4.8 mg/m³) (Dow Chemical Company 1957).

Oral MRLs

An MRL of 0.02 mg/kg/day has been derived for acute-duration oral exposure (14 days or less) to acrylamide.

Available human data are limited to case reports that include findings of persistent peripheral neuropathy in a subject who intentionally ingested 18 g of acrylamide crystals (Donovan and Pearson 1987) and signs of central and peripheral neurological deficits in family members exposed (likely via oral and dermal routes) to acrylamide in well water at a concentration of 400 ppm (Igisu and Matsuoka 2002; Igisu et al. 1975). Epidemiologic studies designed to evaluate noncancer health effects in groups of orally-exposed subjects have not been conducted.

Single oral doses elicit clinical signs of acrylamide-induced neurological effects in laboratory animals at lethal or near-lethal doses (American Cyanamid Company 1953c; Fullerton and Barnes 1966; McCollister et al. 1964; Tilson and Cabe 1979). Repeated oral dosing elicits clinical signs at lower daily dose levels. For example, convulsions and ataxia were noted as early as day 14 in rats administered acrylamide by daily gavage at 25 mg/kg/day for 21 days (Dixit et al. 1981). Daily exposure of male mice to acrylamide in the drinking water for 12 days at an estimated dose of 25.8 mg/kg/day caused decreased rotarod performance and increased hindlimb splay as early as days 6 and 8, respectively (Gilbert and Maurissen 1982).

Histopathological evaluations of nervous tissues following acute-duration oral exposure to acrylamide are limited to electron microscope results from peripheral nerve preparations in subgroups of control and high-dose (20 mg/kg/day) male rats administered acrylamide in the drinking water for up to 93 days (Burek et al. 1980). No signs of acrylamide-induced degenerative effects were observed after 7 days of treatment, but degeneration in axons and myelin sheath were noted after 33 days of treatment. These

2. RELEVANCE TO PUBLIC HEALTH

interim assessments were not performed at lower dose levels and the effect of treatment following 14 days (maximum acute-duration exposure period as defined by ATSDR) was not assessed.

Acrylamide-induced effects on body weight have been reported in rats following acute-duration oral exposure to acrylamide. In one study, repeated-dose oral exposure of rats to doses as low as 15 mg/kg/day for 5 consecutive days resulted in significantly depressed body weight gain (approximately 40% less than controls); higher doses (30–60 mg/kg/day) caused actual weight loss, whereas no effect on body weight was seen at a dose level of 5 mg/kg/day (Tyl et al. 2000b). Another study reported significantly lower mean body weight (magnitude 8%) as early as treatment day 13 in male rats administered acrylamide in the drinking water at a concentration resulting in an estimated dose level of 20 mg/kg/day, but no effect on body weight at an estimated dose level of 5 mg/kg/day (Burek et al. 1980).

Daily oral administration of acrylamide to male rats at 15 mg/kg/day for as little as 5 days followed by mating with unexposed female rats resulted in reduced mating, decreased fertility, and male-mediated increased pre- and postimplantation losses; these effects were not seen at 5 mg/kg/day (Sublet et al. 1989; Tyl et al. 2000b). A similarly-designed study (Tyl et al. 2000b) confirmed the findings of acrylamide-induced increased implantation loss, although, based on pairwise comparisons with controls, statistical significance was achieved only at the highest dose level (60 mg/kg/day). Tyl et al. (2000b) also noted significantly decreased body weight gain during the 5 days of dosing at 15 mg/kg/day and actual weight loss at higher doses, findings not included in the study report of Sublet et al. (1989). Based on the reproducible results for male-mediated implantation loss in the studies of Sublet et al. (1989) and Tyl et al. (2000b) and the lack of supporting information regarding the body weight effects, the male-mediated implantation loss was selected as the critical effect for deriving an acute-duration oral MRL for acrylamide. As discussed in detail in Appendix A, benchmark dose (BMD) analysis was applied to provide a point of departure (BMCL₁₀) of 1.78 mg/kg/day; an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability) was applied, resulting in an acute-duration oral MRL of 0.02 mg/kg/day for acrylamide.

An MRL of 0.002 mg/kg/day has been derived for intermediate-duration oral exposure (15–364 days) to acrylamide.

No human data are available regarding health effects associated with intermediate-duration oral exposure to acrylamide.

2. RELEVANCE TO PUBLIC HEALTH

Results of available animal studies demonstrate that neurological effects in males and females and reproductive effects in males are the most sensitive noncancer effects associated with intermediate-duration oral exposure to acrylamide (Burek et al. 1980; Chapin et al. 1995; Johnson et al. 1984, 1985, 1986; Sakamoto and Hashimoto 1986; Smith et al. 1986; Tyl et al. 2000a, 2000b; Zenick et al. 1986). The lowest dose level reported to induce male-mediated implantation losses was 2.8 mg/kg/day in male Long-Evans rats receiving acrylamide from the drinking water for 80 days; this study identified a no-observed-adverse effect level (NOAEL) of 1 mg/kg/day (Smith et al. 1986). Ultrastructural degenerative peripheral nerve changes were observed at a dose level as low as 1 mg/kg/day in male F344 rats receiving acrylamide from the drinking water for up to 93 days; this study identified a NOAEL of 0.2 mg/kg/day (Burek et al. 1980). Available data suggest that neurological effects represent a more sensitive point of departure than reproductive effects for deriving intermediate- and chronic-duration oral MRLs for acrylamide. Therefore, degenerative nerve change was selected as the critical effect for deriving an intermediate-duration oral MRL for acrylamide. The study of Burek et al. (1980) was selected as the principal study because it identified the lowest lowest-observed-adverse-effect-level (LOAEL) for the critical effect. A NOAEL/LOAEL approach was selected because results of the ultrastructural evaluations included only 3 of 10 rats/group and were reported only as the total numbers of fields (per group) with ultrastructural changes as axolemma invaginations or Schwann cells without axons and/or with degenerating myelin. The lack of adequate quantitative data precludes the utilization of BMD analysis to derive an intermediate-duration oral MRL for acrylamide. The NOAEL of 0.2 mg/kg/day was selected as the point of departure and was divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability) to yield an intermediate-duration oral MRL of 0.002 mg/kg/day for acrylamide.

No chronic-duration oral MRL was derived for acrylamide.

No human data are available to evaluate neurological effects following chronic-duration oral exposure to acrylamide. Two chronic toxicity and carcinogenicity bioassays are available for male and female F344 rats receiving acrylamide from the drinking water for up to 2 years (Friedman et al. 1995; Johnson et al. 1984, 1985, 1986). One study included dose levels of 0, 0.01, 0.1, 0.5, and 2.0 mg/kg/day (Johnson et al. 1984, 1985, 1986); the other study included dose levels of 0, 0.1, 0.5, or 2 mg/kg/day in males and 0, 1.0, or 3.0 mg/kg/day in females (Friedman et al. 1995). Both studies identified a NOAEL of 0.5 mg/kg/day and a LOAEL of 2 mg/kg/day for histopathological evidence of degenerative nerve changes. These values are slightly higher than the NOAEL of 0.2 mg/kg/day and the LOAEL of 1 mg/kg/day for degenerative nerve changes in the intermediate-duration oral study used to derive an intermediate-

2. RELEVANCE TO PUBLIC HEALTH

duration oral MRL for acrylamide. These differences may be solely the result of differences in dose spacing. Because the NOAEL of 0.5 mg/kg/day used as a potential point of departure for deriving a chronic-duration oral MRL for acrylamide is slightly higher than the NOAEL of 0.2 mg/kg/day that serves as the point of departure for the intermediate-duration oral MRL of 0.002 mg/kg/day, the intermediate-duration oral MRL of 0.002 mg/kg/day is expected to be protective of chronic-duration oral exposure to acrylamide.

3. HEALTH EFFECTS

3.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of acrylamide. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals and others address the needs of persons living or working near hazardous waste sites, the information in this section is organized first by route of exposure (inhalation, oral, and dermal) and then by health effect (death, systemic, immunological, neurological, reproductive, developmental, genotoxic, and carcinogenic effects). These data are discussed in terms of three exposure periods: acute (14 days or less), intermediate (15–364 days), and chronic (365 days or more).

Levels of significant exposure for each route and duration are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. "Serious" effects are those that evoke failure in a biological system and can lead to morbidity or mortality (e.g., acute respiratory distress or death). "Less serious" effects are those that are not expected to cause significant dysfunction or death, or those whose significance to the organism is not entirely clear. ATSDR acknowledges that a considerable amount of judgment may be required in establishing whether an end point should be classified as a NOAEL, "less serious" LOAEL, or "serious" LOAEL, and that in some cases, there will be insufficient data to decide whether the effect is indicative of significant dysfunction. However, the Agency has established guidelines and policies that are used to classify these end points. ATSDR believes that there is sufficient merit in this approach to warrant an attempt at distinguishing between "less serious" and "serious" effects. The distinction between "less serious" effects and "serious" effects is considered to be important because it helps the users of the profiles to identify levels of exposure at which major health effects start to appear. LOAELs or NOAELs should also help in determining whether or not

3. HEALTH EFFECTS

the effects vary with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown in the Levels of Significant Exposure (LSE) tables and figures may differ depending on the user's perspective. Public health officials and others concerned with appropriate actions to take at hazardous waste sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAELs) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans (Minimal Risk Levels or MRLs) may be of interest to health professionals and citizens alike.

Levels of exposure associated with carcinogenic effects (Cancer Effect Levels, CELs) of acrylamide are indicated in Table 3-4 and Figure 3-2. Because cancer effects could occur at lower exposure levels, Figure 3-2 also shows a range for the upper bound of estimated excess risks, ranging from a risk of 1 in 10,000 to 1 in 10,000,000 (10^{-4} to 10^{-7}), as developed by EPA.

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

3.2.1 Inhalation Exposure

3.2.1.1 Death

Human data are available from two cohort mortality studies of occupational exposure to acrylamide, one by Collins et al. (1989) with most recent follow up by Marsh et al. (2007) and one by Sobel et al. (1986) with follow up by Swaen et al. (2007). In these studies, no significant associations were found between occupational exposure to acrylamide and incidences of death from all causes. See Section 3.1.2.7 (Cancer) for more detailed information regarding these cohorts and assessments of death due to cancers.

Reliable information regarding death in animals following inhalation exposure to acrylamide is limited. In a study performed for the American Cyanamid Company (1953a), two dogs, seven rats, and seven guinea pigs (sex and strain unspecified) were exposed to acrylamide dust at 15.6 mg/m^3 for 6 hours/day, 5 days/week for up to 12 exposures in a 16-day period. Four of the seven rats died overnight following the first exposure period and two of the remaining rats died a few days later. One of the dogs died on

3. HEALTH EFFECTS

study day 15; there were no deaths among the guinea pigs. The study authors stated that >90% of the particles were in the respirable range (0.3–1.2 μm).

Reliable inhalation mortality data for each species are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.2 Systemic Effects

No human or animal data were located regarding cardiovascular, gastrointestinal, musculoskeletal, hepatic, renal, endocrine, or dermal effects following inhalation exposure to acrylamide.

Respiratory Effects. Available information regarding acrylamide-associated respiratory effects is restricted to complaints of nose and throat irritation in a group of tunnel workers who had been occupationally exposed to acrylamide and N-methylolacrylamide in a chemical grouting agent for 2 months (Hagmar et al. 2001). Increasing incidences of complaints were associated with increasing levels of hemoglobin adducts of acrylamide.

Hematological Effects. No data were located regarding hematological effects in humans following inhalation of acrylamide. Available information in animals is limited to a study in which four rats were exposed to acrylamide as a “saturated” vapor for 6 hours/day, 5 days/week for 3 months at a mean analytical concentration of 1.65 ppm (4.8 mg/m³); the exposures did not affect hematology results (American Cyanamid Company 1954).

Ocular Effects. Available information regarding acrylamide-associated ocular effects is restricted to complaints of eye irritation in a group of tunnel workers who had been occupationally exposed to acrylamide and N-methylolacrylamide in a chemical grouting agent for 2 months (Hagmar et al. 2001). Increasing incidences of complaints were associated with increasing levels of hemoglobin adducts of acrylamide.

Body Weight Effects. No data were located regarding body weight effects in humans following inhalation of acrylamide. Available information in animals is limited to a study in which four rats were exposed to acrylamide as a “saturated” vapor for 6 hours/day, 5 days/week for 3 months at a mean

Table 3-1 Levels of Significant Exposure to Acrylamide - Inhalation

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/m ³)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/m ³)	Serious (mg/m ³)		
ACUTE EXPOSURE								
Death								
1	Rat (NS)	4 d 6 hr/d				15.6	(death of 4/7 rats in 1 day and 2 others before day 8)	American Cyanamid Company 1953a
Neurological								
2	Rat (NS)	4 d 6 hr/d				15.6	(loss of coordination and equilibrium on exposure day 4)	American Cyanamid Company 1953a
INTERMEDIATE EXPOSURE								
Death								
3	Dog (NS)	16 d 5 d/wk 6 hr/d				15.6	(death of 1/2 dogs on day 15)	American Cyanamid Company 1953a
Systemic								
4	Cat	3 mo 5 d/wk 6 hr/d	Hemato	4.8				American Cyanamid Company 1954 Acrylamide
			Bd Wt	4.8				
Neurological								
5	Gn Pig (NS)	16 d 5 d/wk 6 hr/d		15.6				American Cyanamid Company 1953a
6	Dog (NS)	16 d 5 d/wk 6 hr/d				15.6	(CNS effects including loss of equilibrium and coordination)	American Cyanamid Company 1953a

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Acrylamide - Inhalation

(continued)


Key to Figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Route)	System	NOAEL (mg/m ³)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/m ³)	Serious (mg/m ³)		
7	Cat	3 mo 5 d/wk 6 hr/d		4.8			American Cyanamid Company 1954 Acrylamide	

a The number corresponds to entries in Figure 3-1.

Bd Wt = body weight; d = day(s); Hemato = hematological; hr = hour(s); LOAEL = lowest-observed-adverse-effect level; mo = month(s); NOAEL = no-observed-adverse-effect level; NS = not specified; week(s)

DRAFT FOR PUBLIC COMMENT

Figure 3-1 Levels of Significant Exposure to Acrylamide - Inhalation



DRAFT FOR PUBLIC COMMENT

3. HEALTH EFFECTS

analytical concentration of 1.65 ppm (4.8 mg/m³); the exposures did not affect body weights (American Cyanamid Company 1954).

3.2.1.3 Immunological and Lymphoreticular Effects

No data were located regarding immunological or lymphoreticular effects in humans or animals following inhalation exposure to acrylamide.

3.2.1.4 Neurological Effects

Information in humans is available from numerous case reports in which acrylamide exposure has been associated with signs of impaired neurological performance in central and peripheral nervous systems that include impaired motor function and muscle weakness (Auld and Bedwell 1967; Davenport et al. 1976; Dumitru 1989; Fullerton 1969; Garland and Patterson 1967; Gjerløff et al. 2001; Igisu et al. 1975; Kesson et al. 1977; Mapp et al. 1977; Mulloy 1996; Takahashi et al. 1971). Human data are also available from cross-sectional studies that included self-reported symptoms and neurological evaluations of acrylamide-exposed workers with potential for inhalation and dermal (and possibly oral) exposure (Bachmann et al. 1992; Calleman et al. 1994; Hagmar et al. 2001; He et al. 1989; Myers and Macun 1991). Although the case reports and cross-sectional studies provide supportive evidence of acrylamide-induced neurotoxicity, they lack information regarding relative contributions of natural exposure routes (inhalation, oral, dermal), exposure-response relationships, and other confounding exposures. They are therefore unsuitable for meaningful quantitative risk analysis.

He et al. (1989) evaluated health end points in workers employed for 1–18 months at a factory in China that began producing acrylamide and polyacrylamide in 1984. A referent group consisted of unexposed workers from the same town. Concentrations of acrylamide in the workplace air (determined by gas chromatography) reached 5.56–9.02 mg/m³ between March and June 1985 during polymerization when there was an exceptional increase in production, and decreased to an average of 0.0324 mg/m³ (range not specified) after July 1985. The workers were evaluated in October 1985. The study authors reported that heavy skin contamination by aqueous acrylamide monomer was common among the workers. An acrylamide level of 410 mg/L was measured in the water in which three of the workers washed their hands. Personal interviews were conducted to obtain information on demographic factors, occupational history, symptoms, past illnesses, and family history. Physical and neurological examinations, visual acuity and visual field testing, skin temperature measurements, electrocardiography, and

3. HEALTH EFFECTS

electroencephalography were performed. Sixty-nine of the exposed workers and 48 of the referent workers were subjected to electroneuromyographic examinations.

As shown in Table 3-2, significantly greater percentages of the acrylamide-exposed group reported skin peeling from the hands, anorexia, numbness and coldness in hands and feet, lassitude, sleepiness, muscle weakness, clumsiness of the hands, unsteady gait, difficulty in grasping, and stumbling and falling. The authors stated that initial symptoms of skin peeling were the result of dermal exposure to aqueous acrylamide and that other symptoms appeared following 3–10 months of occupational exposure. Greater percentages of acrylamide-exposed workers exhibited erythema of the hands, sensory impairments (vibration, pain, and touch sensation), diminished reflexes, and intention tremor (Table 3-2). Electrical activity, monitored in both the abductor pollicis brevis and abductor digiti minimi muscles of the hand, revealed electromyographic abnormalities in the acrylamide-exposed workers that included denervation potentials (3/69 exposed workers), prolonged duration of motor units (40/69), increased polyphasic potentials (29/69), and discrete pattern of recruitment (9/69). These abnormalities were not seen in referent workers, with the exception of prolonged duration of motor units (4/48 referents). Significantly increased mean duration and mean amplitude of motor unit potentials and significantly decreased mean amplitude of sensory unit potentials were seen in the acrylamide-exposed group compared to the referent group. Assessment of visual acuity and visual field, nerve conduction velocity, electrocardiography, and electroencephalography revealed no significant exposure-related effects.

Calleman et al. (1994) performed a cross-sectional analysis of hemoglobin adduct formation and neurological effects in a group of 41 factory workers who were exposed to acrylamide (and acrylonitrile, from which acrylamide is formed) for 1 month to 11.5 years (mean 3 years) during the production of acrylamide and polyacrylamide at a factory in China. As determined by station sampling and gas chromatography, mean acrylamide air concentrations were 1.07 and 3.27 mg/m³ in the synthesis and polymerization rooms, respectively, during the summer of 1991. Mean exposure concentrations during the time of collection of biomarker data (September 1991) were lower, averaging 0.61 and 0.58 mg/m³ in synthesis and polymerization rooms, respectively. Information regarding demographic factors, smoking and drinking habits, height and weight, occupational history, past illnesses, current symptoms, and reproductive history were collected by questionnaire. Neurological examinations were performed approximately 1 hour after a work shift. Vibration sensitivity thresholds were measured in fingers and

3. HEALTH EFFECTS

Table 3-2. Self-Reported Neurological Symptoms and Observed Clinical Signs Among Acrylamide Workers and Nonexposed Workers

Symptom	Acrylamide group (n=71)		Reference group (n=51)	
	Number	Percent	Number	Percent
Skin peeling from the hands	38	53.5 ^a	2	3.9
Numbness in the hands and feet	15	21.1 ^b	2	3.9
Lassitude	14	19.7 ^b	1	1.9
Sleepiness	12	16.9 ^b	0	0
Muscle weakness	11	15.4 ^b	0	0
Clumsiness of the hands	8	11.2 ^a	0	0
Anorexia	8	11.2 ^a	1	1.9
Unsteady gait	6	8.4 ^a	0	0
Coldness of the hands and feet	6	8.4 ^a	0	0
Difficulty in grasping	5	7.0 ^a	0	0
Stumbling and falling	5	7.0 ^a	0	0
Sweating	27	38.0	14	27.4
Dizziness	7	9.8	2	3.9
Cramping pain	6	8.4	5	9.8
Sign				
Erythema of hands	16	22.5 ^b	0	0
Skin peeling from the hands	16	22.5 ^b	1	1.9
Sensory impairments				
Vibration sensation	12	16.9 ^b	0	0
Pain sensation	7	9.8 ^a	0	0
Touch sensation	6	8.4 ^a	0	0
Position sensation	1	1.4	0	0
Muscle atrophy in hands	4	5.6	0	0
Diminished reflexes				
Biceps	12	16.9 ^b	0	0
Triceps	10	14.0	4	7.8
Knee	11	15.4 ^a	1	1.9
Ankle	8	11.2 ^a	1	1.9
Loss of reflexes				
Biceps	3	4.2	0	0
Triceps	5	7.0	1	1.9
Knee	5	7.0 ^a	0	0
Ankle	17	23.9 ^b	0	0
Intention tremor	13	18.3 ^a	2	3.9
Positive Romberg's sign	15	21.1	3	5.8

^ap <0.05^bp <0.01 (χ^2 test)

Source: He et al. 1989

3. HEALTH EFFECTS

toes. Physical and neurological examinations and electroneuromyographic (ENMG) testing were performed. For each test, a nonexposed referent group was included. Quantitative assessment of contributions of dermal and inhalation exposure were not made, although in the synthesis area of the factory where neurological symptoms were most severe, dermal contact was considered to have been the major exposure route.

As shown in Table 3-3, a variety of symptoms and signs of adverse health effects were noted in acrylamide-exposed workers (Calleman et al. 1994). Other significant ($p < 0.01$) effects in the exposed workers included increased (magnitude $\geq 60\%$) vibration threshold, decreased (10–20%) conduction velocity in the peroneal and sural nerves, and increased (25–36%) latency in median, ulnar, and peroneal nerves. Neurotoxicity index scores, a quantitative expression of the severity of peripheral neuropathy, decreased with physical distance from the synthesis room where the monomer itself was handled. This relationship was not reflected by results of hand or foot vibration sensitivity measurements.

Hagmar et al. (2001) performed a health examination on a group of 210 tunnel construction workers who had been occupationally exposed for 2 months to a chemical grouting agent containing acrylamide and N-methylolacrylamide. Workers were expected to have experienced dermal exposure as well as inhalation exposure. Venous blood samples were drawn and questionnaires and physical examinations were administered 1–5 weeks after exposure was stopped. Quantitative exposure data were limited to two personal air samples showing concentrations of 0.27 and 0.34 mg/m³ for the sum of acrylamide and N-methylolacrylamide; further analysis suggested that the air contained a 50:50 mixture of these compounds. The health examination included an extensive questionnaire and a physical examination that included unspecified tests of peripheral nerve function. Blood samples for the analysis of adducts of acrylamide with N-terminal valines in hemoglobin were drawn within a month after construction work was completed. A group of 50 subjects who claimed recently developed or deteriorated peripheral nervous function at the initial physical examination was subjected to more detailed neurophysiologic examinations and 6-month follow-up clinical ($n=29$) and neurophysiological ($n=26$) examinations. Those with remaining symptoms were examined for up to 18 months postexposure.

Hemoglobin adduct levels for 18 nonsmoking unexposed referents varied between 0.02 and 0.07 nmol/g globin. Adduct levels in 47 of the 210 tunnel workers did not exceed the highest level of the referents. The remaining workers were divided into three categories according to adduct levels as follows: 89 with

3. HEALTH EFFECTS

Table 3-3. Prevalence of Symptoms and Signs of Adverse Health Effects in Acrylamide-Exposed Workers and Controls

Symptom or sign	Exposed (percent)	Controls (n=10)
Numbness of extremities	29/41 (71) ^a	0
Fatigue	29/41 (71) ^a	0
Sweating of hands and feet	28/41 (71) ^a	0
Skin peeling	24/41 (59) ^a	0
Menstruation disorders	4/7 (57)	NA
Loss of pain sensation	22/41 (54) ^a	0
Loss of touch sensation	19/41 (46) ^b	0
Dizziness	18/41 (44) ^b	0
Anorexia	17/41 (41) ^b	0
Loss of vibration sensation	17/41 (41) ^b	0
Nausea	16/41 (39) ^b	0
Loss of ankle reflexes	12/41 (29)	0
Headache	11/41 (27)	0
Unsteady gait	9/41 (22)	0
Loss of knee jerk	8/41 (20)	0
Unsteady Romberg sign	8/41 (20)	0
Loss of triceps reflexes	4/41 (10)	0
Loss of biceps reflexes	4/41 (10)	0

^ap <0.01 (χ^2 test)^bp <0.05

Source: Calleman et al. 1994

3. HEALTH EFFECTS

0.08–0.29 nmol/g globin, 36 with 0.3–1.0 nmol/g globin, and 38 with 1.0–17.7 nmol/g globin. The study authors noted a significant ($p < 0.05$) association between self-reported exposure categories and adduct levels. Significant positive correlations ($p < 0.05$) between prevalence of self-reported peripheral nervous symptoms, irritant symptoms, and symptoms of general discomfort with adduct levels were found. For example, in the groups with adduct levels < 0.08 nmol/g globin, 0.08–0.29 nmol/g globin, 0.3–1.0 nmol/g globin, and > 1.0 nmol/g globin, incidences of reported numbness or tingling in the feet or legs were 2/47 (4%), 10/89 (11%), 9/36 (25%), and 14/38 (37%), respectively. Irritant symptoms and symptoms of general discomfort typically disappeared following the end of a workday, whereas peripheral nervous symptoms persisted. Follow-up examinations revealed that 58% of the subjects with early signs of impaired peripheral nervous function improved, while only 4% showed signs of deterioration.

Myers and Macun (1991) investigated peripheral neuropathy in a cohort of 66 workers in a South African factory that produced polyacrylamide. The investigation followed clinical diagnosis of peripheral neuropathy in five workers at the factory. The workforce was divided into a number of exposure categories, based on environmental sampling and discussions with workers. Exposure levels for the various tasks ranged from 0.07 to 2.5 times the National Institute of Occupational Safety and Health (NIOSH) recommended exposure limit (REL) of 0.3 mg/m^3 . Workers were then classified as being exposed to airborne acrylamide when exposure levels exceeded the REL ($n=22$), and unexposed when exposure levels were below the REL ($n=41$). Workers completed a questionnaire that was designed to capture social, medical, and occupational history. A standard blind neurological examination was also performed.

The exposed group showed higher prevalences of abnormalities for all symptoms (weakness, sensation, balance, fatigue, visual, loss of weight, urogenital, and fingertip skin), most signs (fingertip effects, light touch, tactile discrimination, pain), and reflexes, coordination, motor weakness, gait, and Rombergism. Statistically significant differences between exposed and unexposed groups for individual effects were seen only for abnormal sensation symptoms and signs in fingertip skin (including color, peeling, and sweating). The overall prevalence of acrylamide-related abnormalities among the exposed was 66.7%, which was statistically significantly higher ($p < 0.05$) than that of the unexposed group (prevalence of 14.3%). The authors stated that most workers observed to have abnormalities (number not reported) were employed in areas where exposures were highest (1.6–2.5 times the REL).

Bachmann et al. (1992) performed a follow-up investigation in July 1990 at the same South African factory that had been examined in 1986 by Myers and Macun (1991). The study design was similar to

3. HEALTH EFFECTS

that of Myers and Macun (1991), but included measurements of vibration sensation threshold. Among 82 workers employed at follow-up, increased prevalences of symptoms of tingling and numbness in hands and feet, weakness and pain in arms and legs, peeling hand skin, and sweating hands were reported by exposed workers, compared with those classified as being unexposed. The symptoms of numbness, limb pain, and peeling and sweating of hands were statistically significantly increased in exposed workers. Results of clinical examinations provided supporting evidence for the reported increased symptoms of peeling and sweating of the hands. No gross neurological abnormalities were found. Mean vibration sensation thresholds were similar among unexposed and exposed groups, even when adjusting for age, and no association was found between vibration thresholds and any symptoms.

Information regarding neurological effects in animals exposed to acrylamide by the inhalation route is limited to a single study report in which seven rats, seven guinea pigs, and two dogs were exposed to dust of acrylamide at an analytical concentration of 15.6 mg/m³ for 6 hours/day, 5 days/week for up to 12 exposures in a 16-day period (American Cyanamid Company 1953a). Reported signs of neurological effects in the three rats that survived to exposure termination on day 4 included loss of equilibrium and coordination. There was no mention of neurological signs in the exposed dogs, although one of the dogs lost weight and died on day 15. No toxic signs were seen in the guinea pigs.

3.2.1.5 Reproductive Effects

No data were located regarding reproductive effects in humans or animals following inhalation exposure to acrylamide.

3.2.1.6 Developmental Effects

No data were located regarding developmental effects in humans or animals following inhalation exposure to acrylamide.

3.2.1.7 Cancer

Human data are available from two cohort mortality studies of occupational exposure to acrylamide, one by Collins et al. (1989) with most recent follow up by Marsh et al. (2007) and one by Sobel et al. (1986) with follow up by Swaen et al. (2007). Exposure to acrylamide was considered to have occurred primarily via inhalation and dermal exposure.

3. HEALTH EFFECTS

Collins et al. (1989) conducted a cohort mortality study of all male workers (8,854, of which 2,293 were exposed to acrylamide) who had been hired between January 1, 1925 and January 31, 1973 at four American Cyanamid factories, three in the United States (Fortier, Louisiana [1295 workers]; Warners, New Jersey [7,153 workers]; and Kalamazoo, Michigan [60 workers]) and one in the Netherlands (Botlek [346 workers]). Estimations of acrylamide exposure were based on available monitoring data and worker knowledge of past jobs and processes. Mortality rates among the factory workers were compared with the expected number of deaths among men of the United States from 1925 to 1980 or the Netherlands from 1950 to 1982 to derive standardized mortality ratios (SMRs) as a measure of relative risk for each cohort. No statistically significantly elevated all cause or cause-specific SMRs were found among acrylamide-exposed workers (including cancer of the digestive or respiratory systems, bone, skin, reproductive organs, bladder, kidney, eye, central nervous system, thyroid, or lymphatic system). Trend tests showed no increased risk of mortality due to cancer at several sites (digestive tract, respiratory system, prostate, central nervous system, or lymphopoietic system) with increasing level of exposure to acrylamide.

The most recent update report of the cohort of Collins et al. (1989) includes study periods of 1925–2002 for the 8,508 workers in the three facilities in the United States and 1965–2004 for the 344 workers at the Botlek plant in the Netherlands (the original cohort of 346 people included 2 females who were excluded in the follow up) (Marsh et al. 2007). Among the workers at the three facilities in the United States (during which 4,650 deaths occurred among the 8,508 workers in the period of 1925–2002), excess and deficit overall mortality risks were observed for cancer sites implicated in oral studies in experimental animals: brain and other central nervous system (SMR 0.67, 95% confidence interval [CI] 0.40–1.05), thyroid gland (SMR 1.38, 95% CI 0.28–4.02), and testis and other male genital organs (SMR 0.64, 95% CI 0.08–2.30); and for sites selected in the original report (Collins et al. 1989) of this cohort: respiratory system cancer (SMR 1.17, 95% CI 1.06–1.27), esophagus (SMR 1.20, 95% CI 0.86–1.63), rectum (SMR 1.25, 95% CI 0.84–1.78), pancreas (SMR 0.94, 95% CI 0.70–1.22), and kidney (SMR 1.01, 95% CI 0.66–1.46). None of the mortality excesses were statistically significant, except for respiratory system cancer, which Collins et al. (1989) attributed to muriatic acid (hydrochloric acid) exposure. No significantly elevated SMRs were found for rectal, pancreatic, or kidney cancers in exploratory exposure-response analyses conducted according to the following exposure parameters and categories: duration of employment (<1, 1–, and 15+ years), time since first employment (<20, 20–, and 30+ years), duration of exposure (unexposed, 0.001–, 5–, and 20+ years), cumulative exposure (<0.001, 0.001–, 0.03–, and 0.30+ mg/m³-years), and estimated mean exposure concentrations (unexposed, 0.001–, 0.02–, and 0.3+ mg/m³).

3. HEALTH EFFECTS

Sobel et al. (1986) conducted a mortality study on a cohort of 371 workers assigned to acrylamide and polymerization operations at a Dow Chemical facility in the United States between 1955 and 1979. Analysis and review of air monitoring data and job classifications resulted in estimates of personal 8-hour time-weighted average acrylamide concentrations of 0.1–1.0 mg/m³ before 1957, 0.1–0.6 mg/m³ from 1957 to 1970, and 0.1 mg/m³ thereafter. SMRs, calculated for categories in which at least two deaths were observed, were based on mortality of white males in the United States. No significantly increased incidences of cancer-related deaths were observed within the cohort.

Followup to the Sobel et al. (1986) study cohort was expanded to include employees hired through 2001 (Swaen et al. 2007). Exposure to acrylamide was retrospectively assessed based on personal samples from the 1970s onwards and area samples from the entire study period. Fewer acrylamide workers died (n=141) than expected (n=172.1). No cause-specific SMR for any of the investigated types of cancer was exposure related. The authors reported more total pancreatic cancer deaths (n=5) than expected (n=2.3) (SMR 222.2, 95% CI 72.1–518.5); however, three of the five were in the low-dose group, with no apparent dose-response relationship with acrylamide exposure.

No data were located regarding cancer in animals following inhalation exposure to acrylamide. EPA (IRIS 2009) calculated an inhalation unit risk of $1.3 \times 10^{-3} \text{ (mg/m}^3\text{)}^{-1}$ for acrylamide based on results of a 2-year cancer bioassay in orally-exposed male and female F344 rats (Johnson et al. 1986) and route-to-route extrapolation (see Section 3.2.2.7 for details of the oral study). The air concentrations associated with risk of 1×10^{-4} , 1×10^{-5} , and 1×10^{-6} are 0.08, 0.008, and 0.0008 mg/m³, respectively.

3. HEALTH EFFECTS

3.2.2 Oral Exposure**3.2.2.1 Death**

There are no reports of human deaths associated with oral exposure to acrylamide.

Acrylamide has been demonstrated to be lethal to laboratory animals following a single oral dose. Reported oral LD₅₀ values in rats range from 150 to 413 mg/kg (American Cyanamid Company 1973, 1977; Dow Chemical Company 1957; Fullerton and Barnes 1966; McCollister et al. 1964; Tilson and Cabe 1979; Union Carbide Corporation 1947). Reported LD₅₀ values in mice, guinea pigs, and rabbits range from 107 to 195 mg/kg (American Cyanamid Company 1951; Dow Chemical Company 1957; Hashimoto et al. 1981; McCollister et al. 1964).

Repeated oral exposure to acrylamide has also been associated with death in laboratory animals. In one rat study, a single 100 mg/kg dose was not lethal, but two 100 mg/kg doses administered 24 hours apart resulted in mortalities (Fullerton and Barnes 1966). Sublet et al. (1989) reported death in a group of male Long-Evans hooded rats administered acrylamide by daily gavage for 5 days at a dose of 75 mg/kg/day. Longer repeated-dose exposure periods to lower daily doses are lethal as well. For example, a dose level of 50 mg/kg was lethal to rats receiving 12 daily gavage doses of 50 mg/kg in a 15-day period; the deaths occurred within a few days following the cessation of dosing (Fullerton and Barnes 1966). An acrylamide gavage dose of 30 mg/kg/day resulted in the death of 4/10 male and 2/10 female rats during the third week of daily dosing (Schulze and Boysen 1991). During the last 4 months of a 2-year study in which male and female rats received acrylamide from the drinking water at an estimated dose level of 2 mg/kg/day, decreased survival of both sexes was noted; the increased mortality was statistically significant by study termination (Johnson et al. 1984, 1986).

Reliable acute oral LD₅₀ values for death and other mortality data for each species are recorded in Table 3-4 and plotted in Figure 3-2.

3.2.2.2 Systemic Effects

The highest NOAEL values and all LOAEL values from each reliable study for systemic effects in each species and duration category are recorded in Table 3-4 and plotted in Figure 3-2.

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
ACUTE EXPOSURE								
Death								
1	Rat (Wistar)	Once (GW)				294 M (LD50)	American Cyanamid Company 1973	
2	Rat (Sprague-Dawley)	Once (GW)				413 M (LD50)	American Cyanamid Company 1977	
3	Rat (NS)	Once (GW)				180 (LD50)	Dow Chemical Company 1957; McCollister et al. 1964	
4	Rat (albino)	Once (GW)				203 F (LD50)	Fullerton and Barnes 1966 Acrylamide	
5	Rat (albino)	2 d 1 x/d (GW)				100 (most rats died within 3 days following 2 days of dosing)	Fullerton and Barnes 1966 Acrylamide	
6	Rat (Fischer- 344)	Once (GW)				175 M (LD50)	Tilson and Cabe 1979 Acrylamide	
7	Rat (albino)	Once (GW)				316 M (LD50)	Union Carbide Corporation 1947	
8	Mouse (albino)	Once (G)				195 M (LD50)	American Cyanamid Company 1951	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
9	Mouse	Once (NS)				107 M (LD50)	Hashimoto et al. 1981 Acrylamide	
10	Gn Pig (NS)	Once (GW)				180 (LD50)	Dow Chemical Company 1957; McCollister et al. 1964	
11	Rabbit (NS)	Once (GW)				150 (LD50)	Dow Chemical Company 1957; McCollister et al. 1964	
Systemic								
12	Rat (Fischer- 344)	13 d (W)	Bd Wt	5 M	20 M (8% decreased mean body weight)		Burek et al. 1980 Acrylamide	
13	Rat (Long- Evans)	5 d 1 x/d (GW)	Bd Wt	5 M	15 M (significantly depressed body weight gain during 5 days of acrylamide administration)		Tyl et al. 2000b Acrylamide	
14	Rat (Fischer- 344)	Gd 6-17 (GW)	Bd Wt	20 F			Walden et al. 1981 Acrylamide	Maternal body weight
15	Mouse ddY	Once (NS)	Bd Wt	100 M	150 M (significantly depressed body weight)		Sakamoto et al. 1988 Acrylamide	
16	Gn Pig (NS)	Once (GW)	Bd Wt		126 (very slight initial body weight loss)		Dow Chemical Company 1957; McCollister et al. 1964	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
17	Rabbit (NS)	Once (GW)	Bd Wt		63 (slight initial weight loss)		Dow Chemical Company 1957; McCollister et al. 1964	
Neurological								
18	Rat (Fischer- 344)	7 d (W)		20 M			Burek et al. 1980 Acrylamide	
19	Rat (Wistar)	Up to 21 d 1 x/d (G)				25 M (convulsions and ataxia as early as treatment day 14)	Dixit et al. 1981 Acrylamide	
20	Rat (albino)	Once (GW)				203 F (fine tremors)	Fullerton and Barnes 1966 Acrylamide	
21	Rat (albino)	Once (GW)				100 (fine tremors)	Fullerton and Barnes 1966 Acrylamide	
22	Rat (albino)	2 d 1 x/d (GW)				100 (generalized weakness)	Fullerton and Barnes 1966 Acrylamide	
23	Rat (NS)	Once (GW)				126 F (lethargy)	McCollister et al. 1964 Acrylamide	
24	Rat (Fischer- 344)	Once (GW)		100 M		200 M (decreases in hindlimb grip strength and locomotory performance)	Tilson and Cabe 1979 Acrylamide	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments	
					Less Serious (mg/kg/day)	Serious (mg/kg/day)			
25	Rat (Long- Evans)	5 d 1 x/d (GW)		30 M		45 M (clinical signs of neurotoxicity)	Tyl et al. 2000b Acrylamide		
26	Rat (Fischer- 344)	Gd 6-17 (GW)		20 F			Walden et al. 1981 Acrylamide		
27	Mouse (BALB/c)	12 d (W)				25.8 F (decreased rotarod performance, increased hindlimb splay as early as days 6-8)	Gilbert and Maurissen 1982 Acrylamide		
28	Dog (Mongrel)	Once (C)				100 (severe neurological impairment of the limbs)	American Cyanamid Company 1953c		
29	Rabbit (NS)	Once (GW)				126 (tremors)	McCollister et al. 1964 Acrylamide		
Reproductive									
30	Rat (Long- Evans)	5 d 1 x/d (GW)		^b 5 M		15 M (depressed fertility, increased preimplantation loss)	Sublet et al. 1989 Acrylamide		

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
31	Rat (Long- Evans)	5 d 1 x/d (GW)		30 M		45 M (significantly increased postimplantation losses)	Tyl et al. 2000b Acrylamide	A significant trend for increased postimplantation loss was observed at doses from 15 to 60 mg/kg/day
32	Rat (Fischer- 344)	5 d 1 x/d (GW)				30 M (significantly elevated pre- and post-implantation losses)	Working et al. 1987 Acrylamide	
33	Mouse ddY	Once (NS)				100 M (histologic abnormalities in spermatids)	Sakamoto et al. 1988 Acrylamide	
INTERMEDIATE EXPOSURE								
Death								
34	Rat (albino)	21 d (F)				47 M (4/10 deaths during 21 days of exposure)	American Cyanamid Company 1953b	
35	Rat (Fischer- 344)	28 d (W)				25 M (death of 8/10 male rats during exposure week 4) 24 F (death of 5/10 female rats during exposure week 4)	American Cyanamid Company 1991	

DRAFT FOR PUBLIC COMMENT

ACRYLAMIDE

3. HEALTH EFFECTS

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
36	Rat (albino)	12 doses in 15 d (GW)				50	(death within a few days posttreatment)	Fullerton and Barnes 1966 Acrylamide
37	Rat (Sprague-Dawley)	5 wk 7 d/wk 1 x/d (GW)				30	(death of 4/10 males and 2/10 females during treatment week 3)	Schulze and Boysen 1991 Acrylamide
Systemic								
38	Rat (albino)	21 d (F)	Bd Wt			47 M	(severely depressed body weight gain)	American Cyanamid Company 1953b Magnitude not specified
39	Rat (Sprague-Dawley)	2 wk pre mating and Gd 0-19 (F)	Bd Wt	3.82 F				American Cyanamid Company 1979 Acrylamide
40	Rat (Fischer- 344) (W)	28 d (W)	Endocr	12 M	19 M (decreased serum testosterone, 73% less than controls)			American Cyanamid Company 1991
			Bd Wt	12 M 9 F		19	(emaciation)	
41	Rat (Fischer- 344) (W)	up to 93 d (W)	Hemato	1 F	5 F (decreases in packed cell volume, erythrocytes, hemoglobin)			Burek et al. 1980 Acrylamide

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
42	Rat (Sprague- Dawley)	Gd 6-20 1 x/d (GW)	Bd Wt	2.5 F	7.5 F (12% decreased maternal weight gain)		Field et al. 1990 Acrylamide	Weight gain minus gravid uterine weight
43	Rat (Wistar)	21 d lactation period 1 x/d (G)	Bd Wt			25 F (net maternal weight loss during 21-day lactation treatment period)	Friedman et al. 1999 Acrylamide	
44	Rat (Sprague- Dawley)	5 wk 7 d/wk 1x/d (GW)	Bd Wt			30 (decreased mean body weight, 27% lower than controls)	Schulze and Boysen 1991 Acrylamide	
45	Rat (Wistar)	90 d (W)	Bd Wt	23.7 M			Tanii and Hashimoto 1983 Acrylamide	
46	Rat (Sprague- Dawley)	Gd 6- Ld 10 1 x/d (GW)	Bd Wt	5 F		10 F (approximately 33% depressed maternal body weight gain during 10 days of postpartum exposure)	Wise et al. 1995 Acrylamide	
47	Mouse (CD-1)	Gd 6-20 1 x/d (GW)	Bd Wt	45 F			Field et al. 1990 Acrylamide	
48	Mouse (ICR)	Up to 38 d (W)	Bd Wt			90.8 M (body weight loss)	Ko et al. 1999 Acrylamide	

DRAFT FOR PUBLIC COMMENT

ACRYLAMIDE

3. HEALTH EFFECTS

37

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
49	Dog	8 wk (F)	Bd Wt			7 (up to 12% weight loss)	Satchell and McLeod 1981 Acrylamide	
50	Cat	Up to 16 wk (F)	Bd Wt		15 (body weight loss, magnitude unspecified)		Post and McLeod 1977a Acrylamide	
Neurological								
51	Monkey	44-61 d 5 d/wk 1 x/d				10 F (clinical signs of peripheral neuropathy)	Dow Chemical Company 1981; Maurissen et al. 1983 Acrylamide	Histopathological evaluation of nerve tissue not performed
52	Monkey	6-10 wk 5 d/wk 1 x/d				10 (degenerative changes in visual nerve fibers)	Eskin et al. 1985 Acrylamide	
53	Monkey	NS 1 x/d				30 (peripheral neuropathy)	Leswing and Ribelin 1969 Acrylamide	
54	Monkey (NS)	up to 363 d 5 d/wk (F)		3 F		10 F (clinical signs of peripheral neuropathy)	McCollister et al. 1964 Acrylamide	Only 1 animal per dose group; no clear signs of toxicity at 3 mg/kg/day
55	Monkey	33-47 d 5 d/wk 1 x/d				10 F (ataxia, adverse visual effects)	Merigan et al. 1985 Acrylamide	
56	Rat (albino)	21 d (F)				47 M (paralysis of hind limbs)	American Cyanamid Company 1953b	

DRAFT FOR PUBLIC COMMENT

3. HEALTH EFFECTS

ACRYLAMIDE

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	LOAEL		Reference Chemical Form	Comments	
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)			Serious (mg/kg/day)
57	Rat (albino)	NS 5 d/wk (G)				25 M (clinical signs of peripheral neuropathy)	American Cyanamid Company 1959	
58	Rat (Sprague-Dawley)	2 wk pre mating and Gd 0-19 (F)		3.82 F			American Cyanamid Company 1979 Acrylamide	
59	Rat (Fischer- 344)	28 d (W)		12 M 9 F		19 (clinical signs of peripheral neuropathy)	American Cyanamid Company 1991	Histopathological evaluation of nerve tissue not performed
60	Rat (Fischer- 344)	up to 93 d (W)		0.2 ^c M	1 M (reversible ultrastructural degeneration in sciatic nerve fibers)	20 (progressive signs of peripheral neuropathy)	Burek et al. 1980 Acrylamide	
61	Rat (Wistar)	Up to 21 d 1 x/d (G)				25 M (complete hindlimb paralysis by treatment day 21)	Dixit et al. 1981 Acrylamide	
62	Rat (Wistar)	21 d lactation period 1 x/d (G)				25 F (progressive clinical signs of neuropathy beginning as early as treatment day 4)	Friedman et al. 1999 Acrylamide	electron microscopic examinations were not performed on sciatic nerve

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/ Duration/ Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
63	Rat (albino)	12 doses in 15 d (GW)				50 (severe weakness)	Fullerton and Barnes 1966 Acrylamide	
64	Rat (albino)	variable (F)				6 (slight leg weakness after 40 weeks of exposure)	Fullerton and Barnes 1966 Acrylamide	
65	Rat (albino)	38 d (F)				25 (severe leg weakness)	Fullerton and Barnes 1966 Acrylamide	
66	Rat (albino)	Variable 5 d/wk 1 x/d (GW)				25 (severe leg weakness by 28 days of treatment)	Fullerton and Barnes 1966 Acrylamide	
67	Rat (albino)	55 doses 5 d/wk 1 x/d (GW)		10 F			Fullerton and Barnes 1966 Acrylamide	Histopathological evaluation of nerve tissue not performed
68	Rat (albino)	Variable 1 d/wk 1 x/d (GW)				100 (signs of severe neuropathy by the third dose)	Fullerton and Barnes 1966 Acrylamide	Younger rats appeared to be less severely affected.

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
69	Rat (Fischer- 344)	21 d (W)		3 M 10 F		10 M (clinical and histopathologic evidence of peripheral neuropathy) 30 F (clinical and histopathologic evidence of peripheral neuropathy)	Gorzinski et al. 1979	
70	Rat (Fischer- 344)	3 mo 6 mo (W)		0.5 M	2 M (electron microscopic degenerative effects in sciatic nerve fibers)		Johnson et al. 1984, 1986 Acrylamide	
71	Rat (Sprague-Dawley)	5 wk 7 d/wk 1x/d (GW)				10 (degenerative effects in nerve fibers)	Schulze and Boysen 1991 Acrylamide	
72	Rat (Sprague-Dawley)	GD6-PD21 (W)		3.72 F	7.89 F (abnormal gait)		Takahashi et al. 2009 Acrylamide	
73	Rat (Wistar)	90 d (W)		8.8 M		14.5 M (decreased rotarod performance)	Tanii and Hashimoto 1983 Acrylamide	Electron microscopic evaluations of nerve tissues not performed
74	Rat (Fischer- 344)	4 wk 5 d/wk (GW)				10 M (hindlimb dysfunction)	Tilson and Cabe 1979 Acrylamide	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
75	Rat (Fischer- 344) (W)	16 wk		5 F		5 M (slight axonal fragmentation in peripheral nerves of 6/6 males assessed)	Tyl et al. 2000a Acrylamide	Assessment included clinical signs, histopathological evaluations of peripheral nerves
76	Rat (Sprague-Dawley)	Gd 6- Ld 10 1 x/d (GW)		10 F		15 F (hindlimb splay in 100% of dams during the first few days of postpartum exposure)	Wise et al. 1995 Acrylamide	Histopathology of peripheral nerve tissue was not performed
77	Rat (Long- Evans) (W)	10 wk				7.9 M (increased incidences of hindlimb splay) 14.6 F (increased incidences of foot splay)	Zenick et al. 1986 Acrylamide	NOAEL for neurological effects not established due to lack of histopathologic assessment of peripheral nerve tissue
78	Mouse (CD-1)	Gd 6-20 1 x/d (GW)		15 F		45 F (up to 48% incidence of hindlimb splay)	Field et al. 1990 Acrylamide	
79	Mouse	8 wk 2 x/wk (NS)				36 M (clinical signs of peripheral neuropathy, testicular atrophy)	Hashimoto et al. 1981 Acrylamide	

DRAFT FOR PUBLIC COMMENT

ACRYLAMIDE

3. HEALTH EFFECTS

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
80	Mouse (ICR)	Up to 38 d (W)				90.8 M (clinically and histopathologically confirmed peripheral neuropathy)	Ko et al. 1999 Acrylamide	
81	Mouse (ICR)	NS (W)				91.8 M (clinical signs of peripheral neuropathy; ultrastructural degeneration in cutaneous nerve terminals)	Ko et al. 2000; Ko et al. 2002 Acrylamide	
82	Dog (Mongrel)	19 wk 6 d/wk 1 x/day (C)		1		8 (loss of coordination in the rear extremities)	American Cyanamid Company 1953c	
83	Dog (Mongrel)	29 d 7 d/wk 1 x/day (C)				10 F (incoordination and weakness of the hind legs)	American Cyanamid Company 1953c	
84	Dog	6-7 wk (C)				5.7 (clinical evidence of neuropathy)	Hersch et al. 1989a Acrylamide	
85	Dog	8 wk (F)				7 (clinical signs of peripheral neuropathy)	Satchell and McLeod 1981 Acrylamide	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments	
					Less Serious (mg/kg/day)	Serious (mg/kg/day)			
86	Cat	NS 1 x/d				20 (peripheral neuropathy)	Leswing and Ribelin 1969 Acrylamide		
87	Cat (NS)	up to 367 d 5 d/wk (F)		1		3 (twitching motion in hindquarters at 26 days, slightly unsteady gait at 47 days, definite weakness in the hindquarters at 68 days)	McCollister et al. 1964 Acrylamide	Histopathological evaluation of nerve tissue not performed	
88	Cat	Up to 16 wk (F)				15 (initial weakness of hindlimbs and subsequent paralysis of fore- and hind-limbs)	Post and McLeod 1977a Acrylamide		
89	Baboon	up to 137 d 1 x/d (F)				10 (peripheral neuropathy)	Hopkins 1970 Acrylamide		
Reproductive									
90	Rat (Fischer- 344) (W)	28 d		12 M		19 M (atrophy of the testes and/or seminal vesicles)	American Cyanamid Company 1991		

DRAFT FOR PUBLIC COMMENT

ACRYLAMIDE

3. HEALTH EFFECTS

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
91	Rat (Long- Evans) (W)	80 d		1.5 M		2.8 M (male-mediated increased postimplantation loss)	Smith et al. 1986 Acrylamide	
92	Rat (Fischer- 344) (W)	16 wk		2		5 M (dominant lethal mutation effects) 5 (decreases in implantations and number of live pups)	Tyl et al. 2000a Acrylamide	
93	Rat (Long- Evans) (W)	10 wk		5.1 F	7.9 F (decreased pup body weight)	8.8 M (male-mediated reproductive effects including decreased percentage impregnation of nonexposed females and increased postimplantation loss)	Zenick et al. 1986 Acrylamide	
94	Mouse (CD-1)	16-22 wk (W)		3.1 M		7.5 M (increased early resorptions, total postimplantation loss, decreased numbers of live fetuses, decreased numbers of live pups; apparently male mediated)	Chapin et al. 1995 Acrylamide	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	LOAEL		Reference Chemical Form	Comments
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		
95	Mouse ddY	4 wk (W)		9 M 18.7 F		13.3 M (decreased number of fetuses/dam)	Sakamoto and Hashimoto 1986 Acrylamide
Developmental							
96	Rat (Sprague-Dawley)	2 wk pre mating and Gd 0-19 (F)		3.82			American Cyanamid Company 1979 Acrylamide Assessment included mating and pregnancy indices, litter data, and offspring growth and survival
97	Rat (Sprague-Dawley)	Gd 6-20 1 x/d (GW)		15			Field et al. 1990 Acrylamide Assessment included external, visceral, and skeletal examinations of offspring
98	Rat (Fischer-344)	15 wk Gd 1-21 Ppd 1-84 (GW)		1.3	6 (effects on measures of cognitive motivation)		Garey and Paule 2007 Acrylamide
99	Rat (Wistar)	Throughout lactation via mothers or 5 d 1 x/d (G)				25 M (neurochemical changes in brain regions)	Husain et al. 1987 Acrylamide

DRAFT FOR PUBLIC COMMENT

ACRYLAMIDE

3. HEALTH EFFECTS

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
100	Rat (Sprague-Dawley)	GD6-PD21 (W)		7.89		14.56	(>40% decreased mean pup body weight)	Takahashi et al. 2009 Acrylamide
101	Rat (Sprague-Dawley)	Gd 6- Ld 10 1 x/d (GW)			5 F		(significantly decreased pup body weight during the preweaning period, as much as 9% lower than controls)	Wise et al. 1995 Acrylamide
					15		(increased overall horizontal activity and decreased auditory startle response)	
102	Mouse (CD-1)	Gd 6-20 1 x/d (GW)		15	45		(decreased mean fetal body weight; 15% lower than controls)	Field et al. 1990 Acrylamide Assessment included external, visceral, and skeletal examinations of offspring
CHRONIC EXPOSURE								
Death								
103	Rat (Fischer- 344) (W)	2 yr				2	(significantly decreased survival after 24 months of treatment)	Johnson et al. 1984, 1986 Acrylamide

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
Systemic								
104	Rat (Fischer- 344) (W)	2 yr	Resp	2 M 3 F			Friedman et al. 1995 Acrylamide	
			Cardio	2 M 3 F				
			Gastro	2 M 3 F				
			Hemato	2 M 3 F				
			Musc/skel	2 M 3 F				
			Hepatic	2 M 3 F				
			Renal	2 M 3 F				
			Endocr	2 M 3 F				

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
105	Rat (Fischer- 344) (W)	2 yr	Resp	2			Johnson et al. 1984, 1986 Acrylamide	
			Cardio	2				
			Gastro	2				
			Hemato	2				
			Musc/skel	2				
			Hepatic	2				
			Renal	2				
			Endocr	2				
Neurological								
106	Rat (Fischer- 344) (W)	2 yr		0.5 M	2 M (light microscopic evidence of peripheral nerve degeneration)		Friedman et al. 1995 Acrylamide	electron microscopy not conducted
107	Rat (Fischer- 344) (W)	2 yr		0.5		2 (light microscopic evidence of moderate to severe degeneration in sciatic nerve fibers)	Johnson et al. 1984, 1986 Acrylamide	
108	Cat (NS)	up to 367 d 5 d/wk (F)		1		3 (clinical signs of peripheral neuropathy)	McCollister et al. 1964 Acrylamide	

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
Reproductive								
109	Rat (Fischer- 344)	2 yr (W)		2			Friedman et al. 1995 Acrylamide	Gross and histopathological evaluations of reproductive organs and tissues
110	Rat (Fischer- 344)	2 yr (W)		2			Johnson et al. 1984, 1986 Acrylamide	Gross and histopathological evaluations of reproductive organs and tissues
Cancer								
111	Rat (Fischer- 344)	2 yr (W)				1 F (CEL: mammary gland fibroadenomas and adenomas or carcinomas combined)	Friedman et al. 1995 Acrylamide	Tunica vaginalis mesotheliomas and thyroid gland tumors observed at high dose (2 and 3 mg/kg/day in males and females, respectively)

DRAFT FOR PUBLIC COMMENT

Table 3-4 Levels of Significant Exposure to Acrylamide - Oral

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments	
					Less Serious (mg/kg/day)	Serious (mg/kg/day)			
112	Rat (Fischer- 344)	2 yr (W)					0.5 M (CEL: testicular sac mesotheliomas)	Johnson et al. 1984, 1986 Acrylamide	At 2 mg/kg/day, increased incidences of tumors at other sites in both males and females

a The number corresponds to entries in Figure 3-2.

b Study results used to derive an acute-duration oral minimal risk level (MRL) of 0.02 mg/kg/day for acrylamide, as described in detail in Appendix A. Benchmark dose (BMD) analysis was performed on male-mediated implantation loss data to identify a point of departure (BMDL10 of 1.78 mg/kg/day) which was divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability).


c Used to derive an intermediate-duration oral minimal risk level (MRL) of 0.002 mg/kg/day for acrylamide. The NOAEL of 0.2 mg/kg/day was divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability).

d Differences in levels of health effects and cancer effects between male and females are not indicated in Figure 3-2. Where such differences exist, only the levels of effect for the most sensitive gender are presented.

Bd Wt = body weight; (C) = capsule; Cardio = cardiovascular; CEL = cancer effect level; d = day(s); Endocr = endocrine; (F) = feed; F = Female; (G) = gavage; Gastro = gastrointestinal; Gd = gestation day; Gn Pig = guinea pig; (GW) = gavage in water; Hemato = hematological; Ld = lactation day; LD50 = lethal dose, 50% kill; LOAEL = lowest-observed-adverse-effect level; M = male; mo = month(s); Musc/skel = musculoskeletal; NOAEL = no-observed-adverse-effect level; NS = not specified; ppd = post-parturition day; Resp = respiratory; (W) = drinking water; wk = week(s); x = time(s); yr = year(s)

DRAFT FOR PUBLIC COMMENT

Figure 3-2 Levels of Significant Exposure to Acrylamide - Oral
Acute (≤14 days)




DRAFT FOR PUBLIC COMMENT

Figure 3-2 Levels of Significant Exposure to Acrylamide - Oral (Continued)

Chronic (≥ 365 days)

DRAFT FOR PUBLIC COMMENT



3. HEALTH EFFECTS

Respiratory Effects. No data were located regarding respiratory effects in humans following oral exposure to acrylamide.

No clinical signs or gross or histopathological evidence of respiratory effects were observed in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or for up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986).

Cardiovascular Effects. No data were located regarding cardiovascular effects in humans following oral exposure to acrylamide.

No clinical signs or gross or histopathological evidence of cardiovascular effects were observed in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986).

Gastrointestinal Effects. No data were located regarding gastrointestinal effects in humans following oral exposure to acrylamide.

No clinical signs or gross or histopathological evidence of gastrointestinal effects were observed in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986).

Hematological Effects. No data were located regarding hematological effects in humans following oral exposure to acrylamide.

Burek et al. (1980) reported significant decreases in packed cell volume (PCV), erythrocyte counts, and hemoglobin in male and female F344 rats at day 76 during administration of acrylamide in their drinking water at a dose level of 20 mg/kg/day. These effects were also noted in females (but not males) dosed at 5 mg/kg/day for 93 days. Hematological evaluations at 3, 6, 12, 18, and 24 months on male and female F344 rats receiving acrylamide doses as high as 2 mg/kg/day from the drinking water for up to 2 years revealed no evidence of treatment-related effects (Johnson et al. 1984, 1986).

3. HEALTH EFFECTS

Musculoskeletal Effects. No data were located regarding musculoskeletal effects in humans following oral exposure to acrylamide.

Atrophy of skeletal muscles in the posterior area was noted in F344 rats receiving acrylamide at a dose level of 20 mg/kg/day from the drinking water for up to 93 days; however, this effect was considered the likely result of acrylamide-induced peripheral neuropathy (Burek et al. 1980).

Renal Effects. No data were located regarding renal effects in humans following oral exposure to acrylamide.

Urinalysis and gross and histopathological evaluations revealed no evidence of renal effects in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986). Although distended bladders were occasionally observed in laboratory animals following repeated-dose oral exposure, this effect may be the result of acrylamide-induced effects on nerve fibers that innervate the bladder (Fullerton and Barnes 1966).

Endocrine Effects. No data were located regarding endocrine effects in humans following oral exposure to acrylamide.

Significantly decreased mean serum testosterone levels (27 and 9% of the control mean) were noted in adult male F344 rats administered acrylamide in the drinking water for 28 days at concentrations resulting in estimated acrylamide doses of 19 and 25 mg/kg/day, respectively; there were no statistically significant effects on serum testosterone levels at 14-day interim assessment (American Cyanamid Company 1991). In the same study, serum levels of the thyroid hormones thyroxin (T₄) and triiodothyronine (T₃) and serum prolactin were assessed in both male and female F344 rats receiving acrylamide from the drinking water at estimated doses up to 24–25 mg/kg/day. High-dose males exhibited significantly increased mean T₄ concentration compared to controls at 14-day interim assessment and significantly decreased mean T₃ and T₄ concentrations at study termination. Terminal (28-day) mean T₃ level was significantly decreased in 19 mg/kg/day males as well. Mean T₃ and T₄ levels in treated female rats were not significantly different from controls at either time point. Serum prolactin levels were significantly decreased in 19 and 25 mg/kg/day males at the 14-day interim assessment, but not at terminal assessment. No significant

3. HEALTH EFFECTS

effects on serum prolactin levels were seen in treated female rats. At terminal sacrifice, no significant treatment-related effects on thyroid weights were seen in males or females at any dose level.

Gross and histopathological assessments revealed no evidence of endocrine effects in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986).

Dermal Effects. No data were located regarding dermal effects in humans or animals following oral exposure to acrylamide.

Ocular Effects. No data were located regarding ocular effects in humans following oral exposure to acrylamide.

Ophthalmological evaluations revealed no evidence of ocular effects in male or female F344 rats administered acrylamide in the drinking water for up to 93 days at concentrations resulting in doses up to 20 mg/kg/day (Burek et al. 1980) or up to 2 years at concentrations resulting in doses up to 2 or 3 mg/kg/day (Friedman et al. 1995; Johnson et al. 1984, 1986).

Acrylamide-induced neurological effects on the visual system of orally-exposed primates are discussed in Section 3.2.2.4 (Neurological Effects).

Body Weight Effects. No data were located regarding body weight effects in humans following oral exposure to acrylamide.

Depressed body weight, including actual body weight loss, was consistently reported in laboratory animals following single or repeated oral exposure to acrylamide.

Slight initial weight loss (magnitude not specified) was reported following a single oral dose of 126 mg acrylamide/kg in rats and guinea pigs and 63 mg/kg in rabbits (Dow Chemical Company 1957). Significantly depressed body weight was observed for several days in mice dosed once at 150 mg/kg (lethal dose), but not at 100 mg/kg (sublethal dose) (Sakamoto et al. 1988).

3. HEALTH EFFECTS

Depressed body weight or actual body weight loss was observed in repeat-dosing studies as well. Weight loss was noted in 3- and 6-week-old mice receiving a high daily dose (90.8 mg/kg) of acrylamide for 20 and 38 days, respectively (Ko et al. 1999). Most repeated-dose oral studies, however, identified much lower body weight effect levels. At treatment day 13 of a 3-month study, significantly decreased mean body weights (8% lower than controls) were observed in male and female rats receiving acrylamide at 20 mg/kg/day from the drinking water (Burek et al. 1980). Significantly depressed body weight gain (approximately 40% less than controls) was noted in male Long-Evans rats receiving 15 mg/kg/day of acrylamide for 5 consecutive days; there were no apparent acrylamide-induced effects at 5 mg/kg/day (Tyl et al. 2000b). Severely depressed body weight gain (approximately 50% that of controls) was observed during 21 days of exposure of male albino rats to acrylamide in the food at a concentration resulting in an estimated daily acrylamide dose of 47 mg/kg/day (American Cyanamid Company 1953b). This dose level resulted in the death of 3/10 of the rats by the end of the treatment period; three surviving rats exhibited somewhat greater weight gain than controls during a 10-week recovery period. Severely depressed body weight and emaciation were reported in studies of rats administered acrylamide for periods of 3–5 weeks at doses in the range of 19–30 mg/kg/day (American Cyanamid Company 1953b, 1991; Schulze and Boysen 1991). Tanii and Hashimoto (1983), however, reported no significant effects on body weight in male Wistar rats administered acrylamide in the drinking water for 90 days at concentrations resulting in estimated doses as high as 23.7 mg/kg/day. Dogs and cats are also susceptible to the body weight effects of orally-administered acrylamide. An 8-week dosing period to dogs at 7 mg/kg/day resulted in weight loss (Satchell and McLeod 1981). Cats dosed for 12–16 weeks at 15 mg/kg/day exhibited weight loss of unspecified magnitude (Post and McLeod 1977a).

Several groups of investigators assessed effects of maternal body weight on rat or mouse dams receiving oral acrylamide during gestation and/or lactation. A daily dose as low as 7.5 mg/kg/day during gestation days 6–20 resulted in approximately 12% depressed maternal weight gain in rats, but a NOAEL of 45 mg/kg/day was identified in similarly-treated mouse dams (Field et al. 1990). Net maternal weight loss was observed in rat dams administered acrylamide at 25 mg/kg/day during 21 days of lactation (Friedman et al. 1999). Wise et al. (1995) treated rat dams with 5 or 15 mg/kg/day from gestation day 6 through lactation day 10; no body weight effects were seen at the 5 mg/kg/day dose level, but approximately 33% depressed maternal weight gain was noted during the 10-day lactation period in the 15 mg/kg/day group.

3. HEALTH EFFECTS

3.2.2.3 Immunological and Lymphoreticular Effects

No data were located regarding immunological or lymphoreticular effects in humans or animals following oral exposure to acrylamide.

3.2.2.4 Neurological Effects

Neurological deficits are a hallmark of acrylamide toxicity. Most evidence in humans derives from occupational exposures that predominantly involved inhalation and dermal routes (see Section 3.2.1.4). Available information regarding the neurological effects of oral exposure in humans is limited to a case report of persistent peripheral neuropathy in a subject who intentionally ingested 18 g of acrylamide crystals (Donovan and Pearson 1987) and signs of central and peripheral neurological deficits in family members exposed (likely via oral and dermal routes) to acrylamide in well water at a concentration of 400 ppm (Igisu and Matsuoka 2002; Igisu et al. 1975). Epidemiologic studies designed to evaluate noncancer health effects in groups of orally-exposed subjects have not been conducted.

Neurological effects associated with oral exposure to acrylamide have been well characterized in laboratory animals and include clinical signs such as twitching, loss of balance, tremors, lethargy, and general weakness and more subtle indicators of functional deficits such as decreased rotarod performance and increased limb or foot splay. Evidence of degenerative lesions in peripheral nerve fibers, as observed by light and electron microscopy, have been detected at oral doses lower than those eliciting clinical signs and other overt indications of functional deficit.

Clinical signs have been elicited following single oral exposure of rats, rabbits, and dogs to doses in the range of 100–200 mg/kg (American Cyanamid Company 1953c; Fullerton and Barnes 1966; McCollister et al. 1964; Tilson and Cabe 1979). Five daily doses to rats at 45 mg/kg (Tyl et al. 2000b) or 75 mg/kg (Sublet et al. 1989) elicited typical clinical signs of peripheral neuropathy. Ko et al. (1999) noted the onset of altered gait as early as treatment day 5 in 3- and 8-week-old mice receiving acrylamide from the drinking water at approximately 90 mg/kg/day. Friedman et al. (1999) and Dixit et al. (1981) observed clinical signs of neurological effects as early as treatment days 4 and 14, respectively, in rats receiving acrylamide at 25 mg/kg/day. Gilbert and Maurissen (1982) reported decreased rotarod performance and increased hindlimb splay as early as days 6–8 of a 12-day exposure of mice receiving acrylamide from the drinking water at an estimated dose of 25.8 mg/kg/day.

3. HEALTH EFFECTS

Clinical signs and other indicators of acrylamide-induced functional neurological deficit, such as increased hindlimb splay and decreased rotarod performance, during intermediate-duration (15–364 days) exposure have been observed in several animal species. In rats and cats, dose levels in the range of 8–25 mg/kg/day elicited overt signs of neurological deficit as early as 3–8 weeks following the initiation of treatment (American Cyanamid Company 1991; Burek et al. 1980; Fullerton and Barnes 1966; Gorzinski et al. 1979; Leswing and Ribelin 1969; McCollister et al. 1964; Post and McLeod 1977a; Takahashi et al. 2009; Tanii and Hashimoto 1983; Tilson and Cabe 1979; Wise et al. 1995; Zenick et al. 1986). Slight leg weakness was reported after 40 weeks of oral dosing at 6 mg/kg/day in one rat study (Fullerton and Barnes 1966). In dogs, dose levels of 5.7–10 mg/kg/day elicited clinical signs within 3–4 weeks (American Cyanamid Company 1953c; Hersch et al. 1989a; Satchell and McLeod 1981). Hashimoto et al. (1981) noted typical signs of peripheral neuropathy and diminished rotarod performance in male mice treated at 36 mg/kg/day, 2 days/week for 8 weeks. In primates (monkeys, baboons), clinical signs were elicited by doses of 10–30 mg/kg/day for periods of 42–61 days (Eskin et al. 1985; Hopkins 1970; Leswing and Ribelin 1969; Maurissen et al. 1983; Merigan et al. 1985). Higher levels of oral dosing typically result in earlier onset of clinical signs and more severe effects with continued dosing.

Histopathological assessment of acrylamide-induced neurological effects includes studies in which light and electron microscope evaluations were performed on peripheral nerve fibers of rats administered acrylamide in the drinking water (Burek et al. 1980; Johnson et al. 1984, 1985, 1986). In the study of Burek et al. (1980), the rats received estimated doses of 0.05–20 mg/kg/day for up to 93 days. This study identified a NOAEL of 0.2 mg/kg/day and a LOAEL of 1 mg/kg/day for increased incidences of ultrastructural degeneration (axolemma invaginations with cell organelles and/or dense bodies) in the sciatic nerve; higher doses elicited more pronounced degenerative effects that could also be detected by light microscopy. Clinical signs of peripheral neuropathy were elicited at the 20 mg/kg/day dose level. The study of Johnson et al. (1984, 1985, 1986) employed estimated dose levels of 0.01, 0.1, 0.5, and 2.0 mg/kg/day for up to 2 years. Interim evaluations of peripheral nerve fibers were performed at 3, 6, 12, and 18 months; however electron microscope evaluations at interims >12 months were inconclusive due to high incidences of degenerative results attributed in part to aging. The studies of Johnson et al. (1984, 1985, 1986) identified a NOAEL of 0.5 mg/kg/day and a LOAEL of 2 mg/kg/day for increased incidences of ultrastructural degeneration (axolemma invaginations) in the sciatic nerve at 3, 6, and 12 months and for light microscopy evidence of degenerative effects in peripheral nerve fibers at 12–24 months; there were no clinical signs of neurological effects at any dose level. Friedman et al. (1995) duplicated the study design of Johnson et al. (1986), but excluded electron microscopy in histopathological evaluations of peripheral nerve fibers. The study of Friedman et al. (1995) also identified a NOAEL of 0.5 mg/kg/day

3. HEALTH EFFECTS

and a LOAEL of 2 mg/kg/day for peripheral nerve degeneration as revealed by light microscopy in the absence of clinical signs of neurological effects. Histopathological evidence (light microscope) of degenerative effects was also noted in fibers from cervical and lumbar spinal cord, gasserian and dorsal root ganglia, and sciatic, tibial, and sural nerves of male and female Sprague-Dawley rats that had received acrylamide by gavage for 5 weeks at doses of 10 and 30 mg/kg/day (Schulze and Boysen 1991). The most severe changes were observed in the large fibers of the tibial and sural nerves of high-dose rats. In a 2-generation study of F-344 rats, Tyl et al. (2000a) reported degenerative effects in peripheral nerve fibers of F₀ (but not female or F₁ male or female) rats that had received acrylamide from the drinking water at 5 mg/kg/day for 16 weeks.

Information regarding the neurodevelopmental toxicity of acrylamide is discussed in Section 3.2.2.6.

The highest NOAEL values and all LOAEL values from each reliable study for neurological effects in each species and duration category are recorded in Table 3-4 and plotted in Figure 3-2.

3.2.2.5 Reproductive Effects

No studies were located regarding acrylamide-induced reproductive effects in humans.

Animal studies designed to assess the reproductive toxicity of orally-administered acrylamide revealed pre- and postimplantation losses and decreased numbers of live fetuses in rats and mice at repeated doses in the range of 3–60 mg/kg/day (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Smith et al. 1986; Sublet et al. 1989; Tyl et al. 2000a, 2000b; Working et al. 1987; Zenick et al. 1986).

Results of dominant lethality testing (Chapin et al. 1995; Smith et al. 1986; Tyl et al. 2000a, 2000b) and crossover trials (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Zenick et al. 1986) indicate that acrylamide induces male-mediated reproductive effects at repeated oral doses in the range of 2.8–19 mg/kg/day. Sublet et al. (1989) reported statistically significant moderately decreased sperm mobility in Long-Evans rats administered acrylamide at a gavage dose of 45 mg/kg/day for 5 days, but suggested that this effect was not solely responsible for poorer reproductive performance. In apparent contrast, Tyl et al. (2000b) found no significant effects on sperm parameters in Long-Evans hooded rats following repeated oral dosing at levels as high as 60 mg/kg/day and suggested that indicators of acrylamide-induced reproductive toxicity such as pre- and post-implantation loss and decreased numbers of live fetuses may be at least partly the result of impaired mating performance due to acrylamide neurotoxicity.

3. HEALTH EFFECTS

Histologic indicators of degenerative effects were reported in spermatids of ddY mice administered acrylamide by daily gavage for 5 days at dose levels of 100 or 150 mg/kg/day (Sakamoto et al. 1988). Other investigators reported evidence of acrylamide-induced testicular atrophy in F344 rats receiving acrylamide in the drinking water for 28 or 90 days at concentrations resulting in estimated acrylamide doses of 19 or 5 mg/kg/day, respectively (American Cyanamid Company 1991; Burek et al. 1980). Gross and histopathologic examinations of reproductive organs and tissues from male rats receiving acrylamide from the drinking water for up to 2 years at estimated doses as high as 2 mg/kg/day revealed no signs of acrylamide-induced effects (Friedman et al. 1995; Johnson et al. 1984, 1986).

Prebreeding exposure of female mice to acrylamide at 18.7 mg/kg/day (Sakamoto and Hashimoto 1986) or female Long-Evans rats to doses up to 14.6 mg/kg/day (Zenick et al. 1986) did not adversely affect reproductive performance variables such as fertility or implantation when the animals were bred with nonexposed males. Gross and histopathologic examinations of reproductive organs and tissues from female rats receiving acrylamide from the drinking water for up to 2 years at estimated doses as high as 2–3 mg/kg/day revealed no signs of acrylamide-induced effects (Friedman et al. 1995; Johnson et al. 1984, 1986).

Atrophy of the testes and/or seminal vesicles was reported in F-344 rats receiving acrylamide at 19 or 25 mg/kg/day from the drinking water for 28 days; this effect was not seen at 12 mg/kg/day (American Cyanamid Company 1991).

The highest NOAEL values and all LOAEL values from each reliable study for reproductive effects in each species and duration category are recorded in Table 3-4 and plotted in Figure 3-2.

3.2.2.6 Developmental Effects

No studies were located regarding acrylamide-induced developmental effects in humans.

Several reports are available in which developmental toxicity was assessed in the offspring of rat or mouse dams administered acrylamide via daily gavage during gestation and/or lactation (American Cyanamid Company 1979; Field et al. 1990; Friedman et al. 1999; Garey and Paule 2007; Husain et al. 1987; Takahashi et al. 2009; Walden et al. 1981; Wise et al. 1995).

3. HEALTH EFFECTS

Body weight decreases and decreased auditory startle response were noted in offspring of female Sprague-Dawley rats exposed to 5 and 15 mg/kg-day, respectively, on gestation days 6–10 (Wise et al. 1995). No exposure-related fetal malformations or variations (gross, visceral, or skeletal) were found in offspring of Sprague-Dawley rats administered acrylamide at doses of 2.5, 7.5, or 15 mg/kg/day on gestation days 6–20 or in CD-1 mice at doses of 3, 15, or 45 mg/kg/day on gestation days 6–17 (Field et al. 1990). The highest dose in each species was maternally toxic, as evidenced by depressed maternal weight gain in the rats and mice and increased hindlimb splay in the mice. There were no indications of treatment-related developmental effects in the offspring of Sprague-Dawley rat dams administered acrylamide at doses up to nearly 4 mg/kg/day for 2 weeks pre mating and throughout gestation (American Cyanamid Company 1979). Decreased mean fetal body weight (approximately 15% lower than controls) was noted in the offspring of CD-1 mouse dams administered acrylamide by gavage at 45 mg/kg/day during gestation days 6–20; a NOAEL for this effect was 15 mg/kg/day (Field et al. 1990). Significantly depressed mean body weights were reported in offspring of Sprague-Dawley rat dams administered acrylamide in the drinking water at 100 ppm (mean acrylamide intake 14.56 mg/kg/day) from gestation day 6 through postnatal day 21; the male and female pups exhibited approximately 42 and 46% lower mean body weights, respectively, than unexposed control pups (Takahashi et al. 2009). At lower mean maternal exposure levels doses (25 and 50 ppm; mean doses of 3.72 and 7.89 mg/kg/day, respectively), pup body weights did not differ significantly from those of controls. There were no signs of acrylamide-induced neurotoxicity or testicular toxicity in pups of any exposure group. The 100 ppm dams exhibited increasing severity of gait abnormalities from postnatal day 2 onward; abnormal gait was observed in 50 ppm dams from postnatal day 18 onward.

Periodic significantly decreased brain levels of selected catecholamines (noradrenaline, dopamine, 5-hydroxytryptamine) were noted in pups of rat dams administered acrylamide at 25 mg/kg/day during lactation only (Husain et al. 1987). Similar effects on brain catecholamines were observed in rat pups 12–21 days of age at the beginning of a 5-day period in which they were administered acrylamide by gavage at 25 mg/kg/day; these effects were not seen in rat pups that were 60 days of age at the initiation of dosing (Husain et al. 1987). Decreased performance in an operant test of cognitive motivation was reported in adolescent F344 rats exposed during gestation and lactation and extending through 12 weeks of age at an average dose of 6 mg/kg/day, but not at 1.3 mg/kg/day (Garey and Paule 2007). Significant decreases in whole brain levels of noradrenaline, dopamine, and 5-hydroxytryptamine were observed in pups of rat dams administered acrylamide at 25 mg/kg/day during lactation. Walden et al. (1981) found significant alterations in intestine enzyme levels in young Sprague-Dawley rat pups whose mothers had received acrylamide via gavage on gestation days 6–17 at 20 mg/kg/day; the toxicological significance of these

3. HEALTH EFFECTS

findings is uncertain. Friedman et al. (1999) reported increased mortality and reduced body weights in pups of rat dams dosed at 25 mg/kg/day during lactation; however, serious maternal toxicity was noted as well.

The highest NOAEL values and all LOAEL values from each reliable study for developmental effects in each species and duration category are recorded in Table 3-4 and plotted in Figure 3-2.

3.2.2.7 Cancer

Available epidemiology studies on increased risk of cancer from acrylamide in food include case-control studies (Michels et al. 2006; Mucci et al. 2003, 2004, 2005; Pelucchi et al. 2006, 2007; Wilson et al. 2009a) and reports from ongoing prospective studies (Hogervorst et al. 2007, 2008a, 2008b; Larsson et al. 2009a, 2009b, 2009c, 2009d; Mucci et al. 2006; Wilson et al. 2009b). These studies provide little evidence for acrylamide-induced cancer in humans exposed to acrylamide in the diet. See Section 3.2.1.7 (Cancer) for information regarding cancer among acrylamide workers primarily exposed via inhalation and dermal routes.

Case-control Studies. No statistically significant associations were found between self-reported consumption of foods with high (300–1,200 µg/kg) or moderate (30–299 µg/kg) acrylamide concentrations and increased risk of large bowel, kidney, or bladder cancer in a population-based case-control study (692 controls and 875, 391, and 186 large bowel, kidney, and bladder cancer cases, respectively) (Augustsson et al. 1999; Mucci et al. 2003). Information on intake of various foods and nutrients was assessed by questionnaire. Cancer cases were born in Sweden between 1918 and 1942 and resided in Stockholm for at least 1 month between November 1992 and December 1994. Controls were selected from a Swedish national population registry and matched to cases by age and gender.

Mucci et al. (2005) assessed acrylamide intake of >43,000 women, including 667 breast cancer cases, who were enrolled in the Swedish Women's Lifestyle and Health Cohort. The estimated average daily acrylamide intake among the participants was 25.9 µg/day and was based on results of food frequency questionnaires (FFQs) and the Swedish National Food Administration database of information on acrylamide content of selected food items. After ranking the women into quintiles of estimated acrylamide intake (means of 12, 20, 25, 31, and 44 µg/day), there was no significant increased risk of breast cancer in the higher quintiles compared to the lowest quintile.

3. HEALTH EFFECTS

Within an integrated network of Italian and Swiss hospital-based case-control studies to investigate the relation between dietary acrylamide intake and cancers at several sites, no significant associations were found between estimated dietary acrylamide intake and cancers of the oral cavity and pharynx, esophagus, large bowel, larynx, breast, ovary, or prostate (Pelucchi et al. 2006). Dietary acrylamide intake was estimated based on results of FFQs and average content of acrylamide in foods from resources of the World Health Organization and Swiss Federal Office of Public Health. In another case-control study that included four areas of Italy, no significant association was found between total dietary acrylamide intake and renal cell cancer (Pelucchi et al. 2007). In this study, statistically significantly elevated odds ratios (ORs: 1.49, 95% CI 1.18–1.87; 1.70, 95% CI 1.25–2.30) were noted for weekly white bread portions of 7–<21 and \geq 21, respectively; however, the study authors indicated that the relationship between white bread consumption and renal cell cancer might be explained by a high glycemic content and consequent effect on levels of insulin-like growth factors.

Wilson et al. (2009a) conducted a case-control study to assess possible associations between acrylamide and prostate cancer risk using two measures of acrylamide exposure: intake from FFQs and acrylamide-hemoglobin adduct levels in blood samples. Dietary data were available for 1,499 prostate cancer cases and 1,118 controls from a Cancer of the Prostate in Sweden (CAPS) population-based case-control study. Acrylamide-hemoglobin adduct levels were measured in blood samples from a subset of 170 prostate cancer cases and 161 controls. Controls were randomly selected from the Swedish Population Registry and were frequency matched to cases by 5-year age groups and region of residence. No significant association was found between acrylamide exposure (as measured by FFQ or acrylamide-hemoglobin adduct levels) and risk of prostate cancer.

Michels et al. (2006) conducted a case-control study to evaluate whether diet during preschool age affected a woman's risk of breast cancer later in life. Cases and controls were selected from participants in two prospective cohort studies, the Nurses' Health Study and the Nurses' Health Study II. Information concerning childhood diet of the nurses at ages 3–5 years was obtained from FFQs filled out by the mothers of the participants. The median year of birth of the mothers was 1914 for case mothers and 1913 for control mothers. The results indicated an increased risk of breast cancer among woman who had frequently consumed French fries at preschool age. For one additional serving of French fries per week, the OR for breast cancer adjusted for adult life breast cancer risk factors was 1.27 (95% CI=1.12–1.44). Consumption of whole milk was associated with a slightly decreased risk of breast cancer (covariate-adjusted OR for every additional glass of milk per day=0.90; 95% CI=0.82–0.99). Intake of none of the nutrients calculated was related to the breast cancer risk in this study. The authors noted that they did not

3. HEALTH EFFECTS

observe a similar association of breast cancer with frequent consumption of hot dogs or ground beef, suggesting that French fry consumption was not a marker of “fast food” habits. The study results suggest a possible association between diet before puberty and the subsequent risk of breast cancer, but the conclusions and the study are of limited use. No information is available on cooking methods or acrylamide content in the foods being evaluated, and the ability of mothers to accurately recall preschool diets of their daughters is questionable.

No significant associations were found between acrylamide-hemoglobin or glycidamide-hemoglobin adduct levels and total breast cancer in a Danish nested case-control study that examined breast cancer and acrylamide exposure using acrylamide-hemoglobin and glycidamide-hemoglobin adduct levels in red blood cells as presumed biomarkers for oral exposure to acrylamide (Olesen et al. 2008). After adjusting for confounding factors including smoking behavior, the study authors noted that a 10-fold increase in acrylamide-hemoglobin adduct levels was associated with a 1.9 (95% CI 0.9–4.0) times higher risk of breast cancer and a 5-fold increase (which corresponds to the range in acrylamide-hemoglobin adduct levels among nonsmokers) was associated with a 1.6 (95% CI 0.9–2.6) times higher risk. A significant positive association was observed between acrylamide-hemoglobin adduct level and ER+ breast cancer; a 10-fold increase in adduct level was associated with a 4.9 (95% CI 1.2–20) times increased risk in smokers and 2.7 (95% CI 1.1–6.6) times increased risk after adjustment for smoking. However, this study is limited by the relatively small number of subjects (374 cases and 374 controls) and uncertainty regarding extrapolation of acrylamide exposure as assessed by a few months of acrylamide-hemoglobin adduct measurements to a lifetime of exposure.

Mucci et al. (2004) analyzed data from a large population-based Swedish case-control study of renal cell cancer. FFQs were used to collect information on intake of 11 food items with elevated acrylamide levels as ascertained through extensive food databases in Sweden and the United States and quartiles of daily food and acrylamide intake were created. This study found no evidence that food items with elevated acrylamide were associated with a higher risk of renal cell cancer risk.

Prospective Studies. Recent and ongoing prospective studies designed to evaluate possible associations between acrylamide in food and risk of cancers at various sites include cohorts from Sweden (Larsen et al. 2009a, 2009b, 2009c, 2009d; Mucci et al. 2006), the United States (Wilson et al. 2009b), and the Netherlands (Hogervorst et al. 2007, 2008a, 2008b). Most studies found no statistically significant associations between acrylamide in food and risks of cancers of the esophagus, stomach, or pancreas (Hogervorst et al. 2008b); colon or rectum (Hogervorst et al. 2008b; Larsen et al. 2009c; Mucci et al.

3. HEALTH EFFECTS

2006); bladder or prostate (Hogervorst et al. 2008a); breast (Hogervorst et al. 2007; Larsson et al. 2009d; Wilson et al. 2009b); endometrium (Hogervorst et al. 2007; Larsson et al. 2009a), or ovarian epithelium (Larsson et al. 2009b). Exceptions include increased risks of postmenopausal endometrial and ovarian cancer (Hogervorst et al. 2007) and renal cell cancer (Hogervorst et al. 2008a) with increasing dietary acrylamide in prospective studies of a Dutch population, but in these studies, estimations of dietary acrylamide levels in foods on the market at baseline in 1986 were based on food samples analyzed since 2001 and questionnaires did not include details regarding specifics of food preparation. Some of the tumor sites observed in animal studies (thyroid, testis, central nervous system) have not been evaluated in humans, and there are limitations in some of the study methods and cohort sizes in the prospective studies.

Information regarding the carcinogenicity of acrylamide in orally-exposed animals is available from two similarly-designed 2-year bioassays (Friedman et al. 1995; Johnson et al. 1984, 1985, 1986) in which groups of male and female F344 rats were administered acrylamide in the drinking water at concentrations calculated to provide doses up to 2 mg/kg/day (3 mg/kg/day for high-dose female rats of the Friedman et al. 1995 study). Selected tumor types with significantly increased incidences in acrylamide-treated rats from both studies are summarized in Table 3-5. Significantly increased incidences of mesotheliomas of the tunica vaginalis testis were observed by Johnson et al. (1984, 1986) in male rats at the two highest dose levels (0.5 and 2 mg/kg/day). Additional findings included significantly increased incidences of thyroid (follicular cell) adenomas (no carcinomas), mesotheliomas of the tunica vaginalis testis, and benign adrenal pheochromocytoma in high-dose males and mammary gland benign tumors (adenoma, fibroadenomas, or fibroma), central nervous system tumors of glial origin, thyroid (follicular cell) adenomas or adenocarcinomas, squamous papillomas of the oral cavity, uterine adenocarcinomas, benign clitoral gland adenomas, and pituitary gland adenomas in high-dose females. The study of Friedman et al. (1995) noted increased incidences of tunica vaginalis mesothelioma and thyroid gland (follicular cell) adenoma (and adenoma or carcinoma) in high-dose males and thyroid gland follicular cell neoplasms (adenomas and carcinomas combined) in high-dose females and mammary gland tumors (fibroadenomas or combined fibroadenomas and carcinomas) in low- and high-dose (1 and 3 mg/kg/day) females. The findings of statistically significant increased incidences of adrenal pheochromocytomas in male rats, oral cavity tumors in female rats, central nervous system tumors of glial origin, and clitoral or uterine tumors in female rats in the earlier bioassay (Johnson et al. 1986) were not replicated in the second bioassay (Friedman et al. 1995); however, Rice (2005) reported that the study of Friedman et al. (1995) did not include examination of all the brains or spinal cords in the treatment groups

3. HEALTH EFFECTS

Table 3-5. Incidence of Tumors with Statistically Significant Increases in 2-Year Bioassays with F344 Rats Exposed to Acrylamide in Drinking Water

Reference/tumor type	Dose (mg/kg/day)							
	0	0	0.01	0.1	0.5	1.0	2.0	3.0
Johnson et al. 1986; males								
Follicular cell adenoma	1/60	–	0/58	2/59	1/59	–	7/59 ^a	–
Tunica vaginalis mesothelioma	3/60	–	0/60	7/60	11/60 ^a	–	10/60 ^a	–
Adrenal pheochromocytoma	3/60	–	7/59	7/60	5/60	–	10/60 ^a	–
Johnson et al. 1986; females								
Follicular cell adenoma/ carcinoma	1/58	–	0/59	1/59	1/58	–	5/60 ^b	–
Mammary adenocarcinoma	2/60	–	1/60	1/60	2/58	–	6/61	–
Mammary benign	10/60	–	11/60	9/60	19/58	–	23/61 ^a	–
Mammary benign + malignant ^c	12/60	–	12/60	10/60	21/58	–	29/61 ^a	–
Central nervous system tumors of glial origin	1/60	–	2/59	1/60	1/60	–	9/61 ^a	–
Oral cavity malignant + benign	0/60	–	3/60	2/60	3/60	–	8/60 ^a	–
Uterus adenocarcinoma	1/60	–	2/60	1/60	0/59	–	5/60 ^b	–
Clitoral adenoma, benign	0/2	–	1/3	3/4	2/4	–	5/5 ^b	–
Pituitary gland adenoma	25/59	–	30/60	32/60	27/60	–	32/60 ^b	–
Friedman et al. 1995; males ^d								
Follicular cell adenoma/carcinoma		2/102 ^f	–	12/203	5/101	–	17/75 ^a	–
Tunica vaginalis mesothelioma ^e	4/102	4/102	–	9/204	8/102	–	13/75 ^a	–
Friedman et al. 1995; females ^d								
Follicular cell adenoma/ carcinoma	1/50 7/46	1/50 4/50	– –	– –	– –	10/100 21/94 ^a	– –	23/100 ^a 30/95 ^a
Mammary benign + malignant								

^aStatistically significantly ($p < 0.05$) different from control, Fisher's Exact test.

^bStatistically significantly ($p < 0.05$) different from control, after Mantel-Haenszel mortality adjustment.

^cIncidences of benign and adenocarcinoma were added herein, based on an assumption that rats assessed with adenocarcinoma were not also assessed with benign mammary gland tumors.

^dTwo control groups were included in the study design to assess variability in background tumor responses.

^eIncidences reported herein are those originally reported by Friedman et al. (1995) and not those reported in the reevaluation study by Iatropoulos et al. (1998).

^fThe data reported in Table 4 in Friedman et al. (1995) noted one follicular cell adenoma in the second control group; however, the raw data obtained in the Tegeris Laboratories (1989) report (and used in the time-to-tumor analysis) listed no follicular cell adenomas in this group. The corrected number for adenomas (0) and the total number (2) of combined adenomas and carcinomas in the second control group are used in the tables of this assessment.

Sources: Friedman et al. 1995; Johnson et al. 1986

3. HEALTH EFFECTS

and that seven reported cases of a morphologically distinctive category of primary brain tumor described as “malignant reticulosis” were excluded from the analysis.

The cancer effect levels (CELs) from the rat cancer bioassays of Johnson et al. (1984, 1986) and Friedman et al. (1995) are recorded in Table 3-4 and plotted in Figure 3-2.

The potential for acrylamide to initiate skin tumors was examined in two studies of female mice administered acrylamide at oral dose levels of 0, 12.5, 25, or 50 mg/kg/day, 6 times during a 2-week period, followed 2 weeks later by dermal applications of 12-O-tetradecanoylphorbol-13-acetate (TPA, a tumor promoter) to the shaved back 3 times/week for 20 weeks (Bull et al. 1984a, 1984b). Incidences of skin papillomas and skin carcinomas were significantly elevated in the mice receiving acrylamide at oral dose levels of 25 and/or 50 mg/kg/day, indicating that acrylamide initiated skin tumors under the conditions of the studies.

EPA, IARC, and the Department of Health and Human Services have concluded that acrylamide is likely to be carcinogenic to humans. This conclusion is based on lack of adequate human data and sufficient evidence of carcinogenicity in the animal studies summarized above. The Department of Health and Human Services (NTP 2005) assigned the cancer descriptor “*reasonably anticipated to be a human carcinogen*”. IARC (1994) assigned acrylamide to Group 2A (probably carcinogenic to humans). EPA (IRIS 2009) assigned the classification of B2; probable human carcinogen. Based on the results of Johnson et al. (1986), EPA derived an oral slope factor of $4.5 \text{ (mg/kg/day)}^{-1}$ for acrylamide. For cancer risks of 1×10^{-4} – 1×10^{-6} , the corresponding dose levels range from 2.2×10^{-5} to 2.2×10^{-7} mg/kg/day. These risk levels are presented in Figure 3-2.

3.2.3 Dermal Exposure

3.2.3.1 Death

There are no reports of human deaths associated with dermal exposure to acrylamide.

Acrylamide has been demonstrated to be lethal to laboratory animals following a single dermal dose. Reported dermal LD₅₀ values are 252 mg/kg in rats (American Cyanamid Company 1973) and 941 mg/kg in rabbits (American Cyanamid Company 1977). Dow Chemical Company (1957) reported that one of two rabbits administered a single 24-hour dermal dose of 1,000 mg/kg died 2 days following dosing.

3. HEALTH EFFECTS

Reliable acute dermal LD₅₀ values for death and other mortality data for each species are recorded in Table 3-6.

3.2.3.2 Systemic Effects

No human or animal data were located regarding respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, or endocrine effects following dermal exposure to acrylamide.

The highest NOAEL values and all LOAEL values from each reliable study for systemic effects in each species and duration category are recorded in Table 3-6.

Dermal Effects. Peeling of the skin was a common complaint among workers in factories with measurable acrylamide levels in the air and probable dermal exposure as well (Bachmann et al. 1992; Calleman et al. 1994; He et al. 1989; Myers and Macun 1991). Available information in animals is limited. Slight dermal irritation was reported in rats and rabbits following single dermal application of acrylamide in the range of 200–1,120 mg/kg (American Cyanamid Company 1951, 1973, 1977; Dow Chemical Company 1957). There was no evidence of dermal irritation in male mice receiving dermal applications of acrylamide for 5 days at doses ranging from 25 to 125 mg/kg/day (Gutierrez-Espeleta et al. 1992).

Ocular Effects. No data were located regarding ocular effects in humans following dermal exposure to acrylamide. Mild to moderate irritation was noted in the eyes of rabbits following ocular instillation of acrylamide (in water); irritation resolved during the ensuing 24 hours (American Cyanamid Company 1951; Dow Chemical Company 1957).

Body Weight Effects. No data were located regarding body weight effects in humans following dermal exposure to acrylamide. Available information in animals is restricted to a report of very slight initial weight loss (magnitude not specified) among a group of two rabbits administered a single 500 mg/kg dermal dose of acrylamide (Dow Chemical Company 1957) and another report of weight loss Table 3-6 in rabbits administered acrylamide dermally at 50 mg/kg/day for 5 weeks (Rohm and Haas Company 1975).

Table 3-6 Levels of Significant Exposure to Acrylamide - Dermal

Species (Strain)	Exposure/ Duration/ Frequency (Route)	System	NOAEL	LOAEL		Reference Chemical Form	Comments
				Less Serious	Serious		
ACUTE EXPOSURE							
Death							
Rat (albino)	Once				252 M mg/kg	(LD50)	American Cyanamid Company 1973
Mouse	5 d 1 x/d				200 M mg/kg/day	(4/8 died during 30 days following cessation of dosing)	Gutierrez-Espeleta et al. 1992 Acrylamide
Rabbit (New Zealand)	Once				941 M mg/kg	(LD50)	American Cyanamid Company 1977
Rabbit (NS)	Once				1000 mg/kg	(death of 1/2 rabbits)	Dow Chemical Company 1957; McCollister et al. 1964
Systemic							
Mouse	5 d 1 x/d	Dermal	125 M mg/kg/day				Gutierrez-Espeleta et al. 1992 Acrylamide Histopathological evaluations of peripheral nerve fibers not performed
Rabbit (albino)	Once 18 h	Dermal		1120 M mg/kg	(slight dermal irritation in 1/3 rabbits)		American Cyanamid Company 1951

DRAFT FOR PUBLIC COMMENT

Table 3-6 Levels of Significant Exposure to Acrylamide - Dermal

(continued)

Species (Strain)	Exposure/Duration/Frequency (Route)	System	LOAEL			Reference Chemical Form	Comments
			NOAEL	Less Serious	Serious		
Neurological							
Rat (albino)	Once				200 M (unspecified CNS symptoms)	American Cyanamid Company 1973	
Reproductive							
Mouse	5 d 1 x/d		25 M mg/kg		50 M (significantly increased number of dead implants)	Gutierrez-Espeleta et al. 1992 Acrylamide	
INTERMEDIATE EXPOSURE							
Neurological							
Rabbit (New Zealand)	5 wk 3 x/d				50 (clinical signs of peripheral neuropathy)	Rohm and Haas Company 1975 Acrylamide	

d = day(s); hr = hour(s); LD50 = lethal dose, 50% kill; LOAEL = lowest-observed-adverse-effect level; M = male; NOAEL = no-observed-adverse-effect level; wk = week(s); x = time(s)

DRAFT FOR PUBLIC COMMENT

3. HEALTH EFFECTS

3.2.3.3 Immunological and Lymphoreticular Effects

No human or animal data were located regarding immunological or lymphoreticular effects following dermal exposure to acrylamide.

3.2.3.4 Neurological Effects

Available human data consist of occupational exposure scenarios with suspected inhalation and dermal exposure. See Section 3.2.1.4 for information regarding neurological effects in occupationally-exposed workers. Unspecified clinical signs of central nervous system effects were reported in rats that died following single dermal application of acrylamide at doses in the range of 200–800 mg/kg (American Cyanamid Company 1973). Clinical signs that included shaking and loss of coordination were reported in rabbits following single dermal application at doses in the range of 784–1,568 mg/kg (American Cyanamid Company 1977). Clinical signs of hindlimb neuropathy became apparent during treatment week 4 in a study of rabbits administered acrylamide dermally for 5 weeks at 50 mg/kg/day (Rohm and Haas Company 1975).

The highest NOAEL values and all LOAEL values from each reliable study for neurological effects in each species and duration category are recorded in Table 3-6.

3.2.3.5 Reproductive Effects

No data were located regarding reproductive effects in humans following dermal exposure to acrylamide. Dermal applications of acrylamide to male mice for 5 days at doses ranging from 25 to 125 mg/kg/day followed by matings to unexposed female mice resulted in significantly increased numbers of dead implants (Gutierrez-Espeleta et al. 1992). At doses ≥ 50 mg/kg/day, significantly decreased numbers of living embryos were noted as well; this effect is indicative of dominant lethality.

The highest NOAEL values and all LOAEL values from each reliable study for reproductive effects in each species and duration category are recorded in Table 3-6.

3. HEALTH EFFECTS

3.2.3.6 Developmental Effects

No data were located regarding developmental effects in humans or animals following dermal exposure to acrylamide.

3.2.3.7 Cancer

No data were located regarding cancer associated with dermal exposure to acrylamide in humans.

The potential for acrylamide to initiate skin tumors was examined in a study of female mice administered acrylamide at dermal dose levels of 0, 12.5, 25, or 50 mg/kg/day, 6 times during a 2-week period, followed 2 weeks later by dermal applications of 12-O-tetradecanoylphorbol-13-acetate (TPA, a tumor promoter) to the shaved back 3 times/week for 20 weeks (Bull et al. 1984a). Incidences of skin papillomas and skin carcinomas were not significantly elevated at any acrylamide dose level. It was concluded that acrylamide did not act as a skin tumor initiator under the conditions of the study.

3.3 GENOTOXICITY

The genotoxicity of acrylamide has been studied both *in vivo* and *in vitro*. Studies are limited almost exclusively to laboratory rodents and nonmammalian species with the exception of a few *in vitro* assays of human cells. Results indicate that acrylamide is genotoxic and most potent in its ability to induce clastogenic effects (including heritable translocations in offspring of acrylamide-exposed male rodents mated with untreated females), deoxyribonucleic acid (DNA) damage, and gene mutations (including male-germ-cell-mediated dominant lethal mutations and heritable specific-locus mutations).

Table 3-7 summarizes the results of *in vivo* assays for acrylamide. Numerous assays have been performed in rodents. Dominant lethal mutations were induced consistently following administration of acrylamide via oral, dermal, or intraperitoneal injection in rodents (Adler et al. 2000; Chapin et al. 1995; Gutierrez-Espeleta et al. 1992; Sakamoto and Hashimoto 1986; Shelby et al. 1987; Smith et al. 1986; Sublet et al. 1989; Tyl et al. 2000a, 2000b; Working et al. 1987; Zenick et al. 1986). Exposure via intraperitoneal injection was also associated with specific locus mutations in offspring of male mice exposed to 50–125 mg/kg acrylamide before mating to untreated females (Ehling and Neuhäuser-Klaus

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mammalian gene mutation				
Mouse	Dominant lethal mutation	+	1x125 mg/kg, intraperitoneal injection of males before mating with untreated females	Adler et al. 2000
Mouse	Dominant lethal mutation	+	5x40 or 50 mg/kg, intraperitoneal injection of males before mating with untreated females	Shelby et al. 1987
Mouse	Dominant lethal mutation	+	6 weeks; 0.81, 3.19, or 7.22 mg/kg/day, drinking water of males before mating with untreated females	Chapin et al. 1995
Mouse	Dominant lethal mutation	+	4 weeks; 3.3–16.3 mg/kg/day estimated drinking water doses of males before mating with untreated females	Sakamoto and Hashimoto 1986
Rat	Dominant lethal mutation	+	64 days; 0.5, 5.0 mg/kg/day drinking water of males before mating with untreated females	Tyl et al. 2000a, 2000b
Rat	Dominant lethal mutation	+	5 days; 5, 15, 30, 45, or 60 mg/kg drinking water of males before mating with untreated females	Sublet et al. 1989
Rat	Dominant lethal mutation	+	5 days; 30 mg/kg gavage of males before mating with untreated females	Working et al. 1987
Rat	Dominant lethal mutation	+	80 days; 1.5, 2.8, or 5.8 mg/kg/day, drinking water of males before mating with untreated females	Smith et al. 1986
Rat	Dominant lethal mutation	+	10 weeks; 4.65, 7.86, or 11.53 mg/kg/day estimated drinking water doses of males before mating with untreated females	Zenick et al. 1986
Mouse	Dominant lethal mutation	+	5 days 25–125 mg/kg/day, dermal exposure of males before mating with untreated females	Gutierrez-Espeleta et al. 1992
Mouse (spleen lymphocytes, tk and HPRT loci)	Gene mutation	–	0, 0.14, or 0.70 mmol/kg, intraperitoneal injection on postnatal days 1, 8, or 15	Von Tungeln et al. 2009

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mouse (spleen lymphocytes, tk and HPRT loci)	Gene mutation	+	0, 0.14, or 0.70 mmol/kg, intraperitoneal injection on postnatal days 1–8	Von Tungeln et al. 2009
Mouse (liver cells, lymphocyte HPRT locus)	Gene mutation	+	3–4 weeks; 100 or 500 mg/L drinking water	Manjanatha et al. 2006
Muta® Mouse (lac z loci)	Gene mutation	+	5x50 mg/kg intraperitoneal injection	Hoorn et al. 1993
Muta® Mouse (lac z loci)	Gene mutation	–	1x50–100 mg/kg, intraperitoneal injection	Krebs and Favor 1997
Mouse (offspring coat color loci)	Gene mutation	+	1x50 or 75 mg/kg, intraperitoneal injection of pregnant females	Neuhäuser-Klaus and Schmahl 1989
Mouse (offspring coat color loci)	Gene mutation	+	3x50 or 75 mg/kg, intraperitoneal injection of pregnant females	Neuhäuser-Klaus and Schmahl 1989
Mouse (offspring coat color loci)	Gene mutation	+	5x50 mg/kg, intraperitoneal injection of males before mating with untreated females	Russell et al. 1991
Mouse (offspring coat color loci)	Gene mutation	+	1x100–125 mg/kg, intraperitoneal injection of males before mating with untreated females	Ehling and Neuhäuser-Klaus 1992
Chromosomal alterations in mammalian cells				
Mouse (bone marrow)	Chromosomal aberration	+	1x50–150 mg/kg, intraperitoneal injection	Adler et al. 1988
Mouse (bone marrow)	Chromosomal aberration	+	1x100 mg/kg, intraperitoneal injection	Čihák and Vontorková 1988
Mouse (bone marrow)	Chromosomal aberration	–	1x100–200 mg/kg, intraperitoneal injection	Shiraishi 1978
Mouse (bone marrow)	Chromosomal aberration	–	7–21 days 78 mg/kg-day diet	Shiraishi 1978
Rat (bone marrow)	Chromosomal aberration	–	1x100 mg/kg, intraperitoneal injection	Krishna and Theiss 1995
Mouse (spleen lymphocyte)	Chromosomal aberration	–	1x50–125 mg/kg, intraperitoneal injection	Backer et al. 1989
Mouse (splenocyte)	Chromosomal aberration	–	1x100 mg/kg, intraperitoneal injection	Kligerman et al. 1991
Mouse (spermatogonia)	Chromosomal aberration	–	1x50–150 mg/kg, intraperitoneal injection	Adler et al. 1988
Mouse (spermatogonia)	Chromosomal aberration	–	1x50–125 mg/kg, intraperitoneal injection	Backer et al. 1989
Mouse (spermatogonia)	Chromosomal aberration	–	5x50 mg/kg/day, intraperitoneal injection	Adler 1990

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mouse (spermatocyte)	Chromosomal aberration	+	1x100 mg/kg, intraperitoneal injection	Adler 1990
Mouse (first cleavage one-cell zygote)	Chromosomal aberration	+	1x75 or 125 mg/kg or 5x50 mg/kg/day, intraperitoneal injection of males before mating with untreated females	Pacchierotti et al. 1994
Mouse (first cleavage zygote)	Chromosomal aberration	+	5x50 mg/kg/day, intraperitoneal injection of males before mating with untreated females	Marchetti et al. 1997
Mouse (offspring spermatocyte)	Heritable translocation	+	5x40–50 mg/kg/day, intraperitoneal injection of males before mating with untreated females	Shelby et al. 1987
Mouse (offspring spermatid)	Heritable translocation	+	1x50–100 mg/kg, intraperitoneal injection of males before mating with untreated females	Adler et al. 1994
Mouse (offspring spermatocyte)	Heritable translocation	+	5x50 mg/kg/day, dermal exposure of males before mating with untreated females	Adler et al. 2004
Mouse (offspring spermatocyte)	Heritable translocation	+	5x50 mg/kg/day, intraperitoneal injection	Adler 1990
Mouse (bone marrow)	Polyploidy or aneuploid	+	1x100–200 mg/kg, intraperitoneal injection	Shiraishi 1978
Mouse (bone marrow)	Polyploidy or aneuploid	+	7–21 days 78 mg/kg/day, diet	Shiraishi 1978
Mouse (bone marrow)	Spindle disturbance	–	1x120 mg/kg, intraperitoneal injection	Adler et al. 1993
Mouse (bone marrow)	Micronucleus	+	1x50–125 mg/kg, intraperitoneal injection	Adler et al. 1988
Mouse (bone marrow)	Micronucleus	+	1x100 mg/kg, intraperitoneal injection	Čihák and Vontorková 1988
Mouse (bone marrow)	Micronucleus	+	2 days 25–100 mg/kg/day, intraperitoneal injection	Čihák and Vontorková 1988
Mouse (bone marrow)	Micronucleus	+	1x136 mg/kg, intraperitoneal injection	Knaap et al. 1988
Mouse (bone marrow)	Micronucleus	+	1, 2, or 3 days 42.5–100 mg/kg/day, intraperitoneal injection	Čihák and Vontorková 1990
Sprague-Dawley Rat (bone marrow)	Micronucleus	–	1x100 mg/kg, intraperitoneal injection	Paulsson et al. 2002
Rat (bone marrow)	Micronucleus	–	1x100 mg/kg, intraperitoneal injection	Krishna and Theiss 1995

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mouse (reticulocyte)	Micronucleus	+	1x50–100 mg/kg, intraperitoneal injection	Russo et al. 1994
Mouse (reticulocyte)	Micronucleus	+	1x25–100 mg/kg, intraperitoneal injection	Paulsson et al. 2002
Mouse (reticulocyte and nonchromatic erythrocyte)	Micronucleus	–	0, 0.14, 0.70 mmol/kg, postnatal day 1, 8, 15, intraperitoneal injection	Von Tungeln et al. 2009
Mouse (reticulocyte and nonchromatic erythrocyte)	Micronucleus	–	0, 0.14, 0.70 mmol/kg, postnatal day 1–8, intraperitoneal injection	Von Tungeln et al. 2009
Mouse (reticulocyte)	Micronucleus	+	28 days 0–24 mg/kg/day, gavage	Zeiger et al. 2009
Mouse (normochromatic erythrocyte)	Micronucleus	+	28 days 0–24 mg/kg/day, gavage	Zeiger et al. 2009
Mouse (erythrocyte)	Micronucleus	+	5 days 0, 25, 50 mg/kg/day, intraperitoneal injection	Ghanayem et al. 2005b
Mouse (spleen lymphocyte)	Micronucleus	+	1x50–125 mg/kg, intraperitoneal injection	Backer et al. 1989
Mouse (splenocyte)	Micronucleus	+	1x100 mg/kg, intraperitoneal injection	Kligerman et al. 1991
Mouse (spermatid)	Micronucleus	+	1x10–100 mg/kg, intraperitoneal injection	Collins et al. 1992
Mouse (spermatid)	Micronucleus	+	1x50–100 mg/kg or 4x50 mg/kg/day, intraperitoneal injection	Russo et al. 1994
Lewis Rat (spermatid)	Micronucleus	+	1x50–100 mg/kg or 4x50 mg/kg/day, intraperitoneal injection	Xiao and Tates 1994
Sprague-Dawley rat (spermatid)	Micronucleus	+	1x50–100 mg/kg or 4x50 mg/kg/day, intraperitoneal injection	Lähdetie et al. 1994
Mouse (germ cell)	Synaptonemal complex aberration	–	1x50–150 mg/kg intraperitoneal injection	Backer et al. 1989
Mouse (germ cell)	Synaptonemal complex asynapsis	+	1x50–150 mg/kg, intraperitoneal injection Asynapsis in meiotic prophase	Backer et al. 1989
Sister chromatid exchange				
Mouse (spleen lymphocytes)	Chromatid exchange	+	1x50–125 mg/kg, intraperitoneal injection	Backer et al. 1989

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mouse (splenocytes)	Chromatid exchange	+	1x100 mg/kg, intraperitoneal injection	Kligerman et al. 1991
DNA damage				
Mouse (spermatocytes and early spermatids)	DNA breakage	+	1x0–125 mg/kg, intraperitoneal injection	Sega and Generoso 1990
Mouse (bone marrow, spleen, liver, kidney, lungs, testes)	DNA breakage	+	1x0–125 mg/kg, intraperitoneal injection	Dobrzynska 2007
Mouse (leukocytes, liver, lung)	DNA breakage	+	5 day 0, 25, 50 mg/kg/day, intraperitoneal injection (wild type mice)	Ghanayem et al. 2005b
Rat (hepatocyte)	Unscheduled DNA synthesis	–	1x100 mg/kg or 5x30 mg/kg/day, gavage	Butterworth et al. 1992
Rat (spermatocyte)	Unscheduled DNA synthesis	+	1x100 mg/kg or 5x30 mg/kg/day, gavage	Butterworth et al. 1992
Mouse (germ cell)	Unscheduled DNA synthesis	+	1x7.8–125 mg/kg, intraperitoneal injection	Sega et al. 1990
Mouse (testis)	DNA adduct (Alkylation)	+	1x46 mg/kg, intraperitoneal injection	Sega et al. 1990
Mouse (liver)	DNA adduct (Alkylation)	+	1x46 mg/kg, intraperitoneal injection	Sega et al. 1990
Sprague-Dawley rat (liver, lung, kidney, brain, testis)	DNA adduct (N7-GA-Gua)	+	1x46 mg/kg, intraperitoneal injection	Segeberäck et al. 1995
Mouse (liver, kidney, brain)	DNA adduct (N7-GA-Gua)	+	1x53 mg/kg, intraperitoneal injection	Segeberäck et al. 1995
Neonatal mouse (whole body)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	1x50 mg/kg, intraperitoneal injection	Gamboa da Costa et al. 2003
Mouse (liver, lung, kidney)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	1x50 mg/kg, intraperitoneal injection	Gamboa da Costa et al. 2003
Mouse (liver, lung)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	1x1–10 mg/kg, intraperitoneal injection	Gamboa da Costa et al. 2003
Mouse (lung, liver, spleen, bone marrow)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	0, 0.14, 0.70 mmol/kg intraperitoneal injection, postnatal day 1, 8, 15	Von Tungeln et al. 2009

3. HEALTH EFFECTS

Table 3-7. Genotoxicity of Acrylamide *In Vivo*

Species (test system)	End point	Result	Test conditions	Reference
Mouse (lung, liver, spleen)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	0, 0.14, 0.70 mmol/kg postnatal day 1–8, intraperitoneal injection	Von Tungeln et al. 2009
Mouse (liver)	DNA adduct (GA-Gua)	+	28 days 0–24 mg/kg/day, gavage	Zeiger et al. 2009
Mouse (liver, lung, kidney, leukocyte, testis)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	1x50 mg/kg, intraperitoneal injection	Doerge et al. 2005a
Mouse (liver)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	14 days 1 mg/kg/day, drinking water	Doerge et al. 2005a
Rat (liver, brain, thyroid, leukocyte, mammary gland, testis)	DNA adduct (N7-GA-Gua, N3-GA-Ade)	+	1x50 mg/kg, intraperitoneal injection	Doerge et al. 2005a
Rat (liver)	DNA adduct (N7-GA-Gua)	+	14 days 1 mg/kg/day, drinking water	Doerge et al. 2005a
Nonmammalian gene mutation				
<i>Drosophila melanogaster</i>	Sex-linked recessive lethal	–	1x40–50 mM, abdominal injection	Knaap et al. 1988
<i>D. melanogaster</i>	Sex-linked recessive lethal	+	48 hour 0.25–5 mM, larvae feeding	Tripathy et al. 1991
<i>D. melanogaster</i>	Somatic mutation and recombination	+	1–1.5 mM, larvae feeding until pupation	Knaap et al. 1988
<i>D. melanogaster</i>	Somatic mutation and recombination	+	10–30 mM, larvae feeding until pupation	Batiste-Alentorn et al. 1991
<i>D. melanogaster</i>	Somatic mutation and recombination	+	48 hour 0.25–5 mM, larvae feeding	Tripathy et al. 1991

– = negative result; + = positive result; DNA = deoxyribonucleic acid

3. HEALTH EFFECTS

1992; Russell et al. 1991) and in offspring of pregnant female mice exposed to 50 or 75 mg/kg (Neuhäuser-Klaus and Schmahl 1989). Heritable or reciprocal translocations were noted in offspring of male mice exposed to 50–100 mg/kg acrylamide via intraperitoneal injection or dermal application before mating to untreated females (Adler 1990; Adler et al. 1994, 2004; Shelby et al. 1987). Intraperitoneal exposure to acrylamide also increased mutations at the tk and HPRT loci in lymphocytes of mice (Majanatha et al. 2006; von Tungeln et al. 2009) and at lac z loci in transgenic mice (Hoom et al. 1993; Krebs and Favor 1997).

Prominent clastogenic effects consistently associated with *in vivo* exposure include sister chromatid exchanges (Backer et al. 1989; Kligerman et al. 1991), micronucleus formation (Backer et al. 1989; Čihák and Vontorková 1988, 1990; Collins et al. 1992; Ghanayem et al. 2005b; Kligerman et al. 1991; Knaap et al. 1988; Lähdetie et al. 1994; Paulsson et al. 2002; Russo et al. 1994; Xiao and Tate 1994; Zeiger et al. 2009), and aneuploidy or polyploidy (Shiraishi 1978). Acrylamide induced DNA damage in various rodent tissues (including testes and male germ cells), evidenced as DNA breakage (Dobrzynska 2007; Ghanayem et al. 2005b; Sega and Generoso 1990), unscheduled DNA synthesis (Butterworth et al. 1992; Sega et al. 1990), and DNA adduct formation (Doerge et al. 2005a; Gamboa da Costa et al. 2003; Sega et al. 1990; Segerbäck et al. 1995; Von Tungeln et al. 2009; Zeiger et al. 2009). Increases in chromosomal aberrations were observed in the first cleavage zygote of acrylamide-treated male mice mated with untreated females (Marchetti et al. 1997; Pacchierotti et al. 1994).

Assays for acrylamide-induced chromosomal aberrations in bone marrow, spleen, or spermatogonia of acrylamide-treated rodents produced both positive (Adler 1990; Adler et al. 1988; Čihák and Vontorková 1988) and negative (Adler 1990; Adler et al. 1988; Backer et al. 1989; Kligerman et al. 1991; Krishna and Theiss 1995; Shiraishi 1978) results. Acrylamide did not induce micronuclei in a small number of rat and mouse assays (Krishna and Theiss 1995; Paulsson et al. 2002; Von Tungeln et al. 2009). Results for synaptonemal complex aberrations in male mouse germ cells were negative or only weakly positive (Backer et al. 1989). Acrylamide did not induce spindle disturbances in the bone marrow of intraperitoneally-injected mice (Adler et al. 1993).

Somatic mutations and recombination (Batiste-Alentorn et al. 1991; Knaap et al. 1988; Tripathy et al. 1991) and sex-linked recessive lethal mutations (Tripathy et al. 1991) were induced in drosophila larval feeding assays. Negative results were obtained for sex-linked recessive lethal mutations in another assay that employed abdominal injection (Knaap et al. 1988).

3. HEALTH EFFECTS

Table 3-8 summarizes the results of *in vitro* assays. Acrylamide induced mutations at the tk and HPRT loci in several assays with mammalian cells including mouse lymphoma L5178Y cells (Barfknecht et al. 1988; Knaap et al. 1988; Mei et al. 2008b; Moore et al. 1987) and human promyelocytic leukemia HL-60 and NB4 cells (Ao et al. 2008). In Chinese hamster V79H3 cells (which do not express genes for CYP enzymes), acrylamide did not induce mutations at the HPRT locus (Tsuda et al. 1993). Acrylamide did not induce reverse mutations in multiple strains of *Salmonella typhimurium* or in *Escherichia coli* WP2 *uvrA* with or without metabolic activation (Hashimoto and Tanii 1985; Jung et al. 1992; Knaap et al. 1988; Lijinsky and Andrews 1980; Müller et al. 1993; Tsuda et al. 1993; Zeiger et al. 1987). One exception was a weakly positive result in a few trials of TA98 and TA100, but only with S9 activation (Zeiger et al. 1987).

Clastogenic effects associated with *in vitro* exposure to acrylamide include: chromosomal aberrations in mammalian cells (Knaap et al. 1988; Moore et al. 1987; Tsuda et al. 1993); polyploidy, sister chromatid exchanges, and spindle disturbances in Chinese hamster cells (Adler et al. 1993; Knaap et al. 1988; Martins et al. 2007; Tsuda et al. 1993; Warr et al. 1990); and micronuclei in human hepatoma G2 cells (Jiang et al. 2007). Although acrylamide did not induce micronuclei in one *in vitro* assay using seminiferous tubule segments from rats (Lähdetie et al. 1994), micronuclei were induced in spermatids of rodents exposed to acrylamide *in vivo* (Collins et al. 1992; Lähdetie et al. 1994; Russo et al. 1994; Xiao and Tate 1994).

DNA damage associated with *in vitro* exposure to acrylamide includes DNA damage in *Bacillus subtilis* (Tsuda et al. 1993); DNA breakage and oxidative DNA damage in human hepatoma G2 cells (Jiang et al. 2007); unscheduled DNA synthesis in human mammary epithelial cells (Butterworth et al. 1992); and glycidamide-DNA adducts in Chinese hamster V79 cells (Martins et al. 2007), mouse embryonic fibroblasts (Besaratina and Pfeifer 2004), and human bronchial epithelial cells (Besaratina and Pfeifer 2004). DNA adducts were not detected when mouse lymphoma cells were incubated with up to 20 mM acrylamide, but glycidamide concentrations in the range of 0.5–4 mM induced DNA adducts in these cells (Mei et al. 2008b). Acrylamide *in vitro* exposure also caused cell morphological transformations in several mouse cell lines and Syrian hamster embryo cells (Abernethy and Boreiko 1987; Banerjee and Segal 1986; Park et al. 2002; Tsuda et al. 1993).

3. HEALTH EFFECTS

Table 3-8. Genotoxicity of Acrylamide *In Vitro*

Species (test system)	End point	Result		Test conditions	Reference
		Activation			
		With	out		
Prokaryotic					
Bacterial gene mutation					
<i>Salmonella typhimurium</i> TA98, TA100, TA1535, TA1537	Reverse mutation	-	-	10–10,000 µg/plate +/- S9 activation Weakly positive in TA98 and TA100 in only a few trials with activation; all other strains negative	Zeiger et al. 1987
<i>S. typhimurium</i> TA97, TA98, TA100, TA1535	Reverse mutation	-	-	100–10,000 µg/plate +/- S9 activation	Zeiger et al. 1987
<i>S. typhimurium</i> TA98, TA100, TA102, TA1535, TA1537	Reverse mutation	-	-	1–100 mg/plate +/- S9 activation	Knaap et al. 1988
<i>S. typhimurium</i> TA98, TA100, TA1535, TA1537	Reverse mutation	-	-	0.5–50 mg/plate +/- S9 activation	Tsuda et al. 1993
<i>S. typhimurium</i> TA1535	Reverse mutation	-	-	Up to 5 mg/plate +/- S9 activation	Müller et al. 1993 ; Jung et al. 1992
<i>S. typhimurium</i> TA98, TA100, TA1535, TA1537, TA1538	Reverse mutation	-	-	Up to 1 mg/plate +/- S9 activation	Lijinsky and Andrews 1980
<i>S. typhimurium</i> TA98, TA100, TA 1535, TA1537, TA1538	Reverse mutation	-	-	0.5–5,000 µg/plate +/- S9 activation	Hashimoto and Tanii 1985
<i>Escherichia coli</i> WP2 <i>uvrA</i>	Reverse mutation	-	-	0.5–50 mg/plate +/- S9 activation	Tsuda et al. 1993
<i>Klebsiella pneumoniae</i> ur ⁻ pro ⁻	Fluctuation	No data	-	2–10 mg/L	Knaap et al. 1988
DNA damage and repair and DNA adduct formation					
<i>Bacillus subtilis</i>	DNA damage	+	+	1–50 mg/ disk +/- S9 activation	Tsuda et al. 1993
Mammalian gene mutation assays					
Mouse lymphoma L5178Y TK ^{+/+} , tk locus	Gene mutation	+	+	10 mM	Barfknecht et al. 1988
Mouse lymphoma L5178Y TK ^{+/+} , tk locus	Gene mutation	No data	+	0–0.85 mg/mL No activation Clastogenic response	Moore et al. 1987
Mouse lymphoma L5178Y TK ^{+/+} , tk locus	Gene mutation	No data	+	0–18 mM No activation	Mei et al. 1988

3. HEALTH EFFECTS

Table 3-8. Genotoxicity of Acrylamide *In Vitro*

Species (test system)	End point	Result		Test conditions	Reference
		Activation			
		With	out		
Mouse lymphoma L5178Y TK ^{+/+} , tk and HPRT loci	Gene mutation	-	-	0.5–7.5 mg/mL +/- S9 activation Mutation only at cytotoxic concentrations	Knaap et al. 1988
Mouse lymphoma L5178Y TK ^{+/+} , HPRT locus	Gene mutation	+	+	0.1–0.5 mg/mL Cocultivated activation	Knaap et al. 1988
Chinese hamster V79H3, HPRT locus	Gene mutation	No data	-	1–7 mM No activation	Tsuda et al. 1993
Human promyelocytic leukemia HL-60 and NB4, HPRT locus	Gene mutation	No data	+	0–700 mg/L	Ao et al. 2008
Chromosomal alteration in mammalian cells					
Chinese hamster V79H3	Chromosomal aberration	No data	+	0.5–5 mM No activation	Tsuda et al. 1993
Chinese hamster V79	Chromosomal aberration	+	+	0.1–3 mg/mL +/- S9 activation	Knaap et al. 1988
Chinese hamster V79	Chromosomal aberration	No data	-	0–2,000 µM No activation	Martins et al. 2007
Mouse lymphoma L5178Y TK ^{+/+}	Chromosomal aberration	No data	+	0.65–0.85 mg/mL No activation	Moore et al. 1987
Chinese hamster V79H3	Polyploidy	No data	+	0.5–5 mM	Tsuda et al. 1993
Chinese hamster LUC2 p5	Polyploidy	No data	+	0.0125–0.5 mg/mL	Warr et al. 1990
Chinese hamster V79	Spindle disturbance	No data	+	0.01–1 mg/mL	Adler et al. 1993
Chinese hamster DON:Wg3h	Spindle disturbance	No data	+	0.2–2 mg/mL	Warr et al. 1990
Chinese hamster LUC2 p5	Spindle disturbance	No data	+	0.01–1 mg/mL	Warr et al. 1990
Rat seminiferous tubular segments	Micronucleus	No data	-	5–50 µg/mL	Lähdetie et al. 1994
Human hepatoma G2	Micronucleus	No data	+	0–2.5 mM	Jiang et al. 2007

3. HEALTH EFFECTS

Table 3-8. Genotoxicity of Acrylamide *In Vitro*

Species (test system)	End point	Result		Test conditions	Reference
		Activation			
		With	out		
Sister chromatid exchange					
Chinese hamster V79	Chromatid exchange	+	+	0.1–1 mg/mL	Knaap et al. 1988
Chinese hamster V79	Chromatid exchange	No data	+	0.5–2.5 mg/mL	Tsuda et al. 1993
Chinese hamster V79	Chromatid exchange	No data	+	0–2,000 µM	Martins et al. 2007
DNA damage and repair and DNA adduct formation					
Human hepatoma G2	DNA breakage	No data	+	0–20 mM	Jiang et al. 2007
Human hepatoma G2	Oxidative DNA damage	No data	+	0–20 mM	Jiang et al. 2007
F344 Rat primary hepatocytes	Unscheduled DNA synthesis	No data	–	0.01–1 mM	Butterworth et al. 1992
Human mammary epithelial	Unscheduled DNA synthesis	No data	+	1–10 mM	Butterworth et al. 1992
Chinese hamster V79	DNA adducts (N7-GA-Gua, N3-GA-Ade)	No data	+	0–2,000 µM	Martins et al. 2007
Mouse lymphoma L5178Y TK ^{+/–}	DNA adducts (N7-GA-Gua, N3-GA-Ade)	No data	–	0–20 mM	Mei et al. 2008b
Big Blue mouse embryonic fibroblasts	DNA adducts (N7-GA-Gua, N1-GA-Ade, N3-GA-Ade)	No data	+	0, 0.0032, 0.320. and 16 mM	Besaratinia and Pfeifer 2004
Human bronchial epithelial	DNA adducts (N7-GA-Gua, N1-GA-Ade, N3-GA-Ade)	No data	+	0, 0.320 and 3.2 mM	Besaratinia and Pfeifer 2004
Cell transformation					
Mouse C3H/10T1/2	Morphological transformation	No data	+	25–200 µg/mL	Banerjee and Segal 1986
Mouse NIH/3T ₃	Morphological transformation	No data	+	2–200 µg/mL	Banerjee and Segal 1986

3. HEALTH EFFECTS

Table 3-8. Genotoxicity of Acrylamide *In Vitro*

Species (test system)	End point	Result		Test conditions	Reference
		Activation			
		With	out		
Mouse C3H/10T1/2	Morphological transformation	No data	–	10–300 µg/mL	Abernethy and Boreiko 1987
Mouse BALB/c 3T ₃	Morphological transformation	No data	+	0.5–2 mM	Tsuda et al. 1993
Syrian hamster embryo	Morphological transformation	No data	+	0.1–0.7 mM	Park et al. 2002
Syrian hamster embryo	Morphological transformation	No data	–	0.001–10 mM	Kaster et al. 1998

– = negative result; + = positive result; DNA = deoxyribonucleic acid

3. HEALTH EFFECTS

3.4 TOXICOKINETICS

Acrylamide and its principal and toxicologically significant (epoxide) metabolite, glycidamide, react with various biologically significant targets. The chemical basis for these interactions is strongly associated with the degree of electrophilicity (electron deficiency) with nucleophilic centers (i.e., unshared electrons). Electrophiles and nucleophiles are generally characterized as being either “hard” or “soft” corresponding to a spectral range of high or low charge densities or electronegativity for reactivity (Pearson and Songstad 1967). Due to its α,β -unsaturated structure and ready capacity to undergo Michael-type additions, acrylamide may be classified as a “soft” electrophile. Soft electrophiles like acrylamide react readily with soft nucleophiles such as the thiol groups of proteins or glutathione. Glycidamide, on the other hand, has a relatively high positive charge density, and acts as a hard electrophile, more capable of reacting with centers of high electronegativity (i.e., hard nucleophiles) such as the purine and pyrimidine bases in DNA (Dearfield et al. 1995; Lopachin and DeCaprio 2005;). Hemoglobin adducts have been used as biomarkers of exposure to acrylamide and are based on the assumption that a measured adduct level represents a steady-state level from a continuous exposure to acrylamide over the previous 120 days, which is the average life span of a red blood cell. Hemoglobin adduct levels provide a direct measure of the total amount of acrylamide and its reactive metabolite, glycidamide, in the blood over a given time period, which is quantified as the area under the curve (AUC in amount-unit time/volume). AUC is the integral of “concentration” (e.g., mg or mmol/L) \times “time” (e.g., minutes or hours). Under the reasonable assumption that the amount of parent or reactive toxicant in blood indicates the amount available to bind to tissue macromolecules or DNA, hemoglobin adducts provide a relevant internal metric for use in estimating the risk of acrylamide toxicity.

3.4.1 Absorption

Detection of hemoglobin adducts of acrylamide in exposed workers and volunteers provides qualitative evidence of absorption. Controlled human studies of the formation of hemoglobin adducts of acrylamide following oral and dermal exposure provide some information regarding the extent of absorption via these exposure routes. Available animal data indicate that acrylamide is readily and rapidly absorbed following inhalation and oral exposure, and somewhat less readily absorbed following dermal exposure.

3. HEALTH EFFECTS

3.4.1.1 Inhalation Exposure

Hagmar et al. (2001) measured hemoglobin adducts of acrylamide in a group of 210 tunnel construction workers who were occupationally exposed for 2 months to a chemical grouting agent containing acrylamide and N-methylolacrylamide. Within 1 month after construction work was completed, blood samples were drawn for the analysis of adducts of acrylamide with N-terminal valines (acrylamideVal) in hemoglobin. Workers were expected to have experienced both inhalation and dermal exposure.

Quantitative exposure data were limited to two personal air samples showing concentrations of 0.27 and 0.34 mg/m³ for the sum of acrylamide and N-methylolacrylamide; further analysis suggested that the air contained a 50:50 mixture of these compounds. Hemoglobin adduct levels for 18 nonsmoking unexposed reference subjects varied between 0.02 and 0.07 nmol/g globin. The frequency distribution of adduct levels in the 210 tunnel workers was as follows: 47 with <0.08 nmol/g globin; 89 with 0.08–0.29 nmol/g; 36 with 0.3–1.0 nmol/g; and 38 with 1.0–17.7 nmol/g. Adduct levels were determined in blood samples collected at intervals up to 5 months after cessation of exposure from five workers with initial levels ranging from about 2.2 to 4.4 nmol/g. Adduct levels decreased to background levels within 120 days, consistent with the approximate 120-day life of red blood cells.

Hemoglobin adduct levels were measured in 41 Chinese workers who were exposed to acrylamide for 0.1–8 years (Bergmark et al. 1993). AcrylamideVal hemoglobin levels were measured. Workers were involved in the production of acrylamide (via the hydration of acrylonitrile) and polyacrylamide. The adduct levels in exposed workers ranged from 0.3 to 34 nmol acrylamide/g hemoglobin. AcrylamideVal levels were not detected in blood samples of 10 control workers from the same city who had not been occupationally exposed to acrylamide (or acrylonitrile). Blood samples from 5 of the 41 exposed workers were also analyzed for hemoglobin adducts of glycidamide (glycidamideVal), a principal metabolite of acrylamide in animals (see also Section 3.4.3, Metabolism). A statistically significant linear relationship was observed between levels of acrylamideVal and glycidamideVal in these five workers; the ratio between acrylamideVal and glycidamideVal was approximately 3:10. Average levels of acrylamide in air samples were 1.52 and 0.73 mg/m³ for workplaces involved with polymerization and synthesis processes, respectively. Workers involved in these processes showed average acrylamideVal levels of 7.3±3.4 (n=12, polymerization) and 14.7±10.6 nmol/g hemoglobin (n=14, synthesis).

Bergmark et al. (1993) made the following assumptions based on a central assumption that these workers may have experienced only inhalation exposure: (1) adducts are stable during the life of erythrocytes; (2) the lifespan of human erythrocytes is about 120 days (17 weeks); (3) the second-order reaction rate

3. HEALTH EFFECTS

constant for the reaction of acrylamideVal in human hemoglobin is $4.4 \times 10^{-6} \text{ [L (g Hb)}^{-1} \text{ hour}^{-1}]$ (based on *in vitro* experiments); (4) the human ventilation rate is $0.2 \text{ L minute}^{-1} \text{ kg bw}^{-1}$; and (5) inhaled acrylamide is 100% absorbed. Using these assumptions, the predicted levels of acrylamideVal were only 0.93 and 0.44 nmol/g hemoglobin, respectively. The large disparity between the observed and predicted adduct levels resulted in the conclusion that dermal contact may have been the predominant source of absorption in these workers.

Animal studies indicate that inhaled acrylamide is readily absorbed (Sumner et al. 2003). Male F344 rats and B6C3F1 mice were exposed to approximately 3 ppm of a mixture of [^{13}C]-labeled acrylamide and [^{14}C]-labeled acrylamide vapor via nose-only inhalation for 6 hours. Selected rats and mice were sacrificed immediately following the exposure period for determination of [^{14}C] content in tissues, an indicator of the extent of absorption of inhaled acrylamide. The remaining rats and mice were monitored for 24-hour elimination of acrylamide and metabolites via urine, feces, and expired air. Immediately following the 6-hour exposure period, approximately 18 and 8 μmol of [^{14}C]-equivalents were recovered from tissues and carcasses of the rats and mice, respectively. At the end of the 24-hour postexposure period, 42% of the total recovered radioactivity was in urine, feces, and nose-tube and cage washes of rats; <3% was in exhaled air; and 56% remained in the body. In mice, 51% was recovered in urine, feces, and nose-tube and cage washes; <3% was in exhaled air, and 46% remained in the body. Fractional absorption could not be determined from the presented data because ventilation rates were apparently not measured.

3.4.1.2 Oral Exposure

Fennell and coworkers evaluated metabolism and hemoglobin adduct formation (Fennell et al. 2005b) and kinetics of elimination of urinary metabolites of acrylamide (Fennell et al. 2006) following oral administration of [$1,2,3\text{-}^{13}\text{C}_3$]-acrylamide to 24 adult male volunteers exposed under controlled conditions. All volunteers were aspermic (i.e., clinically sterile because of the potential for adverse effects of acrylamide on sperm), and had not used tobacco products for the prior 6 months. The health of the volunteers was continually monitored. Acrylamide was administered in an aqueous solution (single dose of 0.5, 1, or 3 mg/kg) to the volunteers. For the 3 mg/kg dose group, approximately 40% of the administered dose was recovered as urinary metabolites in the 24-hour urine. Approximately 86% of the urinary metabolites were derived from GSH conjugation and excreted as N-acetyl-S-(3-amino-3-oxopropyl)cysteine and its S-oxide; the remainder included glycidamide, glyceramide, and low levels of N-acetyl-S-(3-amino-2-hydroxy-3-oxopropyl)cysteine. Mean levels of hemoglobin adducts of acrylamide

3. HEALTH EFFECTS

($^{13}\text{C}_3$ -acrylamideVal) and glycidamide ($^{13}\text{C}_3$ -glycidamideVal) in the 3 mg/kg group at 24 hours postadministration were 2,479 and 1,076 fmol/mg globin, respectively. These findings demonstrate that orally-administered acrylamide is rapidly and readily absorbed by the gastrointestinal tract.

Fuhr et al. (2006) evaluated the toxicokinetics of acrylamide in six young healthy volunteers after the consumption of a meal containing 0.94 mg of acrylamide. Urine was collected up to 72 hours thereafter. Unchanged acrylamide; its mercapturic acid metabolite, N-acetyl-S-(2-carbamoylethyl)cysteine (AAMA), its epoxy derivative, glycidamide; and the respective metabolite of glycidamide, N-acetyl-S-(2-hydroxy-2-carbamoylethyl)cysteine (GAMA), were quantified in the urine by liquid chromatography-mass spectrometry. Toxicokinetic variables were obtained by noncompartmental methods. Overall, approximately 60% of the dose was recovered in the urine. Although no glycidamide was found, unchanged acrylamide, AAMA, and GAMA accounted for urinary excretion of approximately 4.5, 50, and 6% of the dose, respectively. These results indicate that most of the acrylamide ingested with food is absorbed in humans.

Boettcher et al. (2006a) investigated the human metabolism of acrylamide to AAMA and GAMA in a healthy male volunteer who received a single dose of about 1 mg deuterium-labeled (d(3)) acrylamide, representing 13 $\mu\text{g}/\text{kg}$ body weight, in drinking water. Urine samples before dosing and within 46 hours after the dose were analyzed for d(3)-AAMA and d(3)-GAMA. Total recovery of AAMA and GAMA in the 24-hour urine was about 51% of the administered dose, which provides additional demonstration of the rapid and extensive absorption of ingested acrylamide.

Boettcher et al. (2006b) reported the influence of an acrylamide-free diet on the excretion of urinary mercapturic acid metabolites derived from acrylamide in three healthy volunteers who fasted for 48 hours. Urinary acrylamide mercapturic acid metabolites were considerably reduced after 48 hours of fasting, with levels even well below the median level in nonsmokers. These results indicate that the acrylamide in the diet is the main source of environmental acrylamide exposure in humans, apart from smoking.

Bjellaas et al. (2007a) reported urinary mercapturic acid derivatives of acrylamide in a clinical study comprised of 53 subjects. Median intakes (range) of acrylamide were estimated based on 24-hour dietary recall as 21 (13–178) μg for nonsmokers and 26 (12–67) μg for smokers. The median dietary exposure to acrylamide was estimated to be 0.47 (range 0.17–1.16) $\mu\text{g}/\text{kg}$ body weight per day. The median (range) total excretion of acrylamide in urine during 24 hours was 16 (7–47) μg for nonsmokers and 74 (38–106)

3. HEALTH EFFECTS

µg for smokers. In a multiple linear regression analysis, a statistically significant correlation was found between the urinary excretion of acrylamide metabolites and intake of aspartic acid, protein, starch, and coffee.

Studies in rats indicate that orally administered acrylamide is rapidly and extensively absorbed by the gastrointestinal tract (Dixit et al. 1982; Doerge et al. 2005b; Fennell et al. 2005a; Kadry et al. 1999; Miller et al. 1982; Ramsey et al. 1984). The time course of urinary elimination of radioactivity from male F344 rats during a 7-day period following single oral doses of 10 mg/kg [2,3-¹⁴C]-acrylamide was essentially the same as that observed with male F344 rats following single intravenous dosing at 10 mg/kg [2,3-¹⁴C]-acrylamide (Miller et al. 1982). This observation suggests that 100% of the oral dose was absorbed. The time courses of urinary elimination of radioactivity were similar for groups of rats given single 1, 10, or 100 mg/kg oral doses of radiolabeled acrylamide, suggesting that the extent of absorption was not affected by dose level in the tested range. The rapidity of absorption was demonstrated by observations that peak plasma levels of radioactivity were attained by 1 hour after administration and that 53–67% of administered radioactivity was detected in the urine collected within 24 hours of administration (Miller et al. 1982). Similar results indicating rapid and extensive oral absorption were reported for studies in which male Sprague-Dawley rats were given single 50 mg/kg oral doses of [1-¹⁴C]-acrylamide (Kadry et al. 1999). Radioactivity was detected in blood 5 minutes after administration, and peak plasma levels of radioactivity occurred at 38 minutes after administration. Approximately 51% of administered radioactivity was detected in urine collected within 24 hours of administration (Kadry et al. 1999). Fennell et al. (2005a) administered 3 mg/kg of [1,2,3-¹³C₃]-acrylamide by gavage to male F344 rats. The total amount of acrylamide metabolites recovered in urine by 24 hours after dosing was 50%, which is similar to that reported by Miller et al. (1982) and Kadry et al. (1999).

Doerge et al. (2005c) compared the toxicokinetics of acrylamide and glycidamide in serum and tissues of male and female B6C3F1 mice following a single 0.1 mg/kg acrylamide dose by intravenous injection or gavage or a comparable dose of acrylamide from a feeding exposure for 30 minutes. Study groups also received an equimolar amount of glycidamide from either intravenous injection or gavage dosing. Following oral dosing, acrylamide was rapidly absorbed, widely distributed to tissues, and efficiently converted to glycidamide. Liver levels of glycidamide-DNA adducts were increased at 8 hours postdosing when acrylamide had been eliminated from the serum. Oral glycidamide dosing also resulted in rapid absorption, wide distribution to tissues, and liver DNA adduct levels that were approximately 40% higher than those from an equimolar dose of orally-administered acrylamide. Based on the kinetics

3. HEALTH EFFECTS

of acrylamide following intravenous injection, oral administration from the diet attenuated acrylamide bioavailability to 23% of the intravenous dose, and aqueous gavage attenuated acrylamide bioavailability to 32–52%. In contrast, oral exposure resulted in higher relative internal glycidamide levels compared with levels following an intravenous injection, likely due to a first-pass effect but possibly the result of some other kinetic change.

3.4.1.3 Dermal Exposure

Fennell and coworkers evaluated metabolism and hemoglobin adduct formation (Fennell et al. 2005b) and kinetics of elimination of urinary metabolites of acrylamide (Fennell et al. 2006) following dermal administration of [1,2,3-¹³C₃]-acrylamide to 24 adult male volunteers exposed under controlled conditions. All volunteers were aspermic (i.e., clinically sterile because of the potential for adverse effects of acrylamide on sperm) and had not used tobacco products for the prior 6 months. The health of the volunteers was continually monitored. Acrylamide was administered as three daily 24-hour occluded dermal doses of 3 mg/kg. Mean levels of hemoglobin adducts of acrylamide (¹³C₃-acrylamideVal) and glycidamide (¹³C₃-glycidamideVal) increased from 116 and 55 fmol/mg globin, respectively, following the first exposure period to 464 and 316 fmol/mg globin, respectively, following the last exposure period. Based on total amount administered, formation of acrylamideVal after dermal exposure was much lower than after oral administration (4.9 vs. 74.7 nmol/g globin/mmol acrylamide/kg). Approximately 4.5% of the administered dose was recovered as urinary metabolites through day 4. Dermal exposure also resulted in much lower formation of glycidamideVal (9.7% of that formed following oral exposure).

Animal studies confirm that acrylamide is readily absorbed following dermal exposure (Ramsey et al. 1984; Sumner et al. 2003). In male F344 rats, an average of 22% of an occluded dermal dose of 162 mg/kg [2,3-¹⁴C]-labeled acrylamide was absorbed during a 24-hour exposure period (Sumner et al. 2003). Following dermal administration of [1,3-¹⁴C]-labeled acrylamide to other F344 rats, peak plasma concentrations of acrylamide occurred at approximately 2 and 5 hours postadministration, respectively (Ramsey et al. 1984). The peak concentration at the high dose was about 20-fold higher than that of the low dose.

Results of an *in vitro* study describe dermal absorption of acrylamide. Marty and Vincent (1998) applied [¹⁴C]-labeled acrylamide (in an aqueous gel of 2% polyacrylamide) to biopsied human abdominal skin for 24 hours at acrylamide concentrations of 1.28 or 2 ppm. Approximately 28 and 21% of the applied doses, respectively, was recovered in the receptor fluid. Between 1.6 and 3.4% of applied doses was recovered

3. HEALTH EFFECTS

in dermis and epidermis. The authors estimated total absorption of acrylamide to be 33.2 and 26.7% at low and high concentration, respectively, based on radioactivity recovered from the receptor phase, epidermis, and dermis.

3.4.2 Distribution

Results from several animal studies indicate that, following absorption, radioactivity from radiolabeled acrylamide is widely distributed with no specific accumulation in any tissues other than red blood cells (Barber et al. 2001a; Crofton et al. 1996; Edwards 1975; Hashimoto and Aldridge 1970; Ikeda et al. 1985; Kadry et al. 1999; Marlowe et al. 1986; Miller et al. 1982; Ramsey et al. 1984) and late-staged spermatids (Sega et al. 1989). Results of intravenous injection studies in pregnant animals indicate that acrylamide and/or its metabolites readily cross the placenta and are distributed within the developing fetus in a manner similar to that of the pregnant mother (Ikeda et al. 1983, 1985; Marlowe et al. 1986). In a recent study designed to assess hemoglobin adduct levels of acrylamide in blood samples of pregnant mothers and umbilical cord blood of neonates, the concentration in umbilical cord blood was approximately 50% of that found in the blood of the mother, indicating that acrylamide readily passes from mother to developing fetus (Schettgen et al. 2004a).

3.4.2.1 Inhalation Exposure

Immediately following a 6-hour inhalation exposure of male F344 rats to 3 ppm of [¹⁴C]/[¹³C]-acrylamide vapor, the rank of acrylamide equivalent concentrations was: blood cells (~7 µg/g) > testes, skin, liver, and kidneys (~6 µg/g) > brain, spleen, lung, and epididymis (~4 µg/g) (Sumner et al. 2003). Acrylamide equivalent concentrations in similarly-treated male B6C3F1 mice were: testes (~14 µg/g) > skin and liver (~11 µg/g) > epididymis (~8 µg/g) > brain (~7 µg/g) > lung and blood (~6 µg/g) > fat (~5 µg/g) (Sumner et al. 2003).

3.4.2.2 Oral Exposure

Following 13 daily oral doses of [1,3-¹⁴C]-labeled acrylamide at 0.05 or 30 mg/kg/day, tissue concentrations of acrylamide in male F344 rats were similar among tissues with the exception of red blood cells, which showed higher concentrations, presumably due to the formation of acrylamideVal and/or glycidamideVal hemoglobin adducts (Ramsey et al. 1984). Mean concentrations (µg equivalents [¹⁴C]-acrylamide per gram of tissue) at the high dose were 383.7 µg/g in red blood cells, 87.74 µg/g in liver, 70.43 µg/g in kidneys, 70.60 µg/g in epididymis, 67.14 µg/g in testis, 54.00 µg/g in sciatic nerve,

3. HEALTH EFFECTS

53.52 µg/g in brain, 47.56 µg/g in carcass, 39.11 µg/g in skin, and 16.45 µg/g in plasma. At the low dose, the mean concentration in red blood cells was 1.26 µg/g (approximately 61% of the total recovered dose) compared with a range of 0.07–0.13 µg/g in other tissues.

In Sprague-Dawley rats given single 50 mg/kg oral doses of [¹⁴C]-labeled acrylamide, tissue concentrations of radioactivity at 28 and 144 hours postadministration were indicative of wide distribution of acrylamide metabolites among tissues with no evidence for accumulation in toxicity targets (Kadry et al. 1999). These results suggest the binding of acrylamide in the absence of its accumulation in erythrocytes or neural tissue. At 28 hours, brain, thyroid, testes, adrenal, pancreas, thymus, liver, kidney, heart, and spleen showed a narrow range of mean concentrations, approximately 0.05–0.10% of the initial dose/g. Higher concentrations were noted in the skin, bone marrow, stomach, and lung (0.15–0.18% of the initial dose/g); only the gastric contents showed a markedly higher concentration (1.37% of the initial dose/g). At 144 hours after administration, tissue concentrations were uniformly low for tissues including the gastric contents, ranging from 0.01 to 0.05% of the initial dose/g, with the exception of skin, bone marrow, and lung, which exhibited mean concentrations of 0.06, 0.08, and 0.19% of the initial dose/g, respectively.

3.4.2.3 Dermal Exposure

Following 24-hour dermal exposure of male F344 rats to 162 mg/kg [¹⁴C]-labeled acrylamide, the highest concentration of acrylamide equivalents (excluding skin at the application site) was found in blood cells (71 µg/g), followed by skin at nondosing sites (~28 µg/g); liver, spleen, testes, and kidneys (~21 µg/g); lungs, thymus, brain, and epididymis (~14 µg/g); and fat (<4 µg/g) (Sumner et al. 2003).

3.4.3 Metabolism

Results from rat and mouse studies indicate that acrylamide is rapidly metabolized and excreted predominantly in the urine as metabolites (Dixit et al. 1982; Edwards 1975; Miller et al. 1982; Ramsey et al. 1984; Sumner et al. 1992, 1999, 2003; Twaddle et al. 2004). Figure 3-3 depicts a metabolic scheme for acrylamide adapted from reports of Calleman (1996), IARC (1994), and Sumner et al. (1992, 1999). According to the metabolic scheme, acrylamide reacts readily with glutathione to form a glutathione conjugate, which is further metabolized to N-acetyl-S-(3-amino-3-oxopropyl)cysteine or S-(3-amino-3-oxopropyl)cysteine. Another initial step, catalyzed by CYP2E1, involves oxidation of acrylamide to the epoxide derivative, glycidamide. Glycidamide can react with glutathione to form conjugates that are


3. HEALTH EFFECTS

further metabolized to N-acetyl-S-(3-amino-2-hydroxy-3-oxopropyl)cysteine or N-acetyl-S-(1-carbamoyl-2-hydroxyethyl)cysteine. Glycidamide may also undergo hydrolysis, perhaps catalyzed by epoxide hydrolases, leading to the formation of 2,3-dihydroxypropionamide and 2,3-dihydroxypropionic acid.

Both acrylamide and glycidamide react with nucleophilic sites in macromolecules (including hemoglobin and DNA) in Michael-type additions (Bergmark et al. 1991, 1993; Segerbäck et al. 1995; Solomon et al. 1985). In rats, the binding index of acrylamide to cysteine in hemoglobin was extremely high (6,400 pmol/g Hb/ μ mol acrylamide/kg); the binding index of glycidamide to cysteine was 1,820 pmol/g Hb/ μ mol glycidamide/kg. The lower binding index for glycidamide may be the result of lower reactivity toward hemoglobin-cysteine and shorter half-life in blood. Rate constants of 0.0054 and 0.021 $M^{-1}s^{-1}$ for reactions of acrylamide with human serum albumin and glutathione, respectively, were reported by Tong et al. (2004), suggesting that these reactions account for most of the elimination of absorbed acrylamide. Doerge et al. (2005a) measured DNA adducts following a single intraperitoneal administration of acrylamide to adult B6C3F1 mice and F344 rats at 50 mg/kg. Glycidamide-derived DNA adducts of adenine and guanine formed in all tissues examined, including target tissues identified in rodent carcinogenicity bioassays and nontarget tissues. Dosing rats and mice with an equimolar amount of glycidamide (61 mg/kg) typically produced higher levels of DNA adducts than those observed from the acrylamide dose. Gamboa da Costa et al. (2003) measured DNA adduct formation in selected tissues of adult and whole body DNA of 3-day-old neonatal mice treated with acrylamide and glycidamide. In adult mice, DNA adduct formation was observed in liver, lung, and kidney; glycidamide treatment produced modestly higher adduct levels than acrylamide treatment. However, glycidamide-treated neonates exhibited 5–7-fold higher whole-body DNA adduct levels than the acrylamide-treated neonates.

Results from studies of CYP2E1 null and wild-type mice demonstrate the importance of CYP2E1 in catalyzing the oxidative formation of glycidamide from acrylamide. Following oral administration of single 50 mg/kg doses of [1,2,3- ^{13}C]-acrylamide to wild-type mice, considerable amounts of metabolites derived from glycidamide were found in the 24-hour urine (Sumner et al. 1999). Approximately 22% of the excreted metabolites were derived from glutathione conjugation with glycidamide (N-acetyl-S-[3-amino-2-hydroxy-3-oxopropyl]cysteine and N-acetyl-S-[1-carbamoyl-2-hydroxyethyl]cysteine) and 28% were derived from glycidamide and its hydrolysis products (2,3-dihydroxypropionamide and 2,3-dihydroxypropionic acid). In

3. HEALTH EFFECTS

Figure 3-3. Metabolic Scheme for Acrylamide and Its Metabolite Glycidamide*

*Processes involving several steps are represented with broken arrows.

GSH = reduced glutathione; Hb = hemoglobin; N-AcCys = N-acetylcysteine

Sources: Calleman 1996; Fennell et al. 2006; IARC 1994a; Sumner et al. 1992, 1999

3. HEALTH EFFECTS

contrast, no evidence was found for the formation of these metabolites in the 24-hour urine of CYP2E1 null mice or wild-type mice treated with the CYP2E1 inhibitor, aminobenzotriazole. The wild-type and CYP2E1-null mice excreted a similar percentage of the administered dose in the urine within 24 hours (about 30%), suggesting that the CYP2E1-null mice compensated for the CYP2E1 deficiency by metabolizing more of the administered acrylamide via direct conjugation with glutathione.

Ghanayem et al. (2005c) investigated the formation of hemoglobin adducts of acrylamide and glycidamide and DNA adducts of glycidamide in wild-type and CYP2E1-null mice following the administration of a single 50 mg/kg dose of acrylamide via intraperitoneal injection. At 6 hours posttreatment, mean plasma levels of acrylamide and glycidamide were 0.84 and 33.0 μM , respectively, in the wild-type mice and 115 and 1.7 μM , respectively, in the CYP2E1-null mice. Levels of hemoglobin adducts of acrylamide were approximately 2-fold higher in the CYP2E1-null mice compared to the wild-type mice. Although only traces of DNA adducts of glycidamide were seen in the CYP2E1-null mice, levels were 52–66 times higher in the wild type mice. These results demonstrate the importance of CYP2E1 in the epoxidation of acrylamide to glycidamide and the formation of hemoglobin adducts and glycidamide-DNA adducts.

N-Acetyl-S-(3-amino-3-oxopropyl)cysteine has been identified as the major urinary metabolite of acrylamide in male F344 rats exposed to oral doses of 1–100 mg/kg [2,3- ^{14}C]-labeled acrylamide (Miller et al. 1982) and in male F344 rats and B6C3F1 mice exposed to oral doses of 50 mg/kg [1,2,3- ^{13}C]-acrylamide (Sumner et al. 1992).

Following oral administration of [1,2,3- ^{13}C]-acrylamide (50 mg/kg) to rats and mice, glycidamide and glycidamide-derived metabolites accounted for about 33% (rats) and 59% (mice) of the total metabolites excreted in the urine within 24 hours, indicating that acrylamide is transformed to glycidamide to a greater extent in mice than in rats (Sumner et al. 1992). Similar results were reported in a study of metabolites in urine collected for 24 hours after 6-hour inhalation exposure (nose only) of rats and mice to 3 ppm [^{14}C]/[^{13}C]-acrylamide where glycidamide and glycidamide-derived metabolites accounted for 36% (rats) and 73% (mice) of total metabolites excreted in the urine within 24 hours (Sumner et al. 2003). Following dermal application of 138 mg/kg [1,2,3- ^{13}C]-acrylamide, male F344 rats excreted approximately 2% of the applied dose in the 24-hour urine as GSH-acrylamide derived metabolites (~50% of the urinary metabolites), glycidamide (~17%), and GSH-glycidamide derived metabolites (~31%) (Sumner et al. 2003).

3. HEALTH EFFECTS

Figure 3-3 does not include a possible minor pathway in which CO₂ may be released from the hydrolysis products of glycidamide, because of conflicting results from several studies. Following intravenous administration of 100 mg/kg [1-¹⁴C]-labeled acrylamide to male albino Porton rats, about 6% of the injected dose of radioactivity was exhaled as CO₂ in 8 hours (Hashimoto and Aldridge 1970), but following administration of [2,3-¹⁴C]-labeled acrylamide to male Fischer 344 rats, no radioactivity was detected in exhaled breath (Miller et al. 1982). Sumner et al. (1992) hypothesized that these results may be consistent with the existence of a minor pathway involving metabolism of 2,3-dihydroxypropionamide to glycerate and hydroxypyruvate with the subsequent release of CO₂ and production of glycoaldehyde, but they did not detect two-carbon metabolites in urine of mice exposed to [1,2,3-¹³C]-acrylamide. In other experiments, no exhaled ¹⁴CO₂ was detected following 50 mg/kg oral dosing of [1-¹⁴C]-labeled acrylamide to male Sprague-Dawley rats (Kadry et al. 1999), whereas 3–4% of intravenously injected [1,3-¹⁴C]-acrylamide (2 or 100 mg/kg) was detected as ¹⁴CO₂ in exhaled breath of male Fischer 344 rats (Ramsey et al. 1984).

Species differences in acrylamide metabolism are apparent. Twaddle et al. (2004) administered acrylamide at approximately 50 mg/kg via gavage to adult male and female B6C3F1 mice. Serum concentrations of acrylamide and glycidamide were taken at 0.5, 1, 2, 4, and 8 hours postdosing. Livers were removed from control and acrylamide-treated mice and analyzed for glycidamide-derived DNA adducts. The results indicated no systematic sex differences in acrylamide or glycidamide serum levels at each time point. Twaddle et al. (2004) estimated an acrylamide half-life of elimination from plasma at 0.73 hours. This value in mice can be compared to an estimate of 2 hours in F344 rats following a subchronic oral administration of 2.8 mM acrylamide in drinking water for 34 days or subacute intraperitoneal doses at 50 mg/kg/day for 11 days (Barber et al. 2001). Miller et al. (1982) estimated a 1.7-hour half-life for acrylamide in rat blood following a 10 mg/kg intravenous dose. Twaddle et al. (2004) reported an elimination half-life of 1.9 hours for the acrylamide-treated mice, which is identical to that measured by Barber et al. (2001) in rats. Barber et al. (2001) also reported a glycidamide/acrylamide-areas under the curve (AUC) ratio of 0.18 for Sprague-Dawley rats treated with 20 mg/kg acrylamide by gavage. This contrasts to the observation of equal AUCs for glycidamide and acrylamide in B6C3F1 mice (Twaddle et al. 2004). Since rats and mice had a comparable glycidamide elimination half-life, this approximately 5-fold difference in internal exposure to glycidamide for mice compared with rats is considered to be the result of an increased rate of glycidamide formation in the mouse.

3. HEALTH EFFECTS

The metabolism of acrylamide has also been investigated in controlled human studies (Boettcher et al. 2006a; Fennell et al. 2005b, 2006; Fuhr et al. 2006).

Fennell and coworkers evaluated metabolism and hemoglobin adduct formation (Fennell et al. 2005b) and kinetics of elimination of urinary metabolites (Fennell et al. 2006) following oral and dermal administration of [1,2,3-¹³C₃]-acrylamide and/or [2,3-¹⁴C]-acrylamide to 24 adult male volunteers.

Metabolism of the administered acrylamide was investigated by ¹³C nuclear magnetic resonance (NMR) spectroscopy (Fennell et al. 2005b) and by liquid chromatography-tandem mass spectroscopy (Fennell et al. 2006). Urinary metabolites accounted for approximately 40% of a 3 mg/kg oral dose of acrylamide and 4.5% of a 3 mg/kg/day repeated (3 consecutive days) dermal dose (Fennell et al. 2006).

Approximately 86% of the recovered urinary metabolites were derived from glutathione conjugation and excreted as N-acetyl-S-(3-amino-3-oxopropyl)cysteine and its S-oxide. Glycidamide, glyceramide (2,3-dihydroxypropionamide), and low levels of N-acetyl-S-(3-amino-2-hydroxy-3-oxopropyl)cysteine were also detected in urine (Fennell et al. 2005b). On oral administration, a linear dose response was observed for acrylamideVal and glycidamideVal in hemoglobin. The main pathway of metabolism in humans was via direct glutathione conjugation, forming N-acetyl-S-(3-amino-3-oxopropyl)cysteine and its S-oxide, which has not been reported previously (Fennell et al. 2006). Epoxidation to glycidamide was the other important pathway, with glyceramide formed as a major metabolite in humans.

Glycidamide was detected in low amounts. The glutathione conjugation of glycidamide, which is a major pathway in rodents, appeared to occur at very low levels in humans. Metabolism via glycidamide in humans was approximately 12% of the total urinary metabolites. This is considerably lower than the amount of glycidamide-derived metabolites reported for oral administration of acrylamide in rats (28% at 50 mg/kg, [Sumner et al. 2003]) and in mice (59% at 50 mg/kg [Sumner et al. 1992]). It should be noted that glyceramide has only been quantitated in human urine by NMR spectroscopy in combination with administration of [1,2,3-¹³C₃]-acrylamide.

The Fennell et al. (2005b) study also provided data on the amount of hemoglobin adducts derived from acrylamide and glycidamide following administration of a defined dose of acrylamide to adult male volunteers. Both acrylamideVal and glycidamideVal increased linearly with increasing dose of acrylamide administered orally, suggesting that saturation of the conversion of acrylamide to glycidamide is not reached in the dosing range of 0.5–3.0 mg/kg. The ratio of glycidamideVal:acrylamideVal produced by administration of acrylamide was similar to the ratio of the background adducts prior to exposure. Compared with the equivalent oral administration in rats (3 mg/kg), the ratio of glycidamideVal:acrylamideVal was lower in humans (0.44 compared to 0.84 for the rat), and the absolute

3. HEALTH EFFECTS

amounts of acrylamideVal and glycidamideVal formed were approximately 2.7- and 1.4-fold higher, respectively, in humans compared to rats.

Fennell et al. (2005b) calculated the expected amount of adduct that would accumulate in men from continuous exposure based on the amount of adduct formed/day of exposure, and from the lifespan of the erythrocyte. Oral intake of 1 µg/kg acrylamide/day (1.05 fmol acrylamideVal/mg globin/day) for the lifespan of the erythrocyte (120 days) was estimated to result in the accumulation of adducts to 63 fmol/mg globin. Daily dermal exposure to 1 µg/kg acrylamide (0.18 fmol acrylamideVal/mg globin/day) for the lifespan of the erythrocyte (120 days) would result in the accumulation of adducts to 10.8 fmol acrylamideVal/mg globin. With workplace exposure of 5 days/week, this would decrease to approximately 7.8 fmol acrylamideVal/mg globin.

Boettcher et al. (2006a) investigated the human metabolism of acrylamide to AAMA and GAMA in a healthy male volunteer who received a single dose of about 1 mg deuterium-labeled (d(3)) acrylamide, representing 13 µg/kg body weight, in drinking water. Urine samples before dosing and within 46 hours after the dose were analyzed for d(3)-AAMA and d(3)-GAMA using liquid chromatography-mass spectrometry (LC-MS). Total recovery in urine after 24 hours was about 51% as the sum of AAMA and GAMA and was similar to recoveries in rats (53–66%) given a gavage dose of 0.1 mg/kg (Doerge et al. 2007). After 2 days, AAMA accounted for 52% of the total acrylamide dose, and was the major metabolite of acrylamide in humans. GAMA accounted for 5%, and appeared as a minor metabolite of acrylamide. A urinary ratio of 0.1 was observed for GAMA/AAMA compared to previously reported values of 0.2 for rats and 0.5 for mice (Doerge et al. 2005b, 2005c). The authors concluded that the metabolic fate of acrylamide in humans was more similar to that in rats than in mice as previously demonstrated in terms of hemoglobin adducts.

Fuhr et al. (2006) evaluated the urinary levels of acrylamide, AAMA, glycidamide, and GAMA (using LC-MS) in six young healthy volunteers after the consumption of a meal containing 0.94 mg of acrylamide. Urine was collected up to 72 hours thereafter. No glycidamide was found. Unchanged acrylamide, AAMA, and GAMA accounted for urinary excretion of approximately 4.5, 50, and 6% of the dose, respectively. Conjugation with glutathione exceeded the formation of the reactive metabolite glycidamide. The data suggest 2- and 4-fold lower relative internal exposures for glycidamide from dietary acrylamide in humans compared with rats and mice, respectively.

3. HEALTH EFFECTS

Hemoglobin adducts of acrylamide and glycidamide and urinary metabolites have been used as biomarkers of exposure to acrylamide (Bergmark 1997; Bergmark et al. 1993; Boettcher et al. 2005; Calleman et al. 1994). Refer to Section 3.8. for detailed information regarding biomarkers of exposure to acrylamide.

3.4.4 Elimination and Excretion

Available human and animal data indicate that urinary excretion of acrylamide metabolites is the primary route of elimination of absorbed acrylamide (Barber et al. 2001a; Boettcher et al. 2006a; Doerge et al. 2007; Fennell et al. 2005b, 2006; Fuhr et al. 2006; Hashimoto and Aldridge 1970; Kadry et al. 1999; Miller et al. 1982; Ramsey et al. 1984; Sumner et al. 1992, 1999, 2003).

3.4.4.1 Inhalation Exposure

No human data were located regarding excretion and elimination following inhalation exposure to acrylamide. Sumner et al. (2003) exposed male F344 rats and B6C3F1 mice nose-only to a mixture of [1,2,3-¹³C]-acrylamide (90%) and [2,3-¹⁴C]-acrylamide (10%) vapors at an average analytical concentration of 1.17 ppm for 6 hours. During a 24-hour postexposure period, the distribution of the inhaled dose in the rats was: 31% in the urine, 56% retained in the body, 3% in the feces, and 2% as exhaled organic volatiles and ¹⁴CO₂. The distribution of the inhaled dose in the mice was: 27% in the urine, 46% in tissues, 5% in feces, 2% as organic volatiles, and 1% as ¹⁴CO₂.

3.4.4.2 Oral Exposure

Approximately 34% of an orally-administered 3 mg/kg dose of acrylamide to adult male volunteers was recovered in the total urinary metabolites within 24 hours of administration (Fennell et al. 2005b). In a male volunteer who received a single dose of about 1 mg deuterium-labeled acrylamide (representing 13 µg/kg body weight) in drinking water, elimination via urinary AAMA and GAMA followed a two-phase pattern (Boettcher et al. 2006a). Elimination half-lives of both AAMA and GAMA were estimated to be approximately 3.5 hours for the first phase and >10 hours up to few days for the second phase. After 2 days, urinary AAMA and GAMA accounted for 52 and 5%, respectively, of the total acrylamide dose. The ratio GAMA/AAMA was approximately 0.2.

Fuhr et al. (2006) measured acrylamide and metabolite levels in a 72-hour urine collection from six young healthy volunteers after the consumption of a meal containing 0.94 mg of acrylamide. Unchanged

3. HEALTH EFFECTS

acrylamide, AAMA, and GAMA accounted for urinary excretion of approximately 4, 50, and 10%, respectively, of the administered dose; glycidamide was not detected. Estimated elimination half-lives for the unchanged acrylamide, AAMA, and GAMA were 2.4, 17.4, and 25.1 hours, respectively.

Heudorf et al. (2009) reported median levels of 36 µg AAMA/L and 13 µg GAMA/L in the urine of 110 children (63 boys and 47 girls; 5–6 years of age). Children who reported higher regular consumption of French fries had significantly higher urinary levels of acrylamide metabolites. Based on the urinary levels of AAMA and GAMA, mean estimated acrylamide dietary intakes were 1.13 µg/kg/day (creatinine excretion based model) and 0.81 µg/kg/day (volume based model). The ratio GAMA/AAMA was approximately 0.4, which is 2-fold higher than that observed by Boettcher et al. (2006a) in adults. Based on this finding, Heudorf et al. (2009) suggest that acrylamide may undergo oxidative metabolism to a greater extent in children than adults.

In male Fischer 344 rats given oral (1, 10, or 100 mg/kg) doses of [2,3-¹⁴C]-acrylamide, about 60 and 70% of the administered radioactivity was excreted in urine collected within 24 hours and 7 days, respectively (Miller et al. 1982). Less than 2% of radioactivity in the urine was accounted for by acrylamide. Elimination of radioactivity from tissues was biphasic with half-lives of about 5 and 8 hours, respectively. The elimination time course of parent compound from tissues exhibited single phase exponential elimination with a half-life of about 2 hours. Fecal excretion accounted for 4.8 and 6% of administered radioactivity at 24 hours and 7 days, respectively. Bile-duct cannulated rats given single intravenous 10 mg/kg doses of [2,3-¹⁴C]-acrylamide excreted about 15% of the administered radioactivity as metabolites within about 6 hours; <1% of radioactivity in the bile was in the form of acrylamide (Miller et al. 1982). These results are consistent with the existence of enterohepatic circulation of metabolites.

Doerge et al. (2007) administered acrylamide to male and female rats and mice at 0.1 mg/kg by oral gavage or via the diet and measured the percentages of parent compound and metabolites in the 24-hour urine. For rats and mice combined, the percentage of total dose excreted as the sum of acrylamide and its metabolites (glycidamide, AAMA, and GAMA) averaged 49%. Excretion of glycidamide and glycidamide-derived species was typically greater in mice than rats.

No radiolabeled CO₂ was captured from rats given [2,3-¹⁴C]-acrylamide orally (Kadry et al. 1999; Miller et al. 1982). However, intravenous injection of rats with a 100 mg/kg dose of [1-¹⁴C]-labeled acrylamide (Hashimoto and Aldridge 1970) or a 2 mg/kg dose of [1,3-¹⁴C]-labeled acrylamide (Ramsey et al. 1984)

3. HEALTH EFFECTS

resulted in the capture of about 6 and 4%, respectively, of the radioactivity as CO₂ in the expired air during the subsequent 6–8 hours.

3.4.4.3 Dermal Exposure

Fennell et al. (2006) exposed 24 adult male volunteers to dermal applications of [1,2,3-¹³C₃]-acrylamide under controlled conditions. Acrylamide was administered as three daily 24-hour occluded dermal doses of 3 mg/kg. Approximately 4.5% of the administered dose (12.35% of the absorbed dose) was recovered as urinary metabolites through day 4. Urinary recovery included ¹³C₃-acrylamide (~4% of total urinary recovery) and metabolites ¹³C₃-cysteine-S-propionamide (¹³C₃-CP; ~0.8%), ¹³C₃-N-acetyl cysteine-S-propionamide (¹³C₃-NACP; ~70%), ¹³C₃-NACP sulfoxide (~22.5%), ¹³C₃-GAMA3 (~2.6%), and ¹³C₃-GAMA2 (~0.4%). Glycidamide was not detected in the majority of the urine samples. ¹³C₃-Acrylamide and ¹³C₃-CP were detected in 2–4 hour urine samples collected on the first day. ¹³C₃-NACP and its sulfoxide were first detectable in the urine between 4 and 8 hours posttreatment. The mercapturic acids of glycidamide (¹³C₃-GAMA2 and ¹³C₃-GAMA2) were first detectable in the urine between 8 and 16 hours posttreatment. The metabolite, glycidamide, was not detectable in the majority of the urine samples collected over the 4-day period.

Sumner et al. (2003) exposed male F344 rats to [2,3-¹⁴C]-labeled acrylamide under occluded dermal conditions for 24 hours. During a 24-hour postexposure period, approximately 8% of the applied dose (36% of the absorbed dose) was excreted in the urine, 3% as volatiles and ¹⁴CO₂ in exhaled air, <1% in feces, and 53% remained in tissues.

3.4.5 Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models

Physiologically based pharmacokinetic (PBPK) models use mathematical descriptions of the uptake and disposition of chemical substances to quantitatively describe the relationships among critical biological processes (Krishnan et al. 1994). PBPK models are also called biologically based tissue dosimetry models. PBPK models are increasingly used in risk assessments, primarily to predict the concentration of potentially toxic moieties of a chemical that will be delivered to any given target tissue following various combinations of route, dose level, and test species (Clewell and Andersen 1985). Physiologically based pharmacodynamic (PBPD) models use mathematical descriptions of the dose-response function to quantitatively describe the relationship between target tissue dose and toxic end points.

3. HEALTH EFFECTS

PBPK/PD models refine our understanding of complex quantitative dose behaviors by helping to delineate and characterize the relationships between: (1) the external/exposure concentration and target tissue dose of the toxic moiety, and (2) the target tissue dose and observed responses (Andersen and Krishnan 1994; Andersen et al. 1987). These models are biologically and mechanistically based and can be used to extrapolate the pharmacokinetic behavior of chemical substances from high to low dose, from route to route, between species, and between subpopulations within a species. The biological basis of PBPK models results in more meaningful extrapolations than those generated with the more conventional use of uncertainty factors.


The PBPK model for a chemical substance is developed in four interconnected steps: (1) model representation, (2) model parameterization, (3) model simulation, and (4) model validation (Krishnan and Andersen 1994). In the early 1990s, validated PBPK models were developed for a number of toxicologically important chemical substances, both volatile and nonvolatile (Krishnan and Andersen 1994; Leung 1993). PBPK models for a particular substance require estimates of the chemical substance-specific physicochemical parameters, and species-specific physiological and biological parameters. The numerical estimates of these model parameters are incorporated within a set of differential and algebraic equations that describe the pharmacokinetic processes. Solving these differential and algebraic equations provides the predictions of tissue dose. Computers then provide process simulations based on these solutions.

The structure and mathematical expressions used in PBPK models significantly simplify the true complexities of biological systems. If the uptake and disposition of the chemical substance(s) are adequately described, however, this simplification is desirable because data are often unavailable for many biological processes. A simplified scheme reduces the magnitude of cumulative uncertainty. The adequacy of the model is, therefore, of great importance, and model validation is essential to the use of PBPK models in risk assessment.

PBPK models improve the pharmacokinetic extrapolations used in risk assessments that identify the maximal (i.e., the safe) levels for human exposure to chemical substances (Andersen and Krishnan 1994). PBPK models provide a scientifically sound means to predict the target tissue dose of chemicals in humans who are exposed to environmental levels (for example, levels that might occur at hazardous waste sites) based on the results of studies where doses were higher or were administered in different species. Figure 3-4 shows a conceptualized representation of a PBPK model.

3. HEALTH EFFECTS

Figure 3-4. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance



Note: This is a conceptual representation of a physiologically based pharmacokinetic (PBPK) model for a hypothetical chemical substance. The chemical substance is shown to be absorbed via the skin, by inhalation, or by ingestion, metabolized in the liver, and excreted in the urine or by exhalation.

Source: adapted from Krishnan and Andersen 1994

3. HEALTH EFFECTS

If PBPK models for acrylamide exist, the overall results and individual models are discussed in this section in terms of their use in risk assessment, tissue dosimetry, and dose, route, and species extrapolations.

Physiologically based pharmacokinetic (PBPK) models for acrylamide are available (Calleman et al. 1992, 1993; Kirman et al. 2003; Walker et al. 2007; Young et al. 2007).

Calleman et al. (1992, 1993) Model

Calleman et al. (1992, 1993) developed a nonlinear dosimetric model to simultaneously determine hemoglobin adduct formation by acrylamide and its genotoxic metabolite, glycidamide, in rats. The model is not useful for human health risk assessment.


Kirman et al. (2003) Model

Kirman et al. (2003) developed a PBPK model (Figure 3-5) to predict the behavior of acrylamide and its epoxide metabolite, glycidamide, in the rat for intravenously-, intraperitoneally-, or orally-administered acrylamide. The model includes components for both acrylamide and glycidamide, each consisting of five compartments (arterial blood, venous blood, liver, lung, and all other tissues). The acrylamide component is linked to the glycidamide portion of the model via metabolism in the liver. Model parameters are listed in Table 3-9. Rat physiological parameters were selected from measured data (Brown et al. 1997; Kedderis et al. 1996). Partition coefficients for acrylamide were estimated based on an algorithm derived by Poulin and Krishnan (1995, 1996a, 1996b) using specific chemical properties. Partition coefficients for glycidamide were derived by application of a proportionality constant of 3.2 to the values calculated for acrylamide (Kedderis et al. 1996). Metabolism of acrylamide and glycidamide are represented only in the liver. Estimated values for metabolism and tissue binding were based on fitting of the model to metabolism and urinary elimination data from Miller et al. (1982), Ramsey et al. (1984), Raymer et al. (1993), and Sumner et al. (1992). Depletion and resynthesis of glutathione were included in the model structure (D'Souza et al. 1988).

The PBPK model of Kirman et al. (2003) was developed to predict the behavior of acrylamide and glycidamide in the rat. The model has only been partially validated in rats and is not useful for human risk assessment.

3. HEALTH EFFECTS

Figure 3-5. Schematic of the Kirman et al. (2003) Physiologically Based Pharmacokinetic (PBPK) Model for Acrylamide



EH = epoxide hydrolase; GST = glutathione *S*-transferase; i.p. = intraperitoneal injection; i.v. = intravenous injection;

Source: Kirman et al. 2003

3. HEALTH EFFECTS

Table 3-9. Original Model Parameter Values for Rats in the Kirman et al. (2003) Physiologically Based Toxicokinetic (PBTK) Model

Parameter group	Parameter	Symbol (units)	Value	Reference/source
Rat physiology	Body weight	BW (kg)	Study specific	Study specific
	Cardiac output	QCC (L/hour-kg)	14	Kedderis et al. 1996
	Alveolar ventilation	QPC (L/hour-kg)	14	Kedderis et al. 1996
	Liver blood flow	QLC (fraction QCC)	0.25	Kedderis et al. 1996
	Fraction arterial	FABC (fraction VB)	0.35	Kedderis et al. 1996
	Fraction venous	FVBC (fraction VB)	0.65	Kedderis et al. 1996
	Liver volume	VLC (fraction BW)	0.04	Brown et al. 1997
	Tissue volume	VTC (fraction BW)	0.87	Calculated (0.91-VLC)
	Tissue blood flow	QTC (fraction QCC)	0.75	Calculated (1-QLC)
	Volume blood	VBC (fraction BW)	0.06	Brown et al. 1997
	Hemoglobin concentration	HGB (g/L)	1.5	Kedderis et al. 1996
	Fraction blood cells	FBC (fraction VB)	0.44	Kedderis et al. 1996
	Fraction blood serum	FBS (fraction VB)	0.56	Kedderis et al. 1996
Absorption	Absorption rate from gastrointestinal tract (oral dose) or intraperitoneal cavity (intraperitoneal dose)	KA (/hour)	5	Model simulations fit to Miller et al. 1982; Ramsey et al. 1984; Raymer et al. 1993
	Infusion time (intravenous dose)	TINF (hour)	0.003	Model simulations fit to Miller et al. 1982; Ramsey et al. 1984; Raymer et al. 1993

3. HEALTH EFFECTS

Table 3-9. Original Model Parameter Values for Rats in the Kirman et al. (2003) Physiologically Based Toxicokinetic (PBTK) Model

Parameter group	Parameter	Symbol (units)	Value	Reference/source
Partition coefficients	Blood:air, AA	PB1 (unitless)	31,000,000	Estimated
	Liver:blood, AA	PL1 (unitless)	0.83	Poulin and Krishnan 1995, 1996a, 1996b
	Tissue:blood, AA	PL1	0.95	Poulin and Krishnan 1995, 1996a, 1996b
	Blood:air, GA	PB2 (unitless)	98,000,000	Poulin and Krishnan 1995, 1996a, 1996b
	Liver:blood, GA	PL2 (unitless)	2.7	Poulin and Krishnan 1995, 1996a, 1996b
	Tissue:blood, GA	PT2 (unitless)	3.0	Poulin and Krishnan 1995, 1996a, 1996b
Metabolism	Cytochrome P-450 oxidation rate, AA	VMAXC1 (mg/hour-kg)	1.6	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992
	Cytochrome P-450, Michaelis-Menten constant, AA	KMC1 (mg/L)	10	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992
	Epoxide hydrolase hydrolysis rate, GA	VMAXC2 (mg/hour-kg)	1.9	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992
	Epoxide hydrolase, Michaelis-Menten constant, GA	KMC2 (mg/L)	100	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992
	Reaction with glutathione, AA	KGSTC1 (L/mmol GSH-hr)	0.55	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992
	Reaction with glutathione, GA	KGSTC2 (L/mmol GSH-hour)	0.8	Model simulations fit to Miller et al. 1982; Raymer et al. 1993; Sumner et al. 1992

3. HEALTH EFFECTS

Table 3-9. Original Model Parameter Values for Rats in the Kirman et al. (2003) Physiologically Based Toxicokinetic (PBTK) Model

Parameter group	Parameter	Symbol (units)	Value	Reference/source
Tissue binding	Binding to hemoglobin, AA	KHGB1 (L/gHGB)	0.5	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to hemoglobin, GA	KHGB2 (L/gHGB-hour)	0.25	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to liver macromolecules, acrylamide	KFEEL1 (/hour)	0.2	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to liver macromolecules, GA	KFEEL2 (/hour)	0.1	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to tissue macromolecules, AA	KFEET1 (/hour)	0.08	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to tissue macromolecules, GA	KFEET1 (/hour)	0.04	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to blood macromolecules, AA	KFEEB1 (/hour)	0.01	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Binding to blood macromolecules, GA	KFEEB2 (/hour)	0.005	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
	Protein turnover	KPT (/hour)	0.008	Model simulations fit to Miller et al. 1982; Raymer et al. 1993
Glutathione	GSH production rate	KGSHP (mmol/hour)	0.025	D'Souza et al. 1988
	GSH loss rate	KGSHL (/hour)	0.35	D'Souza et al. 1988
	Initial GSH concentration in liver	GSHL0 (mmol/L)	7.0	D'Souza et al. 1988

AA = acrylamide; GA = glycidamide; GSH = glutathione

Source: Kirman et al. 2003

3. HEALTH EFFECTS

Young et al. (2007) Model


Young et al. (2007) produced a PBPK/toxicodynamic (TD) model to predict the behavior of acrylamide, glycidamide, and their respective glutathione conjugates in rats, mice, and humans (Figure 3-6). The model was developed using PostNatal, a windows-based program from the U.S. FDA's National Center for Toxicological Research (NCTR). The program controls up to four PBPK model units (depicted as PBPK 1–4 in Figure 3-6) under one shell with multiple input and output options. Each PBPK unit is comprised of 28 organ/tissue/fluid components maintained independently or connected through metabolic pathways. Pharmacodynamic components for liver glycidamide-DNA adducts and hemoglobin adducts with acrylamide and glycidamide are included in the model as well. Dose administration can be either acrylamide or glycidamide. All metabolism and urinary elimination are considered first order.

Physiological parameters were assigned values from within the PostNatal program based on animal species, gender, and total body weight. Serum acrylamide and glycidamide concentrations and urinary elimination levels for male and female rats and mice were simulated from intravenous and oral administration of 0.1 or 0.12 mg/kg glycidamide. Adduct formation and decay rates were determined from a 6-week exposure to 1 mg/kg acrylamide in the drinking water followed by 6 weeks of nontreatment.

The data used to calibrate the Young et al. (2007) model for rats and mice include acrylamide serum levels in rats following intraperitoneal administration (Raymer et al. 1993); plasma acrylamide and glycidamide levels, and acrylamide and glycidamide hemoglobin adduct levels following relatively high (50 mg/kg bw) repeat intraperitoneal dosing in rats for 11 days or 2.8 mM of acrylamide in drinking water for 47 days (Barber et al. 2001); urinary excretion profile and acrylamide and glycidamide hemoglobin adduct levels following dosing via intraperitoneal injection (50 mg/kg), gavage (50 mg/kg), dermal application (150 mg/kg), or inhalation (3 ppm for 6 hours) (Sumner et al. 2003); and serum and tissue levels of acrylamide and glycidamide, and liver glycidamide-DNA adduct data in rats and mice following relatively low-dose administration via intravenous injection (acrylamide and glycidamide at 0.1–0.12 mg/kg), gavage (acrylamide and glycidamide at 0.12 and 50 mg/kg), diet (~0.1 mg/kg over 30 minutes), and drinking water (~1 mg/kg acrylamide over 42 days) (Doerge et al. 2005a, 2005b, 2005c). The single and multiple oral data from Barber et al. (2001) were combined with the urinary elimination data of Sumner et al. (1992, 2003) and simulated with the model. The Raymer et al. (1993)

3. HEALTH EFFECTS

Figure 3-6. Schematic of the Young et al. (2007) Physiologically Based Pharmacokinetic/Toxicodynamic (PBPK/TD) Model For Acrylamide and Glycidamide*



*Block diagram for the metabolism of AA to GA and further metabolism of both to their glutathione conjugates. All pharmacokinetic and pharmacodynamic (PD) processes are first order.

3. HEALTH EFFECTS

data were also combined with the urinary elimination data of Sumner et al. (1992, 2003) and simulated in a similar manner. The NCTR tissue data (Doerge et al. 2005a, 2005b, 2005c) were used to develop partition coefficients. Only those tissues specifically analyzed for acrylamide or glycidamide were partitioned differently from the blood compartment. Values for the human parameters were calibrated against urinary excretion data (Fennell et al. 2005b; Fuhr et al. 2006) and hemoglobin adduct data from a dietary exposure (Boettcher et al. 2005).

Values for the metabolism and elimination of acrylamide or glycidamide, for acrylamide or glycidamide binding to hemoglobin, and for glycidamide-DNA adduct formation were derived by optimizing the fit of the simulation results to individual animal data. All rate constants for the metabolic and elimination processes, the binding and decay of acrylamide or glycidamide to hemoglobin, and the binding of glycidamide to liver macromolecule are represented as first order. Although Young et al. (2007) calibrated their model parameter values in a logical sequence against the data identified in the paper, a number of sensitive parameters were allowed to vary when fitting the individual animal data so as to optimize the model fit to each set of data. The authors evaluated the resulting differences among the model parameter values relative to gender and study conditions for insights into the toxicokinetics of acrylamide and glycidamide, and to assess the uncertainty in the model parameter values. Although there are statistically significant differences in some cases in the fitted model parameter values for basic physiological functions such as excretion of acrylamide-GSH conjugates in urine (which varies as much as 4–6 for model fits to different studies), the authors argued that the ranges of values are not exceedingly wide considering that different routes of administration for different chemicals are all being compared, and that there is very little difference for each metabolic rate constant when comparing across gender, dose, and route.

For the purpose of quantitative risk assessment, a PBPK model is generally developed to produce a single set of parameter values that fits at least the most relevant data for a particular application. Evaluating the importance of uncertainty in parameter values also depends upon the choice of the dose metric and its sensitivity to parameters of interest. For useful application of the PBPK model of Young et al. (2007) to human health risk, additional studies are needed to identify a single set of parameters, and to evaluate the sensitivity of various dose metrics to the parameters that are the most uncertain.

3. HEALTH EFFECTS

Walker et al. (2007) Model

Walker et al. (2007) described an adaptation of the PBPK model of Kirman et al. (2003) to account for: (1) hemoglobin adduct data for rats (Fennell et al. 2003) and (2) extrapolation to adult humans using human adduct data (Fennell et al. 2004, 2005b). The Walker et al. (2007) model incorporates available information regarding children's changing physiology and metabolic capacity for selected processes involved in acrylamide disposition (i.e., CYP2E1, glutathione conjugation, epoxide hydrolase). However, this model has not been calibrated or validated with respect to acrylamide dosimetry in children.

3.5 MECHANISMS OF ACTION**3.5.1 Pharmacokinetic Mechanisms**

Acrylamide is readily absorbed via all natural routes of exposure (inhalation, oral, dermal) and is distributed throughout the body via the blood. No studies were located regarding mechanisms for acrylamide absorption across lung, gut, or skin; processes may include passive and/or facilitative mechanisms. Acrylamide and its reactive metabolite, glycidamide, readily bind to hemoglobin, but do not accumulate appreciably in any tissues. Acrylamide metabolism is relatively rapid; both parent compound and the epoxide, glycidamide, appear to be involved in acrylamide toxicity. Animal data demonstrate the importance of CYP2E1 in acrylamide metabolism. Metabolism is assumed to take place primarily in the liver. Urinary excretion of conjugated acrylamide derivatives represents the major excretory pathway for acrylamide.

3.5.2 Mechanisms of Toxicity

Neurotoxic Effects. The neurotoxicity of acrylamide has been assessed since the 1950s in numerous animal studies; notable signs and symptoms include hindfoot splay, decreased grip strength, and increasingly impaired mobility with continued exposure (see Sections 3.2.1.4, 3.2.2.4, and 3.2.3.4 for detailed information regarding neurological effects in animals administered acrylamide by inhalation, oral, and dermal exposure routes, respectively). Observations of neurological effects in acrylamide-exposed humans include muscle weakness and other signs of functional impairments (Calleman et al. 1994; Hagmar et al. 2001; He et al. 1989). Early histopathologic investigations performed on acrylamide-intoxicated animals revealed degenerative effects in distal portions of large peripheral nerve fibers and associated myelin sheath (Spencer and Schaumberg 1974, 1977). The degeneration was observed to progress proximally in a process termed “dying back”. Early hypotheses to explain this “dying back”

3. HEALTH EFFECTS

effect involved damage to axonal neurofilaments, impairment of cellular metabolism, effects on axonal transport mechanisms, and disruption of axon cytoskeleton (Cavanaugh 1964; Harris et al. 1994; Harry 1992; Lapadula et al. 1989; LoPachin and Lehning 1994; Padilla et al. 1993; Pleasure et al. 1969; Sickles 1991; Spencer et al. 1979).

Specific mechanisms of acrylamide neurotoxicity have not been clearly elucidated. However, results of numerous studies designed to assess mechanisms of acrylamide-induced neurological effects have led to two major hypotheses: (1) acrylamide-induced disruption of fast axonal transport (Sickles et al. 2002a) and (2) acrylamide-induced disruption of nitric oxide signaling at nerve terminals (LoPachin and Barber 2006; LoPachin et al. 2008). In addition, recent work by Zhu et al. (2008) provides some support to a hypothetical mode of action whereby acrylamide exposure results in increases in reactive oxygen species, damage to cellular macromolecules, and subsequent degeneration of neural tissues.

Disruption of Fast Axonal Transport. The hypothesis that acrylamide induces peripheral neuropathy via disruption of fast axonal transport proposes the binding of acrylamide to kinesin, which leads to impairment of the fast axonal transport system responsible for the distal delivery of macromolecules. This would result in deficiencies in proteins responsible for maintaining axonal structure and function. The particular vulnerability of distal axons and nerve terminals is based on the large axonal volume and transport distance from cell body to distal regions. Kinesin, which plays an integral role in intracellular transport along microtubules, is inhibited by neurotoxic doses of acrylamide. Evidence was provided from results of a microtubule motility assay in which preincubation of purified kinesin with acrylamide produced a dose-dependent loss of numbers of microtubules moving over a bed of kinesin and less-steady locomotory activity (Sickles et al. 1996). These effects were proposed to result from covalent adduction through sulfhydryl alkylation by acrylamide. Comparable effects were produced by glycidamide.

Disruption of Nitric Oxide Signaling at Nerve Terminals. The hypothesis that acrylamide induces peripheral neuropathy via disruption of nitric oxide signaling at nerve terminals was proposed by LoPachin and coworkers (LoPachin and Barber 2006; LoPachin et al. 2008) based on critical review of available data for conjugated α,β -unsaturated carbonyl derivatives, a group of type-2 alkenes that includes acrylamide. The hypothesis is based on evidence that: (1) acrylamide inhibits neurotransmission in peripheral and central synapses and (2) soft electrophiles such as acrylamide form Michael-type adducts with soft nucleophilic sulfhydryl groups on protein cysteine residues.

3. HEALTH EFFECTS

According to the review of LoPachin and Barber (2006), the functional status of many synaptic processes is determined by proteins whose activity is regulated by the redox state of highly nucleophilic sulfhydryl groups within corresponding catalytic triads. Acrylamide (a soft electrophile) forms adducts with soft nucleophilic sulfhydryl groups on cysteine residues (reviewed in LoPachin and DeCaprio 2005). Results of early electrophysiological studies demonstrated acrylamide-induced neurotransmitter inhibition at peripheral and central synapses (reviewed in LoPachin et al. 2002, 2003). Results of subsequent mechanistic studies indicate that acrylamide causes decreases in neurotransmitter release, uptake, and storage (Barber and LoPachin 2004; LoPachin et al. 2004, 2006) and that these effects may be mediated by acrylamide-sulfhydryl adduction of specific cysteine residues on functionally critical proteins (Barber and LoPachin 2004). Because the cysteine-acrylamide adduction sites are also sites of nitric oxide nitrosylation (Jaffrey et al. 2001; Stamler et al. 2001), it has been suggested that acrylamide inhibits synaptic activity by disrupting nitric oxide signaling (LoPachin and Barber 2006). LoPachin and coworkers suggest that acrylamide itself, and not its major electrophilic metabolite glycidamide, is responsible for the neurotoxicity because glycidamide is a hard electrophile that forms adducts with hard nucleophiles (nitrogen, carbon, oxygen) consistent with glycidamide adducts on adenine and guanine bases.

Reactive Oxygen Species Hypothesis. Results of Zhu et al. (2008) provide some indication that acrylamide-induced neurotoxicity may involve enhancement of lipid peroxidation and decreased antioxidative capacity, depletion of neural glutathione levels and antioxidant enzyme activities, leading to increased levels of reactive oxygen species, damage to cellular macromolecules, and subsequent degeneration of neural tissues. Time-dependent decreased glutathione levels and anti-reactive oxygen species activities and increased malondialdehyde levels in sciatic nerve preparations were highly correlated with changes in electrophysiological indices of acrylamide-induced neurotoxicity in rats receiving repeated intraperitoneally-injected doses of 40 mg/kg acrylamide.

Reproductive Effects. Mechanisms of acrylamide-induced reproductive toxicity are poorly understood. There is some indication that mutagenic effects on male germ cells may play a significant role (see Section 3.3 Genotoxic Effects). Available data provide suggestions that acrylamide-induced male dominant lethal mutations may involve clastogenic events from binding of acrylamide and/or glycidamide to spermatid protamines or spindle fiber proteins and/or direct alkylation of DNA by glycidamide (Adler et al. 2000; Perrault 2003; Sega et al. 1989; Tyl and Friedman 2003; Tyl et al. 2000b). Tyl and Friedman (2003) also suggested that adverse effects on mounting, sperm motility, and intromission could be related to distal axonopathy resulting from binding of acrylamide to motor proteins.

3. HEALTH EFFECTS

Results of one study using male CYP2E1-null and wild-type mice demonstrate the importance of metabolism in acrylamide-induced germ cell mutations (Ghanayem et al. 2005a). The male mice were administered acrylamide via intraperitoneal injection for 5 consecutive days at dose levels of 0, 12.5, 24, or 50 mg/kg/day and subsequently mated with untreated B6C3F1 female mice. Dose-related increased incidences of dominant lethal mutations and decreases in numbers of pregnant females and proportion of living fetuses were observed in females mated to wild-type male mice. No significant changes in any fertility parameters were seen in females mated to CYP2E1-null male mice. These results demonstrate the importance of CYP2E1-mediated epoxidation to acrylamide-induced germ cell mutations in male mice.

Genotoxic/Carcinogenic Effects. Specific mechanisms whereby acrylamide induces tumors in laboratory animals are not understood at present. However, the weight of evidence supports a mutagenic mode of action (Besaratnia and Pfeiffer 2005, 2007; Dearfield et al. 1995; Moore et al. 1987; Segerbäck et al. 1995). Evidence for a mutagenic mode of action includes findings that: (1) acrylamide is metabolized by CYP2E1 to DNA-reactive glycidamide; (2) acrylamide and glycidamide induce mutations in lymphocyte HPRT and liver cII cells; (3) DNA adducts of glycidamide have been detected in tissues of all relevant tumor targets of acrylamide-and glycidamide-exposed male and female rats and mice; (4) glycidamide is mutagenic to bacteria and male mouse germ cells and male and female mouse somatic cells *in vivo*; (5) acrylamide induces heritable translocations and specific locus mutations in germ cells of exposed male mice; (6) acrylamide induces clastogenic effects in mouse lymphoma assays; and (7) dominant lethal effects in rodents occur at subchronic oral exposure levels comparable to those associated with carcinogenic effects in chronically-exposed rats. These findings support a proposed mode of action whereby acrylamide is metabolized to the relatively long-lived epoxide, glycidamide, which reacts with proteins and DNA, causing mutations that persist in viable somatic cells, resulting in tumor formation. Ghanayem et al. (2005b) observed significant dose-related increases in micronucleated erythrocytes and DNA damage in somatic cells (leukocytes, liver, lung) of acrylamide-treated wild-type mice, but not in CYP2E1-null mice, indicating that genetic damage in somatic cells is dependent on metabolism of acrylamide by CYP2E1. Allen et al. (2005) investigated dose-response relationships from *in vivo* genotoxicity data (chromosomal damage, gene mutations, and recombinations in somatic and germ cells) for acrylamide using three different mathematical models according to end point; the investigation included benchmark dose calculations. The results were not consistent with a genotoxic mode of action for the thyroid tumors reported in 2-year cancer bioassays in rats (Friedman et al. 1995; Johnson et al. 1984, 1986). Based on these results, Allen et al. (2005) suggested that a nongenotoxic mode of action may be primarily responsible for acrylamide-induced thyroid tumors in rats.

3. HEALTH EFFECTS

Another hypothetical mode of action involves disruption of hormone levels or hormone signaling for acrylamide-induced tumors in hormonally-sensitive tissues including mammary gland and thyroid or tissues adjacent to hormonally sensitive tissue, such as tunica vaginalis mesothelium (DFG 2009, Dourson et al. 2008; Environ 2002; Haber et al. 2009; Klaunig 2008; Shipp et al. 2006). In support of this hypothetical mode of action, American Cyanamid Company (1991) reported decreases in serum testosterone and thyroid hormone (T₃ and T₄) levels in rats following oral exposure to acrylamide. The induction of cellular proliferation is yet another proposed mode of action for acrylamide carcinogenicity in selected target tissues, although limited supporting data are available. Lafferty et al. (2004) demonstrated that tumorigenic doses of acrylamide administered to male rats induced cell proliferation in thyroid, tunica vaginalis, and adrenal medulla (tumor target tissues), but not in liver or adrenal cortex (nontarget tissues).

3.5.3 Animal-to-Human Extrapolations

Available data from rats and mice indicate that acrylamide is transformed to glycidamide to a greater extent in mice than rats. Following oral administration of radiolabeled acrylamide (50 mg/kg), glycidamide and glycidamide-derived metabolites accounted for about 33% (rats) and 59% (mice) of the total metabolites excreted in the urine within 24 hours (Sumner et al. 1992). Similar results were reported in a study of metabolites in urine collected for 24 hours after 6-hour inhalation exposure (nose only) of rats and mice to 3 ppm acrylamide where glycidamide and glycidamide-derived metabolites accounted for 36 % (rats) and 73 % (mice) of total metabolites excreted in the urine within 24 hours (Sumner et al. 2003).

Available PBPK models for acrylamide (Kirman et al. 2003; Walker et al. 2009; Young et al. 2007) have not been adequately calibrated and validated for useful extrapolation from animals to humans.

3.6 TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS

Recently, attention has focused on the potential hazardous effects of certain chemicals on the endocrine system because of the ability of these chemicals to mimic or block endogenous hormones. Chemicals with this type of activity are most commonly referred to as *endocrine disruptors*. However, appropriate terminology to describe such effects remains controversial. The terminology *endocrine disruptors*, initially used by Thomas and Colborn (1992), was also used in 1996 when Congress mandated the EPA to develop a screening program for "...certain substances [which] may have an effect produced by a

3. HEALTH EFFECTS

naturally occurring estrogen, or other such endocrine effect[s]...”. To meet this mandate, EPA convened a panel called the Endocrine Disruptors Screening and Testing Advisory Committee (EDSTAC), and in 1998, the EDSTAC completed its deliberations and made recommendations to EPA concerning *endocrine disruptors*. In 1999, the National Academy of Sciences released a report that referred to these same types of chemicals as *hormonally active agents*. The terminology *endocrine modulators* has also been used to convey the fact that effects caused by such chemicals may not necessarily be adverse. Many scientists agree that chemicals with the ability to disrupt or modulate the endocrine system are a potential threat to the health of humans, aquatic animals, and wildlife. However, others think that endocrine-active chemicals do not pose a significant health risk, particularly in view of the fact that hormone mimics exist in the natural environment. Examples of natural hormone mimics are the isoflavonoid phytoestrogens (Adlercreutz 1995; Livingston 1978; Mayr et al. 1992). These chemicals are derived from plants and are similar in structure and action to endogenous estrogen. Although the public health significance and descriptive terminology of substances capable of affecting the endocrine system remains controversial, scientists agree that these chemicals may affect the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body responsible for maintaining homeostasis, reproduction, development, and/or behavior (EPA 1997). Stated differently, such compounds may cause toxicities that are mediated through the neuroendocrine axis. As a result, these chemicals may play a role in altering, for example, metabolic, sexual, immune, and neurobehavioral function. Such chemicals are also thought to be involved in inducing breast, testicular, and prostate cancers, as well as endometriosis (Berger 1994; Giwercman et al. 1993; Hoel et al. 1992).

No studies were located regarding endocrine disruption in humans after exposure to acrylamide.

One repeated-dose oral study in rats reported acrylamide-induced decreases in serum testosterone levels, thyroid hormone (T₃ and T₄) levels, and prolactin (American Cyanamid Company 1991).

No *in vitro* studies were located regarding endocrine disruption of acrylamide.

3.7 CHILDREN'S SUSCEPTIBILITY

This section discusses potential health effects from exposures during the period from conception to maturity at 18 years of age in humans, when all biological systems will have fully developed. Potential effects on offspring resulting from exposures of parental germ cells are considered, as well as any indirect

3. HEALTH EFFECTS

effects on the fetus and neonate resulting from maternal exposure during gestation and lactation. Relevant animal and *in vitro* models are also discussed.

Children are not small adults. They differ from adults in their exposures and may differ in their susceptibility to hazardous chemicals. Children's unique physiology and behavior can influence the extent of their exposure. Exposures of children are discussed in Section 6.6, Exposures of Children.

Children sometimes differ from adults in their susceptibility to hazardous chemicals, but whether there is a difference depends on the chemical (Guzelian et al. 1992; NRC 1993). Children may be more or less susceptible than adults to health effects, and the relationship may change with developmental age (Guzelian et al. 1992; NRC 1993). Vulnerability often depends on developmental stage. There are critical periods of structural and functional development during both prenatal and postnatal life, and a particular structure or function will be most sensitive to disruption during its critical period(s). Damage may not be evident until a later stage of development. There are often differences in pharmacokinetics and metabolism between children and adults. For example, absorption may be different in neonates because of the immaturity of their gastrointestinal tract and their larger skin surface area in proportion to body weight (Morselli et al. 1980; NRC 1993); the gastrointestinal absorption of lead is greatest in infants and young children (Ziegler et al. 1978). Distribution of xenobiotics may be different; for example, infants have a larger proportion of their bodies as extracellular water, and their brains and livers are proportionately larger (Altman and Dittmer 1974; Fomon 1966; Fomon et al. 1982; Owen and Brozek 1966; Widdowson and Dickerson 1964). The infant also has an immature blood-brain barrier (Adinolfi 1985; Johanson 1980) and probably an immature blood-testis barrier (Setchell and Waites 1975). Many xenobiotic metabolizing enzymes have distinctive developmental patterns. At various stages of growth and development, levels of particular enzymes may be higher or lower than those of adults, and sometimes unique enzymes may exist at particular developmental stages (Komori et al. 1990; Leeder and Kearns 1997; NRC 1993; Vieira et al. 1996). Whether differences in xenobiotic metabolism make the child more or less susceptible also depends on whether the relevant enzymes are involved in activation of the parent compound to its toxic form or in detoxification. There may also be differences in excretion, particularly in newborns who all have a low glomerular filtration rate and have not developed efficient tubular secretion and resorption capacities (Altman and Dittmer 1974; NRC 1993; West et al. 1948). Children and adults may differ in their capacity to repair damage from chemical insults. Children also have a longer remaining lifetime in which to express damage from chemicals; this potential is particularly relevant to cancer.

3. HEALTH EFFECTS

Certain characteristics of the developing human may increase exposure or susceptibility, whereas others may decrease susceptibility to the same chemical. For example, although infants breathe more air per kilogram of body weight than adults breathe, this difference might be somewhat counterbalanced by their alveoli being less developed, which results in a disproportionately smaller surface area for alveolar absorption (NRC 1993).

Specific information regarding acrylamide-induced health effects in children was not located.

Neurotoxic end points have been examined in acrylamide-exposed mature and immature animals; however, with respect to possible age-related differences in susceptibility to acrylamide neurotoxicity, results are conflicting. Some reports indicate that young animals may be less susceptible than older ones (Fullerton and Barnes 1966; Kaplan et al. 1973), whereas other reports present evidence that young animals may be more sensitive (Ko et al. 1999; Suzuki and Pfaff 1973). In rats administered acrylamide orally at 100 mg/kg/day, Fullerton and Barnes (1966) noted that 26-week-old rats experienced earlier and more severe neurotoxic effects than 5-week-old rats. In contrast, Ko et al. (1999) reported that oral administration of acrylamide at 91 mg/kg/day resulted in earlier onset and more rapid progression of neuropathy in 3-week-old mice compared to 8-week-old mice. Studies that employed intraperitoneal injection of acrylamide also yielded conflicting results (Kaplan et al. 1973; Suzuki and Pfaff 1973). In rats repeatedly injected with 50 mg/kg acrylamide, earlier and more prominent degenerative histopathologic changes were noted in peripheral nerves of 1-day-old pups compared to adults. Kaplan et al. (1973) found the opposite in rats repeatedly injected with the same dose; 14-week-old rats experienced impaired rotarod performance earlier than 5-week-old rats (although the younger rats recovered more slowly).

Cancer is also a possible human health effect based on evidence that carcinogenic responses in rats chronically exposed to acrylamide throughout adulthood are likely mediated through a mutagenic mode of action that could occur in humans. In the absence of direct evidence that early-life exposure leads to increased risk for cancer, EPA (2005b) assumes that increased risk occurs with early-life exposure to agents that act through a mutagenic mode of action. For acrylamide, no human or animal studies were located that examined whether early-life exposure to acrylamide increased the risk for cancer, compared with exposure during adulthood alone; however, there is evidence that acrylamide acts through a mutagenic carcinogenic mode of action. CYP2E1, which catalyzes acrylamide to its DNA-reactive metabolite glycidamide, is not expressed in the developing fetus, but attains levels of expression during early postnatal periods that are similar to levels in adults (Johnsrud et al. 2003).

3. HEALTH EFFECTS

There is no information regarding developmental health effects as a result of acrylamide exposure in humans. Developmental effects observed in acrylamide-exposed rats include decreased brain levels of catecholamines, decreased auditory startle response, decreased performance in an operant test of cognitive motivation, and decreased body weight (Field et al. 1990; Garey and Paule 2007; Husain et al. 1987; Wise et al. 1995). Decreased brain levels of catecholamine resulted from various lengths of exposure via lactation or gavage (Husain et al. 1987). Decreased auditory startle response followed exposure during gestation (Wise et al. 1995), and decreased performance in an operant test of cognitive motivation followed either exposure during gestation, lactation, or through 12 weeks of age (Garey and Paule 2007). Acrylamide has been shown to distribute across the placenta in exposed pregnant animals. Two hours following intravenous administration of acrylamide to pregnant beagle dogs, concentrations of radioactivity in blood, brain, heart, and lung were similar in both maternal and fetal tissues (Ikeda et al. 1983, 1985; Marlowe et al. 1986). Humans are unlikely to be exposed by intravenous routes, but these results show that acrylamide could cross the placenta if exposure was great enough to achieve comparable maternal blood levels.

Sörgeel et al. (2002) reported the detection of low levels of acrylamide (3.17–18.8 ng/mL) in breast milk samples taken from mothers (n=2) during 8 hours following the ingestion of potato chips (estimated acrylamide dose of 0.8–1 mg); predose acrylamide in the breast milk was below the level of quantification (5 ng/mL).

In studies of offspring of nursing rat dams receiving acrylamide orally at 25 mg/kg/day throughout lactation, the dams exhibited severe toxic effects (some mortalities, body weight loss, and hindlimb splay in 90% of the dams) (Friedman et al. 1999). Starting at lactation day 4, offspring showed progressively decreased body weight, compared with control offspring (at lactation day 21, mean body weight of exposed pups was about 43% of the control pup weight); many of the pups died or became moribund during the lactational period without exhibiting signs of peripheral neuropathy (Friedman et al. 1999). In surviving acrylamide-exposed male pups, grip strength was decreased at postnatal day (PND) 30, compared with controls, but was not significantly different from control values at PNDs 60 and 90 (Friedman et al. 1999). The absence of milk in the stomachs of exposed pups that died or became moribund during lactation suggests that inadequate milk supply caused these effects in the offspring. An earlier study by Husain et al. (1987) reported complete hindlimb paralysis in male offspring of comparably-exposed rat dams of the same strain, but neither of the studies examined breast milk for the presence of acrylamide. Free acrylamide was not detected in the serum of Sprague-Dawley rat dams

3. HEALTH EFFECTS

receiving acrylamide from the drinking water from gestation day 6 through postnatal day 21 at concentrations resulting in mean daily acrylamide doses as high as 14.56 mg/kg (Takahashi et al. 2009). No free acrylamide was detected in stomach contents and serum of the pups at postnatal day 14. However, dose-related increases in acrylamide-Hb adduct levels of both dams and their pups were noted; the levels in the pups were ≥ 10 -fold lower than those of their respective dams.

3.8 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility (NAS/NRC 1989).

A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule(s) or cell(s) that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself, substance-specific metabolites in readily obtainable body fluid(s), or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic substance (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time samples can be taken. It may be difficult to identify individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc, and selenium). Biomarkers of exposure to acrylamide are discussed in Section 3.8.1.

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction (e.g., increased liver enzyme activity or pathologic changes in female genital epithelial cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are not often substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by acrylamide are discussed in Section 3.8.2.

3. HEALTH EFFECTS

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic substance. It can be an intrinsic genetic or other characteristic or a preexisting disease that results in an increase in absorbed dose, a decrease in the biologically effective dose, or a target tissue response. If biomarkers of susceptibility exist, they are discussed in Section 3.10, Populations That Are Unusually Susceptible.

3.8.1 Biomarkers Used to Identify or Quantify Exposure to Acrylamide

Several biomarkers of exposure to acrylamide have been reported in the literature. Unchanged acrylamide, its mercapturic acid metabolite, AAMA, its epoxy derivative, glycidamide, and the respective metabolite of glycidamide, GAMA, were quantified in the urine of six volunteers after the consumption of a meal containing 0.94 mg of acrylamide (Fuhr et al. 2006). Urinary mercapturic acid derivatives of acrylamide and/or glycidamide were quantified in other human studies as well (Bjellaas et al. 2007a; Boettcher et al. 2005, 2006a, 2006b). Results of epidemiological studies support the use of hemoglobin adducts of acrylamide and/or glycidamide as biomarkers of exposure to acrylamide (Bergmark et al. 1993; Boettcher et al. 2005; Calleman et al. 1994; Fennell et al. 2005b, 2006; Hagmer et al. 2001; Olesen et al. 2008). Results of animal studies indicate similar biomarkers of exposure to acrylamide. Metabolites can be measured for assessment of recent exposure. Hemoglobin adducts provide a biomarker of exposure for longer periods. See the introductory paragraph in Section 3.4 (Toxicokinetics) for a more detailed discussion of hemoglobin adduct levels as biomarkers of exposure.

It should be noted that hemoglobin adducts of N-methylolacrylamide are indistinguishable from hemoglobin adducts of acrylamide.

3.8.2 Biomarkers Used to Characterize Effects Caused by Acrylamide

Glycidamide-derived DNA adduct formation has been quantified in rats and mice exposed to acrylamide (Doerge et al. 2005a; Gamboa da Costa et al. 2003). There are no other known biomarkers of effect that are considered to be specific to acrylamide exposure.

3.9 INTERACTIONS WITH OTHER CHEMICALS

Nesterova et al. (1999) demonstrated an enhanced effect of acrylamide-induced clastogenicity in male mice administered acrylamide in combination with Verapamil (a calcium antagonist). No other

3. HEALTH EFFECTS

information was located regarding health effects attributed to interactions between acrylamide and other substances.

3.10 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE

A susceptible population will exhibit a different or enhanced response to acrylamide than will most persons exposed to the same level of acrylamide in the environment. Reasons may include genetic makeup, age, health and nutritional status, and exposure to other toxic substances (e.g., cigarette smoke). These parameters result in reduced detoxification or excretion of acrylamide, or compromised function of organs affected by acrylamide. Populations who are at greater risk due to their unusually high exposure to acrylamide are discussed in Section 6.7, Populations with Potentially High Exposures.

No human data were located regarding populations that would be particularly sensitive to acrylamide toxicity.

Available animal data demonstrate increased susceptibility of males to acrylamide-induced reproductive effects expressed as male-mediated implantation loss, reduced number of fetuses, and testicular atrophy. The dominant lethal effects observed in male, but not female, rodents may be caused by acrylamide-induced alkylation of sperm protamine during spermiogenesis (Adler et al. 2000; Generoso et al. 1996; Perrault 2003; Segal et al. 1989; Sublet et al. 1989). Key determinants of male reproductive performance such as copulatory behavior (Zenick et al. 1986) and sperm motility (Sublet et al. 1989; Tyl et al. 2000b) may also be adversely affected by acrylamide.

Available animal data do not suggest gender-related differences in susceptibility to acrylamide neurotoxicity. Male and female rats experience similar neurological effects at comparable dose levels (Burek et al. 1980; Friedman et al. 1995; Fullerton and Barnes 1996; Johnson et al. 1984, 1986). Results of animal cancer bioassays do not indicate any gender-related differences in susceptibility to acrylamide carcinogenicity. Chronic exposure of F344 rats to acrylamide in drinking water induced increased incidences of thyroid follicular cell tumors in both genders, scrotal sac mesotheliomas in males, and mammary gland fibroadenomas in females (Friedman et al. 1995; Johnson et al. 1986).

No human data were located regarding age-related differences in susceptibility to acrylamide toxicity in humans. Conflicting reports are available from animal studies. Some reports indicate that young animals may be more susceptible than older ones (Ko et al. 1999; Suzuki and Pfaff 1973), whereas other reports

3. HEALTH EFFECTS

suggest decreased susceptibility in young animals (Fullerton and Barnes 1966; Kaplan et al. 1973). It should be noted that CYP2E1, which catalyzes acrylamide to its DNA-reactive metabolite glycidamide, is not expressed in the developing fetus, but attains levels of expression during early postnatal periods that are similar to levels in adults (Johnsrud et al. 2003); however, the toxicological significance of this finding has not been demonstrated. Refer to Section 3.7 (Children's Susceptibility) for a detailed discussion of children's susceptibility to acrylamide toxicity.

Genetic polymorphisms in the acrylamide metabolizing P-450 enzyme CYP2E1 have been identified in humans (Hanioka et al. 2003). Polymorphisms in CYP2E1 could conceivably confer a differential risk to acrylamide toxicity and carcinogenicity. There is currently no quantitative estimate of differences in acrylamide or glycidamide tissue or blood levels that might result from CYP2E1 polymorphisms at high or low levels of acrylamide exposure. It is also noted that, since both acrylamide and glycidamide can exert toxic effects, different catalytic activities of CYP2E1 may result in different spectra of adverse effects.

Because the acrylamide metabolite, glycidamide, is DNA reactive, individual differences in DNA repair and detoxification mechanisms might influence susceptibility to acrylamide toxicity.

3.11 METHODS FOR REDUCING TOXIC EFFECTS

This section will describe clinical practice and research concerning methods for reducing toxic effects of exposure to acrylamide. However, because some of the treatments discussed may be experimental and unproven, this section should not be used as a guide for treatment of exposures to acrylamide. When specific exposures have occurred, poison control centers and medical toxicologists should be consulted for medical advice. The following texts provide specific information about treatment following exposures to acrylamide:

Currance PL, Clements B, Bronstein AC. 2007. Tri-ortho-cresyl phosphate (TOCP) and related compounds. In: Emergency care for hazardous materials exposure. 3rd ed. St. Louis, MO: MosbyJems, 482-484.

Goldfrank LR, Flomenbaum NE, Lewin NA, et al. 1998. Goldfrank's toxicologic emergencies. Stamford, CT: Appleton & Lange, 322-324, 475.

Leikin JB, Paloucek FP. 2002. Leikin & Paloucek's poisoning & toxicology handbook. 3rd ed. Hudson, OH: Lexi-Comp, Inc., 193-194.

3. HEALTH EFFECTS

Palmer RB. 2004. Acrylic acid and derivatives. In: Dart RC, ed. Medical toxicology. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1358-1368.

The following methods for reducing the toxic effects of acrylamide are applicable to numerous organic chemicals; they are not unique to acrylamide.

3.11.1 Reducing Peak Absorption Following Exposure

Rapid absorption of acrylamide can occur following exposure via inhalation, oral, and dermal routes. In the case of inhalation exposure, recommendations include removal from the site of exposure, establishment of a patent airway in the patient, and accompaniment by suction, ventilation, or administration of oxygen by a nonbreathing mask, if necessary (Currance et al. 2007). In the presence of airborne acrylamide, an impervious body protector may shield from absorption via dermal exposure (Leikin and Paloucek 2002). To reduce percutaneous absorption, immediate decontamination with mild liquid soap and large quantities of water has been recommended (Currance et al. 2007; Goldfrank et al. 1998). To reduce absorption resulting from oral exposure, administration of activated charcoal (Leikin and Paloucek 2002; Palmer 2004), particularly within 1 hour of ingestion (Leikin and Paloucek 2002) has been recommended. Because of its potential for central nervous system depression and seizures, ipecac-induced emesis has been discouraged for treatment of acrylamide ingestion (Currance et al. 2007). Use of gastric lavage has not been proven helpful and is not recommended (Palmer 2004).

3.11.2 Reducing Body Burden

No data were located regarding methods for reducing the body burden of absorbed acrylamide.

Animal studies indicate that acrylamide and its metabolites do not accumulate in any tissue other than red blood cells (Barber et al. 2001a; Crofton et al. 1996; Edwards 1975; Hashimoto and Aldridge 1970; Ikeda et al. 1985; Kadry et al. 1999; Marlowe et al. 1986; Miller et al. 1982; Ramsey et al. 1984) and late staged spermatids (Sega et al. 1989) and that elimination of acrylamide and its metabolites is relatively rapid (Fuhr et al. 2006; Leikin and Paloucek 2002).

3.11.3 Interfering with the Mechanism of Action for Toxic Effects

The peripheral nervous system is the primary target of acrylamide toxicity in humans and animals. Neurotoxic effects have been observed following inhalation, oral, and dermal exposures. Administration

3. HEALTH EFFECTS

of pyridoxine has been suggested as a possible treatment to delay the onset of neurotoxic effects (Leikin and Paloucek 2002). N-Acetylcysteine has also been utilized, but is of unproven benefit (Leikin and Paloucek 2002).

Administration of 2-cyclopentyl-5-(5-isoquinolylsulfonyl)-6-nitro-1*H*-benzo[D]imidazole to acrylamide-exposed rodents reduced acrylamide-induced behavioral deficits. This effect is suggestive of a therapeutic potential for peripheral neuropathy (Nakagawa-Yagi et al. 2001).

3.12 ADEQUACY OF THE DATABASE

Section 104(I)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylamide is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylamide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

3.12.1 Existing Information on Health Effects of Acrylamide

The existing data on health effects of inhalation, oral, and dermal exposure of humans and animals to acrylamide are summarized in Figure 3-7. The purpose of this figure is to illustrate the existing information concerning the health effects of acrylamide. Each dot in the figure indicates that one or more studies provide information associated with that particular effect. The dot does not necessarily imply anything about the quality of the study or studies, nor should missing information in this figure be interpreted as a “data need”. A data need, as defined in ATSDR’s *Decision Guide for Identifying Substance-Specific Data Needs Related to Toxicological Profiles* (Agency for Toxic Substances and Disease Registry 1989), is substance-specific information necessary to conduct comprehensive public

3. HEALTH EFFECTS

Figure 3-7. Existing Information on Health Effects of Acrylamide

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation					●					●
Oral		●								
Dermal					●					●

Human

	Systemic									
	Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation	●	●	●			●				
Oral	●	●	●	●		●	●	●	●	●
Dermal	●	●				●	●	●	●	●

Animal

● Existing Studies

3. HEALTH EFFECTS

health assessments. Generally, ATSDR defines a data gap more broadly as any substance-specific information missing from the scientific literature.

3.12.2 Identification of Data Needs

Acute-Duration Exposure. Available human information is limited to a single case in which intentional ingestion of 18 g of acrylamide resulted in clinical signs of peripheral neuropathy (Donovan and Pearson 1987). Limited acute-duration animal data are available for inhalation and dermal exposure routes. Clinical signs of neurological effects were observed in laboratory animals exposed to acrylamide dust at 15.6 mg/m³, a lethal or near-lethal concentration (American Cyanamid Company 1953a). Acute-duration dermal exposure elicited indications of male-mediated decreased fertility in rabbits at 50 mg/kg/day (Gutierrez-Espeleta et al. 1992), clinical signs of neurological effects in rats at doses ≥ 200 mg/kg (American Cyanamid Company 1973), and slight initial weight loss and slight dermal irritation in rabbits at doses ≥ 500 mg/kg (American Cyanamid Company 1951; Dow Chemical Company 1957).

Numerous reports are available regarding the effects of acute-duration oral exposure in animals. Acute oral LD₅₀ values range from 107 to 413 mg/kg (American Cyanamid Company 1951, 1973, 1977; Dow Chemical Company 1957; Fullerton and Barnes 1966; Hashimoto et al. 1981; McCollister et al. 1964; Tilson and Cabe 1979; Union Carbide Corporation 1947). Symptoms of acrylamide-induced neurotoxicity were elicited by single oral doses at lethal or near-lethal levels (100–200 mg/kg) (American Cyanamid Company 1953c; Fullerton and Barnes 1966; McCollister et al. 1964; Tilson and Cabe 1979) and at lower dose levels (25–45 mg/kg/day) during repeated oral dosing for ≥ 14 days (Dixit et al. 1981; Gilbert and Maurissen 1982; Tyl et al. 2000b). Single oral dosing at 63–150 mg/kg resulted in depressed body weight gain or actual weight loss (Dow Chemical Company 1957; Sakamoto et al. 1988); similar effects on body weight were elicited during repeated oral dosing at lower dose levels (15–20 mg/kg/day) (Burek et al. 1980; Tyl et al. 2000b). Male-mediated implantation losses were noted following acute-duration repeated oral dosing of rats at levels as low as 15–45 mg/kg/day (Sublet et al. 1989; Tyl et al. 2000b; Working et al. 1987).

Additional acute-duration studies of animals exposed by inhalation could be designed to assess exposure concentration-response relationships and provide a basis for an acute-duration inhalation MRL for acrylamide, although the general population is not likely to encounter acutely hazardous airborne

3. HEALTH EFFECTS

concentrations of acrylamide. Additional dermal studies using multiple dose levels could be designed to more extensively characterize the hazards of dermal exposure to acrylamide.

Intermediate-Duration Exposure. Information in humans is available from numerous case reports in which acrylamide exposure was associated with signs of impaired neurological performance in central and peripheral nervous systems that include impaired motor function and muscle weakness (Auld and Bedwell 1967; Davenport et al. 1976; Dumitru 1989; Fullerton 1969; Garland and Patterson 1967; Gjerloff et al. 2001; Igisu et al. 1975; Kesson et al. 1977; Mapp et al. 1977; Mulloy 1996; Takahashi et al. 1971). Human data are also available from cross-sectional studies that included self-reported symptoms and neurological evaluations of acrylamide-exposed workers with potential for inhalation and dermal (and possibly oral) exposure (Bachmann et al. 1992; Calleman et al. 1994; Hagmar et al. 2001; He et al. 1989; Myers and Macun 1991). In the cross-sectional studies, workers were exposed for time periods as short as 1 month (but predominantly for more than 1 year); typical signs and symptoms of acrylamide-induced neuropathy were reported. However, the human studies do not include meaningful exposure-response data, and relative contributions of inhalation, dermal, and oral exposure routes could not be determined.

Most available intermediate-duration animal studies employed the oral exposure route. Treatment-related deaths were noted at repeated doses in the range of 25–50 mg/kg/day (American Cyanamid Company 1953b, 1991; Fullerton and Barnes 1966; Schulze and Boysen 1991). Several studies reported acrylamide-induced adverse effects on body weight (American Cyanamid Company 1953b, 1979, 1991; Field et al. 1990; Friedman et al. 1999; Ko et al. 1999; Post and McLeod 1977a; Satchell and McLeod 1981; Schulze and Boysen 1991; Tanii and Hashimoto 1983; Wise et al. 1995). One study reported 12% decreased maternal weight gain in rats administered acrylamide at a dose as low as 7.5 mg/kg/day during gestation days 6–20 (Field et al. 1990). A 28-day drinking water study reported acrylamide-associated decreases in serum testosterone and thyroid hormones T₃ and T₄ in male rats (American Cyanamid Company 1991). Male-mediated implantation losses were noted following intermediate-duration repeated oral dosing of rats or mice at levels as low as 2.8–13.3 mg/kg/day (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Smith et al. 1986; Tyl et al. 2000a; Zenick et al. 1986). The developmental toxicity of acrylamide has been assessed to some extent (American Cyanamid Company 1979; Field et al. 1990; Garey and Paule 2007; Husain et al. 1987; Sakamoto and Hashimoto 1986; Wise et al. 1995); reported effects include neurochemical changes in the neonatal rat brain (Husain et al. 1987), decreased fetal body weight (Field et al. 1990), increased activity and decreased startle response (Wise et al. 1995), and subtle effects on measures of cognitive motivation (Gary and Paule 2007).

3. HEALTH EFFECTS

Numerous studies reported clinical signs and other indicators (e.g., hindlimb splay) of neurological effects associated with intermediate-duration oral exposure in laboratory animals at doses in the range of 5.7–100 mg/kg/day (American Cyanamid Company 1953b, 1953c, 1959, 1991; Burek et al. 1980; Dixit et al. 1981; Dow Chemical Company 1981; Field et al. 1990; Friedman et al. 1999; Fullerton and Barnes 1966; Gorzinski et al. 1979; Hashimoto et al. 1981; Hersch et al. 1989a, 1989b; Hopkins 1970; Ko et al. 1999, 2000, 2002; Leswing and Ribelin 1969; Maurissen et al. 1983; McCollister et al. 1964; Merigan et al. 1985; Post and McLeod 1977a; Satchell and McLeod 1981; Tanii and Hashimoto 1983; Tilson and Cabe 1979; Wise et al. 1995; Zenick et al. 1986). In general, the higher the dose level, the earlier the appearance of neurological signs. Some studies included histopathological evaluation of peripheral nerve fibers in acrylamide-treated animals (Burek et al. 1980; Eskin et al. 1985; Gorzinski et al. 1979; Johnson et al. 1985, 1986; Schulze and Boysen 1991). One study reported ultrastructural evidence of degenerative effects in sciatic nerve fibers from male F344 rats that had received acrylamide from the drinking water at 1 mg/kg/day for up to 93 days and served as the basis for deriving an intermediate-duration oral MRL for acrylamide (Burek et al. 1980).

Available intermediate-duration animal data for inhalation and dermal exposure are limited. Death and clinical signs of neurotoxicity were reported in dogs repeatedly exposed to acrylamide dust at 15.6 mg/m³, for up to 16 days (American Cyanamid Company 1953a) and rabbits administered acrylamide dermally at 50 mg/kg/day for 5 weeks (Rohm and Haas 1975). No adverse effects were observed in cats repeatedly exposed to acrylamide dust at 4.8 mg/m³ for up to 3 months (American Cyanamid Company 1954).

Additional intermediate-duration studies of animals exposed by inhalation designed to assess exposure concentration-response relationships could serve as the basis for deriving an intermediate-duration inhalation MRL for acrylamide. Additional dermal studies using multiple dose levels could be designed to better characterize the hazards of intermediate-duration dermal exposure to acrylamide.

Chronic-Duration Exposure and Cancer. Reports of chronic occupational exposure in humans describe a spectrum of neurological effects resulting from inhalation (Calleman et al. 1994; Myers and Macun 1991) and dermal effects such as peeling of skin as a result of dermal exposure (Bachmann et al. 1992; Calleman et al. 1994; He et al. 1989; Myers and Macun 1991). Questionnaires, physical examinations, neurological examinations, and ENMG tests provide support from a cross-sectional analysis on factory workers in China (Calleman et al. 1994). This study also utilized hemoglobin adduct levels as a biomarker (Calleman et al. 1994). Neurological effects in rats from chronic oral exposures to

3. HEALTH EFFECTS

acrylamide are documented and provide NOAEL and LOAEL for increased incidence of ultrastructural degeneration in the sciatic nerve and degenerative effects in peripheral nerve fibers (Johnson et al. 1984, 1985, 1986).

One case-control human study suggests a possible association between diet before puberty and subsequent risk of breast cancer; this particular study, however, is of limited use due to lacking documentation of significant factors such as cooking methods and accurate recall (Michels et al. 2006). Olesen et al. (2008) discovered a significant positive correlation between acrylamide-hemoglobin adduct levels and ER+ breast cancer, but the conclusions are also only of limited use due to the small study size and uncertainty regarding extrapolation of exposure. Most prospective studies found no statistically significant associations between acrylamide in food and risks of cancers of the esophagus, stomach, or pancreas (Hogervorst et al. 2008b); colon or rectum (Hogervorst et al. 2008b; Larsen et al. 2009c; Mucci et al. 2006); bladder or prostate (Hogervorst et al. 2008a); breast (Hogervorst et al. 2007; Larsson et al. 2009d; Wilson et al. 2009b); endometrium (Hogervorst et al. 2007; Larsson et al. 2009a); or ovarian epithelium (Larsson et al. 2009b). Exceptions include increased risks of postmenopausal endometrial and ovarian cancer (Hogervorst et al. 2007) and renal cell cancer (Hogervorst et al. 2008a) with increasing dietary acrylamide. The conclusions of these studies are only somewhat reliable as the estimations of dietary acrylamide levels in food on the market at baseline in 1986 are based on food samples analyzed in 2001 and details regarding the specifics of food preparation are lacking.

Conclusions of a 2-year bioassay of male and female F344 rats include significant increases in thyroid (follicular cell) adenomas (no carcinomas), mesotheliomas of the tunica vaginalis testis, mammary gland benign tumors (adenoma, fibroadenomas, or fibroma), and pituitary gland adenomas in high dose females (Johnson et al. 1984, 1986).

The NCTR (2009) recently performed cancer bioassays on Fischer 344 rats and B6C3F1 mice administered acrylamide or glycidamide in the drinking water for 2 years. The carcinogenicity of acrylamide and glycidamide has also been assessed in neonatally-exposed B6C3F1 mice. Post-public comment drafts of this ATSDR Toxicological Profile for Acrylamide will include available results from these carcinogenicity bioassays.

Genotoxicity. The genotoxicity of acrylamide has been studied both *in vivo* and *in vitro*. Studies are limited almost exclusively to laboratory rodents and nonmammalian species with the exception of a few *in vitro* assays of human cells. Results indicate that acrylamide is genotoxic and most potent in its ability

3. HEALTH EFFECTS

to induce clastogenic effects (including heritable translocations in offspring of acrylamide-exposed male rodents mated with untreated females), DNA damage, and gene mutations (including male-germ-cell-mediated dominant lethal mutations and heritable specific-locus mutations).

Genotoxicity assessments for acrylamide and its metabolite, glycidamide, in rats and mice are currently underway at NTCR (2009). Post-public comment drafts of this ATSDR Toxicological Profile for Acrylamide will include available results from these assessments.

Reproductive Toxicity. The reproductive toxicity of acrylamide has been studied almost exclusively in orally-exposed rats and mice. Pre- and postimplantation losses and decreased numbers of live fetuses were noted at repeated doses in the range of 3–60 mg/kg/day (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Smith et al. 1986; Sublet et al. 1989; Tyl et al. 2000a, 2000b; Working et al. 1987; Zenick et al. 1986).

Results of dominant lethality testing and crossover trials indicate that acrylamide induces male-mediated reproductive effects at repeated oral doses in the range of 2.8–19 mg/kg/day (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Smith et al. 1986; Tyl et al. 2000a, 2000b; Zenick et al. 1986). Other reported effects include decreased sperm mobility in Long-Evans rats receiving acrylamide at 45 mg/kg/day for 5 days (Sublet et al. 1989), degenerative effects in spermatids of ddY mice dosed at 100 or 150 mg/kg/day for 5 days (Sakamoto et al. 1988), and testicular atrophy in F344 rats at doses as low as 5 mg/kg/day for 90 days (American Cyanamid Company 1991; Burek et al. 1980). In apparent contrast, Tyl et al. (2000b) found no significant effects on sperm parameters in Long-Evans hooded rats following repeated oral dosing at dose levels as high as 60 mg/kg/day. Gross and histopathologic examinations of reproductive organs and tissues from male and female rats receiving acrylamide from the drinking water for up to 2 years at estimated doses as high as 2 mg/kg/day revealed no signs of acrylamide-induced effects (Friedman et al. 1995; Johnson et al. 1984, 1986). Prebreeding exposure of female rats and mice to acrylamide at doses of approximately 14 and 19 mg/kg/day did not adversely affect reproductive performance (Sakamoto and Hashimoto 1986; Zenick et al. 1986).

The reproductive toxicity of acrylamide appears to have been adequately characterized in laboratory animals. Continued assessment of possible reproductive effects in humans with potential for significant exposure to acrylamide is recommended.

3. HEALTH EFFECTS

Developmental Toxicity. The developmental toxicity of acrylamide has been assessed only in oral studies of rats and mice. Effects include body weight decreases and decreased auditory startle response in offspring of female Sprague-Dawley rats exposed to 5 and 15 mg/kg-day, respectively, on gestation days 6–10 (Wise et al. 1995), decreased mean fetal body weight in offspring of CD-1 mouse dams administered acrylamide by gavage at 45 mg/kg/day during gestation days 6–20 (Field et al. 1990), decreased mean fetal body weight in offspring of Sprague-Dawley rat dams receiving acrylamide from the drinking water (estimated acrylamide intake as high as 14.56 mg/kg/day) from gestation day 6 throughout lactation (Takahashi et al. 2009), decreased brain levels of selected catecholamines (noradrenaline, dopamine, 5-hydroxytryptamine) in pups of rat dams administered acrylamide at 25 mg/kg/day during lactation only or in young pups receiving the same dose for 5 days (Husain et al. 1987), and decreased cognitive motivation in adolescent F344 rats exposed during gestation and lactation and extending through 12 weeks postpartum (Garey and Paule 2007). No exposure-related fetal malformations or variations (gross, visceral, or skeletal) were found in offspring of Sprague-Dawley rats administered acrylamide at doses of 2.5, 7.5, or 15 mg/kg/day on gestation days 6–20 or in CD-1 mice at doses of 3, 15, or 45 mg/kg/day on gestation days 6–17 (Field et al. 1990).

Section 3.12.3 (Ongoing Studies) lists information regarding an ongoing developmental neurotoxicity study in rats. The results of this study should be evaluated prior to assessing a need for additional developmental toxicity studies of acrylamide.

Immunotoxicity. No human or animal data were located in which acrylamide was considered to be a potential immunological concern. Immunotoxicity studies do not appear necessary at this time.

Neurotoxicity. Information in humans is available from numerous case reports in which acrylamide exposure has been associated with signs of impaired neurological performance in central and peripheral nervous systems that include impaired motor function and muscle weakness (Auld and Bedwell 1967; Davenport et al. 1976; Dumitru 1989; Fullerton 1969; Garland and Patterson 1967; Gjerløff et al. 2001; Igisu et al. 1975; Kesson et al. 1977; Mapp et al. 1977; Mulloy 1996; Takahashi et al. 1971). Human data are also available from cross-sectional studies that included self-reported symptoms and neurological evaluations of acrylamide-exposed workers with potential for inhalation and dermal (and possibly oral) exposure (Bachmann et al. 1992; Calleman et al. 1994; Hagmar et al. 2001; He et al. 1989; Myers and Macun 1991).

3. HEALTH EFFECTS

Neurological effects associated with oral exposure to acrylamide have been well characterized in laboratory animals and include clinical signs such as twitching, loss of balance, tremors, lethargy, and general weakness and more subtle indicators of functional deficits such as decreased rotarod performance and increased limb or foot splay. Evidence of degenerative lesions in peripheral nerve fibers, as observed by light and electron microscopy, have been detected at oral doses lower than those eliciting clinical signs and other overt indications of functional deficit. See Sections 3.2.1.4, 3.2.2.4, and 3.2.3.4 for detailed information regarding neurological effects in acrylamide-exposed animals. Available animal data appear to adequately characterize acrylamide neurotoxicity; additional neurotoxicity studies do not appear necessary at this time.

Refer to Section 3.12.2 (Developmental Effects) for a summary of information regarding acrylamide-induced neurodevelopmental effects.

Epidemiological and Human Dosimetry Studies. Early epidemiological studies focused on neurological signs and symptoms in workers employed in the manufacture and/or use of acrylamide (Bachmann et al. 1992; Calleman et al. 1994; Hagmar et al. 2001; He et al. 1989; Myers and Macun 1991). Although these studies, as well as available case reports, provide supportive evidence of acrylamide-induced neurotoxicity, they lack information regarding relative contributions of natural exposure routes (inhalation, oral, dermal), exposure-response relationships, and other confounding exposures.

More recent investigations have focused on examining possible associations between acrylamide dietary intake and various cancer end points (Hogervorst et al. 2007, 2008a, 2008b; Larsson et al. 2009a, 2009b, 2009c, 2009d; Michels et al. 2006; Mucci et al. 2003, 2004, 2005, 2006; Pelucchi et al. 2006, 2007; Wilson et al. 2009a, 2009b). These studies provide little evidence of acrylamide-induced carcinogenicity from estimated intakes through the diet. However, a major deficiency of these studies is the use of questionnaires to estimate prior dietary intake of acrylamide.

Continued epidemiological research should focus on improving methods for estimating acrylamide intake. Additional information regarding biomarkers of exposure to acrylamide and improved PBPK modeling could be beneficial.

3. HEALTH EFFECTS

Biomarkers of Exposure and Effect.

Exposure. Biomarkers of exposure to acrylamide include unchanged acrylamide, its mercapturic acid metabolite, AAMA, its epoxy derivative, glycidamide, and the respective metabolite of glycidamide, GAMA in urine (Bjellaas et al. 2007a; Boettcher et al. 2005, 2006a, 2006b; Fuhr et al. 2006) and hemoglobin adducts of acrylamide and glycidamide (Bergmark et al. 1993; Boettcher et al. 2005; Calleman et al. 1994; Fennell et al. 2005b; Hagmer et al. 2001; Olesen et al. 2008). However, N-methylolacrylamide forms hemoglobin adducts that are indistinguishable from hemoglobin adducts of acrylamide. The development of a method to measure levels of the metabolite, glyceramide, in urine could help to determine the relative importance of glycidamide via glutathione conjugation versus hydrolysis.

See Section 3.12.3 (Ongoing Studies) for information regarding assessments currently underway to assess biomarkers of exposure to acrylamide.

Effect. Glycidamide-DNA adducts may be considered a biomarker of effect from exposure to acrylamide (Doerge et al. 2005a; Gamboa da Costa et al. 2003). It is not likely that additional biomarkers of effect specific to acrylamide would be identified.

Absorption, Distribution, Metabolism, and Excretion. Available human data indicate that acrylamide is readily absorbed following oral or dermal exposure; significant absorption of inhaled acrylamide is expected. In animals, acrylamide is readily absorbed following inhalation, oral, or dermal exposure. Absorbed acrylamide is widely distributed (accumulates only in red blood cells), rapidly metabolized in the liver, and excreted in the urine mainly as acrylamide-derived conjugates. Existing PBPK models for acrylamide are not adequate for human health risk assessment. However, as described in Section 3.12.3 (Ongoing Studies) a new PBPK model for acrylamide is presently being developed. Additional pharmacokinetic studies may be necessary to assist in calibration and validation of PBPK models for acrylamide.

Comparative Toxicokinetics. Available data in rats and mice indicate species differences in acrylamide metabolism. Additional comparative toxicokinetic studies may be warranted pending the outcome of the new PBPK model being developed for acrylamide.

3. HEALTH EFFECTS

Methods for Reducing Toxic Effects. Suggested methods for reducing absorption of acrylamide include removal from exposure source, decontamination of exposed skin with soap and water, and administration of activated charcoal within 1 hour following ingestion. There are no known methods for reducing the body burden following absorption of acrylamide and no proven methods for interfering with the mechanism of action for toxic effects.

Children's Susceptibility. No specific data were located regarding acrylamide-induced health effects in children. The database of information in animals provides conflicting results. Children are not expected to be less susceptible than adults to acrylamide toxicity.

Data needs relating to both prenatal and childhood exposures, and developmental effects expressed either prenatally or during childhood, are discussed in detail in the Developmental Toxicity subsection above.

Child health data needs relating to exposure are discussed in Section 6.8.1, Identification of Data Needs: Exposures of Children.

3.12.3 Ongoing Studies

Search of the NCTR website revealed the following (NCTR 2009):

Dr. Frederick Beland (principal investigator), Division of Biochemical Toxicology, is comparing the carcinogenicity of acrylamide and its metabolite glycidamide in B6C3F1 mice treated neonatally and in B6C3F1 mice and Fischer 344 rats treated for 2 years. The 2-year study is funded by the National Toxicology Program (NTP).

Dr. Daniel Doerge (principal investigator), Division of Biochemical Toxicology, in collaboration with the Division of Genetic and Reproductive Toxicology and Division of Personalized Nutrition and Medicine and funding from the University of Maryland, is working to (1) develop a PBPK/PD model for acrylamide and glycidamide and (2) determine mutagenicity of acrylamide and glycidamide in Big Blue® rats. Dr. Doerge, in collaboration with the Department of genetic and Reproductive Toxicology and funding from NTP, is studying genotoxicity, mutagenicity, and biomarkers of exposure for acrylamide and glycidamide in rodents.

3. HEALTH EFFECTS

Dr. Merle Paule (principal investigator), Division of Neurotoxicology, in collaboration with the Division of Biochemical Toxicology and funding from NTP, is assessing developmental neurotoxicity of acrylamide in rats.

Dr. Eden Tareke (principal investigator), Division of Neurotoxicology, is investigating the effects of acrylamide on normal human brain cortical neuronal (HCN-1), PC12, and HepG2 cells *in vitro*.

Search of the Federal Research in Progress database (FEDRIP 2009) revealed the following:

Dr. John Essigmann (principal investigator), Massachusetts Institute of Technology, Cambridge, Massachusetts, is investigating the genotoxic effects of several compounds, including acrylamide, with the goal of developing new biomarkers. The sponsoring organization is the National Institute of Environmental Health Sciences.

Dr. Richard LoPachin (principal investigator), Montefiore Medical Center, Bronx, New York, is performing ongoing research to elucidate specific mechanisms responsible for acrylamide toxicity.

4. CHEMICAL AND PHYSICAL INFORMATION

4.1 CHEMICAL IDENTITY

Acrylamide is an unsaturated amide that is produced mainly for use as an intermediate in the production of polyacrylamide (Abdelmagic 1982; Haberman 2002). Information regarding the chemical identity of acrylamide is located in Table 4-1.

4.2 PHYSICAL AND CHEMICAL PROPERTIES

Acrylamide is a white or colorless, odorless crystalline solid. Information regarding the physical and chemical properties of acrylamide is located in Table 4-2. Acrylamide is stable at room temperature but can violently polymerize at its melting point or under UV light (Lewis 2000, 2007; O'Neil et al. 2006). Acrid fumes, as well as NO_x , may be released when it is heated to decomposition (Lewis 2000). Acrylamide is soluble in water, alcohol, and acetone, and is insoluble in benzene and heptanes (Lewis 2007).

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-1. Chemical Identity of Acrylamide

Characteristic	Information	Reference
Chemical name	Acrylamide	
Synonym(s)	AAM; acrylagel; acrylic acid amide; acrylic amide; akrylamid (Czech); amresco acryl-40; amid kyseliny akrylove; ethylenecarboxamide; optimum; propenamide; 2-propenamide; propereamide; propenoic acid amide; vinyl amide	HSDB 2009; Lewis 2000
Registered trade name(s)	No data	
Chemical formula	C ₃ H ₅ NO	HSDB 2009; WHO 2003
Chemical structure		
Identification numbers:		
CAS registry	79-06-1	
NIOSH RTECS	NIOSH/AS3325000	NIOSH 2005
EPA hazardous waste	U007	HSDB 2009; Lewis 2000
OHM/TADS	No data	
DOT/UN/NA/IMDG shipping	UN 2074; IMO 6.1	HSDB 2009; Lewis 2000
HSDB	191	HSDB 2009
NCI	No data	
STCC	49 091 83 49 131 87	HSDB 2009 HSDB 2009

CAS = Chemical Abstracts Service; DOT/UN/NA/IMDG = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances; STCC = Standard Transport Commodity Code

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-2. Physical and Chemical Properties of Acrylamide

Property	Information	Reference
Molecular weight	71.08	Haberman 2002; HSDB 2009
Color	White Colorless	HSDB 2009; Lewis 2000 Lewis 2007; WHO 2003
Physical state	Crystalline solid	HSDB 2009; Lewis 2000 Lewis 2007; WHO 2003
Melting point	84.5 °C	Lewis 2007
Boiling point	192.6 °C 87 °C (2 mm) 103 °C (5 mm) 125 °C (25 mm)	Lide 2008 O'Neil et al. 2006 O'Neil et al. 2006 Lewis 2007; O'Neil et al. 2006
Density at 30 °C/4 °C	1.122	HSDB 2009; Lewis 2007
Odor	Odorless	Lewis 2007
Odor threshold:		
Water	No data	
Air	No data	
Solubility:		
Water at 0 °C	No data	
at 20 °C	3.711x10 ⁵ mg/L	HSDB 2009
at 30 °C	4.048x10 ⁵ mg/L	HSDB 2009
Organic solvents	Soluble in water, alcohol, acetone Insoluble in benzene and heptanes	Lewis 2007 Lewis 2007
Partition coefficients:		
Log K _{ow}	-0.67	HSDB 2009
Log K _{oc}	No data	
Vapor pressure		
at 25 °C	0.9 Pa (7x10 ⁻³ mm Hg)	Haberman 2002; HSDB 2009
at 40 °C	4.4 Pa (3.3x10 ⁻² mm Hg)	Haberman 2002
at 50 °C	9.3 Pa (7.0x10 ⁻² mm Hg)	Haberman 2002
Henry's law constant:		
at 25 °C	1.7x10 ⁻⁹ atm·m ³ /mol (estimated)	HSDB 2009
Autoignition temperature	424 °C	HSDB 2009
Flash point	138 °C (closed cup)	HSDB 2009
Flammability limits	No data	
Conversion factors (25 °C, 1 atm)	1 mg/m ³ =0.34 ppm 1 ppm=2.95 mg/m ³	HSDB 2009

4. CHEMICAL AND PHYSICAL INFORMATION

This page is intentionally blank.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.1 PRODUCTION

No information is available in the TRI database on facilities that manufacture or process acrylamide because this chemical is not required to be reported under Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986) (EPA 1998).

Acrylamide can be produced from acrylonitrile treated with sulfuric acid or hydrochloric acid, followed by base neutralization or use of an ion exclusion column to separate the chemical from its sulfate salt (Haberman 2002; HSDB 2009; Lewis 2007; O'Neil et al. 2006). Although historically the primary method of acrylamide production, this method results in sulfate byproducts and significant waste streams, and thus, commercial acrylamide producers no longer use this process (Haberman 2002). A newer process was developed in which a solution of acrylonitrile is passed over a fixed bed copper catalyst at 85 °C, resulting in an acrylamide solution. This method has been modified by various companies since its development. A Raney copper catalyst can also be used in both slurry and fixed-bed reactors (Haberman 2002). American Cyanamid uses a proprietary catalytic direct hydration process to produce acrylamide from acrylonitrile, using a reaction of water of solid surfaces of metal, metallic salts, or metallic oxides, which avoids the formation of unwanted byproducts (HSDB 2009).

Numerous other methods of acrylamide production have been developed. Acryloyl chloride or acrylic anhydride can be reacted with ammonia to yield acrylamide. Microorganisms can also be used to convert acrylonitrile to acrylamide via an enzymatic hydration process (Haberman 2002).

Table 5-1 lists the facilities in each state that manufacture, process, or use acrylamide as well as the intended use and the range of amounts of acrylamide that are stored on site. There are currently 216 facilities that produce, process, or use acrylamide in the United States. The data listed in Table 5-1 are derived from the Toxics Release Inventory (TRI 07 2009). These data should be used with caution, however, since only certain types of facilities are required to report. Therefore, this is not an exhaustive list. Commercial production of acrylamide in 1983 was reported as 86,233 pounds (HSDB 2009). Currently, there are four major producers of acrylamide in the United States, with a combined production capacity of approximately 141,000 metric tons (311 million pounds) (SRI 2008). These producers and their respective plant locations are provided in Table 5-2.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Acrylamide

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
AL	4	10,000	9,999,999	6, 14
AR	5	0	99,999	6, 7, 12
CA	15	0	9,999,999	1, 2, 3, 4, 6, 7, 8, 9, 10
CO	2	0	99,999	6
CT	6	10,000	9,999,999	2, 3, 4, 6
DE	1	0	0	0
FL	3	100,000	999,999	2, 3, 6
GA	8	0	9,999,999	1, 4, 6, 7
IL	14	0	9,999,999	2, 3, 6, 7, 12
IN	1	10,000	99,999	12
KY	4	10,000	999,999	6, 12
LA	13	0	9,999,999	1, 2, 3, 4, 6, 9, 10, 12, 13
MA	4	1,000	99,999	6, 7
MD	4	0	999,999	6, 7
MI	21	0	49,999,999	1, 2, 3, 4, 6, 7, 8, 10, 11, 12
MO	5	100	99,999	1, 2, 3, 6, 7, 8
NC	7	100	999,999	6, 7
NE	4	0	99,999	3, 7, 8, 11, 12
NJ	8	0	999,999	1, 6, 10, 12, 13
NY	11	100	99,999	6, 7, 9, 10, 12
OH	10	0	999,999	1, 6, 7, 12, 13
OR	1	1,000	9,999	6
PA	6	0	499,999,999	6
SC	12	0	999,999	6, 7, 12
TN	8	0	9,999,999	6, 12
TX	18	0	9,999,999	1, 5, 6, 7, 12, 13
UT	1	1,000	9,999	12
VA	7	100,000	49,999,999	1, 2, 3, 4, 6
WA	2	100,000	999,999	6, 12
WI	5	0	99,999	2, 3, 6, 7

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Acrylamide

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
WV	5	0	499,999,999	1, 5, 6, 12
WY	1	1,000,000	9,999,999	9

^aPost office state abbreviations used

^bAmounts on site reported by facilities in each state

^cActivities/Uses:

- | | | |
|--------------------------|--------------------------|-----------------------------|
| 1. Produce | 6. Impurity | 11. Chemical Processing Aid |
| 2. Import | 7. Reactant | 12. Manufacturing Aid |
| 3. Onsite use/processing | 8. Formulation Component | 13. Ancillary/Other Uses |
| 4. Sale/Distribution | 9. Article Component | 14. Process Impurity |
| 5. Byproduct | 10. Repackaging | |

Source: TRI07 2009 (Data are from 2007)

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-2. 2008 Acrylamide Production in the United States

Company	Location	Annual capacity (metric tons ^a)	Remarks
Chemtall, Inc.	Riceboro, Georgia	65,000	From purchased acrylonitrile; captive consumption for polyacrylamide
Ciba Specialty Chemicals Corporation Water & Paper Treatment Business Segment	Suffolk, Virginia	15,000	From purchased acrylonitrile; primarily captive consumption for polyacrylamide
Kemira Water Solutions, Inc.	Waggaman, Louisiana	41,000	From captive acrylonitrile; captive consumption for polyacrylamide
Nalco Company	Garyville, Louisiana	20,000	From purchased acrylonitrile; primarily captive consumption for polyacrylamide
Total		141,000	

^aAll production is as solution.

Source: SRI 2008; SRI Consulting estimates as of February 1, 2008.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Acrylamide is produced as a byproduct when foods are cooked at high temperatures, such as by frying, roasting, and baking (Muttucumaru et al. 2008). This occurs by the Maillard reaction, where thermal degradation of amino acids occurs in the presence of reducing sugars at temperatures above 120 °C. The major precursors for acrylamide formation by the Maillard reaction are free asparagines and reducing sugars (Arisseto et al. 2007; Mottram 2002; Muttucumaru et al. 2008; Stadler et al. 2002). Acrylamide concentrations in the foods rise with temperature and length of heating, and appear to be affected by water content, food composition, and processing conditions (WHO 2002).

5.2 IMPORT/EXPORT

The demand for acrylamide in the United States was reported as 245 and 253 million pounds in 2006 and 2007, respectively. U.S. imports were 61 and 55 million pounds in 2006 and 2007, respectively, while U.S. exports were 57 and 53 million pounds in 2006 and 2007, respectively (CMR 2008).

5.3 USE

Commercially-produced acrylamide is used mainly as an intermediate in the production of polyacrylamides (EU 2002; Haberman 2002; O'Neil et al. 2006; WHO 2003). This accounts for 94% of manufactured acrylamide (Haberman 2002). Polyacrylamides are then primarily used as flocculants for clarifying drinking and treating municipal and industrial effluents (Abdelmagid 1982; EPA 2006c; EU 2002; Haberman 2002; WHO 2003;). They aid in dewatering sludge from sewage treatment plant effluent as well as industrial waste water from pulp and paper plants (Abdelmagid 1982; Haberman 2002). Polyacrylamides are also found in cosmetics and toiletries and are used to prepare polyacrylamide gels for use in biotechnology laboratories (EU 2002; Lewis 2007). In the oil industry, acrylamide is used as a flow control agent to enhance oil production from wells. Acrylamide and polyacrylamides are also used in the production of dyes and organic chemicals such as N-methylacrylamide, in copolymers for contact lenses, in permanent-press fabrics, for sizing paper and textiles, as a binder and retention aid for pulp and paper production, in the processing of ore, in sugar refining, and as a chemical grouting agent and soil stabilizer for the construction of tunnels, sewers, wells, and reservoirs (Abdelmagid 1982; EPA 2006c; EU 2002; Haberman 2002; Lewis 2007; O'Neil et al. 2006; WHO 2003).

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.4 DISPOSAL

Acrylamide is typically produced as an intermediate for the production of polyacrylamides (Haberman 2002; O'Neil et al. 2006), and is therefore consumed in the process. No information was found regarding the disposal of acrylamide.

6. POTENTIAL FOR HUMAN EXPOSURE

6.1 OVERVIEW

Acrylamide has been identified in at least 3 of the 1,699 hazardous waste sites that have been proposed for inclusion on the EPA National Priorities List (NPL) (HazDat 2007). However, the number of sites evaluated for acrylamide is not known. The frequency of these sites can be seen in Figure 6-1.


Acrylamide is an industrial chemical used mainly in the production of polyacrylamides, which are primarily used as flocculants for clarifying drinking and treating municipal and industrial effluents (Abdelmagid 1982; EPA 2006c; EU 2002; Haberman 2002; O'Neil et al. 2006; WHO 2003). Acrylamide may be released to the environment during production and use of polyacrylamides, which are used as clarifiers in water treatment. Residual monomer released from the polyacrylamide coagulants is the main source of acrylamide contamination of drinking water (Abdelmagid 1982; Cavalli et al. 2004; EPA 2006c; WHO 2003), though it can also be released from plastics and dye industries and from acrylamide-containing grouting agents used in reservoirs and wells (Cavalli et al. 2004; EPA 2006c). Acrylamide can be released to the environment during formulation of cosmetics or other consumer products and in the laboratory while gel chromatography is being performed (Boettcher and Angerer 2005; EU 2002). Due to its low vapor pressure and high water solubility, acrylamide is rarely identified in atmospheric samples (WHO 2003).

Acrylamide is expected to be highly mobile in soil and water (EPA 2006c; HSDB 2009; WHO 2003). It is highly susceptible to biodegradation in both soils and surface water (Abdelmagid 1982; EPA 2006c; Haberman 2002; WHO 2003). It is not typically present in the atmosphere (HSDB 2009; Pratt 2000). Acrylamide is not expected to significantly bioconcentrate (EPA 2006c; Haberman 2002; WHO 2003).

Acrylamide is sometimes present in drinking water due to leaching of monomer during treatment processes (Abdelmagid 1982; Cavalli et al. 2004; EPA 2006c; van Dijk-Looijaard and van Genderen 2000; WHO 2003), as well as release from grouting agents in wells and dams and releases from plastics and dye industries (Cavalli et al. 2004). It is rarely found in soil samples (HSDB 2009). Acrylamide is not a common air pollutant, due to its low vapor pressure and high water solubility. It is rarely identified in atmospheric samples (HSDB 2009; Pratt 2000; WHO 2003).

6. POTENTIAL FOR HUMAN EXPOSURE

Figure 6-1. Frequency of NPL Sites with Acrylamide Contamination



Derived from HazDat 2007

6. POTENTIAL FOR HUMAN EXPOSURE

In 2002, acrylamide was first identified in samples of food cooked at high temperatures (Tareke et al. 2002). Concentrations of acrylamide in food vary with the type of food and method of cooking, and typically increase with temperature and length of heating (Sorgel et al. 2002; WHO 2002, 2003). Carbohydrate-rich foods typically contain the highest levels of acrylamide (Muttucumararu et al. 2008; Tareke et al. 2002; WHO 2003), whereas protein-based foods contain smaller amounts (Tareke et al. 2002). As acrylamide content in food appears to be affected by temperature, water content, food thickness, and length of heating, various steps can be taken to minimize the exposure of acrylamide from food sources (Sorgel et al. 2002; WHO 2002, 2003).

Acrylamide is a carcinogen with the potential to cause nervous system damage (Arisetto et al. 2007; EPA 2006c; Lewis 2000). Exposure occurs mainly via ingestion, dermal contact, and inhalation routes (Lewis 2000; Sorgel et al. 2002). Ingestion of foodstuffs containing acrylamide appears to be one of the most common methods of exposure for the general public. Average estimated intake of acrylamide from food sources ranged from 0.8 to 6.0 µg/kg bw/day for short-term exposure and 0.3 to 0.8 µg/kg bw/day for long-term exposure (WHO 2002, 2003). Children may be susceptible to food-borne exposure 2–3 times that of adults on a body weight basis (WHO 2002, 2003).

Ingestion of polyacrylamide-treated drinking water containing residual monomer as well as water in contact with acrylamide-containing products, such as grouting agents, can result in exposure to acrylamide (EPA 2006c; EU 2002; WHO 2003). The presence of acrylamide in tobacco smoke can result in inhalation exposure for both adults and children (EU 2002; Moreno Navarro et al. 2007). Dermal exposure can result from contact with cosmetics and toiletries containing polyacrylamides (EU 2002). Once in the body, acrylamide is widely dispersed by body fluids, and can also cross the placental barrier (WHO 2003), resulting in exposure to unborn children. Breast milk of mothers with diets high in acrylamide-containing foods can contain high amounts of acrylamide (Sorgel et al. 2002). Occupational exposure to acrylamide is primarily due to dermal contact when handling bags and drums of the chemical or preparing polyacrylamide gels, followed by inhalation of dust or aerosols (EU 2002; Lewis 2000).

6.2 RELEASES TO THE ENVIRONMENT

The Toxics Release Inventory (TRI) data should be used with caution because only certain types of facilities are required to report (EPA 2005a). This is not an exhaustive list. Manufacturing and processing facilities are required to report information to the TRI only if they employ 10 or more full-time employees; if their facility is included in Standard Industrial Classification (SIC) Codes 10 (except 1011,

6. POTENTIAL FOR HUMAN EXPOSURE

1081, and 1094), 12 (except 1241), 20–39, 4911 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4931 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4939 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4953 (limited to facilities regulated under RCRA Subtitle C, 42 U.S.C. section 6921 et seq.), 5169, 5171, and 7389 (limited S.C. section 6921 et seq.), 5169, 5171, and 7389 (limited to facilities primarily engaged in solvents recovery services on a contract or fee basis); and if their facility produces, imports, or processes $\geq 25,000$ pounds of any TRI chemical or otherwise uses $>10,000$ pounds of a TRI chemical in a calendar year (EPA 2005a).

6.2.1 Air

Estimated releases of 11,791 pounds (~5.3 metric tons) of acrylamide to the atmosphere from 81 domestic manufacturing and processing facilities in 2007, accounted for about 0.2% of the estimated total environmental releases from facilities required to report to the TRI (TRI07 2009). These releases are summarized in Table 6-1.

Acrylamide may be released to the atmosphere during production of polymers, formulation of cosmetics or other consumer products, and in the laboratory while gel chromatography is being performed (Boettcher and Angerer 2005). Release of tobacco smoke, which was shown to contain acrylamide (Schumacher et al. 1977), may also contribute to airborne acrylamide in confined spaces. However, acrylamide is not anticipated to be a common air contaminant due to its low vapor pressure and high water solubility. Acrylamide is not expected to be removed from soils or water by volatilization (WHO 2003). Limited data indicate that atmospheric acrylamide concentrations are very low, when it is identified at all (Pratt 2000; HSDB 2009).

The total national baseline National Toxics Inventory (NTI) emissions for acrylamide during 1900–1998 were 35.4 tons/year. Total urban emissions were 33.5 tons/year while total rural emissions were 1.9 tons/year (EPA 2000). The National Emissions Inventory database, maintained by the EPA, reports total U.S. acrylamide emissions of 12.12 tons per year in 2005 from various sources, including degreasing, waste disposal, pulp and paper, solvents, oil and gas production, industrial surface coatings, and oil and gasoline production among others (EPA 2009f).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Acrylamide^a

State ^c	RF ^d	Reported amounts released in pounds per year ^b						Total release		
		Air ^e	Water ^f	UI ^g	Land ^h	Other ⁱ	On-site ^j	Off-site ^k	On- and off-site	
AL	1	90	0	0	4	4,555	90	4,559	4,649	
AR	3	55	0	0	1	580	55	581	636	
CA	2	1	No data	0	0	0	1	0	1	
CO	1	0	No data	0	0	0	0	0	0	
CT	2	2	7	0	83	0	9	83	92	
DE	1	0	No data	0	0	0	0	0	0	
GA	6	5,629	0	0	798	0	5,629	798	6,427	
IL	6	99	0	0	0	0	99	0	99	
KY	1	0	No data	0	0	0	0	0	0	
LA	5	1,980	216	873,489	2,936	0	876,491	2,130	878,621	
MA	1	0	No data	0	0	5	0	5	5	
MD	2	2	0	0	0	0	2	0	2	
MI	3	6	0	0	0	0	6	0	6	
NC	8	1,860	0	0	0	0	1,860	0	1,860	
NE	1	2	0	0	0	0	2	0	2	
NJ	2	19	0	0	0	0	19	0	19	
OH	6	1,299	0	960,000	10	0	961,299	10	961,309	
PA	5	19	5	0	0	0	24	0	24	
SC	8	500	250	0	0	0	750	0	750	
TN	5	98	No data	0	0	357	98	357	455	
TX	6	26	0	4,303,938	8	0	4,303,971	0	4,303,971	
UT	1	1	0	0	0	0	1	0	1	
VA	1	64	0	0	176	0	64	176	240	
WA	1	21	No data	0	0	0	21	0	21	

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Acrylamide^a

State ^c	RF ^d	Reported amounts released in pounds per year ^b							
		Air ^e	Water ^f	UI ^g	Land ^h	Other ⁱ	Total release		
							On-site ^j	Off-site ^k	On- and off-site
WI	3	18	No data	0	0	2,038	18	2,038	2,056
Total	81	11,791	478	6,137,427	4,016	7,535	6,150,510	10,737	6,161,247

^aThe TRI data should be used with caution since only certain types of facilities are required to report. This is not an exhaustive list. Data are rounded to nearest whole number.

^bData in TRI are maximum amounts released by each facility.

^cPost office state abbreviations are used.

^dNumber of reporting facilities.

^eThe sum of fugitive and point source releases are included in releases to air by a given facility.

^fSurface water discharges, waste water treatment-(metals only), and publicly owned treatment works (POTWs) (metal and metal compounds).

^gClass I wells, Class II-V wells, and underground injection.

^hResource Conservation and Recovery Act (RCRA) subtitle C landfills; other on-site landfills, land treatment, surface impoundments, other land disposal, other landfills.

ⁱStorage only, solidification/stabilization (metals only), other off-site management, transfers to waste broker for disposal, unknown

^jThe sum of all releases of the chemical to air, land, water, and underground injection wells.

^kTotal amount of chemical transferred off-site, including to POTWs.

RF = reporting facilities; UI = underground injection

Source: TRI07 2009 (Data are from 2007)

6. POTENTIAL FOR HUMAN EXPOSURE

6.2.2 Water

Estimated releases of 478 pounds (~0.22 metric tons) of acrylamide to surface water from 81 domestic manufacturing and processing facilities in 2007, accounted for about 0.01% of the estimated total environmental releases from facilities required to report to the TRI (TRI07 2009). These releases are summarized in Table 6-1.

Acrylamide in drinking water is typically a result of the release of residual monomer from polyacrylamide coagulants that are used as clarifiers in the treatment of raw water (Abdelmagid 1982; Cavalli et al. 2004; EPA 2006c; WHO 2003). The clarifiers served to coagulate and trap suspended solids such that they may be removed more easily from the water. Acrylamide that does not coagulate is released into the environment as a drinking water contaminant (Abdelmagid 1982; EPA 2006c). Use of polyacrylamides as grouting agents in reservoirs and wells can result in release to drinking water supplies. They can also be released in water from plastics and dye industries (Cavalli et al. 2004; EPA 2006c). When released to land, acrylamide does not bind to soil and will thus move rapidly through the soil column (EPA 2006c), which can result in increased risk of surface or groundwater contamination.

Total release of acrylamide to water between 1987 and 1993 was 36,287 pounds. Major industries found to release acrylamide to water include plastics and resins (19,002 pounds), pulp mills (8,000 pounds), industrial organics (3,107 pounds), and industrial inorganics (2,510 pounds) (EPA 2006c). According to the EPA's Toxic Chemical Release Inventory, release of acrylamide to land and water from 1987 to 1993 was over 40,000 pounds. Acrylamide releases were typically from plastics industries, with the largest releases occurring in Michigan (EPA 2006c).

As required by the 1974 Safe Drinking Water Act, EPA developed a maximum contaminant level (MCL) for acrylamide, which specifies the concentration at which it is not expected to cause health problems. The MCL for acrylamide is zero (EPA 2006c). EPA requires water suppliers to control the amount of acrylamide added to water during the treatment process. This can be accomplished by limiting the amount of acrylamide in the polyacrylamide flocculants or by limiting the dose of flocculants (WHO 2003). Uncoagulated acrylamide in drinking water must be <0.5 µg/L (ppb) (Cavalli et al. 2004; EPA 2006c). This concentration corresponds to a maximum authorized dose of polymer of 1 mg/L (for a monomer content of 0.05%), as this corresponds to 0.5 µg/L of monomer in the water (Cavalli et al. 2004;

6. POTENTIAL FOR HUMAN EXPOSURE

WHO 2003). In European Union countries, the maximum allowable concentration in drinking water is 0.1 µg/L (Cavalli et al. 2004).

If acrylamide is found to be present in raw water, the concentration can be decreased through ozonation or by treating the water with potassium permanganate. Acrylamide is not removed by conventional water treatment processes (WHO 2003).

6.2.3 Soil

Estimated releases of 4,016 pounds (~1.8 metric tons) of acrylamide to soils from 81 domestic manufacturing and processing facilities in 2007, accounted for about 0.06% of the estimated total environmental releases from facilities required to report to the TRI (TRI07 2009). An additional 6,137,427 pounds (~2,783 metric tons), constituting about 99.6% of the total environmental emissions, were released via underground injection (TRI07 2009). These releases are summarized in Table 6-1.

Polyacrylamides releases to land can occur from plastics and dye industries (Cavalli et al. 2004; EPA 2006c). When released to land, acrylamide does not bind to soil and will thus move rapidly through the soil column (EPA 2006c; WHO 2003). Total release of acrylamide to land between 1987 and 1993 was 5,818 pounds. Major industries found to release acrylamide to land include plastics and resins (2,177 pounds), industrial organics (2,200 pounds), and industrial inorganics (500 pounds) (EPA 2006c). According to the EPA's Toxic Chemical Release Inventory, release of acrylamide to land and water from 1987 to 1993 was over 40,000 pounds. Acrylamide releases were typically from plastics industries, with the largest releases occurring in Michigan (EPA 2006c).

6.3 ENVIRONMENTAL FATE

6.3.1 Transport and Partitioning

Acrylamide is expected to be highly mobile in soil and water. When released to land, acrylamide does not bind to soil and will thus move rapidly through the soil column and into groundwater, where it is also expected to have high mobility (EPA 2006c; HSDB 2009; WHO 2003). Acrylamide has a higher mobility in sandy soils than in clay soils, which can result in increased risk of surface or groundwater contamination (Abdelmagid 1982; EPA 2006c; WHO 2003). Acrylamide is not expected to be removed from soils or water by volatilization (WHO 2003).

6. POTENTIAL FOR HUMAN EXPOSURE

Acrylamide is not expected to significantly bioconcentrate in aquatic organisms due to its high water solubility and its ability to be degraded by microorganisms (EPA 2006c; Haberman 2002; WHO 2003). Acrylamide was determined to have a bioconcentration factor (BCF) of around 1 in both the carcass and viscera (0.86–1.44 and 1.12–1.65, respectively) of rainbow trout exposed to acrylamide concentrations of 0.338 and 0.710 mg/L. This indicates that acrylamide did not bioaccumulate significantly in the trout (Petersen et al. 1985).

6.3.2 Transformation and Degradation

Acrylamide is susceptible to biodegradation in both soils and surface water (Abdelmagid 1982; EPA 2006c; Haberman 2002; HSDB 2009; WHO 2003). As it does not bind to soil, acrylamide moves rapidly through the soil column where it is quickly degraded (EPA 2006c; WHO 2003).

In sandy soils, acrylamide has a higher mobility and lower rate of degradation than in clay soils (Abdelmagid 1982; EPA 2006c; WHO 2003). Acrylamide has been shown to biodegrade in effluent from a sludge dewatering process (WHO 2003).

Enzyme-catalyzed hydrolysis is a dominant mechanism for removal of acrylamide from soils (WHO 2003). Abdelmagid (1982) showed that acrylamide is readily hydrolyzed in soils under both aerobic and anaerobic conditions, resulting in the release of NH_4^+ . The rate of decomposition in soil is influenced by temperature and incubation time. Soil type did not appear to be a factor, as acrylamide decomposed in both sandy and heavy-textured soils (Abdelmagid 1982).

Acrylamide was shown to degrade by an average of 41.5% by BOD in 14 days at 25 °C and 100 mg/L test substance, with 30 mg/L activated sludge (CERI 1999).

6.3.2.1 Air

Limited data indicate that acrylamide concentrations in the atmosphere are very low (HSDB 2009; Pratt 2000). In the atmosphere, acrylamide is susceptible to degradation via photochemically generated hydroxyl radicals and ozone. The rate constant for acrylamide's reaction with hydroxyl radicals has been estimated as 1.1×10^{-11} cm³/molecule-second, and its reaction with ozone is estimated as 1.7×10^{-18} cm³/molecule-second using a structure estimation method (Meylan and Howard 1993). Using an average atmospheric hydroxyl radical concentration of 1.5×10^6 molecules/cm³ and ozone concentration

6. POTENTIAL FOR HUMAN EXPOSURE

of 7×10^{11} molecules/cm³, estimated half-lives of approximately 12 hours (hydroxyl radical reaction) and 6.5 days (ozone reaction) can be calculated.

6.3.2.2 Water

Acrylamide is expected to be quickly degraded in water by biological processes (Abdelmagid 1982; EPA 2006c; Haberman 2002; WHO 2003). At an initial concentration of 10 ppm, acrylamide was completely degraded in about 12 days using water obtained from the Hackensack River (Cherry et al. 1956).

Acrylamide, at an initial concentration of 8 µg/L, was rapidly degraded following a lag period of approximately 9 days in well-aerated, sunlit river water obtained from the Thames River, England (Croll et al. 1974). Subsequent re-seeding of the water with acrylamide resulted in rapid degradation with little or no lag period.

6.3.2.3 Sediment and Soil

When released to land, acrylamide does not bind to soil and will thus move rapidly through the soil column, where it is expected to be quickly degraded (EPA 2006c; Haberman 2002; WHO 2003).

Acrylamide has a higher mobility and lower rate of degradation in sandy soils than in clay soils (Abdelmagid 1982; EPA 2006c; WHO 2003).

Enzyme-catalyzed hydrolysis is a dominant mechanism for removal of acrylamide from soils (WHO 2003). Abdelmagid (1982) showed that acrylamide is readily hydrolyzed in soils, producing NH₄⁺. The rate of decomposition in soil is influenced by temperature and incubation time. Soil type did not appear to be a factor, as acrylamide decomposed in both sandy and heavy-textured soils. Acrylamide, therefore, does not appear to accumulate in soils (Abdelmagid 1982).

6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT

Reliable evaluation of the potential for human exposure to acrylamide depends in part on the reliability of supporting analytical data from environmental samples and biological specimens. Concentrations of acrylamide in unpolluted atmospheres and in pristine surface waters are often so low as to be near the limits of current analytical methods. In reviewing data on acrylamide levels monitored or estimated in the environment, it should also be noted that the amount of chemical identified analytically is not necessarily equivalent to the amount that is bioavailable. The analytical methods available for monitoring acrylamide in a variety of environmental media are detailed in Chapter 7.

6. POTENTIAL FOR HUMAN EXPOSURE

6.4.1 Air

Acrylamide is not anticipated to be a common air contaminant due to its low vapor pressure and high water solubility. Limited data indicate that acrylamide concentrations in the atmosphere are very low (HSDB 2009; Pratt 2000). Air samples were collected at 25 sites throughout Minnesota over various periods of time (up to 8 years) from 1991 to 1998. Acrylamide was not identified in any of the samples (Pratt 2000). As of 1978, acrylamide concentrations in the air near six U.S. acrylamide/polyacrylamide producers or users averaged $<0.2 \mu\text{g}/\text{m}^3$ in vapor or particulate form. Concentrations ranged from <0.1 to $1.1 \mu\text{g}/\text{m}^3$ (EPA 1978).

The total national baseline NTI emissions for acrylamide during 1900–1998 were 35.4 tons/year. Total urban emissions were 33.5 tons/year, while total rural emissions were 1.9 tons per year (EPA 2000).

6.4.2 Water

Acrylamide in drinking water is typically a result of leaching during treatment processes (van Dijk-Looijaard and van Genderen 2000). Acrylamide results from the release of residual monomer from polyacrylamide coagulants that are used as clarifiers in the treatment of raw water (Abdelmagid 1982; Cavalli et al. 2004; EPA 2006c; WHO 2003). Polyacrylamides can enter water supplies due to their use as grouting agents in reservoirs and wells in addition to being released in water from plastics and dye industries (Cavalli et al. 2004; EPA 2006c).

Acrylamide concentrations of $<5 \mu\text{g}/\text{L}$ were found in river and tap water. In the sampling area, polyacrylamides were known to be used in treating potable water (WHO 2003). Public drinking water supply wells in West Virginia contained acrylamide concentrations of 0.024 – $0.041 \mu\text{g}/\text{L}$ (WHO 2003).

One water sample, taken from downstream of the effluent of a producer of polyacrylamide, contained acrylamide at $1,500 \mu\text{g}/\text{L}$, while samples from other industrial sites (acrylamide and polyacrylamide production and use locations) contained $<0.8 \mu\text{g}/\text{L}$ acrylamide (EPA 1978).

Various effluents in the United Kingdom have been found to contain acrylamide, including effluent from a clay pit ($16.0 \mu\text{g}/\text{L}$), a tailings lagoon (39 – $42 \mu\text{g}/\text{L}$), coal washing lagoon ($1.8 \mu\text{g}/\text{L}$), colliery/cooking plant, ($0.74 \mu\text{g}/\text{L}$), treated paper mill (0.47 – $14.4 \mu\text{g}/\text{L}$), and paper mill process water ($45.4 \mu\text{g}/\text{L}$) (Croll et

6. POTENTIAL FOR HUMAN EXPOSURE

al. 1974; IPCS 1985). In Devon, England, 17.4 ppb of acrylamide was detected in sewage effluent (Brown and Rhead 1979).

6.4.3 Sediment and Soil

Soil samples obtained near six U.S. acrylamide and/or polyacrylamide producers contained <0.02 µg/g of acrylamide. Concentrations ranged from <0.02 to <0.08 µg/g (EPA 1978).

6.4.4 Other Environmental Media

Acrylamide was detected in waste materials and containers at 3 of the 1,699 hazardous waste sites that have been proposed for inclusion on the EPA NPL (HazDat 2007). Acrylamide was also identified in tobacco smoke samples, obtained from nonfiltered cigarettes smoked under typically conditions. The presence of acrylamide was identified in the smoke condensate by infrared (IR), mass spectrometry (MS), and nuclear magnetic resonance (NMR) (Schumacher et al. 1977).

In 2002, acrylamide was first shown to be produced when foods are cooked at high temperatures (Tareke et al. 2002). Concentrations of acrylamide in food vary with the type of food and method of processing and cooking. Acrylamide concentrations in food typically increase with temperature and length of heating (Sorgel et al. 2002; WHO 2002, 2003). Starchy foods, such as potato-based products, typically contain the highest levels of acrylamide (Muttucumaru et al. 2008; Tareke et al. 2002; WHO 2003), whereas protein-based foods contain smaller amounts (Tareke et al. 2002).

Tareke et al. (2002) analyzed acrylamide content of various heated foods. Acrylamide concentrations were determined by gas chromatography (GC)-MS and liquid chromatography (LC)-MS/MS. In heated protein-rich foods, acrylamide concentrations of 5–50 µg/kg were found. Carbohydrate-rich foods, such as potato, beetroot, potato products, and crispbread, contained much higher concentrations, ranging from 150 to 4,000 µg/kg. The median acrylamide concentration in fried foods, including beef, chicken, soy meal, grated potatoes, boiled mashed potatoes, and grated beetroot were 17, 28, 16, 447, 172, and 850 µg/kg, respectively. In microwave-heated grated potatoes, the median acrylamide concentration was found to be 551 µg/kg, while microwave-heated cod was less than the detection limit (<5 µg/kg). In restaurant-prepared or purchased foods, the median acrylamide concentrations for hamburger, French fries, potato crisps, and three types of crispbread were 18, 424, 1,739, and 208, respectively. Unheated

6. POTENTIAL FOR HUMAN EXPOSURE

controls and boiled foods contained very little acrylamide, with all results being less than the detection limits for the methods ($<5 \mu\text{g}/\text{kg}$ by GC-MS and $<10 \mu\text{g}/\text{kg}$ by LC-MS/MS) (Tareke et al. 2002).

Acrylamide concentrations were analyzed in food samples in Norway, Sweden, Switzerland, the United Kingdom, and the United States in various studies (Table 6-2). Almost all items analyzed contained some level of acrylamide. Of the food tested, chips and crisps contained the highest average concentrations, ranging from not detectable to $3.5 \text{ mg}/\text{kg}$ (WHO 2002).

Acrylamide concentrations were determined for various foods prepared using home cooking methods, including the use of a household oven and microwave oven. The data are presented in Tables 6-3 and 6-4. The authors concluded that longer cooking times and higher temperatures appear to cause increased acrylamide concentrations in food (Sorgel et al. 2002).

The acrylamide content of various carbohydrate-rich Brazilian foods was determined by LC-MS/MS. Samples were obtained from grocery stores, restaurants, and fast food restaurants in Campinas, Sao Paulo, Brazil between September 2004 and April 2006. The foods sampled included French fries, potato chips, bread, crispbread, crackers, breakfast cereals, coffee, beer, and other high carbohydrate foods typically processed at high temperatures. Of 111 samples from 19 product categories, acrylamide concentrations ranged from <20 to $2,528 \text{ mg}/\text{kg}$. Considerable differences in concentration existed between individual foods within the same product class. Potato chips, processed at high temperatures, and instant coffee had the highest concentrations of acrylamide, while cassava- and maize-based foods, bread, and beer were found to have the lowest levels. The detection limit and limit of quantification were 10 and $20 \text{ mg}/\text{kg}$, respectively (Arisseto et al. 2007).

An ongoing U.S. Food and Drug Administration (FDA) survey collects acrylamide data on U.S. food products under the Total Diet Study (TDS). Data on approximately 280 core foods (or TDS foods) were collected to determine the nutrient and contaminant levels in the foods from each of the geographic regions in the United States (West, North Central, South, and Northeast). Food samples are collected from grocery stores and fast food restaurants from three cities in each region, prepared for consumption, and then analyzed (FDA 2009). The summary results for the 2003–2006 TDS data on acrylamide are presented in Table 6-5. Foods within certain food groups (i.e., grains/starches/baked goods) have been found to have very high levels of acrylamide. It should be noted that the high levels within the vegetable group were primarily due to potato chips and French fries.

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-2. Acrylamide Levels in Different Food and Food Product Groups from Norway, Sweden, Switzerland, the United Kingdom, and the United States

Food/product group	Acrylamide levels (µg/kg) ^a			Number of Samples
	Mean ^b	Median ²	Minimum-Maximum	
Crisps, potato/sweet potato ^c	1,312	1,343	170–2,287	38
Chips, potato ^d	537	330	<50–3,500	39
Batter based products	36	36	<30–42	2
Bakery products	112	<50	<50–450	19
Biscuits, crackers, toast, bread crisps	423	142	<30–3,200	58
Breakfast cereals	298	150	<30–1346	29
Crisps, corn	218	167	34–416	7
Bread, soft	50	30	<30–162	41
Fish and seafood products, crumbled, battered	35	35	30–39	4
Poultry or game, crumbed, battered	52	52	39–64	2
Instant malt drinks	50	50	<50–70	3
Chocolate powder	75	75	<50–100	2
Coffee powder	200	200	170–230	3
Beer	<30	<30	<30	1

^aThe limits of detection and quantification varied among laboratories; values reported as less than a value are below the limit reported by the laboratory.

^bMean and median values were calculated where individual data were available; sample sizes were extremely small, particularly for some food categories; where the mean and median are different, it reflects the skewed distribution of the underlying data that were collected in different countries and may represent different food items within the larger category.

^cProducts that are thinly sliced and fried (such as potato chips).

^dProducts that are more thickly sliced (such as French fries).

Source: WHO 2002

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-3. Acrylamide in Potato Products

Sample	Size (mm)	Cooking temperature (°C)	Cooking time (minutes)	Acrylamide concentration (µg/kg)
Potato chips	3	180	2	2,557.9
	3	140	2	36.3
	3	180	4	7,678.3
	3	140	4	53.3
	1	180	1	121.1
	1	140	2	20.0
	1	180	3	9,670.2
	1	140	4	35.3
	French fries	5	180	2
5		160	2	53.8
5		140	2	11
5		180	4	2,687.0
5		160	4	1,049.5
5		140	4	77.0
10		180	2	674.2
10		160	2	<8
10		140	2	<8
10		180	4	1,084.3
10		160	4	87.3
10		140	4	10.0
13		180	4	476.2
13		140	2	<8
13		180	6	880.0
13		140	4	<8

Source: Sorgel et al. 2002

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-4. Acrylamide Content of Food

Sample	Description	Cooking method	Cooking time (minutes)	Acrylamide concentration (µg/kg)
Walnuts		Roasted		<8
Pine seeds		Roasted		16.4
Candied chestnuts		Baked		8.3
Crème caramel		Boiled		<8
Whiskey				<8
Liver of beef		Baked		<8
Breaded fish		Baked		<8
Goose skin		Oven		<8
Goose meat		Oven		<8
Gingerbread	Lightly baked	200 °C		12.1
Gingerbread	Dark baked	200 °C		113.0
Gingerbread				180.7
Gingerbread balls				639.5
Spiced cookies				143.6
Popcorn		Microwave, 800 W	1	29.7
Popcorn		Microwave, 600 W	2	50.0
Popcorn		Microwave, 800 W	2	132.8
Popcorn		Microwave, 600 W	3	250.3
Popcorn		Microwave, 600 W	4	307.1
Croquettes		180 °C	2	354.7
Croquettes		180 °C	4	453.4
Hash browns		180 °C	2	446.1
Hash browns		180 °C	4	751.7
Fried potatoes				543.1
Croutons		Baked		27.9
Bread crust				370.6
Toast		Toasted		54.1

Source: Sorgel et al. 2002

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-5. Acrylamide Levels in Food Products Sampled for the 2003–2006 Total Diet Studies (TDS) Summary

TDS food category	Acrylamide range (ppb) for 2003	Acrylamide range (ppb) for 2004	Acrylamide range (ppb) for 2005	Acrylamide range (ppb) for 2006
Dairy	ND–16	ND	ND–16	ND–18
Eggs	ND	No data	ND	No data
Baby food	ND–267	ND–407	ND–442	ND–381
Meat/poultry/fish	ND–26	ND–27	ND–30	ND–19
Legumes	ND–93	ND–93	ND–84	ND–54
Grains/starches/ baked goods	ND–647	ND–946	ND–616	ND–470
Fruits	ND–202	109–355 ^a	ND–250	326 ^a
Vegetables	ND–536 ^b	ND–829 ^b	ND–529	ND–393
Mixtures (e.g., casseroles, sandwiches, soups, pizza)	ND–187	ND–210	ND–93	ND–55
Candy/sweets/ Sugars/syrups	ND–29	ND–42	13–50	17–32
Fats/oils	ND	No data	No data	No data
Beverages	ND–12	ND–21	ND–28	ND

^aOnly bottle prune juice was included in this category for 2004 and 2006.

^bThe highest levels in the vegetable category are for potato chips and French fries; the majority of the vegetables had low or non-detectable levels.

ND = Not detected

Source: U.S. Food and Drug Administration 2006

6. POTENTIAL FOR HUMAN EXPOSURE

Temperature, water content, food thickness, and length of heating appear to be important factors affecting the amount of acrylamide that is formed while cooking foods (Sorgel et al. 2002; WHO 2002, 2003). Reversing these factors could help minimize acrylamide formation (e.g., heating to <180 °C, making products very thick, etc.) (Sorgel et al. 2002). In many countries, food manufacturers have been requested to take steps to decrease the formation of acrylamide in their products, which has become an important area of research (Amrein et al. 2007).

With potato products, controlling reducing sugars, processing temperature, and moisture content can help limit formation of acrylamide (Amrein et al. 2007). It is thought that the amount of asparagine is also an important factor in acrylamide formation in potatoes (Muttucumaru et al. 2008). With bakery products, the amount of free asparagines as well as the type of baking agent can affect acrylamide concentration. Replacing the baking agent NH_4HCO_3 , for example, with NaHCO_3 in sweet bakery products can help reduce acrylamide content. In almonds, temperature and free asparagine content are thought to affect the amount of acrylamide formed during roasting. High concentrations of acrylamide can form during heating of olives and dried fruit (Amrein et al. 2007). Additional research is needed to determine the acrylamide reduction strategies for each type of food.

6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE

Acrylamide is a probable or confirmed carcinogen with the potential to cause nervous system damage, weakness, and incoordination of the legs following short-term exposure to high levels. Long-term effects of acrylamide include nervous system damage, paralysis, and cancer (Arisetto et al. 2007; EPA 2006c; Lewis 2000). Exposure can occur via ingestion, dermal contact, inhalation, and intraperitoneal routes (Lewis 2000; Sorgel et al. 2002). Once in the body, acrylamide is widely dispersed by body fluids. It can also cross the placental barrier (WHO 2003).

Exposure to high levels of acrylamide can result from ingestion of various foodstuffs. Acrylamide is produced when certain foods are cooked at high temperatures (Tareke et al. 2002). Concentrations of acrylamide in food vary with the type of food and method of processing and cooking, and typically increase with temperature and length of heating (Sorgel et al. 2002; WHO 2002, 2003). Foodstuffs, particularly those rich in carbohydrates, may develop high levels of acrylamide when cooked at high temperatures (Moreno Navarro et al. 2007). Potato products, including chips and other potatoes cooked at high temperatures, may comprise a large percentage of the total acrylamide intake from food. However,

6. POTENTIAL FOR HUMAN EXPOSURE

foods with lower acrylamide content but eaten at a higher frequency, such as bread, may contribute significantly to the total exposure from foods (WHO 2002).

Food was found to contribute significantly to the overall exposure of the general population to acrylamide. Short-term dietary intake of acrylamide was estimated using Monte Carlo techniques based on data from The Netherlands, United States, and Sweden. The short-term exposure estimates ranged from 0.8 to 6.0 $\mu\text{g}/\text{kg}\text{-bw}/\text{day}$ for the average consumer to the 98th percentile consumer (WHO 2002). Average estimated long-term intake of acrylamide from food sources was determined based on data from Australia, Norway, The Netherlands, Sweden, and the United States, as well as data from the International Agency for Research on Cancer (IARC). Long-term intake estimates ranged from 0.3 to 0.8 $\mu\text{g}/\text{kg}\text{-bw}/\text{day}$ (WHO 2002, 2003). Tareke et al. (2002) concluded that the acrylamide levels found in heated foods could result in a daily acrylamide intake of a few tens of micrograms.

Due to the presence of acrylamide in tobacco smoke (EU 2002; Schumacher et al. 1977), the general population can be exposed to acrylamide via inhalation (Moreno Navarro et al. 2007). This may include second-hand smoke, and is thought to be a significant source of exposure for the general population (WHO 2002). Due to its low vapor pressure and high water solubility, acrylamide is not anticipated to be a common air contaminant (WHO 2003), and thus, inhalation exposure is likely limited to tobacco smoke exposure.

Nonfood exposure to acrylamide is thought to be low for nonsmokers. Exposure can result from ingestion of polyacrylamide-treated drinking water containing residual monomer (EPA 2006c; EU 2002; WHO 2003), and from water in contact with acrylamide-containing products (e.g., grouting agents in dams) (Sorgel et al. 2002, van Dijk-Looijaard and van Genderen 2000). The MCL for acrylamide in drinking water, which specifies the concentration at which it is not expected to cause health problems, is zero (EPA 2006c). Dermal exposure can result from contact with cosmetics and toiletries containing polyacrylamides (EU 2002).

Acrylamide and its metabolite, glycidamide, can react with biomolecules such as hemoglobin. Concentrations of adducts in the blood relevant to dietary exposure can be determined, and thus, the adducts can be used as biomarkers to determine a time-averaged estimate of acrylamide exposure. Urinary biomarker use may also be possible (WHO 2002).

6. POTENTIAL FOR HUMAN EXPOSURE

Occupational exposure occurs mainly through dermal contact of acrylamide from solution, followed by inhalation of dry monomer or aerosols of solution during manufacture of acrylamide and polyacrylamide as well as during acrylamide grouting or preparation of polyacrylamide gels in the laboratory. Ingestion is a minimal concern in the workplace (EU 2002; Lewis 2000). A unique danger of acrylamide is its ability to be absorbed through unbroken skin (Lewis 2000). In the workplace, handling of dry acrylamide, such as emptying bags and drums, results in an inhalation and dermal hazard from dust (Haberman 2002).

6.6 EXPOSURES OF CHILDREN

This section focuses on exposures from conception to maturity at 18 years in humans. Differences from adults in susceptibility to hazardous substances are discussed in Section 3.7, Children's Susceptibility.

Children are not small adults. A child's exposure may differ from an adult's exposure in many ways. Children drink more fluids, eat more food, breathe more air per kilogram of body weight, and have a larger skin surface in proportion to their body volume. A child's diet often differs from that of adults. The developing human's source of nutrition changes with age: from placental nourishment to breast milk or formula to the diet of older children who eat more of certain types of foods than adults. A child's behavior and lifestyle also influence exposure. Children crawl on the floor, put things in their mouths, sometimes eat inappropriate things (such as dirt or paint chips), and spend more time outdoors. Children also are closer to the ground, and they do not use the judgment of adults to avoid hazards (NRC 1993).

Acrylamide exposure to children can occur via ingestion and inhalation. Food was found to contribute significantly to the overall exposure of the general population to acrylamide. The short-term exposure estimates ranged from 0.8 to 6.0 $\mu\text{g}/\text{kg}$ bw/day, while the estimated long-term intake of acrylamide from food sources ranged from 0.3 to 0.8 $\mu\text{g}/\text{kg}$ bw/day for adults. Exposure in children is expected to be 2–3 times that of adults on a body weight basis (WHO 2002, 2003). Children can also be exposed to acrylamide via inhalation of second-hand smoke (Moreno Navarro et al. 2007; WHO 2002).

In the body, acrylamide is widely dispersed by body fluids, and can also cross the placental barrier, exposing the fetus to high levels of acrylamide (Sorgel et al. 2002, WHO 2003). Fetuses and newborns do not have a fully developed blood-brain barrier, and thus, women who consume large amounts of acrylamide-containing foods risk significant acrylamide transfer to the fetus. Postnatally, acrylamide can be transferred via breast milk. According to Sorgel et al. (2002), when mothers consume foods with high

6. POTENTIAL FOR HUMAN EXPOSURE

acrylamide concentrations, daily consumption of 500 mL of breast milk can result in up to 10 µg of acrylamide transferred to the baby. Women who eat lesser amounts of acrylamide-containing foods, such as potato chips, may still transfer 2 µg of acrylamide to the baby. For these two scenarios, the doses for a 3 kg infant would be 3.3 and 0.66 µg/kg, respectively (Sorgel et al. 2002).

6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES

In addition to the individuals who are occupationally exposed to acrylamide (see Section 6.5), there are several groups within the general population that may receive potentially high exposures (higher than background levels) to acrylamide. These populations include individuals living in proximity to sites where acrylamide is produced or used in manufacturing or sites where acrylamide is disposed, and includes individuals living near the three NPL hazardous waste sites where acrylamide has been detected in some environmental media (HazDat 2007). Smokers and those breathing second-hand smoke are subject to high acrylamide exposures (Moreno Navarro et al. 2007). Children of mothers whose diets are high in acrylamide-containing foods can be exposed to high amounts of acrylamide through breast milk (Sorgel et al. 2002).

6.8 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylamide is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylamide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

6. POTENTIAL FOR HUMAN EXPOSURE

6.8.1 Identification of Data Needs

Physical and Chemical Properties. The physical and chemical properties of acrylamide are well understood and have been discussed in Chapter 4 (Table 4-2). There are no data needs.

Production, Import/Export, Use, Release, and Disposal. According to the Emergency Planning and Community Right-to-Know Act of 1986, 42 U.S.C. Section 11023, industries are required to submit substance release and off-site transfer information to the EPA. The TRI, which contains this information for 2007, became available in February of 2009. This database is updated yearly and should provide a list of industrial production facilities and emissions.

Methods of manufacturing, production volumes, and uses of acrylamide are available and have been discussed in Chapter 5. Demand and import/export volumes are also available. No data needs are identified.

Environmental Fate. The environmental fate of acrylamide is reasonably well understood. If released to the environment, acrylamide has a low potential to volatilize to air from water or soil surfaces; however, its low soil adsorption coefficient suggests that it has potential to leach into groundwater. Acrylamide appears to degrade fairly rapidly by biological processes in both soil (Abdelmagid 1982) and water (Cherry et al. 1956; Croll et al. 1974). If released to air, acrylamide is expected to degrade through reaction with photochemically generated hydroxyl radicals and ozone molecules. No data needs are identified.

Bioavailability from Environmental Media. No data exist regarding acrylamide's bioavailability from environmental media such as soil or drinking water. Acrylamide is produced when certain foods are cooked at high temperatures (Tareke et al. 2002). Foodstuffs, particularly those rich in carbohydrates, may develop high levels of acrylamide when cooked at high temperatures (Moreno Navarro et al. 2007).

Food Chain Bioaccumulation. Acrylamide is not expected to significantly bioconcentrate due to its high water solubility and its ability to be degraded by microorganisms (EPA 2006c; Haberman 2002; WHO 2003). Therefore, bioaccumulation through the food chain is expected to be low. No data needs are identified.

6. POTENTIAL FOR HUMAN EXPOSURE

Exposure Levels in Environmental Media. Reliable monitoring data for the levels of acrylamide in contaminated media at hazardous waste sites are needed so that the information obtained on levels of acrylamide in the environment can be used in combination with the known body burden of acrylamide to assess the potential risk of adverse health effects in populations living in the vicinity of hazardous waste sites.

Exposure Levels in Humans. Acrylamide was found to be present in various foods at high concentrations (Tareke et al. 2002). In order to better characterize the degree of exposure from food sources, further information is necessary to determine the background levels of acrylamide in both the environment and in humans. Additional information concerning methods to decrease acrylamide content in foods, and thus decrease exposure, is needed.

This information is necessary for assessing the need to conduct health studies on these populations.

Exposures of Children. Children are exposed to acrylamide by the same routes as adults. Food was found to contribute to high levels of exposure in children, possibly 2–3 times that of an adult (WHO 2002, 2003). Inhalation of second-hand smoke can also result in exposure (Moreno Navarro et al. 2007; WHO 2002). Acrylamide has been detected in breast milk, which can result in significant exposure to infants (Sorgel et al. 2002). Additional data would be useful to determine the exposures of children to acrylamide via these methods.

Child health data needs relating to susceptibility are discussed in Section 3.12.2, Identification of Data Needs: Children's Susceptibility.

Exposure Registries. No exposure registries for acrylamide were located. This substance is not currently one of the compounds for which a sub-registry has been established in the National Exposure Registry. The substance will be considered in the future when chemical selection is made for sub-registries to be established. The information that is amassed in the National Exposure Registry facilitates the epidemiological research needed to assess adverse health outcomes that may be related to exposure to this substance.

6. POTENTIAL FOR HUMAN EXPOSURE

6.8.2 Ongoing Studies

The Federal Research in Progress (FEDRIP 2009) database provides additional information obtainable from a few ongoing studies that may fill data needs identified in Section 6.8.1. Titles for these studies are listed in Table 6-6.

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-6. Ongoing Research Regarding the Environmental Fate and Exposure to Acrylamide

Investigator	Affiliation	Description	Sponsor
Lehotay SJ, Handel A	Drexel University	Development of new methods to detect and control acrylamide formation in deep-fat fried foods	USDA
Chen H	University of Delaware	Improvement of thermal and alternative processes for foods	USDA
Calder BL, Perkins LB, Bushway AA, et al.	University of Maine	Improving the post-harvest quality of fresh-cut and processed Maine potatoes	USDA
Daley LS	Oregon State University	Regulation of photosynthetic processes	USDA

USDA = U.S. Department of Agriculture

Source: FEDRIP 2009

6. POTENTIAL FOR HUMAN EXPOSURE

This page is intentionally blank.

7. ANALYTICAL METHODS

The purpose of this chapter is to describe the analytical methods that are available for detecting, measuring, and/or monitoring acrylamide, its metabolites, and other biomarkers of exposure and effect to acrylamide. The intent is not to provide an exhaustive list of analytical methods. Rather, the intention is to identify well-established methods that are used as the standard methods of analysis. Many of the analytical methods used for environmental samples are the methods approved by federal agencies and organizations such as EPA and the National Institute for Occupational Safety and Health (NIOSH). Other methods presented in this chapter are those that are approved by groups such as the Association of Official Analytical Chemists (AOAC) and the American Public Health Association (APHA). Additionally, analytical methods are included that modify previously used methods to obtain lower detection limits and/or to improve accuracy and precision.

7.1 BIOLOGICAL MATERIALS

Methods for the detection of acrylamide in biological materials are summarized in Table 7-1. Acrylamide can be detected in biological samples using gas chromatography (GC) with electron-capture detection. Acrylamide is converted to its 2,3-dibromopropionamide derivation in aqueous solution, plasma, or tissue homogenates by ionic bromination. The limits of detection for this method correspond to 9.5×10^{-12} g of acrylamide on the column or 8.4×10^{-9} g in the final biological extract of 0.5 mL. Acrylamide recovery by this method exceeds 80% at nanogram levels (HSDB 2009).

A liquid chromatography/tandem mass spectrometry (LC-MS/MS) assay was developed to examine unchanged acrylamide concentrations in bodily fluids, specifically urine, breast milk, and placental perfusion medium. A bioanalytical method was developed to analyze acrylamide in bodily fluids using LC-MS/MS following liquid/liquid extraction of acrylamide and evaporation under a stream of N_2 at 35 °C. A reversed-phase column eluted with an isocratic solvent system consisting of water, acetic acid, and an organic modifier was used for chromatographic separation. Method detection limits were 1 ng/mL for urine, 2 ng/mL for placenta perfusate, and 5 ng/mL for breast milk (Sorgel et al. 2002).

GC-MS methods have been used to determine the adducts of acrylamide and its metabolites (such as glycidamide) with hemoglobin in blood samples. These methods are sensitive enough to measure adducts

7. ANALYTICAL METHODS

Table 7-1. Analytical Methods for Determining Acrylamide in Biological Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Urine	Liquid/liquid extraction, evaporation under N ₂ stream at 35 °C	LC-MS/MS	1 ng/mL		Sorgel et al. 2002
Breast milk	Liquid/liquid extraction, evaporation under N ₂ stream at 35 °C	LC-MS/MS	5 ng/mL		Sorgel et al. 2002
Placental perfusion medium	Liquid/liquid extraction, evaporation under N ₂ stream at 35 °C	LC-MS/MS	2 ng/mL		Sorgel et al. 2002
Biological samples	Ionic bromination of acrylamide in sample	GC with electron-capture device	Corresponds to 9.5x10 ⁻¹² g acrylamide (column), 8.4x10 ⁻⁹ g (extract)		HSDB 2009

GC = gas chromatography; LC = liquid chromatography; MS = mass spectrometry

7. ANALYTICAL METHODS

at blood levels relevant to potential dietary acrylamide exposure, which allows for use of the adducts as biomarkers of exposure (WHO 2002, 2003). A more recent method of detection for hemoglobin adducts of acrylamide utilizes LC-MS/MS (Fennell et al. 2003).

Acrylamide and glycidamide were measured in serum and tissues of mice following dosing by intravenous, gavage, and dietary routes at 0.1 mg/kg acrylamide and by intravenous and gavage routes at equimolar amounts of glycidamide. LC-electrospray (ES)/MS/MS was utilized to measure acrylamide and glycidamide in serum and tissues with recoveries of 70% for acrylamide and 63% for glycidamide. The limit of detection was below 0.1 pmol/mg tissue for both acrylamide and glycidamide (Doerge et al. 2005b). Doerge et al. (2005a) detected DNA adducts derived from administration of acrylamide and glycidamide to mice and rats. A single intraperitoneal injection containing an aqueous solution of 100 μ L of 50 mg/kg acrylamide or 61 mg/kg glycamide was given to the mice. Rats were treated with a single intraperitoneal injection of the same concentration using a volume of 100 μ L per 100 g body weight. Six hours after dosing, the tissues were removed following gassing by carbon dioxide. Tissues were frozen and stored at -80 °C. DNA adducts were analyzed using a Quattro Ultima triple stage quadrupole mass spectrometer with an electrospray source, with an ion source temperature of 120 °C, desolvation gas temperature of 400 °C, and constant cone voltage of 35 V. DNA was isolated using a Blood and Cell Culture Maxi kit and adducts were quantified using LC-/MS/MS. The limit of quantification ranged from 1 to 1.5 adducts in 10^8 nucleotides and the limit of detection was 0.5 adducts in 10^8 nucleotides (Doerge et al. 2005a).

7.2 ENVIRONMENTAL SAMPLES

Methods for the detection of acrylamide in environmental samples are summarized in Table 7-2.

Acrylamide detection in water samples can be achieved using direct injection and a reversed-phase high performance liquid chromatography (HPLC)-ultraviolet (UV) absorption procedure, which has a limit of detection of 5 μ g/L (Cavalli et al. 2004). An HPLC method can be used to determine the amount of acrylamide monomer in natural and polluted aqueous environments. Acrylamide undergoes bromination and the resulting dibromopropionamide is assayed. The detection limit for the method was found to be 0.2 μ g/L in river, sea, and estuarine waters as well as potable waters, sewage, and china clay works effluents (HSDB 2009).

Solid phase extraction, extraction with activated carbon filter, and GC-MS analyses can also be used (Cavalli et al. 2004). A water sample is brominated to form 2,3-dibromopropionamide, which is then

7. ANALYTICAL METHODS

Table 7-2. Analytical Methods for Determining Acrylamide in Environmental Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Water	Direct injection	Reversed-phase HPLC-UV absorption	5 µg/L		Cavalli et al. 2004
Natural and polluted water	Bromination of acrylamide in sample	HPLC	0.20 µg/L		HSDB 2009
Organic-free reagent water		HPLC	10 µg/L		HSDB 2009
Aqueous matrices		GC	0.032 µg/L		HSDB 2009
Drinking water	Direct injection of 500 µL	Ion-exclusion chromatographic separation/MS	0.20 ppb		Cavalli et al. 2004
Airborne dust, vapors		Differential pulse polarographic method			HSDB 2009
Food	Water extract food, brominates sample	GC-MS	5–10 µg/kg		WHO 2002
Food	Bromination of sample	GC-MS	5 µg/kg		Tareke et al. 2002
Food	Water extract food	LC-MS/MS	20–50 µg/kg		WHO 2002
Food		LC-MS/MS	10 µg/kg		Tareke et al. 2002
Food		LC-MS/MS	10 µg/kg	100–115%	Arisseto et al. 2007

GC = gas chromatography; HPLC = high performance liquid chromatography; LC = liquid chromatography; MS = mass spectrometry; SIM = selected ion monitoring; UV = ultraviolet

7. ANALYTICAL METHODS

analyzed by GC-MS using methacrylamide as an internal standard (Ahn et al. 2002). Additional methods include polarography and electron capture GC (WHO 2003). EPA-OSW 8316 method determines acrylamide concentrations by HPLC in organic-free reagent water, with a limit of detection of 10 µg/L. The EPA-OSW 8032A method analyzes acrylamide in aqueous matrices using GC, with a detection limit of 0.032 µg/L (HSDB 2009).

Cavalli et al. (2004) utilized a method for detecting acrylamide in drinking water using a combination of ion-exclusion chromatographic separation and MS detection. Drinking water samples are injected directly into the microbore ICE-AS1 column and are then detected in the selected-ion monitoring mode by a single quadrupole system with electrospray ionization. The detection limit for the method was determined to be 0.20 ppb with an injection volume of 500 µL (Cavalli et al. 2004).

Airborne acrylamide vapors and dust can be determined in air by a differential pulse polarographic method (HSDB 2009). Two additional methods of detection in air were found (called Method 21 and OSHA PV2004), though only very limited details were provided (HSDB 2009).

Various methods are available to determine the acrylamide content of food. The most common methods utilize either GC-MS or LC-MS/MS (Ahn et al. 2002; Arisseto et al. 2007). Water is often used as the extraction solvent. Solid-phase extraction (SPE) is typically used to prepare samples. SPE phases that have been used for this method include graphitized carbon black, mixed mode anion and cation exchange, and polymeric materials (Arisseto et al. 2007)

The GC-MS method of determining acrylamide content in food is well established. A water extract of the food is brominated to form 2,3-dibromopropionamide, a derivative of acrylamide with enhanced GC properties. The derivative is then analyzed by GC-MS using methacrylamide as an internal standard (Ahn et al. 2002; WHO 2002). Detection limits for the GC-MS method are typically in the 5–10 µg/kg range (WHO 2002). Tareke et al. (2002) analyzed food samples for acrylamide using an improved GC-MS method involving bromination. A detection limit of 5 µg/kg was achieved.

The LC-MS/MS method was developed over concerns that artifacts were forming during the bromination procedure used with the GC-MS method. LC-MS/MS allows for direct acrylamide analysis without the need for a derivative (WHO 2002). In this method, a water extract of food is tested by LC-MS/MS using deuterated acrylamide as an internal standard (Ahn et al. 2002). Acrylamide is then identified by its retention time and the relative ion intensities. WHO (2002) reports that the detection limits of this

7. ANALYTICAL METHODS

method are typically 20–50 µg/kg, although lower detection limits have since been obtained for this method.

Tareke et al. (2002) also utilized the LC-MS/MS method to determine underivitized acrylamide concentrations. This method had a detection limit of 10 µg/kg. Sorgel et al. (2002) used a modified LC-MS/MS method, similar to that developed for analysis of acrylamide in bodily fluids, to determine acrylamide content in food. Brazilian foods were analyzed by LC-MS/MS to determine acrylamide content (Arisseto et al. 2007). Detection limits of 10 µg/kg were achieved, along with a limit of quantification of 20 µg/kg and mean recoveries ranging from 100 to 115%.

7.3 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylamide is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylamide.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

7.3.1 Identification of Data Needs

Methods for Determining Biomarkers of Exposure and Effect.

Exposure. Exposure to acrylamide can be determined directly by LC-MS/MS for biological samples, including urine, breast milk, and placental perfusion medium with detection limits of 1, 5, and 2 ng/mL, respectively (Sorgel et al. 2002). GC-MS methods are also used to detect the adducts of acrylamide and its metabolites in blood and tissue samples (HSDB 2009; WHO 2002, 2003). Methods are described in

7. ANALYTICAL METHODS

Table 7-1. Additional methods for the detection of acrylamide and its metabolites in biological samples would be helpful.

Effect. Information on biomarkers of effect for acrylamide would be useful.

Methods for Determining Parent Compounds and Degradation Products in Environmental

Media. There are several variations on methods for identifying acrylamide in water, air, and food. The methods are summarized in Table 7-2. The methods for water, primarily involving the use of HPLC and GC are fairly sensitive (e.g., ~0.03-10 µg/L for water (Cavalli et al. 2004; HSDB 2009). In air, a differential pulse polarographic method was identified, although limited details were provided (HSDB 2009). Additional information concerning the detection of acrylamide in air, including detection limits, would be useful. Various methods are available to determine acrylamide content of food using primarily GC-MS or LC-MS/MS (Ahn et al. 2002; Ariseto et al. 2007). GC-MS involved bromination, while LC-MS/MS can determine acrylamide content directly. Detection limits for GC-MS methods typically range from 5 to 10 µg/kg, while limits of detection for the LC-MS/MS method are typically in the range of 10–50 µg/kg (Tareke et al. 2002; WHO 2002). More sensitive methods for direct determination of acrylamide in food via the LC-MS/MS method would use useful.

7.3.2 Ongoing Studies

The Federal Research in Progress (FEDRIP 2009) database contains a study sponsored by the U.S. Department of Agriculture in which new methods are being developed for detecting and controlling acrylamide formation in deep-fat fried foods.

7. ANALYTICAL METHODS

This page is intentionally blank.

8. REGULATIONS, ADVISORIES, AND GUIDELINES

MRLs are substance specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors and other responders to identify contaminants and potential health effects that may be of concern at hazardous waste sites.

ATSDR has derived an acute-duration oral MRL of 0.02 mg/kg/day for acrylamide based on benchmark dose (BMD) analysis of results of fertility testing of male F344 rats administered acrylamide by gavage for 5 days prior to 1-week mating sessions with untreated female rats (Sublet et al. 1989). The resulting BMDL₁₀ of 1.78 mg/kg/day was divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability). See Appendix A for details regarding BMD analysis.

ATSDR has derived an intermediate-duration oral MRL of 0.002 mg/kg/day for acrylamide based on a NOAEL of 0.2 mg/kg/day and a LOAEL of 1 mg/kg/day for ultrastructural changes in peripheral nerve fibers in male F344 rats receiving acrylamide from the drinking water for up to 93 days (Burek et al. 1980). The NOAEL of 0.2 was divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability).

EPA (IRIS 2009) has established an oral reference dose (RfD) for acrylamide of 2×10^{-4} mg/kg/day based on a no-observed-effect-level (NOEL) of 0.2 mg/kg/day for nerve damage in male and female Fischer 344 rats exposed to acrylamide in the drinking water for 90 days (Burek et al. 1980). The uncertainty factor used in this assessment was 1,000 (10 for interspecies extrapolation, 10 for the protection of sensitive human subpopulations, and 10 for the uncertainty in the effect of duration when extrapolating from subchronic to chronic exposure).

EPA (IRIS 2009) has not derived an inhalation reference concentration (RfC) for acrylamide.

The International Agency for Research on Cancer (IARC) has classified acrylamide as a Group 2A carcinogen (probably carcinogenic to humans) (IARC 2009). The National Toxicology Program (NTP) has determined that acrylamide is reasonably anticipated to be a human carcinogen (NTP 2005). EPA has classified acrylamide as a B2 carcinogen (probable human carcinogen) (IRIS 2009). The American Conference of Governmental Industrial Hygienists (ACGIH) has classified acrylamide as an A3 carcinogen (confirmed animal carcinogen with unknown relevance to humans) (ACGIH 2008).

8. REGULATIONS AND ADVISORIES

OSHA has required employers of workers who are occupationally exposed to acrylamide to institute engineering controls and work practices to reduce and maintain employee exposure at or below permissible exposure limits (PELs) (OSHA 2009). The employer must use engineering and work practice controls to reduce exposures to not exceed 0.3 mg/m^3 for acrylamide at any time (OSHA 2009).

EPA has designated acrylamide as a hazardous air pollutant (HAP) under the Clean Air Act (CAA) (EPA 2009a). Acrylamide is on the list of chemicals appearing in "Toxic Chemicals Subject to Section 313 of the Emergency Planning and Community Right-to-Know Act of 1986" and has been assigned a reportable quantity (RQ) limit of 5,000 pounds (EPA 2009b). Acrylamide is also considered to be an extremely hazardous substance (EPA 2009c). The RQ represents the amount of a designated hazardous substance which, when released to the environment, must be reported to the appropriate authority.

The international and national regulations, advisories, and guidelines regarding acrylamide in air, water, and other media are summarized in Table 8-1.

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Acrylamide

Agency	Description	Information	Reference
<u>INTERNATIONAL</u>			
Guidelines:			
IARC	Carcinogenicity classification	Group 2A ^a	IARC 2009
WHO	Air quality guidelines	No	WHO 2000
	Drinking water quality guidelines	0.0005 mg/L ^b	WHO 2006
<u>NATIONAL</u>			
Regulations and Guidelines:			
a. Air			
ACGIH	TLV (8-hour TWA) ^{c,d}	0.03 mg/m ³	ACGIH 2008
	TLV-basis (critical effect)	Central nervous system impairment	
AIHA	ERPG values	No	AIHA 2008
EPA	Hazardous air pollutant	Yes	EPA 2009a 42 USC 7412
	Second AEGL Chemical Priority List	Yes ^e	EPA 2008
NIOSH	REL (10-hour TWA) ^f	0.03 mg/m ³	NIOSH 2005
	IDLH	60 mg/m ³	
	Potential occupational carcinogen	Yes	
OSHA	Target organs	Eyes, skin, central nervous system, peripheral nervous system, and reproductive system	OSHA 2009 29 CFR 1910.1000, Table Z-1
	PEL (8-hour TWA) for general industry ^f	0.3 mg/m ³	
b. Water			
EPA	Drinking water standards and health advisories		EPA 2006a
	1-day health advisory for a 10-kg child	1.5 mg/L	
	10-day health advisory for a 10-kg child	0.3 mg/L	
	DWEL	0.007 mg/L	
	Lifetime	No data	
	10 ⁻⁴ Cancer risk	0.0008 mg/L	

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Acrylamide

Agency	Description	Information	Reference
<u>NATIONAL</u> (cont.)			
EPA	National primary drinking water standards		EPA 2003
	Treatment technique	Yes ^g	
	Potential health effects from exposure above the MCL	Nervous system or blood problems	
	Common sources of acrylamide in drinking water	Added to water during sewage/waste water treatment; increased risk of cancer treatment	
	Public health goal	Zero	
	National recommended water quality criteria	No	EPA 2006b
c. Food			
FDA	EAFUS ^h	No	FDA 2008
d. Other			
ACGIH	Carcinogenicity classification	A3 ⁱ	ACGIH 2008
EPA	Carcinogenicity classification	B2 ^j	IRIS 2009
	RfC	Value not estimated ^k	
	RfD	2x10 ⁻⁴ mg/kg/day	
	Inhalation unit risk	1.3x10 ⁻³ (µg/m ³) ⁻¹	
	Oral slope factor	4.5 (mg/kg/day) ⁻¹	
	Superfund, emergency planning, and community right-to-know		
	Designated CERCLA hazardous substance	Yes ^l	EPA 2009b 40 CFR 302.4
	Reportable quantity	5,000 pounds	
	Effective date of toxic chemical release reporting	01/01/1987	EPA 2009d 40 CFR 372.65
	Extremely hazardous substances and its threshold planning quantity	1,000/10,000 pounds	EPA 2009c 40 CFR 355, Appendix A
	TSCA health and safety data reporting		EPA 2009e 40 CFR 716.120
	Effective date	10/04/1982	
	Sunset date	10/04/1992	

8. REGULATIONS AND ADVISORIES

Table 8-1. Regulations and Guidelines Applicable to Acrylamide

Agency	Description	Information	Reference
NATIONAL (cont.)			
NTP	Carcinogenicity classification	Reasonably anticipated to be a human carcinogen	NTP 2005

^aGroup 2A: probably carcinogenic to humans

^bFor substances that are considered to be carcinogenic, the guideline value is the concentration in drinking-water associated with an upper-bound excess lifetime cancer risk of 10^{-5} (one additional cancer per 100,000 of the population ingesting drinking-water containing the substance at the guideline value for 70 years) (WHO 2006).

^cTWA: inhalable fraction and vapor. Material exerts sufficient vapor pressure such that it may be present in both particle and vapor phases.

^dSkin: refers to the potential significant contribution to the overall exposure by the cutaneous route.

^eAcrylamide is included on the list of 371 priority chemicals that are acutely toxic and represent the selection of chemicals for AEGL development by the NAC/AEGL committee during the next several years.

^fSkin designation

^gEach water system must certify, in writing, to the state (using third-party or manufacturers certification) that when it uses acrylamide to treat water, the combination (or product) of dose and monomer level does not exceed 0.05% of acrylamide dosed at 1 mg/L (or equivalent).

^hThe EAFUS list of substances contains ingredients added directly to food that FDA has either approved as food additives or listed or affirmed as GRAS.

ⁱA3: confirmed animal carcinogen with unknown relevance to humans.

^jB2: probable human carcinogen, based on sufficient evidence of carcinogenicity in animals

^kThe health effects data for acrylamide were reviewed by the U.S. EPA RfD/RfC Work Group and determined to be inadequate for derivation of an inhalation RfC.

^lDesignated CERCLA hazardous substance pursuant to Section 112 of the Clean Air Act and Section 3001 of the Resource Conservation and Recovery Act.

ACGIH = American Conference of Governmental Industrial Hygienists; AEGL = acute exposure guideline levels; AIHA = American Industrial Hygiene Association; CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act; CFR = Code of Federal Regulations; DWEL = drinking water equivalent level; EAFUS = Everything Added to Food in the United States; EPA = Environmental Protection Agency; ERPG = emergency response planning guidelines; FDA = Food and Drug Administration; GRAS = Generally Recognized As Safe; IARC = International Agency for Research on Cancer; IDLH = immediately dangerous to life or health; IRIS = Integrated Risk Information System; MCL = maximum contaminant level; NAC = National Advisory Committee; NIOSH = National Institute for Occupational Safety and Health; NTP = National Toxicology Program; OSHA = Occupational Safety and Health Administration; PEL = permissible exposure limit; REL = recommended exposure limit; RfC = inhalation reference concentration; RfD = oral reference dose; TLV = threshold limit values; TSCA = Toxic Substances Control Act; TWA = time-weighted average; USC = United States Code; WHO = World Health Organization

8. REGULATIONS AND ADVISORIES

This page is intentionally blank.

9. REFERENCES

- *Abdelmagid HM, Tabatabai MA. 1982. Decomposition of acrylamide in soils. *J Environ Qual* 11(4):701-704.
- *Abernethy DJ, Boreiko CJ. 1987. Acrylonitrile and acrylamide fail to transform C3H/10T1/2 cells. *Environ Mutagen* 9(Suppl 8):2.
- Abou-Donia MB, Ibrahim SM, Corcoran JJ, et al. 1993. Neurotoxicity of glycidamide, an acrylamide metabolite, following intraperitoneal injections in rats. *J Toxicol Environ Health* 39(4):447-464.
- Abramsson-Zetterberg L, Vikstrom AC, Tornqvist M, et al. 2008. Differences in the frequency of micronucleated erythrocytes in humans in relation to consumption of fried carbohydrate-rich food. *Mutat Res* 653(1-2):50-56.
- *ACGIH. 2008. Acrylamide. In: *Threshold limit values for chemical substances and physical agents and biological exposure indices*. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- *Adinolfi M. 1985. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 27(4):532-537.
- *Adler ID. 1990. Clastogenic effects of acrylamide in different germ-cell stages of male mice. In: Allen B, Bridges B, Lyon M, eds. *Biology of mammalian germ cell mutagenesis*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 115-131.
- Adler ID, Baumgartner A, Friedman MA. 1999. 1-Aminobenzotriazole reduces acrylamide-induced dominant lethal effects in sperm and spermatids of mice. [Abstract]. *Environ Mol Mutagen* 33:10.
- *Adler ID, Baumgartner A, Gonda H, et al. 2000. 1-Aminobenzotriazole inhibits acrylamide-induced dominant lethal effects in spermatids of male mice. *Mutagenesis* 15(2):133-136.
- Adler ID, Gonda H, Hrabe de Angelis M, et al. 2004. Heritable translocations induced by dermal exposure of male mice to acrylamide. *Cytogenet Genome Res* 104(1-4):271-276.
- *Adler ID, Ingwersen I, Kliesch U, et al. 1988. Clastogenic effects of acrylamide in mouse bone marrow cells. *Mutat Res* 206(3):379-385.
- *Adler ID, Reitmeir P, Schmoller R, et al. 1994. Dose response for heritable translocations induced by acrylamide in spermatids of mice. *Mutat Res* 309(2):285-291.
- *Adler ID, Zouh R, Schmid E. 1993. Perturbation of cell division by acrylamide *in vitro* and *in vivo*. *Mutat Res* 301(4):249-254.

* Cited in text

+ Cited in Supplemental Document

9. REFERENCES

*Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environ Health Perspect Suppl* 103(7):103-112.

*Agency for Toxic Substances and Disease Registry. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry, Division of Toxicology. *Fed Regist* 54(174):37618-37634.

Agency for Toxic Substances and Disease Registry. 1990. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary, and immune systems. Subcommittee on Biomarkers of Organ Damage and Dysfunction. Atlanta, GA: Agency for Toxic Substances and Disease Registry.

Agrawal AK, Squibb RE. 1981. Effects of acrylamide given during gestation on dopamine receptor binding in rat pups. *Toxicol Lett* 7:233-238.

Agrawal AK, Seth PK, Squibb RE, et al. 1981. Neurotransmitter receptors in brain regions of acrylamide-treated rats: 1. Effects of a single exposure to acrylamide. *Pharmacol Biochem Behav* 14(4):527-532.

Aguas PC, Fitzhenry MJ, Giannikopoulos G, et al. 2006. Analysis of acrylamide in coffee and cocoa by isotope dilution liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem* 385(8):1526-1531.

Agut J, Font E, Sacristan A, et al. 1983. Effect of oral CDP-choline on acrylamide-induced lesion. *Arzneimittelforschung* 33(7A):1029-1033.

*Ahn JS, Castle L, Clarke DB, et al. 2002. Verification of the findings of acrylamide in heated foods. *Food Addit Contam* 19(12):1116-1124.

*AIHA. 2008. Emergency Response Planning Guidelines (ERPG). Fairfax, VA: American Industrial Hygiene Association. <http://www.aiha.org/1documents/Committees/ERP-erpglevels.pdf>. May 19, 2009.

Al Deeb S, Al Moutaery K, Arshaduddin M, et al. 2000. Attenuation of acrylamide-induced neurotoxicity in diabetic rats. *Neurotoxicol Teratol* 22(2):247-253.

Aldous CN, Farr CH, Sharma RP. 1983. Evaluation of acrylamide treatment on levels of major brain biogenic amines, their turnover rates, and metabolites. *Fundam Appl Toxicol* 3(3):182-186.

Ali SF. 1983. Acrylamide-induced changes in the monoamines and their acid metabolites in different regions of the rat brain. *Toxicol Lett* 17(1-2):101-105.

Ali SF, Hong JS, Wilson WE, et al. 1983. Effect of acrylamide on neurotransmitter metabolism and neuropeptide levels in several brain regions and upon circulating hormones. *Arch Toxicol* 52(1):35-44.

*Allen B, Zeiger E, Lawrence G, et al. 2005. Dose-response modeling of *in vivo* genotoxicity data for use in risk assessment: Some approaches illustrated by an analysis of acrylamide. *Regul Toxicol Pharmacol* 41(1):6-27.

*Altman PL, Dittmer DS. 1974. In: *Biological handbooks: Biology data book*. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies of Experimental Biology, 1987-2008, 2041.

9. REFERENCES

- +*American Cyanamid Company. 1951. Acute eye, dermal and oral toxicity. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211662. OTS206055.
- +*American Cyanamid Company. 1953a. Inhalation toxicity supplement to reports dated May 2, 1951 and August 13, 1952. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA879211664. OTS206055.
- +*American Cyanamid Company. 1953b. Subacute feeding section I-acrylamide section II-acrylamide with metabolic studies supplement to report dated August 13, 1952. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211665. OTS206055.
- +*American Cyanamid Company. 1953c. Chronic oral administration-dogs oral administration and neurohistology-dogs supplement to reports dated August 13, 1952 and May 8, 1953. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211666. OTS206055.
- +*American Cyanamid Company. 1954. Chronic inhalation exposure-acrylamide. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211670. OTS206055.
- +*American Cyanamid Company. 1959. Acrylamide toxicity: Effect of polyvinylpyrrolidone (PVP). Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211673. OTS206055.
- +*American Cyanamid Company. 1973. Range finding studies single oral single dermal (FHSA) single inhalation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211677. OTS206055.
- +*American Cyanamid Company. 1977. Limited release toxicity tests. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211678. OTS206055.
- +*American Cyanamid Company. 1979. A fetal toxicity study of acrylamide in rats. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878211679. OTS206055.
- +*American Cyanamid Company. 1991. 28-Day subchronic dose toxicity study in rats. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0533854.
- *Amrein TM, Andres L, Escher F, et al. 2007. Occurrence of acrylamide in selected foods and mitigation options. *Food Addit Contam* 24(Suppl 1):13-25.
- *Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, and replacement*. New York: Marcel Dekker Inc., 9-25.
- *Andersen ME, Clewell HJ, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87(2):185-205.
- Andersen ME, Scimeca J, Olin SS. 2005. Kinetic and mechanistic data needs for a human physiologically based pharmacokinetic (PBPK) model for acrylamide: Pharmacokinetic model for acrylamide. *Adv Exp Med Biol* 561:117-125.

9. REFERENCES

- Anderson RJ. 1981. Selective effect on peripheral nerves after subchronic administration of acrylamide. *Bull Environ Contam Toxicol* 27(6):888-893.
- Anderson RJ. 1982. Alterations in nerve and muscle compound action potentials after acute acrylamide administration. *Environ Health Perspect* 44:153-158.
- Annola K, Karttunen V, Keski-Rahkonen P, et al. 2008a. Transplacental transfer of acrylamide and glycidamide are comparable to that of antipyrine in perfused human placenta. *Toxicol Lett* 182:50-56.
- Annola K, Keski-Rahkonen P, Vahakangas K, et al. 2008b. Simultaneous determination of acrylamide, its metabolite glycidamide and antipyrine in human placental perfusion fluid and placental tissue by liquid chromatography-electrospray tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 876:191-197.
- *Ao L, Liu SX, Yang MS, et al. 2008. Acrylamide-induced molecular mutation spectra at *HPRT* locus in human promyelocytic leukemia HL-60 and NB4 cell lines. *Mutagenesis* 23(4):309-315.
- *Arisseto AP, Toledo MC, Govaert Y, et al. 2007. Determination of acrylamide levels in selected foods in Brazil. *Food Addit Contam* 24(3):236-241.
- Arocena M. 2006. Effect of acrylamide on the cytoskeleton and apoptosis of bovine lens epithelial cells. *Cell Biol Int* 30(12):1007-1012.
- Aschner M, Cao CC, Wu Q, et al. 2003. The acute effects of acrylamide on astrocyte functions. *Ann N Y Acad Sci* 993:296-304.
- Aschner M, Wu Q, Friedman MA. 2005. Effects of acrylamide on primary neonatal rat astrocyte functions. *Ann N Y Acad Sci* 1053:444-454.
- Aspengren S, Wielbass L, Wallin M. 2006. Effects of acrylamide, latrunculin, and nocodazole on intracellular transport and cytoskeletal organization in melanophores. *Cell Motil Cytoskeleton* 63(7):423-436.
- +*Augustsson K, Skog K, Jagerstad M, et al. 1999. Dietary heterocyclic amines and cancer of the colon, rectum, bladder, and kidney: A population-based study. *Lancet* 353(9154):703-707.
- *Auld RB, Bedwell SF. 1967. Peripheral neuropathy with sympathetic overactivity from industrial contact with acrylamide. *Can Med Assoc J* 96:652-654.
- Aureli F, Di Pasquale M, Lucchetti D, et al. 2007. An absorption study of dietary administered acrylamide in swine. *Food Chem Toxicol* 45(7):1202-1209.
- Axelrad DA, Morello-Frosch RA, Woodruff TJ, et al. 1999. Assessment of estimated 1990 air toxics concentrations in urban areas in the United States. *Environ Sci Policy* 2(4-5):397-411.
- +*Bachmann M, Myers JE, Bezuidenhout BN. 1992. Acrylamide monomer and peripheral neuropathy in chemical workers. *Am J Ind Med* 21(2):217-222.
- *Backer LC, Dearfield KL, Erexson GL, et al. 1989. The effects of acrylamide on mouse germ-line and somatic cell chromosomes. *Environ Mol Mutagen* 13:218-226.

9. REFERENCES

- Bailey E, Farmer PB, Bird I, et al. 1986. Monitoring exposure to acrylamide by the determination of S-(2-carboxyethyl)cysteine in hydrolyzed hemoglobin by gas chromatography-mass spectrometry. *Anal Biochem* 157(2):241-248.
- *Banerjee S, Segal A. 1986. *In vitro* transformation of C3H/10T1/2 and NIH/3T3 cells by acrylonitrile and acrylamide. *Cancer Lett* 32(3):293-304.
- *Barber DS, LoPachin RM. 2004. Proteomic analysis of acrylamide-protein adduct formation in rat brain synaptosomes. 201:120-136.
- *Barber DS, Hunt JR, Ehrich MF, et al. 2001a. Metabolism, toxicokinetics and hemoglobin adduct formation in rats following subacute and subchronic acrylamide dosing. *Neurotoxicology* 22(3):341-353.
- Barber DS, Hunt J, LoPachin RM, et al. 2001b. Determination of acrylamide and glycidamide in rat plasma by reversed-phase high performance liquid chromatography. *J Chromatogr B Biomed Sci Appl* 758(2):289-293.
- Barber DS, Stevens S, LoPachin RM. 2007. Proteomic analysis of rat striatal synaptosomes during acrylamide intoxication at a low dose rate. *Toxicol Sci* 100(1):156-167.
- *Barfknecht TR, Mecca, DJ, Naismith RW. 1988. The genotoxic activity of acrylamide. *Environ Mol Mutagen* 11(Suppl 11):9.
- *Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486.
- Basile A, Ferranti P, Moccaldi R, et al. 2008. Proteomic approach for the analysis of acrylamide-hemoglobin adducts. Perspectives for biological monitoring. *J Chromatogr A* 1215(1-2):74-81.
- *Batiste-Alentorn M, Xamena N, Creus A, et al. 1991. Genotoxicity studies with the unstable Zeste-White (UZ) System of *Drosophila melanogaster*: Results with ten carcinogenic compounds. *Environ Mol Mutagen* 18:120-125.
- Baum M, Fauth E, Fritzen S, et al. 2005. Acrylamide and glycidamide: Genotoxic effects in V79-cells and human blood. *Mutat Res* 580(1-2):61-69.
- Beiswanger CM, Mandella RD, Graessle TR, et al. 1993. Synergistic neurotoxic effects of styrene oxide and acrylamide: Glutathione-independent necrosis of cerebellar granule cells. *Toxicol Appl Pharmacol* 118(2):233-244.
- Benn TM, Thomas SM. 1997. Acrylamide induces oxidative damage to DNA in *in vitro* neuronal cultures. [Abstract]. *Hum Exp Toxicol* 16(7):389.
- *Berger GS, ed. 1994. Epidemiology of endometriosis. In: *Endometriosis: Modern surgical management of endometriosis*. New York, NY: Springer-Verlag, 3-7.
- *Bergmark E. 1997. Hemoglobin adducts of acrylamide and acrylonitrile in laboratory workers, smokers and nonsmokers. *Chem Res Toxicol* 10(1):78-84.

9. REFERENCES

- *Bergmark E, Calleman CJ, Costa LG. 1991. Formation of hemoglobin adducts of acrylamide and its epoxide metabolite glycidamide in the rat. *Toxicol Appl Pharmacol* 111(2):352-363.
- *Bergmark E, Calleman CJ, He F, et al. 1993. Determination of hemoglobin adducts in humans occupationally exposed to acrylamide. *Toxicol Appl Pharmacol* 120(1):45-54.
- Bermudo E, Nunez O, Moyano E, et al. 2007. Field amplified sample injection-capillary electrophoresis-tandem mass spectrometry for the analysis of acrylamide in foodstuffs. *J Chromatogr A* 1159(1-2):225-232.
- Berti-Mattera LN, Eichberg J, Schrama L, et al. 1990. Acrylamide administration alters protein phosphorylation and phospholipid metabolism in rat sciatic nerve. *Toxicol Appl Pharmacol* 103(3):502-511.
- Besaratinia A, Pfeifer GP. 2003. Weak yet distinct mutagenicity of acrylamide in mammalian cells. *J Natl Cancer Inst* 95(12):889-896.
- Besaratinia A, Pfeifer GP. 2004a. Genotoxicity of acrylamide and glycidamide. *J Natl Cancer Inst* 96(13):1023-1029.
- *Besaratinia A, Pfeifer GP. 2004b. Genotoxicity of acrylamide in humans and mice: Promutagenic glycidamide-DNA adducts in the human p53 gene and the cII transgene. *Proc Am Assoc Cancer Res* 45:452. <http://www.aacrmeetingabstracts.org/cgi/content/abstract/2004/1/452-b>. July 13, 2009.
- Besaratinia A, Pfeifer GP. 2005. DNA adduction and mutagenic properties of acrylamide. *Mutat Res* 580(1-2):31-40.
- *Besaratinia A, Pfeifer GP. 2007. A review of mechanisms of acrylamide carcinogenicity. *Carcinogenesis* 28(3):519-528.
- Beyer DJ, Belsito DV. 2000. Allergic contact dermatitis from acrylamide in a chemical mixer. *Contact Dermatitis* 42(3):181-182.
- Bikales NM. 1973. Preparation of acrylamide polymers. In: Bikales NM, ed. *Water-soluble polymers, polymer science and technology series*. Vol. 2. New York, NY: Plenum Press, 213-225.
- Bishop JB, Chapin RE, Fail P, et al. 1991. Acrylamide induced dominant lethality in mice following low dose chronic administration in drinking water. [Abstract]. *Environ Mol Mutagen* 17(Suppl 19):11.
- Bjellaas T, Olstorn HB, Becher G, et al. 2007c. Urinary metabolites as biomarkers of acrylamide exposure in mice following dietary crisp bread administration or subcutaneous injection. *Toxicol Sci* 100(2):374-380.
- Bjellaas T, Olesen PT, Frandsen H, et al. 2007b. Comparison of estimated dietary intake of acrylamide with hemoglobin adducts of acrylamide and glycidamide. *Toxicol Sci* 98(1):110-117.
- *Bjellaas T, Stolen LH, Haugen M, et al. 2007a. Urinary acrylamide metabolites as biomarkers for short-term dietary exposure to acrylamide. *Food Chem Toxicol* 45(6):1020-1026.

9. REFERENCES

- Blank I. 2005. Current status of acrylamide research in food: measurement, safety assessment, and formation. *Ann N Y Acad Sci* 1043:30-40.
- Blasiak J, Gloc E, Wozniak K, et al. 2004. Genotoxicity of acrylamide in human lymphocytes. *Chem Biol Interact* 149(2-3):137-149.
- *Boettcher MI, Angerer J. 2005. Determination of the major mercapturic acids of acrylamide and glycidamide in human urine by LC-ESI-MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 824(1-2):283-294.
- *Boettcher MI, Bolt HM, Angerer J. 2006b. Acrylamide exposure via the diet: Influence of fasting on urinary mercapturic acid metabolite excretion in humans. *Toxicol* 80:817-819.
- *Boettcher MI, Bolt HM, Drexler H, et al. 2006a. Excretion of mercapturic acids of acrylamide and glycidamide in human urine after single oral administration of deuterium-labelled acrylamide. *Arch Toxicol* 80(2):55-61.
- *Boettcher MI, Schettgen T, Kutting B, et al. 2005. Mercapturic acids of acrylamide and glycidamide as biomarkers of the internal exposure to acrylamide in the general population. *Mutat Res* 580(1-2):167-176.
- Bolger PM. 2005. Overview of dietary acrylamide. [Abstract]. *Toxicol Sci* 84(1-S):60.
- Bondy SC, Tilson HA, Agrawal AK. 1981. Neurotransmitter receptors in brain regions of acrylamide-treated rats: 2. Effects of extended exposure to acrylamide. *Pharmacol Biochem Behav* 14(4):533-538.
- Boon PE, de Mul A, van der Voet H, et al. 2005. Calculations of dietary exposure to acrylamide. *Mutat Res* 580(1-2):143-155.
- Borenfreund E, Babich H. 1987. In-vitro cytotoxicity of heavy metals acrylamide and organotin salts to neural cells and fibroblasts. *Cell Biol Toxicol* 3(1):63-73.
- Bowyer JF, Latendresse JR, Delongchamp RR, et al. 2008a. The effects of subchronic acrylamide exposure on gene expression, neurochemistry, hormones, and histopathology in the hypothalamus-pituitary-thyroid axis of male Fischer 344 rats (Erratum in: *Toxicol Appl Pharmacol* 232:498). *Toxicol Appl Pharmacol* 230(2):208-215.
- Bowyer JF, Latendresse JR, Delongchamp RR, et al. 2008b. Corrigendum to "The effects of subchronic acrylamide exposure on gene expression, neurochemistry, hormones, and histopathology in the hypothalamus-pituitary-thyroid axis of male Fischer 344 rats" (Erratum to: *Toxicol Appl Pharmacol* 230:208-215). *Toxicol Appl Pharmacol* 232:498.
- Brady ST, Pfister KK, Bloom GS. 1990. A monoclonal antibody against kinesin inhibits both anterograde and retrograde fast axonal transport in squid axoplasm. *Neurobiology* 87:1061-1065.
- Brantsaeter AL, Haugen M, Mul A, et al. 2008. Exploration of different methods to assess dietary acrylamide exposure in pregnant women participating in the Norwegian Mother and Child Cohort Study (MoBa). *Food Chem Toxicol* 46(8):2808-2814.

9. REFERENCES

- Brat DJ, Brimijoin S. 1993. Acrylamide and glycidamide impair neurite outgrowth in differentiating N1E.115 neuroblastoma without disturbing rapid bidirectional transport of organelles observed by video microscopy. *J Neurochem* 60(6):2145-2152.
- Brimijoin WS, Hammond PI. 1985. Acrylamide neuropathy in the rat: Effects on energy metabolism in sciatic nerve. *Mayo Clin Proc* 60(1):3-8.
- Brismar T, Hildebrand C, Tegner R. 1987. Nodes of Ranvier in acrylamide neuropathy: Voltage clamp and electron microscopic analysis of rat sciatic nerve fibres at proximal levels. *Brain Res* 423(1-2):135-143.
- *Brown L, Rhead M. 1979. Liquid chromatographic determination of acrylamide monomer in natural and polluted aqueous environments. *Analyst* 104:391-399.
- *Brown RP, Delp MD, Lindstedt SL, et al. 1997. Physiological parameter values for physiologically based pharmacokinetic models. *Toxicol Indl Health* 13(4):407-484.
- *Bull RJ, Robinson M, Laurie RD, et al. 1984a. Carcinogenic effects of acrylamide in Sencar and A/J mice. *Cancer Res* 44(1):107-111.
- *Bull RJ, Robinson M, Stober JA. 1984b. Carcinogenic activity of acrylamide in the skin and lung of Swiss-ICR mice. *Cancer Lett* 24(2):209-212.
- +*Burek JD, Albee RR, Beyer JE, et al. 1980. Subchronic toxicity of acrylamide administered to rats in the drinking water followed by up to 144 days of recovery. *J Environ Pathol Toxicol* 4(5-6):157-182.
- *Butterworth BE, Eldridge SR, Sprankle CS, et al. 1992. Tissue-specific genotoxic effects of acrylamide and acrylonitrile. *Environ Mol Mutagen* 20(3):148-155.
- *Calleman CJ. 1996. The metabolism and pharmacokinetics of acrylamide: Implications for mechanisms of toxicity and human risk estimation. *Drug Metab Rev* 28(4):527-590.
- Calleman CJ, Bergmark E, Costa LG. 1990. Acrylamide is metabolized to glycidamide in the rat: Evidence from hemoglobin adduct formation. *Chem Res Toxicol* 3(5):406-412.
- Calleman CJ, Bergmark E, He F, et al. 1993a. Determination of hemoglobin adducts and adverse health effects in workers occupationally exposed to acrylamide. *J Toxicol Environ Health* 40(2-3):447-448.
- *Calleman CJ, Bergmark E, Stern LG, et al. 1993b. A nonlinear dosimetric model for hemoglobin adduct formation by the neurotoxic agent acrylamide and its genotoxic metabolite glycidamide. *Environ Health Perspect* 99:221-223.
- *Calleman CJ, Stern LG, Bergmark E, et al. 1992. Linear versus nonlinear models for hemoglobin adduct formation by acrylamide and its metabolite glycidamide: Implications for risk estimation. *Cancer Epidemiol Biomarkers Prev* 1(5):361-368.
- +*Calleman CJ, Wu Y, He F, et al. 1994. Relationship between biomarkers of exposure and neurological effects in a group of workers exposed to acrylamide. *Toxicol Appl Pharmacol* 126(2):361-371.

9. REFERENCES

- Cao J, Liu Y, Jia L, et al. 2008. Curcumin attenuates acrylamide-induced cytotoxicity and genotoxicity in HepG2 cells by ROS scavenging. *J Agric Food Chem* 56(24):12059-12063.
- Carere A. 2006. Genotoxicity and carcinogenicity of acrylamide: A critical review. *Ann Ist Super Sanita* 42(2):144-155.
- Carlson GP, Weaver PM. 1985. Distribution and binding of [¹⁴C]acrylamide to macromolecules in SENCAR and BALB/c mice following oral and topical administration. *Toxicol Appl Pharmacol* 79(2):307-313.
- Castle L, Eriksson S. 2005. Analytical methods used to measure acrylamide concentrations in foods. *J AOAC Int* 88(1):274-284.
- *Cavalli S, Polesello S, Sacconi G. 2004. Determination of acrylamide in drinking water by large-volume direct injection and ion-exclusion chromatography-mass spectrometry. *J Chromatogr A* 1039(1-2):155-159.
- *Cavanagh JB. 1964. The significance of the "dying back" process in experimental and human neurological disease. *Int Rev Exp Pathol* 3:219-267
- Cavanagh JB. 1982. The pathokinetics of acrylamide intoxication: A reassessment of the problem. *Neuropathol Appl Neurobiol* 8(4):315-336.
- Cavanagh JB, Gysbers MF. 1980. Dying back above a nerve ligature produced by acrylamide. *Acta Neuropathol* 51(3):169-178.
- Cavanagh JB, Gysbers MF. 1983. Ultrastructural features of the Purkinje cell damage caused by acrylamide in the rat: A new phenomenon in cellular neuropathology. *J Neurocytol* 12(3):413-438.
- *CERI. 1999. Biodegradation database and estimation. Acrylamide. Chemicals Evaluation and Research Institute, Japan. <http://qsar.cerij.or.jp/cgi-bin/QSAR/index.cgi?e>. June 8, 2009.
- +*Chapin RE, Fail PA, George JD, et al. 1995. The reproductive and neural toxicities of acrylamide and three analogues in Swiss mice, evaluated using the continuous breeding protocol. *Fundam Appl Toxicol* 27(1):9-24.
- Chauhan NB, Spencer PS, Sabri MI. 1993. Acrylamide-induced depletion of microtubule-associated proteins (MAP1 and MAP2) in the rat extrapyramidal system. *Brain Res* 602(1):111-118.
- *Cherry AB, Gabaccia AJ, Senn HW. 1956. The assimilation behavior of certain toxic organic compounds in natural water. *Sewage Ind Waste* 28(9):1137-1146.
- Chevolleau S, Jacques C, Canlet C, et al. 2007. Analysis of hemoglobin adducts of acrylamide and glycidamide by liquid chromatography-electrospray ionization tandem mass spectrometry, as exposure biomarkers in French population. *J Chromatogr A* 1167(2):125-134.
- Chico Galdo V, Massart C, Jin L, et al. 2006. Acrylamide, an *in vivo* thyroid carcinogenic agent, induces DNA damage in rat thyroid cell lines and primary cultures. *Mol Cell Endocrinol* 257-258:6-14.

9. REFERENCES

- Chretien M, Patey G, Souyri F, et al. 1981. Acrylamide-induced neuropathy and impairment of axonal transport of proteins: 2. Abnormal accumulations of smooth endoplasmic reticulum as sites of focal retention of fast transported proteins: Electron microscope radioautographic study. *Brain Res* 205(1):15-28.
- Chu S, Metcalfe CD. 2007. Analysis of acrylamide in water using a coevaporation preparative step and isotope dilution liquid chromatography tandem mass spectrometry. *Anal Chem* 79(13):5093-5096.
- *Cihak R, Vontorkova M. 1988. Cytogenetic effects of acrylamide in the bone marrow of mice. *Mutat Res* 209(1-2):91-94.
- *Cihak R, Vontorkova M. 1990. Activity of acrylamide in single-, double-, and triple-dose mouse bone marrow micronucleus assays. *Mutat Res* 234:125-127.
- Clarke CH, Sickles DW. 1996. Decreased GAP-43 accumulation in neurite tips of cultured hippocampal neurons by acrylamide. *Neurotoxicology* 17(2):397-406.
- Clay P. 1991. Procarbazine and acrylamide induce small colony tk^{-/-} mutants in a microwell 15178y mouse lymphoma mutation assay. [Abstract]. *Mutagenesis* 6:432.
- Clement FC, Dip R, Naegeli H. 2007. Expression profile of human cells in culture exposed to glycidamide, a reactive metabolite of the heat-induced food carcinogen acrylamide. *Toxicology* 240(1-2):111-124.
- *Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- *CMR. 2008. Chemical profile: Acrylamide. *Chem Market Rep* July 21-27
- *Collins BW, Howard DR, Allen JW. 1992. Kinetochores-staining of spermatid micronuclei: Studies of mice treated with X-radiation or acrylamide. *Mutat Res* 281(4):287-294.
- +*Collins JJ, Swaen GMH, Marsh GM, et al. 1989. Mortality patterns among workers exposed to acrylamide. *J Occup Med* 31(7):614-617.
- Comblath DR. 2004. Neuropathy in grout workers. (Letter and author response to *Scand J Work Environ Health* 30(1):21-29). *Scand J Work Environ Health* 30(3):253-254.
- Costa LG, Deng H, Calleman CJ, et al. 1995. Evaluation of the neurotoxicity of glycidamide, an epoxide metabolite of acrylamide: Behavioral, neurochemical and morphological studies. *Toxicology* 98(1-3):151-161.
- Costa LG, Deng H, Gregotti C, et al. 1992. Comparative studies on the neuro- and reproductive toxicity of acrylamide and its epoxide metabolite glycidamide in the rat. *Neurotoxicology* 13:219-224.
- *Crofton KM, Padilla S, Tilson HA, et al. 1996. The impact of dose rate on the neurotoxicity of acrylamide: The interaction of administered dose, target tissue concentrations, tissue damage, and functional effects. *Toxicol Appl Pharmacol* 139(1):163-176.
- *Croll BT, Arkell GM, Hodge RP. 1974. Residues of acrylamide in water. *Water Res* 8(11):989-993.

9. REFERENCES

- *Currance PL, Clements B, Bronstein AC. 2007. Tri-ortho-cresyl phosphate (TOCP) and related compounds. In: Emergency care for hazardous materials exposure. 3rd ed. St. Louis, MO: MosbyJems, 482-484.
- Damjanov I, Friedman MA. 1998. Mesotheliomas of tunica vaginalis testis of Fischer 344 (F344) rats treated with acrylamide: A light and electron microscopy study. *In Vivo* 12(5):495-502.
- Das M, Mukhtar H, Seth PK. 1982. Effect of acrylamide on brain and hepatic mixed-function oxidases and glutathione-S-transferase in rats. *Toxicol Appl Pharmacol* 66(3):420-424.
- *Davenport JG, Farrell DF, Sumi SM. 1976. 'Giant axonal neuropathy' caused by industrial chemicals: Neurofilamentous axonal masses in man. *Neurology* 26(10):919-923.
- *Dearfield KL, Douglas GR, Ehling UH, et al. 1995. Acrylamide: A review of its genotoxicity and an assessment of heritable genetic risk. *Mutat Res* 330(1-2):71-99.
- Degrandchamp RL, Lowndes HE. 1990. Early degeneration and sprouting at the rat neuromuscular junction following acrylamide administration. *Neuropathol Appl Neurobiol* 16(3):239-254.
- DeGrandchamp RL, Reuhl KR, Lowndes HE. 1990. Synaptic terminal degeneration and remodeling at the rat neuromuscular junction resulting from a single exposure to acrylamide. *Toxicol Appl Pharmacol* 105(3):422-433.
- Deng H, He F, Zhang S, et al. 1993. Quantitative measurements of vibration threshold in healthy adults and acrylamide workers. *Int Arch Occup Environ Health* 65(1):53-56.
- De Rojas TC, Goldstein BD. 1987. Primary afferent terminal function following acrylamide: Alterations in the dorsal root potential and reflex. *Toxicol Appl Pharmacol* 88(2):175-182.
- *DFG. 2009. Acrylamide. In: Greim H, ed. The MAK-collection for occupational health and safety. Part I: MAK Value Documentations. Vol. 25. Weinheim, Germany: Deutsche Forschungsgemeinschaft. Wiley-VCH Verlag, 1-43.
- Diembeck W, Dusing HJ, Akhiana M. 1998. Dermal absorption and penetration of acrylamide ([¹⁴C] acrylamide as tracer) in different cosmetic formulations and polyacrylamide-solution after topical application to excised pig skin. *Beiersdorf* 9.
- Dixit R, Das M, Seth PK, et al. 1984. Interaction of acrylamide with glutathione in rat erythrocytes. *Toxicol Lett* 23(3):291-298.
- +*Dixit R, Husain R, Mukhtar H, et al. 1981. Effect of acrylamide on biogenic amine levels, monoamine oxidase, and cathepsin D activity of rat brain. *Environ Res* 26(1):168-173.
- Dixit R, Husain R, Seth PK, et al. 1980a. Effect of diethyl maleate on acrylamide-induced neuropathy in rats. *Toxicol Lett* 6(6):417-422.
- Dixit R, Mukhtar H, Seth PK, et al. 1980b. Binding of acrylamide with glutathione s transferases. *Chem Biol Interact* 32(3):353-359.

9. REFERENCES

- *Dixit R, Seth PK, Mukhtar H. 1982. Metabolism of acrylamide into urinary mercapturic-acid and cysteine conjugates in rats. *Drug Metab Dispos* 10(2):196-197.
- *Dobrzynska MM. 2007. Assessment of DNA damage in multiple organs from mice exposed to X-rays or acrylamide or a combination of both using the comet assay. *In Vivo* 21(4):657-662.
- *Doerge DR, da Costa GG, McDaniel LP, et al. 2005a. DNA adducts derived from administration of acrylamide and glycidamide to mice and rats. *Mutat Res* 580(1-2):131-141.
- *Doerge DR, Twaddle NC, Boettcher MI, et al. 2007. Urinary excretion of acrylamide and metabolites in Fischer 344 rats and B6C3F₁ mice administered a single dose of acrylamide. *Toxicol Lett* 169(1):34-42.
- *Doerge DR, Young JF, McDaniel LP, et al. 2005b. Toxicokinetics of acrylamide and glycidamide in B6C3F₁ mice. *Toxicol Appl Pharmacol* 202(3):258-267.
- *Doerge DR, Young JF, McDaniel LP, et al. 2005c. Toxicokinetics of acrylamide and glycidamide in Fischer 344 rats. *Toxicol Appl Pharmacol* 208(3):199-209.
- Doerge DR, Young JF, Chen JJ, et al. 2008. Using dietary exposure and physiologically based pharmacokinetic/pharmacodynamic modeling in human risk extrapolations for acrylamide toxicity. *J Agric Food Chem* 56(15):6031-6038.
- Doerge DR, Young JF, McDaniel L, et al. 2005d. Toxicokinetics of acrylamide and glycidamide in B6C3F₁ mice and Fischer 344 rats. [Abstract]. *Toxicol Sci* 84(1-S):60.
- *Donovan JW, Pearson T. 1987. Ingestion of acrylamide with severe encephalopathy neurotoxicity and hepatotoxicity. [Abstract]. *Vet Hum Toxicol* 29(6):462.
- Doroshenko O, Fuhr U, Kunz D, et al. 2009. *In vivo* role of cytochrome P450 2E1 and glutathione-S-transferase activity for acrylamide toxicokinetics in humans. *Cancer Epidemiol Biomarkers Prev* 18(2):433-443.
- *Dourson M, Hertzberg R, Allen B, et al. 2008. Evidence-based dose-response assessment for thyroid tumorigenesis from acrylamide. *Regul Toxicol Pharmacol* 52(3):264-289.
- +*Dow Chemical Company. 1957. Results of toxicological tests on acrylamide with cover letter. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878210955. OTS206135.
- +*Dow Chemical Company. 1981. Effects of acrylamide monomer on sensory thresholds in monkeys Sept. 15, 1979 to Sept. 18, 1981. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878210964. OTS206135.
- *D'Souza RW, Francis WR, Andersen ME. 1988. Physiological model for tissue glutathione depletion and increased resynthesis after ethylene dichloride exposure. *J Pharmacol Expl Ther* 245:563-568.
- Duale N, Bjellaas T, Alexander J, et al. 2009. Biomarkers of human exposure to acrylamide and relation to polymorphisms in metabolizing genes. *Toxicol Sci* 108(1):90-99.

9. REFERENCES

- *Dumitru DL. 1989. Occupational intoxication with acrylamide: Discussion of cases with acute and chronic intoxication. *Rev Ig Bacteriol Virusol Parazitol Epidemiol Pneumoftiziol Pneumo* 38(4):359-364.
- Dumont J, Van Sande J, Massart C, et al. 2005. Study *in vitro* of the tumorigenic action of acrylamide on thyroid cells. [Abstract]. *Toxicol Sci* 84(1-S):142.
- Durling LJ, Abramsson-Zetterberg L. 2005. A comparison of genotoxicity between three common heterocyclic amines and acrylamide. *Mutat Res* 580(1-2):103-110.
- Dybing E, Sanner T. 2003. Risk assessment of acrylamide in foods. *Toxicol Sci* 75(1):7-15.
- Dybing E, Farmer PB, Andersen M, et al. 2005. Human exposure and internal dose assessments of acrylamide in food. *Food Chem Toxicol* 43(3):365-410.
- *Edwards PM. 1975. The distribution and metabolism of acrylamide and its neurotoxic analogues in rats. *Biochem Pharmacol* 24(20):1277-1282.
- Edwards PM. 1976. Insensitivity of the developing rat fetus to the toxic effects of acrylamide. *Chem Biol Interact* 12:13-18.
- Edwards PM, Sporel-Ozokat RE, Gispens WH. 1991. Peripheral pain fiber function is relatively insensitive to the neurotoxic actions of acrylamide in the rat. *Toxicol Appl Pharmacol* 111(1):43-48.
- *Ehling UH, Neuhaeuser-Klaus A. 1992. Reevaluation of the induction of specific-locus mutations in spermatogonia of the mouse by acrylamide. *Mutat Res* 283(3):185-191.
- El-Din AY, Al-Maskati HA, Mohamed AY, et al. 1993. Acrylamide as an inducer of metabolic activation system (S9) in rats. *Mutat Res* 300(2):91-97.
- Eldridge SR, Sprankle CS, Butterworth BE. 1992. Tissue-specific genotoxic effects of acrylamide and glycidamide. [Abstract]. *Environ Mol Mutagen* 19(20):17.
- El-Hadri L, Ghanayem BI. 2004. Comparative metabolism and disposition of 1-¹⁴C- and 2, 3-¹⁴C-acrylamide in cytochrome P450 2E1-null (KO) and wild-type (WT) mice. [Abstract]. *Toxicologist* 78(1-S):297.
- *Environ. 2002. Toxicological review of acrylamide (CAS No. 79-06-01). Ruston, Louisiana: Environ International Corporation.
- EPA. 1976. Investigation of selected potential environmental contaminants: Acrylamides. Final report. Washington, DC: U.S. Environmental Protection Agency. EPA560276008.
- *EPA. 1978. Environmental monitoring near industrial sites acrylamide. Washington, DC: U.S. Environmental Protection Agency. EPA560678001. PB281879.
- *EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency Office of Health and Environmental Assessment, Office of Research and Development, Environmental Criteria and Assessment Office.

9. REFERENCES

- *EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency Risk Assessment Forum.
- *EPA. 1998. Automated Form R for Windows: User's guide (RY97). Washington, DC: U.S. Environmental Protection Agency Office of Pollution Prevention and Toxics.
- *EPA. 2000. National air pollutant emission trends, 1900-1998. Research Triangle Park, NC: U. S. Environmental Protection Agency. EPA454R00002. <http://www.epa.gov/ttn/chieftrends/trends98/trends98.pdf>. June 8, 2009.
- *EPA. 2003. National primary drinking water regulations. Washington, DC: Office of Ground Water and Drinking Water, U.S. Environmental Protection Agency. <http://www.epa.gov/safewater/contaminants/index.html>. May 19, 2009.
- *EPA. 2005a. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency.
- *EPA. 2005b. Supplemental guidance for assessing susceptibility from early life exposure to carcinogens. Risk assessment forum. Washington, DC: U.S. Environmental Protection Agency. EPA630R03003F.
- *EPA. 2006a. Drinking water standards and health advisories. Washington, DC: Office of Water, U.S. Environmental Protection Agency. <http://epa.gov/waterscience/criteria/drinking/>. May 19, 2009.
- *EPA. 2006b. National recommended water quality criteria. Washington, DC: Office of Water, Office of Science and Technology, U.S. Environmental Protection Agency. <http://www.epa.gov/waterscience/criteria/wqcriteria.html>. May 11, 2009.
- *EPA. 2008. Acute exposure guideline levels (AEGLs). Second AEGL Chemical Priority List. Washington, DC: Office of Pollution Prevention and Toxics, U.S. Environmental Protection Agency. http://www.epa.gov/oppt/aegl/pubs/priority_2.htm. May 19, 2009.
- *EPA. 2009a. Hazardous air pollutants. Clean Air Act. U.S. Environmental Protection Agency. United States Code. 42 USC 7412. <http://www.epa.gov/ttn/atw/orig189.html>. May 19, 2009.
- *EPA. 2009b. Superfund, emergency planning, and community right-to-know programs. Designation, reportable quantities, and notifications. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 20, 2009.
- *EPA. 2009c. Superfund, emergency planning, and community right-to-know programs. Extremely hazardous substances and their threshold planning quantities. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 355, Appendix A. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 20, 2009.
- *EPA. 2009d. Superfund, Emergency Planning, And Community Right-To-Know Programs. Toxic Chemical Release Reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372.65. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 11, 2009.

9. REFERENCES

- *EPA. 2009e. Toxic substances control act. Health and safety data reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 716.120. <http://www.epa.gov/lawsregs/search/40cfr.html>. May 20, 2009.
- *EPA. 2009f. 2005 National emissions inventory data and documentation. National emissions inventory. U.S. Environmental Protection Agency. <http://www.epa.gov/ttn/chief/net/2005inventory.html>. May 26, 2009.
- Ericsson AC, Walum E. 1984. Cytotoxicity of cyclophosphamide and acrylamide in glioma and neuroblastoma cell lines cocultured with liver cells. *Toxicol Lett* 20(3):251-256.
- +*Eskin TA, Lapham LW, Maurissen JP, et al. 1985. Acrylamide effects on the macaque visual system. II. Retinogeniculate morphology. *Invest Ophthalmol Vis Sci* 26(3):317-329.
- *EU. 2002. Opinion of the scientific committee on food on new findings regarding the presence of acrylamide in food. Brussels: European Union. <http://europa.eu.int/comm/food/fs/sc/scf/index.en.html>. June 3, 2009.
- Evans HL, Teal JJ. 1981. Appetitive behaviors as models of the neuro toxicity of acrylamide. [Abstract]. *Fed Proc* 40(3 Part 1):677.
- Exon JH. 2006. A review of the toxicology of acrylamide. *J Toxicol Environ Health B Crit Rev* 9:397-412.
- Fail PA, George JD, Grizzle TB, et al. 1993. Dominant lethality of acrylamide and three of its congeners after chronic exposure in Swiss mice. [Abstract]. *Environ Mol Mutagen* 21(Suppl 22):19.
- Fang M, Boobis AR, Edwards RJ. 2007. Searching for novel biomarkers of centrally and peripherally-acting neurotoxicants, using surface-enhanced laser desorption/ionisation-time-of-flight mass spectrometry (SELDI-TOF MS). *Food Chem Toxicol* 45(11):2126-2137.
- Favor J, Shelby MD. 2005. Transmitted mutational events induced in mouse germ cells following acrylamide or glycidamide exposure. *Mutat Res* 580(1-2):21-30.
- Favor J, Layton D, Sega G, et al. 1995. Genetic risk extrapolation from animal data to human disease. *Mutat Res* 330:23-34.
- *FDA. 2008. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://vm.cfsan.fda.gov/~dms/eafus.html>. May 19, 2009.
- *FDA. 2009. Survey data on acrylamide in food: Total diet study results. U.S. Food and Drug Administration. <http://www.fda.gov/Food/FoodSafety/FoodContaminantsAdulteration/ChemicalContaminants/Acrylamide/ucm053566.htm>. June 8, 2009.
- *FEDRIP. 2009. Acrylamide. Federal Research in Progress database. Springfield, VA: National Technical Information Service.
- Feng Y, Forgac M. 1992. Cysteine 254 of the 73-kDa A subunit is responsible for inhibition of the coated vesicle (H⁺)-ATPase upon modification by sulfhydryl reagents. *J Biol Chem* 267(9):5817-5822.

9. REFERENCES

- Fennell TR, Friedman MA. 2005. Comparison of acrylamide metabolism in humans and rodents. In: Friedman M, Mottram D, eds. Chemistry and safety of acrylamide in food. Vol. 561. New York, NY: Springer, 109-116.
- *Fennell T, Snyder R, Burgess JP, et al. 2004. Metabolism and hemoglobin adducts of [1, 2, 3-¹³C₃] acrylamide in humans. [Abstract]. Toxicologist 78(1-S):173-174.
- *Fennell T, Snyder RW, Friedman MA. 2005a. DNA and hemoglobin adducts from oral administration of acrylamide to male Fischer 344 rats. [Abstract]. Toxicol Sci 84(1-S):103.
- *Fennell T, Snyder RW, Sumner SC, et al. 2006. Kinetics of elimination of urinary metabolites of acrylamide in humans. Toxicol Sci 93(2):256-267.
- Fennell TR, Macneela JP, Sumner SCJ. 1990. Metabolism of 1,2,3-¹³C acrylamide in the rat and mouse determined by ¹³C nuclear magnetic resonance spectroscopy. Proc Annu Meet Am Assoc Cancer Res 31:114.
- *Fennell TR, Snyder RW, Krol WL, et al. 2003. Comparison of the hemoglobin adducts formed by administration of N-methylolacrylamide and acrylamide to rats. Toxicol Sci 71(2):164-175.
- *Fennell TR, Sumner SC, Snyder RW, et al. 2005b. Metabolism and hemoglobin adduct formation of acrylamide in humans. Toxicol Sci 85(1):447-459.
- Ferri GL, Zareh S, Sbraccia M, et al. 1991. Heterogeneous visceral nerve changes in acrylamide intoxication. Exp Brain Res 87(2):363-370.
- Field EA, Price CJ, Sleet RB, et al. 1989. Teratologic evaluation of acrylamide (ACRL) in CD-1 mice and CD rats. [Abstract]. Toxicologist 9(1):273.
- +*Field EA, Price CJ, Sleet RB, et al. 1990. Developmental toxicity evaluation of acrylamide in rats and mice. Fundam Appl Toxicol 14(3):502-512.
- *Fomon SJ. 1966. Body composition of the infant: Part 1: The male reference infant. In: Faulkner F, ed. Human development. Philadelphia, PA: WB Saunders, 239-246.
- *Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. Am J Clin Nutr 35(Suppl 5):1169-1175.
- Frantz SW, Dryzga MD, Fresheur NL, et al. 1995. *In vitro/in vivo* determination of cutaneous penetration of residual acrylamide monomer from polyacrylamide water solutions. Dow Chemical Toxicology Research Laboratory.
- Friedman M. 2003. Chemistry, biochemistry, and safety of acrylamide. A review. J Agric Food Chem 51(16):4504-4526.
- Friedman M, Levin CE. 2008. Review of methods for the reduction of dietary content and toxicity of acrylamide. J Agric Food Chem 56(15):6113-6140.

9. REFERENCES

- +*Friedman MA, Dulak LH, Stedham MA. 1995. A lifetime oncogenicity study in rats with acrylamide. *Fundam Appl Toxicol* 27(1):95-105.
- +*Friedman MA, Tyl RW, Marr MC, et al. 1999. Effects of lactational administration of acrylamide on rat dams and offspring. *Reprod Toxicol* 13(6):511-520.
- Friedman MA, Zeiger E, Marroni DE, et al. 2008. Inhibition of rat testicular nuclear kinesins (krp2; KIFC5A) by acrylamide as a basis for establishing a genotoxicity threshold. *J Agric Food Chem* 56:6024-6030.
- *Fuhr U, Boettcher MI, Kinzig-Schippers M, et al. 2006. Toxicokinetics of acrylamide in humans after ingestion of a defined dose in a test meal to improve risk assessment for acrylamide carcinogenicity. *Cancer Epidemiol Biomarkers Prev* 15(2):266-271.
- +*Fullerton PM. 1969. Electrophysiological and histological observations on peripheral nerves in acrylamide poisoning in man. *J Neurol Neurosurg Psychiatry* 32:186-192.
- *Fullerton PM, Barnes JM. 1966. Peripheral neuropathy in rats produced by acrylamide. *Br J Ind Med* (23):210-221.
- Galesa K, Bren U, Kranjc A, et al. 2008. Carcinogenicity of acrylamide: A computational study. *J Agric Food Chem* 56(18):8720-8727.
- *Gamboa da Costa G, Churchwell MI, Hamilton LP, et al. 2003. DNA adduct formation from acrylamide via conversion to glycidamide in adult and neonatal mice. *Chem Res Toxicol* 16(10):1328-1337.
- +*Garey J, Paule MG. 2007. Effects of chronic low-dose acrylamide exposure on progressive ratio performance in adolescent rats. *Neurotoxicology* 28(5):998-1002.
- Garey J, Ferguson SA, Paule MG. 2004. Behavioral effects of acrylamide exposure during early development in Fischer 344 rats. [Abstract]. *Birth Defects Res Part A Clin Mol Teratol* 70(5):346.
- Garey J, Ferguson SA, Paule MG. 2005. Effect of low-dose acrylamide exposure on preweaning behavior of Fisher 344 rats. [Abstract]. *Neurotoxicol Teratol* 27(3):379-380.
- Garey J, Ferguson SA, Paule MG. 2006. Effects of chronic, low-dose acrylamide treatment on spatial learning and neurodevelopment in Fischer 344 rats. [Abstract]. *Toxicol Sci* 90(1-S):298.
- Gargas ML, Kirman CR, Sweeney LM, et al. 2009. Acrylamide: Consideration of species differences and nonlinear processes in estimating risk and safety for human ingestion. *Food Chem Toxicol* 47(4):760-768.
- *Garland TO, Patterson MWH. 1967. Six cases of acrylamide poisoning. *Br Med J* 4:134-138.
- Gassner P, Adler ID. 1996. Induction of hypoploidy and cell cycle delay by acrylamide in somatic and germinal cells of male mice. *Mutat Res* 367(4):195-202.

9. REFERENCES

- *Generoso WM, Sega GA, Lockhart AM, et al. 1996. Dominant lethal mutations, heritable translocations, and unscheduled DNA synthesis induced in male mouse germ cells by glycidamide, a metabolite of acrylamide. *Mutat Res* 371(3-4):175-183.
- Ghanayem BI, Bai R, Burka LT. 2009. Effect of dose volume on the toxicokinetics of acrylamide and its metabolites and 2-deoxy-D-glucose. *Drug Metab Dispos* 37(2):259-263.
- *Ghanayem BI, McDaniel LP, Churchwell MI, et al. 2005c. Role of CYP2E1 in the epoxidation of acrylamide to glycidamide and formation of DNA and hemoglobin adducts. *Toxicol Sci* 88(2):311-318.
- *Ghanayem BI, Witt KL, El-Hadri L, et al. 2005a. Comparison of germ cell mutagenicity in male CYP2E1-null and wild-type mice treated with acrylamide: Evidence supporting a glycidamide-mediated effect. *Biol Reprod* 72:157-163.
- *Ghanayem BJ, Witt KL, Kissling GE, et al. 2005b. Absence of acrylamide-induced genotoxicity in CYP2E1-null mice: Evidence consistent with a glycidamide-mediated effect. *Mutat Res* 578:284-297.
- Ghetti B, Wisniewski HM, Cook RD, et al. 1973. Changes in the central nervous system after acute and chronic acrylamide intoxication. *Am J Pathol* 70(2):78a.
- Gho M, McDonald K, Ganetzky B, et al. 1992. Effects of kinesin mutations on neuronal functions. *Science* 258:313-316.
- +*Gilbert SG, Maurissen JP. 1982. Assessment of the effects of acrylamide, methylmercury and 2,5-hexanedione on motor functions in mice. *J Toxicol Environ Health* 10(1):31-42.
- Gipon L, Schotman P, Jennekens FG, et al. 1977. Polyneuropathies and CNS protein metabolism: I. Description of the acrylamide syndrome in rats. *Neuropathol Appl Neurobiol* 3(2):115-123.
- *Giwercman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. *Environmental Health Perspectives Supplement* 101(2):65-71.
- *Gjerløff T, Elsborg H, Bonde JP. 2001. [Severe chronic acrylamide intoxication]. *Ugeskr Laeger* 163(32):4204-4205.
- Gobel A, Kliemant A. 2007. The German minimization concept for acrylamide. *Food Addit Contam* 24(Suppl 1):82-90.
- Goffeng LO, Heier MS, Kjuus H, et al. 2008a. Nerve conduction, visual evoked responses and electroretinography in tunnel workers previously exposed to acrylamide and N-methylolacrylamide containing grouting agents. *Neurotoxicol Teratol* 30(3):186-194.
- Goffeng LO, Kjuus H, Heier MS, et al. 2008b. Colour vision and light sensitivity in tunnel workers previously exposed to acrylamide and N-methylolacrylamide containing grouting agents. *Neurotoxicology (Little Rock)* 29:31-39.
- Gold BG, Griffin JW, Price DL. 1985. Slow axonal transport in acrylamide neuropathy: Different abnormalities produced by single-dose and continuous administration. *J Neurosci* 5(7):1755-1768.

9. REFERENCES

- Gold BG, Griffin JW, Price DL. 1992. Somatofugal axonal atrophy precedes development of axonal degeneration in acrylamide neuropathy. *Arch Toxicol* 66(1):57-66.
- Gold BG, Voda J, Yu X, et al. 2004. The immunosuppressant FK506 elicits a neuronal heat shock response and protects against acrylamide neuropathy. *Exp Neurol* 187(1):160-170.
- *Goldfrank LR, Flomenbaum NE, Lewin NA, et al. 1998. Goldfrank's toxicologic emergencies. Stamford, CT: Appleton & Lange, 322-324, 475.
- Goldstein B, Lowndes HE. 1979. Spinal cord defect in the peripheral neuropathy resulting from acrylamide. *Neurotoxicology* 1(1):75-88.
- Goldstein BD. 1985a. Acrylamide preferentially affects slowly adapting cutaneous mechanoreceptors. *Toxicol Appl Pharmacol* 80(3):527-533.
- Goldstein BD. 1985b. Acrylamide neurotoxicity: Altered spinal monosynaptic responses to quipazine, a serotonin agonist, in cats. *Toxicol Appl Pharmacol* 78(3):436-444.
- Goldstein BD, Fincher DR. 1986. Paradoxical changes in spinal cord reflexes following the acute administration of acrylamide. *Toxicol Lett* 31(2):93-99.
- Goldstein BD, Lowndes HE. 1977. Mono- and polysynaptic reflexes in cats with experimental acrylamide neuropathy. [Abstract]. *Pharmacologist* 19(2):224.
- Goldstein BD, Lowndes HE. 1979. Spinal cord defect in the peripheral neuropathy resulting from acrylamide. *Neurotoxicology* 1(1):75-87.
- +*Gorzinski SJ, Morden DC, Albee RR, et al. 1979. Results of papatability (12-day) and tolerance (21-day) studies on acrylamide monomer administered in the drinking water to rats. Dow Chemical Company. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878210959. OTS206135.
- Gould RM, Brady ST. 2004. Neuropathology: Many paths lead to hereditary spastic paraplegia. *Curr Biol* 14:R903-R904.
- Granath F, Ehrenberg L, Paulsson B, et al. 2001. Cancer risk from exposure to occupational acrylamide. *Occup Environ Med* 58(9):608-609.
- Granby K, Fagt S. 2004. Analysis of acrylamide in coffee and dietary exposure to acrylamide from coffee. *Anal Chim Acta* 520(1-2):177-182.
- Graveleau J, Loirat P, Nusinovici V. 1970. Polyneuritis caused by acrylamide. *Rev Neurol (Paris)* 123:62-65.
- Griffin JW, Price DL, Drachman DB. 1977. Impaired axonal regeneration in acrylamide intoxication. *J Neurobiol* 8(4):355-370.
- Guenther H, Anklam E, Wenzl T, et al. 2007. Acrylamide in coffee: Review of progress in analysis, formation and level reduction. *Food Addit Contam* 24 (Suppl 1):60-70.

9. REFERENCES

- Gueuning C, Graff GL, Dictus-Vermeulen C. 1978. Neurotoxicity of acrylamide in the rat: I. Subacute intoxication, absence of effect on incorporation of plasma phosphate in acid-soluble phosphates of the sciatic nerve and the gastrocnemius muscle, in anatomic continuity and after cutting the nerve. *Arch Int Physiol Biochim* 86(2):342-352.
- +*Gutierrez-Espeleta GA, Hughes LA, Piegorsch WW, et al. 1992. Acrylamide: Dermal exposure produces genetic damage in male mouse germ cells. *Fundam Appl Toxicol* 18(2):189-192.
- *Guzelian PS, Henry CJ, Olin SS. 1992. In: Similarities and differences between children and adults: Implications for risk assessment. Washington DC: International Life Sciences and Press Institute Press.
- *Haber LT, Maier A, Kroner OL, et al. 2009. Evaluation of human relevance and mode of action for tunica vaginalis mesotheliomas resulting from oral exposure to acrylamide. *Regul Toxicol Pharmacol* 53(2):134-149.
- *Habermann CE. 2002. Acrylamide. In: Kirk-Othmer encyclopedia of chemical technology. John Wiley & Sons, 1-22.
<http://mrw.interscience.wiley.com/emrw/9780471238966/kirk/article/acryhabe.a01/current/pdf>. June 08, 2009.
- +*Hagmar L, Tornqvist M, Nordander C, et al. 2001. Health effects of occupational exposure to acrylamide using hemoglobin adducts as biomarkers of internal dose. *Scand J Work Environ Health* 27(4):219-226.
- Hagmar L, Wirfalt E, Paulsson B, et al. 2005. Differences in hemoglobin adduct levels of acrylamide in the general population with respect to dietary intake, smoking habits and gender. *Mutat Res* 580(1-2):157-165.
- *Hanioka N, Tanaka-Kagawa T, Miyata Y, et al. 2003. Functional characterization of three human cytochrome P450 2E1 variants with amino acid substitutions. *Xenobiotica* 33(6):575-586.
- *Harris CH, Gulati AK, Friedman MA, et al. 1994. Toxic neurofilamentous axonopathies and fast axonal transport. V. Reduced bidirectional vesical transport in cultured neurons by acrylamide and glycidamide. *J Toxicol Environ Health* 42:343-356.
- *Harry GJ. 1992. Acrylamide-induced alterations in axonal transport. Biochemical and autoradiographic studies. *Mol Neurobiol* 6(2-3):203-216.
- Hartley CL, Anderson VE, Anderson BH, et al. 1997. Acrylamide and 2,5-hexanedione induce collapse of neurofilaments in SH-SY5Y human neuroblastoma cells to form perikaryal inclusion bodies. *Neuropathol Appl Neurobiol* 23(5):364-372.
- Hartmann EC, Boettcher MI, Bolt HM, et al. 2008a. N-Acetyl-S-(1-carbamoyl-2-hydroxy-ethyl)-L-cysteine (iso-GAMA) a further product of human metabolism of acrylamide: Comparison with the simultaneously excreted other mercapturic acids. *Arch Toxicol* 83(7):731-734.
- Hartmann EC, Boettcher MI, Schettgen T, et al. 2008b. Hemoglobin adducts and mercapturic acid excretion of acrylamide and glycidamide in one study population. *J Agric Food Chem* 56(15):6061-6068.

9. REFERENCES

- Hasegawa K, Miwa S, Isomura K, et al. 2008. Acrylamide-responsive genes in the nematode *Caenorhabditis elegans*. *Toxicol Sci* 101(2):215-225.
- Hasegawa K, Miwa S, Tajima T, et al. 2007. A rapid and inexpensive method to screen for common foods that reduce the action of acrylamide, a harmful substance in food. *Toxicol Lett* 175(1-3):82-88.
- Hashimoto K. 1980. Toxicity of acrylamide. *Sangyo Igaku* 22(4):233-248.
- *Hashimoto K, Aldridge WN. 1970. Biochemical studies on acrylamide, a neurotoxic agent. *Biochem Pharmacol* 19:2591-2604.
- Hashimoto K, Ando K. 1975. Studies on the percutaneous absorption of acrylamide. [Abstract]. Abstracts of the XVIII International Congress on Occupational Health, Brighton, England, September 1975:453.
- *Hashimoto K, Tanii H. 1985. Mutagenicity of acrylamide and its analogs in *Salmonella typhimurium*. *Mutat Res* 158:129-133.
- +*Hashimoto K, Sakamoto J, Tanii H. 1981. Neurotoxicity of acrylamide and related compounds and their effects on male gonads in mice. *Arch Toxicol* 47:179-189.
- Hattis D, Shapiro K. 1990. Analysis of dose/time/response relationships for chronic toxic effects: The case of acrylamide. *Neurotoxicology* 11(2):219-236.
- *HazDat. 2007. Acrylamide. HazDat Database: ATSDR's Hazardous Substance Release and Health Effects Database. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- +*He F, Zhang S, Wang H, et al. 1989. Neurological and electroneuromyographic assessment of the adverse effects of acrylamide on occupationally exposed workers. *Scand J Work Environ Health* 15(2):125-129.
- +*Hersch MI, McLeod JG, Satchell PM, et al. 1989a. Breathing pattern, lung inflation reflex and airway tone in acrylamide neuropathy. *Respir Physiol* 76(2):257-276.
- Hersch MI, McLeod JG, Sullivan CE. 1989b. Abnormal cough reflex in canine acrylamide neuropathy. *Ann Neurol* 26(6):738-745.
- Hersch MI, Satchell PM, Sullivan CE, et al. 1986. Abnormal pulmonary slowly adapting receptors in canine acrylamide neuropathy. *J Appl Physiol* 60(2):376-384.
- *Heudorf U, Hartmann E, Angerer J. 2009. Acrylamide in children-exposure assessment via urinary acrylamide metabolites as biomarkers. *Int J Hyg Environ Health* 212:135-14.
- Hilbig A, Freidank N, Kersting M, et al. 2004. Estimation of the dietary intake of acrylamide by German infants, children and adolescents as calculated from dietary records and available data on acrylamide levels in food groups. *Int J Hyg Environ Health* 207(5):463-471.
- Hill EJ, Woehrling EK, Prince M, et al. 2008. Differentiating human NT2/D1 neurospheres as a versatile *in vitro* 3D model system for developmental neurotoxicity testing. 249:243-250.

9. REFERENCES

- Hills BW, Greife AL. 1986. Evaluation of occupational acrylamide exposures. *Appl Ind Hyg* 1(3):148-152.
- Hinson JA, Roberts DW. 1992. Role of covalent and noncovalent interactions in cell toxicity: effects on proteins. *Toxicol Appl Pharmacol* 32:471-510.
- *Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.
- +*Hogervorst JG, Schouten LJ, Konings EJ, et al. 2007. A prospective study of dietary acrylamide intake and the risk of endometrial, ovarian, and breast cancer. *Cancer Epidemiol Biomarkers Prev* 16(11):2304-2313.
- +*Hogervorst JG, Schouten LJ, Konings EJ, et al. 2008a. Dietary acrylamide intake and the risk of renal cell, bladder, and prostate cancer. *Am J Clin Nutr* 87(5):1428-1438.
- +*Hogervorst JG, Schouten LJ, Konings EJ, et al. 2008b. Dietary acrylamide intake is not associated with gastrointestinal cancer risk. *J Nutr* 138(11):2229-2236.
- Holden LJ, Coleman MD. 2007. Assessment of the astroglial responses of three human astrocytoma cell lines to ethanol, trimethyltin chloride and acrylamide. *Toxicology* 241(1-2):75-83.
- Holland N, Ahlborn T, Turteltaub K, et al. 1999. Acrylamide causes preimplantation abnormalities in embryos and induces chromatin-adducts in male germ cells of mice. *Reprod Toxicol* 13(3):167-178.
- *Hoorn AJW, Custer LI, Myhr BC. 1993. Detection of chemical mutagens using Muta Mouse: A transgenic mouse model. *Mutagenesis* 8(1):7-10.
- +*Hopkins A. 1970. The effect of acrylamide on the peripheral nervous system of the baboon. *J Neurol Neurosurg Psychiatry* 33(6):805-816.
- Hopkins AP, Gilliatt RW. 1971. Motor and sensory nerve conduction velocity in the baboon: Normal values and changes during acrylamide neuropathy. *J Neurol Neurosurg Psychiatry* 34(4):415-426.
- Howland RD. 1981. The etiology of acrylamide neuropathy: Enolase, phosphofructokinase, and glyceraldehyde-3-phosphate dehydrogenase activities in peripheral nerve, spinal cord, brain, and skeletal muscle of acrylamide-intoxicated cats. *Toxicol Appl Pharmacol* 60(2):324-333.
- Howland RD. 1985. Biochemical studies of acrylamide neuropathy. *Neurotoxicology* 6(4):7-15.
- Howland RD, Lowndes HE. 1984. Peripheral nerve phospholipids in acrylamide neuropathy. *Arch Toxicol* 55(3):178-181.
- Howland RD, Vyas IL, Lowndes HE, et al. 1980. The etiology of toxic peripheral neuropathies: *In vitro* effects of acrylamide and 2,5-hexanedione on brain enolase and other glycolytic enzymes. *Brain Res* 202(1):131-142.
- *HSDB. 2009. Acrylamide. Hazardous Substances Data Bank. National Library of Medicine. <http://toxnet.nlm.nih.gov>. April 27, 2009.

9. REFERENCES

- HSE. 1987. Acrylamide in air. Methods for the determination of hazardous substances (MDHS) 57. Health and Safety Executive. <http://www.hse.gov.uk/pubns/mdhs/pdfs/mdhs57.pdf>. May 4, 2009.
- Huang CC, Li CM, Wu CF, et al. 2007. Analysis of urinary N-acetyl-S-(propionamide)-cysteine as a biomarker for the assessment of acrylamide exposure in smokers. *Environ Res* 104(3):346-351.
- Hurd DD, Saxton WM. 1996. Kinesin mutations cause motor neuron disease phenotypes by disrupting fast axonal transport in *Drosophila*. *Genetics* 144(3):1075-1085.
- Hurt ME, Bentley KS, Working PK. 1987. Effects of acrylamide and acrylonitrile on unscheduled DNA synthesis (UDS) in rat spermatocytes. *Environ Mutagen* 9(Suppl 8):49-50.
- +*Husain R, Dixit R, Das M, et al. 1987. Neurotoxicity of acrylamide in developing rat brain: Changes in the levels of brain biogenic amines and activities of monoamine oxidase and acetylcholine esterase. *Ind Health* 25(1):19-28.
- *IARC. 1994. Acrylamide. In: IARC monographs of some industrial chemicals, summary of data reported and evaluation. Lyon, France: International Agency for Research on Cancer, 389-433.
- *IARC. 2009. Agents reviewed by the IARC Monographs. Volumes 1-99. Lyon, France: International Agency for Research on Cancer. <http://monographs.iarc.fr/ENG/Classification/index.php>. May 19, 2009.
- Igisu H, Kinoshita Y. 2007. Magnetic resonance for evaluation of toxic encephalopathies: Implications from animal experiments. *Neurotoxicology* 28(2):252-256.
- *Igisu H, Matsuoka M. 2002. Acrylamide encephalopathy. *J Occup Health* 44:63-68.
- *Igisu H, Goto I, Kawamura Y, et al. 1975. Acrylamide encephaloneuropathy due to well water pollution. *J Neurol Neurosurg Psychiatry* 38:581-584.
- *Ikeda GJ, Miller E, Sapienza PP, et al. 1983. Distribution of ^{14}C -labelled acrylamide and betaine in fetuses of rats, rabbits, beagle dogs and miniature pigs. *Food Chem Toxicol* 21:49-58.
- *Ikeda GJ, Miller E, Sapienza PP, et al. 1985. Maternal-foetal distribution studies in late pregnancy. II. Distribution of $[1-^{14}\text{C}]$ acrylamide in tissues of beagle dogs and miniature pigs. *Food Chem Toxicol* 23(8):757-761.
- Ikeda GJ, Miller E, Sapienza PP, et al. 1987. Comparative tissue distribution and excretion of $[1-^{14}\text{C}]$ acrylamide in beagle dogs and miniature pigs. *Food Chem Toxicol* 25(11):871-875.
- Imai T, Cho YM, Hasumura M, et al. 2005. Enhancement by acrylamide of N-methyl-N-nitrosourea-induced rat mammary tumor development-possible application for a model to detect co-modifiers of carcinogenesis. *Cancer Lett* 230(1):25-32.
- Imai T, Takami S, Cho Y, et al. 2006. Inhibitory effects of antioxidants and phase II-enzyme inducers on acrylamide induced rat mammary carcinogenesis. [Abstract]. *Toxicol Sci* 90(1-S):408-409.
- *IPCS. 1985. Acrylamide. Environmental Health Criteria 49. International Programme on Chemical Safety. <http://www.inchem.org/documents/ehc/ehc/ehc49.htm>. June 8, 2009.

9. REFERENCES

- *IRIS. 2009. Acrylamide. Integrated Risk Information System. Washington, DC: <http://www.epa.gov/iris/subst/index.html>. May 11, 2009.
- Jakobsen J, Sidenius P. 1983. Early and dose-dependent decrease of retrograde axonal transport in acrylamide-intoxicated rats. *J Neurochem* 40(2):447-454.
- Jennekens FG, Veldman H, Schotman P, et al. 1979. Sequence of motor nerve terminal involvement in acrylamide neuropathy. *Acta Neuropathol* 46(1-2):57-64.
- Jiang L, Cao J, An Y, et al. 2007. Genotoxicity of acrylamide in human hepatoma G2 (HepG2) cells. *Toxicol In Vitro* 21(8):1486-1492.
- Jin L, Chico-Galdo V, Massart C, et al. 2008. Acrylamide does not induce tumorigenesis or major defects in mice *in vivo*. *J Endocrinol* 198(2):301-307.
- *Johanson CE. 1980. Permeability and vascularity of the developing brain: Cerebellum vs. cerebral cortex. *Brain Res* 190(1):3-16.
- Johnson EC, Murphy SD. 1976. Effect of acrylamide intoxication on biochemical components of brain and sciatic nerve in rats. [Abstract]. *Toxicol Appl Pharmacol* 37(1):131.
- Johnson EC, Murphy SD. 1977. Effect of acrylamide intoxication on pyridine nucleotide concentrations and functions in rat cerebral cortex. *Biochem Pharmacol* 26(22):2151-2156.
- +*Johnson KA, Beyer JE, Bell TJ, et al. 1984. Acrylamide: A two-year drinking water chronic toxicity-oncogenicity study in Fischer 344 rats. American Cyanamid Company. Dow Chemical U.S.A. Nalco Chemical Company. The Standard Oil Company. Submitted to the U.S. Environmental Protection Agency under TSCA Section 4. OTS0507273.
- *Johnson KA, Beyer JE, Bell TJ, et al. 1985. Acrylamide: A two-year drinking water chronic toxicity-oncogenicity study in Fischer 344 rats. Electron microscopy portion. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878216184. OTS0506849.
- +*Johnson KA, Gorzinski SJ, Bodner KM, et al. 1986. Chronic toxicity and oncogenicity study on acrylamide incorporated in the drinking water of Fischer 344 rats. *Toxicol Appl Pharmacol* 85(2):154-168.
- *Johnsrud EK, Kuokouritaki SB, Divakaran K, et al. 2003. Human hepatic CYP2E1 expression during development. *J Pharmacol Exp Ther* 307(1):402-407. Erratum in: *J Pharmacol Exp Ther* 309(1):439.
- Jones HB, Cavanagh JB. 1986. The axon reaction in spinal ganglion neurons of acrylamide-treated rats. *Acta Neuropathol* 71(1-2):55-63.
- Jortner BS, Ehrlich M. 1993. Comparison of toxicities of acrylamide and 2,5-hexanedione in hens and rats on 3-week dosing regimens. *J Toxicol Environ Health* 39(4):417-428.
- *Jung R, Engelhart G, Herbolt B. 1992. Collaborative study of mutagenicity with *Salmonella typhimurium* TA102. *Mutat Res* 278:265-270.

9. REFERENCES

- *Kadry AM, Friedman MA, Abdel-Rahman MS. 1999. Pharmacokinetics of acrylamide after oral administration in male rats. *Environ Toxicol Pharmacol* 7(2):127-133.
- Kaji R, Liu Y, Duckett S, et al. 1989. Slow recovery of central axons in acrylamide neuropathy. *Muscle Nerve* 12(10):816-826.
- Kaplan ML, Murphy SD. 1971. Rotarod performance and sciatic nerve betaglucuronidase activity as indices of acrylamide neurotoxicity in rats. *Toxicol Appl Pharmacol* 19(2):408-409.
- Kaplan ML, Murphy SD. 1972. Modification of acrylamide neuropathy in rats by various factors. *Toxicol Appl Pharmacol* 22(2):302-303.
- *Kaplan ML, Murphy SD, Gilles FH. 1973. Modification of acrylamide neuropathy in rats by selected factors. *Toxicol Appl Pharmacol* 24:564-579.
- Kasamatsu T, Kohda K. 2006. Balancing risks. *Regul Toxicol Pharmacol* 46(1):100-104.
- *Kaster JK, Kamendulls LM, Friedman MA, et al. 1998. Syrian hamsters embryo (SHE) cell transformation by acrylamide and hormones. *Toxicol Sci* 42(1-S):76.
- *Kedderis GL, Teo SK, Batra R, et al. 1996. Refinement and verification of the physiologically based dosimetry description for acrylonitrile in rats. *Toxicol Appl Pharmacol* 140:422-435.
- *Kesson CM, Baird AW, Lawson DH. 1977. Acrylamide poisoning. *Postgrad Med J* 53:16-17.
- Khanna VK, Husain R, Seth PK. 1992. Protein malnourishment: A predisposing factor in acrylamide toxicity in pregnant rats. *J Toxicol Environ Health* 36(4):293-305.
- King DJ, Noss RR. 1989. Toxicity of polyacrylamide and acrylamide monomer. *Rev Environ Health* 8(1-4):3-16.
- Kinoshita Y, Matsumura H, Igisu H, et al. 2000. Brain of rats intoxicated with acrylamide: Observation with 4.7 tesla magnetic resonance. *Arch Toxicol* 74(8):487-489.
- *Kirman CR, Gargas ML, Deskin R, et al. 2003. A physiologically based pharmacokinetic model for acrylamide and its metabolite, glycidamide, in the rat. *J Toxicol Environ Health A* 66(3):253-274.
- Kjuus H, Goffeng LO, Heier MS, et al. 2004. Effects on the peripheral nervous system of tunnel workers exposed to acrylamide and N-methylolacrylamide. *Scand J Work Environ Health* 30(1):21-29.
- Kjuus H, Hansteen IL, Ryberg D, et al. 2005. Chromosome aberrations in tunnel workers exposed to acrylamide and N-methylolacrylamide. *Scand J Work Environ Health* 31(4):300-306.
- *Klaunig JE. 2008. Acrylamide carcinogenicity. *J Agric Food Chem* 56(15):5984-5988.
- Klaunig JE, Kamendulis LM. 2005. Mechanisms of acrylamide induced rodent carcinogenesis. *Adv Exp Med Biol* 561:49-62.
- Kliesch U, El-Tarras A, Ingwersen I, et al. 1989. Clastogenicity of acrylamide in somatic and germinal cells of mice. *Mutagenesis* 4:319-320.

9. REFERENCES

- *Kligerman AD, Atwater AL, Bryant MF, et al. 1991. Cytogenetic studies of ethyl acrylate using C57BL/6 mice. *Mutagenesis* 6(2):137-141
- *Knaap AC, Kramers PN, Voogd CE, et al. 1988. Mutagenic activity of acrylamide in eukaryotic systems but not in bacteria. *Mutagenesis* 3:263-268.
- +Ko MH, Chen WP, Hsieh ST. 2000. Cutaneous nerve degeneration induced by acrylamide in mice. *Neurosci Lett* 293(3):195-198.
- *Ko MH, Chen WP, Hsieh ST. 2002. Neuropathology of skin denervation in acrylamide-induced neuropathy. *Neurobiol Dis* 11(1):155-165.
- +*Ko MH, Chen WP, Lin-Shiau SY, et al. 1999. Age-dependent acrylamide neurotoxicity in mice: Morphology, physiology, and function. *Exp Neurol* 158(1):37-46.
- Kohriyama K. 1992. [The suppression of CK activity in the brain by acrylamide]. *J UOEH* 14(1):47-58.
- *Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29(18):4430-4433.
- Konings EJ, Ashby P, Hamlet CG, et al. 2007. Acrylamide in cereal and cereal products: A review on progress in level reduction. *Food Addit Contam* 24 (Suppl 1):47-59.
- Konings EJ, Baars AJ, van Klaveren JD, et al. 2003. Acrylamide exposure from foods of the Dutch population and an assessment of the consequent risks. *Food Chem Toxicol* 41(11):1569-1579.
- Kopp EK, Dekant W. 2009. Toxicokinetics of acrylamide in rats and humans following single oral administration of low doses. *Toxicol Appl Pharmacol* 235(2):135-142.
- Kopp EK, Sieber M, Kellert M, et al. 2008. Rapid and sensitive HILIC-ESI-MS/MS quantitation of polar metabolites of acrylamide in human urine using column switching with an online trap column. *J Agric Food Chem* 56(21):9828-9834.
- Koyama N, Sakamoto H, Sakuraba M, et al. 2006. Genotoxicity of acrylamide and glycidamide in human lymphoblastoid TK6 cells. *Mutat Res* 603(2):151-158.
- *Krebs O, Favor J. 1997. Somatic and germ cell mutagenesis in lambda lacZ transgenic mice treated with acrylamide or ethylnitrosourea. *Mutat Res* 388(2-3):239-248.
- *Krishna G, Theiss C. 1995. Concurrent analysis of cytogenetic damage *in vivo*: A multiple endpoint-multiple tissue approach. *Environ Mol Mutagen* 25:314-320.
- *Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- *Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures. Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.

9. REFERENCES

- KS Crump Group, Inc. 1999a. Consideration of the potency classification of acrylamide based on the incidence of tunica vaginalis mesotheliomas (TVMs) in male F344 rats. June 11, 1999.
- KS Crump Group, Inc. 1999b. Mechanism of acrylamide induction of benign mammary fibroadenomas in the aging female F344 rat: Relevance to human health risk assessment. December 14, 1999.
- Kurebayashi H, Ohno Y. 2006. Metabolism of acrylamide to glycidamide and their cytotoxicity in isolated rat hepatocytes: Protective effects of GSH precursors. *Arch Toxicol* 80(12):820-828.
- Kutting B, Uter W, Drexler H. 2008. The association between self-reported acrylamide intake and hemoglobin adducts as biomarkers of exposure. *Cancer Causes Control* 19(3):273-281.
- *Lafferty JS, Kamendulis LM, Kaster J, et al. 2004. Subchronic acrylamide treatment induces a tissue-specific increase in DNA synthesis in the rat. *Toxicol Lett* 154(1-2):95-103.
- *Lähdetie J, Suutari A, Sjoblom T. 1994. The spermatid micronucleus test with the dissection technique detects the germ cell mutagenicity of acrylamide in rat meiotic cells. *Mutat Res* 309(2):255-262.
- Lambert J, Matthieu L, Dockx P. 1988. Contact dermatitis from acrylamide. *Contact Dermatitis* 19(1):65.
- Lamy E, Volkel Y, Roos PH, et al. 2008. Ethanol enhanced the genotoxicity of acrylamide in human, metabolically competent HepG2 cells by CYP2E1 induction and glutathione depletion. *Int J Hyg Environ Health* 211(1-2):74-81.
- Lande SS, Bosch SJ, Howard PH. 1979. Degradation and leaching of acrylamide in soil. *J Environ Qual* 8(1):133-137.
- *Lapadula DM, Bowe M, Carrington CD, et al. 1989. *In vitro* binding of [¹⁴C]acrylamide to neurofilament and microtubule proteins of rats. *Brain Res* 481(1):157-161.
- Lapin EP, Maker HS, Weissbarth S, et al. 1984. The influence of systemic factors on acrylamide-induced changes in brain, nerve, and other tissues. *J Neurosci Res* 11(4):395-404.
- +*Larsson SC, Akesson A, Wolk A. 2009d. Long-term dietary acrylamide intake and breast cancer risk in a prospective cohort of Swedish women. *Am J Epidemiol* 169(3):376-381.
- +*Larsson SC, Akesson A, Wolk A. 2009b. Long-term dietary acrylamide intake and risk of epithelial ovarian cancer in a prospective cohort of Swedish women. *Cancer Epidemiol Biomarkers Prev* 18(3):994-997.
- +*Larsson SC, Hakansson N, Akesson A, et al. 2009a. Long-term dietary acrylamide intake and risk of endometrial cancer in a prospective cohort of Swedish women. *Int J Cancer* 124(5):1196-1199.
- +*Larsson SC, Kesson A, Bergkvist L, et al. 2009c. Dietary acrylamide intake and risk of colorectal cancer in a prospective cohort of men. *Eur J Cancer* 45:513-516.

9. REFERENCES

- Lee KY, Shibutani M, Kuroiwa K, et al. 2005. Chemoprevention of acrylamide toxicity by antioxidative agents in rats-effective suppression of testicular toxicity by phenylethyl isothiocyanate. *Arch Toxicol* 79(9):531-541.
- *Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.
- Lehning EJ, Balaban CD, Ross JF, et al. 2002a. Acrylamide neuropathy. I. Spatiotemporal characteristics of nerve cell damage in rat cerebellum. *Neurotoxicology* 23(3):397-414.
- Lehning EJ, Balaban CD, Ross JF, et al. 2002b. Acrylamide neuropathy. II. Spatiotemporal characteristics of nerve cell damage in brainstem and spinal cord. *Neurotoxicology* 23:415-429.
- Lehning EJ, Balaban CD, Ross JF, et al. 2003a. Acrylamide neuropathy. II. Spatiotemporal characteristics of nerve cell damage in brainstem and spinal cord. *Neurotoxicology* 24(1):109-123.
- Lehning EJ, Balaban CD, Ross JF, et al. 2003b. Acrylamide neuropathy. III. Spatiotemporal characteristics of nerve cell damage in forebrain. *Neurotoxicology* 24(1):125-136.
- Lehning EJ, Gaughan CL, LoPachin RM, Jr. 1997. Acrylamide intoxication modifies *in vitro* responses of peripheral nerve axons to anoxia. *J Peripher Nerv Syst* 2(2):165-174.
- Lehning EJ, LoPachin EM, Mathew J, et al. 1994. Changes in Na-K ATPase and protein kinase C activities in peripheral nerve of acrylamide-treated rats. *J Toxicol Environ Health* 42(3):331-342.
- Lehning EJ, Persaud A, Dyer KR, et al. 1998. Biochemical and morphologic characterization of acrylamide peripheral neuropathy. *Toxicol Appl Pharmacol* 151(2):211-221.
- *Leikin JB, Paloucek FP. 2002. Leikin & Paloucek's poisoning and toxicology handbook. 3rd ed. Poisoning and toxicology handbook. 3rd ed. Hudson, OH: Lexi-Comp, Inc., 193-194.
- +*Leswing RJ, Ribelin WE. 1969. Physiologic and pathologic changes in acrylamide neuropathy. *Arch Environ Health* 18(1):23-29.
- *Leung H. 1993. Physiologically-based pharmacokinetic modelling. In: Ballantyne B, Marrs T, Turner P, eds. *General and applied toxicology*. Vol. 1. New York, NY: Stockton Press, 153-164.
- *Lewis RJ. 2000. Acrylamide. In: *Sax's dangerous properties of industrial materials*. 10th ed. New York, NY: John Wiley & Sons, Inc., 66-67.
- *Lewis RJ. 2007. Acrylamide. In: *Hawley's condensed chemical dictionary*. 15th ed. Hoboken, NJ: John Wiley & Sons, Inc., 19-20.
- Lewkowski JP, Yang YY, Orthoefer JG, et al. 1978. Effects of low level exposure to acrylamide in water on spontaneous locomotor activity. *Toxicol Appl Pharmacol* 45(1):251.
- Li SX, Cui N, Zhang CL, et al. 2006. Effect of subchronic exposure to acrylamide induced on the expression of bcl-2, bax and caspase-3 in the rat nervous system. *Toxicology* 217(1):46-53.

9. REFERENCES

- Lide DR, ed. 2008. [Acrylamide]. In: CRC handbook of chemistry and physics. 88th ed. New York, NY: CRC Press, 3-8 to 3-9.
- *Lijinsky W, Andrews AW. 1980. Mutagenicity of vinyl compounds in *Salmonella typhimurium*. *Teratog Carcinog Mutagen* 1:259-267.
- Ling B, Authier N, Balayssac D, et al. 2005. Assessment of nociception in acrylamide-induced neuropathy in rats. *Pain* 119(1-3):104-112.
- *Livingston AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4(2-3):301-324.
- LoPachin RM. 2004. The changing view of acrylamide neurotoxicity. *Neurotoxicology* 25(4):617-630.
- LoPachin RM. 2005. Acrylamide neurotoxicity: Neurological, morphological and molecular endpoints in animal models. *Adv Exp Med Biol* 561:21-37.
- LoPachin RM, Barber DS. 2005. Acrylamide disrupts uptake of dopamine into rat striatal synaptic vesicles. [Abstract]. *Toxicol Sci* 84(1-S):315.
- *LoPachin RM, Barber DS. 2006. Synaptic cysteine sulfhydryl groups as targets of electrophilic neurotoxicants. *Toxicol Sci* 94:240-255.
- *LoPachin RM, Decaprio AP. 2005. Protein adduct formation as a molecular mechanism in neurotoxicity. *Toxicol Sci* 86(2):214-225.
- *LoPachin RM, Gavin T. 2008. Acrylamide-induced nerve terminal damage: Relevance to neurotoxic and neurodegenerative mechanisms. *J Agric Food Chem* 56(15):5994-6003.
- *LoPachin RM, Lehning EJ. 1994. Acrylamide-induced distal axon degeneration: A proposed mechanism of action. *Neurotoxicology* 15(2):247-259.
- *LoPachin RM, Balaban CD, Ross JF. 2003. Acrylamide axonopathy revisited. *Toxicol Appl Pharmacol* 188(3):135-153.
- *LoPachin RM, Barber DS, Gavin T. 2008. Molecular mechanisms of the conjugated alpha, beta-unsaturated carbonyl derivatives: Relevance to neurotoxicity and neurodegenerative diseases. *Toxicol Sci* 104(2):235-249.
- LoPachin RM, Barber DS, Geohagen BC, et al. 2007a. Structure-toxicity analysis of type-2 alkenes: *In vitro* neurotoxicity. *Toxicol Sci* 95(1):136-146.
- LoPachin RM, Barber DS, He D, et al. 2006. Acrylamide inhibits dopamine uptake in rat striatal synaptic vesicles. 89:224-234.
- LoPachin RM, Gavin T, Geohagen BC, et al. 2007b. Neurotoxic mechanisms of electrophilic type-2 alkenes: Soft-soft interactions described by quantum mechanical parameters. *Toxicol Sci* 98:561-570.
- LoPachin RM, Geohagen BC, Gavin T. 2009. Synaptosomal toxicity and nucleophilic targets of 4-hydroxy-2-nonenal. *Toxicol Sci* 107:171-181.

9. REFERENCES

- LoPachin RM, Ross JF, Lehning EJ. 2002a. Nerve terminals as the primary site of acrylamide action: A hypothesis. *Neurotoxicology* 23(1):43-59.
- LoPachin RM, Ross JF, Reid ML, et al. 2002b. Neurological evaluation of toxic axonopathies in rats: Acrylamide and 2,5-hexanedione. *Neurotoxicology* 23(1):95-110.
- LoPachin RM, Schwarcz AI, Gaughan CL, et al. 2004. *In vivo* and *in vitro* effects of acrylamide on synaptosomal neurotransmitter uptake and release. *Neurotoxicology* 25(3):349-363.
- Lowndes HE, Baker T. 1976. Studies on drug-induced neuropathies: III. Motor nerve deficit in cats with experimental acrylamide neuropathy. *Eur J Pharmacol* 35(1):177-184.
- Lowndes HE, Baker T, Cho ES, et al. 1978. Position sensitivity of de-efferented muscle spindles in experimental acrylamide neuropathy. *J Pharmacol Exp Ther* 205(1):40-48.
- Lu ZR, Zou HC, Park SJ, et al. 2009. The effects of acrylamide on brain creatine kinase: Inhibition kinetics and computational docking simulation. *Int J Biol Macromol* 44(2):128-132.
- Mangan D, Snyder IS. 1978. The effect of acrylamide on human polymorphonuclear neutrophils *in vitro*. *Br J Ind Med* 35(4):305-311.
- Maniere I, Godard T, Doerge DR, et al. 2005. DNA damage and DNA adduct formation in rat tissues following oral administration of acrylamide. *Mutat Res* 580(1-2):119-129.
- *Manjanatha MG, Aidoo A, Shelton SD, et al. 2006. Genotoxicity of acrylamide and its metabolite glycidamide administered in drinking water to male and female Big Blue mice. *Environ Mol Mutagen* 47(1):6-17.
- Mannaa F, Abdel-Wahhab MA, Ahmed HH, et al. 2006. Protective role of Panax ginseng extract standardized with ginsenoside Rg3 against acrylamide-induced neurotoxicity in rats. *J Appl Toxicol* 26(3):198-206.
- *Mapp C, Mazzotta M, Bartolucci GB, et al. 1977. [La neuropatia da acrilamide: Prime osservazioni in Italia]. (Italian) *Med Lav* 68(1):1-12.
- Marchetti F, Bishop J, Lowe X, et al. 2009. Chromosomal mosaicism in mouse two-cell embryos after paternal exposure to acrylamide. *Toxicol Sci* 107(1):194-205.
- *Marchetti F, Lowe X, Bishop J, et al. 1997. Induction of chromosomal aberrations in mouse zygotes by acrylamide treatment of male germ cells and their correlation with dominant lethality and heritable translocations. *Environ Mol Mutagen* 30(4):410-417.
- *Marlowe C, Clark MJ, Mast RW, et al. 1986. The distribution of [¹⁴C]acrylamide in male and pregnant Swiss-Webster mice studied by whole-body autoradiography. *Toxicol Appl Pharmacol* 86(3):457-465.
- Marsh GM, Lucas LJ, Youk AO, et al. 1999. Mortality patterns among workers exposed to acrylamide: 1994 follow up. *Occup Environ Med* 56(3):181-190.
- *Marsh GM, Youk AO, Buchanich JM, et al. 2007. Mortality patterns among workers exposed to acrylamide: Updated follow up. *J Occup Environ Med* 49(1):82-95.

9. REFERENCES

- Martenson CH, Sheetz MP, Graham DG. 1995. *In vitro* acrylamide exposure alters growth cone morphology. *Toxicol Appl Pharmacol* 131(1):119-129.
- *Martins C, Oliveira NG, Pingarilho M, et al. 2007. Cytogenetic damage induced by acrylamide and glycidamide in mammalian cells: Correlation with specific glycidamide-DNA adducts. *Toxicol Sci* 95(2):383-390.
- *Marty JP, Vincent CM. 1998. *In vitro* percutaneous absorption of acrylamide across human skin. Paris, France, Faculty of Pharmacy, Universite de Paris Sud.
- Matsuoka M, Igisu H. 1992. Brain energy metabolites in mice intoxicated with acrylamide: Effects of ischemia. *Toxicol Lett* 62(1):39-43.
- +*Maurissen JP, Weiss B, Davis HT. 1983. Somatosensory thresholds in monkeys exposed to acrylamide. *Toxicol Appl Pharmacol* 71(2):266-279.
- Maurissen JP, Weiss B, Cox C. 1990. Vibration sensitivity recovery after a second course of acrylamide intoxication. *Fundam Appl Toxicol* 15(1):93-98.
- *Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74(2-3):135-149.
- +*McCollister DD, Oyen F, Rowe VK. 1964. Toxicology of acrylamide. *Toxicol Appl Pharmacol* 6:172-181.
- McLean JD, Mann JR, Jacoby JA. 1978. A monitoring method for the determination of acrylamide in an industrial environment. *Am Ind Hyg Assoc J* 39(3):247-250.
- Medrano CJ, LoPachin RM. 1989. Effects of acrylamide and 2,5-hexanedione on brain mitochondrial respiration. *Neurotoxicology* 10(2):249-255.
- Mei N, Guo L, Tseng J, et al. 2008a. Gene expression changes associated with xenobiotic metabolism pathways in mice exposed to acrylamide. *Environ Mol Mutagen* 49(9):741-745.
- *Mei N, Hu J, Churchwell MI, et al. 2008b. Genotoxic effects of acrylamide and glycidamide in mouse lymphoma cells. *Food Chem Toxicol* 46(2):628-636.
- Merigan WH, Barkdoll E, Maurissen JP. 1982. Acrylamide-induced visual impairment in primates. *Toxicol Appl Pharmacol* 62(6):342-345.
- +*Merigan WH, Barkdoll E, Maurissen JP, et al. 1985. Acrylamide effects on the macaque visual system. I. Psychophysics and electrophysiology. *Invest Ophthalmol Vis Sci* 26(3):309-316.
- *Meylan WM, Howard PH. 1993. Computer estimation of the atmospheric gas-phase reaction rate of organic compounds with hydroxyl radicals and ozone. *Chemosphere* 26(12):2293-2299.
- +*Michels KB, Rosner BA, Chumlea WC, et al. 2006. Preschool diet and adult risk of breast cancer. *Int J Cancer* 118(3):749-754.

9. REFERENCES

- Miller MJ, McQueen CA. 1986. The effect of acrylamide on hepatocellular DNA repair. *Environ Mutagen* 8(1):99-108.
- *Miller MJ, Carter DE, Sipes IG. 1982. Pharmacokinetics of acrylamide in Fischer-344 rats. *Toxicol Appl Pharmacol* 63:36-44.
- Miller MS, Miller MJ, Burks TF, et al. 1983. Altered retrograde axonal transport of nerve growth factor after single and repeated doses of acrylamide in the rat. *Toxicol Appl Pharmacol* 69(1):96-101.
- Miller MJ, Miller MS, Burks TF, et al. 1984. A simple, sensitive method for detecting early peripheral nerve dysfunction in the rat following acrylamide treatment. *Neurotoxicology* 5(2):15-23.
- Mills C, Tlustos C, Evans R, et al. 2008. Dietary acrylamide exposure estimates for the United Kingdom and Ireland: Comparison between semiprobabilistic and probabilistic exposure models. *J Agric Food Chem* 56(15):6039-6045.
- *Moore MM, Amtower A, Doer C, et al. 1987. Mutagenicity and clastogenicity of acrylamide in 15178y mouse lymphoma cells. *Environ Mutagen* 9(3):261-267.
- *Moreno Navarro IM, Rubio Armendariz C, Gutierrez Fernandez AJ, et al. 2007. La acrilamida, contaminante quimico de procesamiento: Revision. *Revista de Toxicologia (Elche, Spain)* 24(1):1-9.
- Moretto A, Sabri MI. 1988. Progressive deficits in retrograde axon transport precede degeneration of motor axons in acrylamide neuropathy. *Brain Res* 440(1):18-24.
- *Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokinet* 5(6):485-527.
- Moser VC, Anthony DC, Sette WF, et al. 1992. Comparison of subchronic neurotoxicity of 2-hydroxyethyl acrylate and acrylamide in rats. *Fundam Appl Toxicol* 18(3):343-352.
- Mottram DS, Friedman M. 2008. Symposium on the chemistry and toxicology of acrylamide. *J Agric Food Chem* 56(15):5983.
- *Mottram DS, Wedzicha BL, Dodson AT. 2002. Acrylamide is formed in the Maillard reaction. *Nature* 419(6906):448-449.
- Mucci LA, Wilson KM. 2008. Acrylamide intake through diet and human cancer risk. *J Agric Food Chem* 56(15):6013-6019.
- +*Mucci LA, Adami HO, Wolk A. 2006. Prospective study of dietary acrylamide and risk of colorectal cancer among women. *Int J Cancer* 118(1):169-173.
- *Mucci LA, Dickman PW, Steineck G, et al. 2003. Dietary acrylamide and cancer of the large bowel, kidney, and bladder: Absence of an association in a population-based study in Sweden. *Br J Cancer* 88(1):84-89.
- +*Mucci LA, Lindblad P, Steineck G, et al. 2004. Dietary acrylamide and risk of renal cell cancer. *Int J Cancer* 109(5):774-776.

9. REFERENCES

- +*Mucci LA, Sandin S, Balter K, et al. 2005. Acrylamide intake and breast cancer risk in Swedish women. *JAMA* 293(11):1326-1327.
- Mukhtar H, Dixit R, Seth PK. 1981. Reduction in cutaneous and hepatic glutathione contents, glutathione S-transferase and aryl hydrocarbon hydroxylase activities following topical application of acrylamide to mouse. *Toxicol Lett* 9(2):153-156.
- Muller M, Wacker K, Getts D, et al. 2008. Further evidence for a crucial role of resident endoneurial macrophages in peripheral nerve disorders: Lessons from acrylamide-induced neuropathy. *Glia* 56(9):1005-1016.
- *Müller W, Engelhart G, Herbold B, et al. 1993. Evaluation of mutagenicity testing with *Salmonella typhimurium* TA102 in three different laboratories. *Environ Health Perspect* 101(Suppl 3):33-36.
- *Mulloy KB. 1996. Two case reports of neurological disease in coal mine preparation plant workers. *Am J Ind Med* 30(1):56-61.
- *Muttucumaru N, Elmore JS, Curtis T, et al. 2008. Reducing acrylamide precursors in raw materials derived from wheat and potato. *J Agric Food Chem* 56(15):6167-6172.
- Muttucumaru N, Halford NG, Elmore JS, et al. 2006. Formation of high levels of acrylamide during the processing of flour derived from sulfate-deprived wheat. *J Agric Food Chem* 54(23):8951-8955.
- +*Myers JE, Macun I. 1991. Acrylamide neuropathy in a South African factory: An epidemiologic investigation. *Am J Ind Med* 19(4):487-493.
- Nagao T. 1994. Developmental abnormalities due to exposure of mouse paternal germ cells, preimplantation embryos, and organogenic embryos to acrylamide. *Congenit Anom (Kyoto)* 34(1):35-46.
- *Nakagawa-Yagi Y, Choi D, Ogane, N, et al. 2001. Discovery of a novel compound insight into mechanisms for acrylamide-induced axonopathy and colchicine-induced apoptotic neuronal cell death. *Brain Res* 909:8-19.
- Nalco. 1977. Exhaust AAM sampling data sheets with cover letters. Nalco Chem. Co. Submitted to the U.S. Environmental Protection Agency under TSCS Section 8D. OTS0206160.
- *NAS/NRC. 1989. Report of the oversight committee. In: *Biologic markers in reproductive toxicology*. Washington, DC National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- *NCTR. 2009. National Center for Toxicological Research annual report. Research accomplishments and plans. Department of Health and Human Services, National Center for Toxicological Research, 52, 56, 69. <http://www.fda.gov/downloads/AboutFDA/CentersOffices/NCTR/ResearchAccomplishmentsPlans/UCM163319.pdf>. July 16, 2009.
- Nemoto S, Takatsuki S, Sasaki K, et al. 2002. Determination of acrylamide in foods by GC/MS using ¹³C-labeled acrylamide as an internal standard. *Shokuhin Eiseigaku Zasshi* 43(6):371-376.

9. REFERENCES

- *Nesterova EV, Durnev AD, Seredenin SB. 1999. Verapamil contributes to the clastogenic effects of acrylamide, cyclophosphamide, and dioxidine on somatic cells of BALB/C and C57BL/6 mice. *Mutat Res* 440(2):171-179.
- *Neuhauser-Klaus A, Schmahl W. 1989. Mutagenic and teratogenic effects of acrylamide in the mammalian spot test *Mutat Res* 226(3):157-162.
- Newman B, Oblak L, Kumiski D, et al. 2005. Reductions in neuromuscular junction cholinesterase activity by subacute and chronic acrylamide exposure. [Abstract]. *Toxicol Sci* 84(1-S):317.
- Nilsen OG, Toftgard R, Ingelman-Sundberg M, et al. 1978. Qualitative alterations of cytochrome P-450 in mouse liver microsomes after administration of acrylamide and methylmethacrylate. *Acta Pharmacol Toxicol* 43(4):299-305.
- NIOSH. 1985. Health hazard evaluation report No. HETA-84-307-1581, Big Dry Creek Plant, Westminster, Colorado. Cincinnati, OH: National Institute for Occupational Safety and Health. PB86132792.
- *NIOSH. 2005. Acrylamide. In: NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, <http://www.cdc.gov/niosh/npg/>. May 19, 2009.
- Nordin-Andersson M, Walum E, Kjellstrand P, et al. 2003. Acrylamide-induced effects on general and neurospecific cellular functions during exposure and recovery. *Cell Biol Toxicol* 19(1):43-51.
- *NRC. 1993. Pesticides in the diets of infants and children. Washington, DC: National Research Council. National Academy Press.
- *NTP. 2005. Report on carcinogens, 11th edition. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. <http://ntp-server.niehs.nih.gov/ntp/roc/toc11.html>. May 11, 2009.
- Oishi Y, Yamamoto H, Nagano M, et al. 1996. The effects of 2,5-hexanedione and acrylamide on myosin heavy chain isoforms of slow and fast skeletal muscles of the rat. *Toxicol Appl Pharmacol* 139(1):15-21.
- Okazaki S, Takashima H, Yamaguchi M, et al. 2003. [Neurobehavioral toxicity of acrylamide and IDPN (3,3'-iminodipropionitrile) in rats by 28-day oral administration--problems encountered in collaborative study and a commentary on conducting neurobehavioral testing.]. *J Toxicol Sci* 28 (Suppl 1):1-14.
- +*Olesen PT, Olsen A, Frandsen H, et al. 2008. Acrylamide exposure and incidence of breast cancer among postmenopausal women in the Danish Diet, Cancer and Health study. *Int J Cancer* 122(9):2094-2100.
- Olstorn HB, Paulsen JE, Alexander J. 2007. Effects of perinatal exposure to acrylamide and glycidamide on intestinal tumorigenesis in *Min/+* mice and their wild-type litter mates. *Anticancer Res* 27(6B):3855-3864.

9. REFERENCES

- *O'Neil MJ, Heckelman PE, Koch CB, et al. 2006. Acrylamide. In: O'Neil MJ, Heckelman PE, Koch CB, eds. The Merck index. An encyclopedia of chemicals, drugs, and biologicals. 14th ed. Whitehouse Station, NJ: Merck & Co., Inc., 22-23.
- *OSHA. 2009. Toxic and Hazardous Substances. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.1000, Table Z 1. <http://www.osha.gov/comp-links.html>. May 19, 2009.
- O'Shaughnessy DJ, Losos GJ. 1986. Comparison of central and peripheral nervous system lesions caused by high-dose short-term and low-dose subchronic acrylamide treatment in rats. *Toxicol Pathol* 14(4):389-394.
- Otles S, Otles S. 2004. Acrylamide in food. (Chemical structure of acrylamide). *Electron J Environ Agric Food Chem* 3(5):723-730.
- Otles S, Otles S. 2005. Acrylamide in food-formation of acrylamide and its damages to health. *Electron J Pol Agric Univ* 7(2) <http://www.ejpau.media.pl/volume7/issue2/food/art-02.html>. May 28, 2009.
- *Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. Human development. Philadelphia, PA: WB Saunders, 222-238.
- Pabst K, Mathar W, Palavinskas R, et al. 2005. Acrylamide-occurrence in mixed concentrate feed for dairy cows and carry-over into milk. *Food Addit Contam* 22(3):210-213.
- *Pacchierotti F, Tiveron C, D'Archivio M, et al. 1994. Acrylamide-induced chromosomal damage in male mouse germ cells detected by cytogenetic analysis of one-cell zygotes. *Mutat Res* 309(2):273-284.
- *Padilla S, Atkinson MB, Breuer AC. 1993. Direct measurement of fast axonal organelle transport in the sciatic nerve of rats treated with acrylamide. *J Toxicol Environ Health* 39(4):429-445.
- Palazoglu TK, Gokmen V. 2008. Reduction of acrylamide level in french fries by employing a temperature program during frying. *J Agric Food Chem* 56(15):6162-6166.
- *Palmer RB. 2004. Acrylic acid and derivatives. In: Dart RC, ed. Medical toxicology. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1358-1368.
- Pantusa VP, Stock TH, Morandi MT, et al. 2002. Inhalation exposures to acrylamide in biomedical laboratories. *AIHA J (Fairfax, Va)* 63(4):468-473.
- *Park J, Kamendulis LM, Friedman MA, et al. 2002. Acrylamide-induced cellular transformation. *Toxicol Sci* 65(2):177-183.
- Parker RD, Sharma RP, Obersteiner EJ. 1978. Acrylamide toxicity in developing chick embryo. [Abstract]. *Pharmacologist* 20(3):249.
- Parzefall W. 2008. Mini review on the toxicity of dietary acrylamide. *Food Chem Toxicol* 46(4):1360-1364.
- Paule MG, Garey J, Ferguson SA. 2004. Developmental toxicity of acrylamide in Fischer 344 rats. [Abstract]. *Birth Defects Res Part A Clin Mol Teratol* 70(5):312.

9. REFERENCES

- *Paulsson B, Grawe J, Tornqvist M. 2002. Hemoglobin adducts and micronucleus frequencies in mouse and rat after acrylamide or N-methylolacrylamide treatment. *Mutat Res* 516(1-2):101-111.
- Paulsson B, Kotova N, Grawe J, et al. 2003. Induction of micronuclei in mouse and rat by glycidamide, genotoxic metabolite of acrylamide. *Mutat Res* 535(1):15-24.
- Paulsson B, Larsen KO, Tornqvist M. 2006. Hemoglobin adducts in the assessment of potential occupational exposure to acrylamides-Three case studies. *Scand J Work Environ Health* 32(2):154-159.
- Paulsson B, Rannug A, Henderson AP, et al. 2005. *In vitro* studies of the influence of glutathione transferases and epoxide hydrolase on the detoxification of acrylamide and glycidamide in blood. *Mutat Res* 580(1-2):53-59.
- *Pearson RG, Songstad J. 1967. Application of the principle of hard and soft acids and bases to organic chemistry. *J Am Chem Soc* 89:1827-1836.
- +*Pelucchi C, Galeone C, Dal Maso L, et al. 2007. Dietary acrylamide and renal cell cancer. *Int J Cancer* 120(6):1376-1377.
- +*Pelucchi C, Galeone C, Levi F, et al. 2006. Dietary acrylamide and human cancer. *Int J Cancer* 118(2):467-471.
- *Perrault SD. 2003. Distinguishing between fertilization failure and early pregnancy loss when identifying male-mediated adverse pregnancy outcomes. *Adv Exp Med Biol* 518:189-198.
- Peters AK, Steemans M, Hansen E, et al. 2008. Evaluation of the embryotoxic potency of compounds in a newly revised high throughput embryonic stem cell test. *Toxicol Sci* 105(2):342-350.
- *Petersen DW, Kleinow KM, Kraska RC, et al. 1985. Uptake disposition and elimination of acrylamide in rainbow trout *Salmo gairdneri*. *Toxicol Appl Pharmacol* 80(1):58-65.
- Pienta RJ. 1980. Transformation of Syrian hamster embryo cells by diverse chemicals and correlation with their reported carcinogenic and mutagenic activities. *Chem Mutagens* 6:175-202.
- *Pleasure DE, Mishler KC, Engel, WK. 1969. Axonal transport of proteins in experimental neuropathies. *Science* 166:524-525.
- Poole CF, Sye WF, Zlatkis A, et al. 1981. Determination of acrylamide in nerve tissue homogenates by electron-capture gas chromatography. *J Chromatogr* 217:239-245.
- Post EJ. 1978. Unmyelinated nerve fibers in feline acrylamide neuropathy. *Acta Neuropathol* 42(1):19-24.
- +*Post EJ, McLeod JG. 1977a. Acrylamide autonomic neuropathy in the cat: 1. Neurophysiological and histological studies. *J Neurol Sci* 33(3):353-374.
- *Post EJ, McLeod JG. 1977b. Acrylamide autonomic neuropathy in the cat: 2. Effects on mesenteric vascular control. *J Neurol Sci* 33(3):375-386.

9. REFERENCES

- *Poulin P, Krishnan K. 1995. A biologically-based algorithm for predicting human tissue:blood partition coefficients of organic chemicals. *Hum Exp Toxicol* 14:273-280.
- *Poulin P, Krishnan K. 1996a. A mechanistic algorithm for predicting blood:air partition coefficients of organic chemicals with the consideration of reversible binding in hemoglobin. *Toxicol Appl Pharmacol* 136:131-137.
- *Poulin P, Krishnan K. 1996. A tissue composition-based algorithm for predicting tissue:air partition coefficients of organic chemicals. *Toxicol Appl Pharmacol* 136:126-130.
- *Pratt GC, Palmer K, Wu CY, et al. 2000. An assessment of air toxics in Minnesota. *Environ Health Perspect* 108(9):815-825.
- Preston A, Fodey T, Douglas A, et al. 2009. Monoclonal antibody development for acrylamide-adducted human haemoglobin; a biomarker of dietary acrylamide exposure. *J Immunol Methods* 341(1-2):19-29.
- Prineas J. 1969. The pathogenesis of dying back polyneuropathies. Part II. An ultrastructural study of experimental acrylamide intoxication in the cat. *J Neuropathol Exp Neurol* 28:598-621.
- Puppel N, Tjaden Z, Fueller F, et al. 2005. DNA strand breaking capacity of acrylamide and glycidamide in mammalian cells. *Mutat Res* 580(1-2):71-80.
- Rafales LS, Bornschein RL, Caruso V. 1982. Behavioral and pharmacological responses following acrylamide exposure in rats. *Neurobehav Toxicol Teratol* 4(3):355-364.
- Rafales LS, Lasley SM, Greenland RD, et al. 1983. Effects of acrylamide on locomotion and central monoamine function in the rat. *Pharmacol Biochem Behav* 19(4):635-644.
- *Ramsey JC, Young JD, Gorzinsky SJ. 1984. Acrylamide: Toxicodynamics in rats. Submitted to the U.S. Environmental Protection Agency under TSCA Section 4. OTS0507270. [Unpublished study for peer review]
- Rasool CG, Bradley WG. 1978. Studies on axoplasmic transport of individual proteins: 1. Acetylcholinesterase (AChE) in acrylamide neuropathy. *J Neurochem* 31(2):419-426.
- Ravindranath V, Pai KS. 1991. The use of rat brain slices as an *in vitro* model for mechanistic evaluation of neurotoxicity-studies with acrylamide. *Neurotoxicology* 12(2):225-234.
- *Raymer JH, Sparacino CM, Velez GR, et al. 1993. Determination of acrylamide in rat serum and sciatic nerve by gas chromatography-electron-capture detection. *J Chromatogr* 619:223-234.
- Recio L, Caspary W, Tourous D, et al. 2006. Dose-response comparison of micronucleated reticulocyte frequencies in rodent peripheral blood with four genotoxic agents by flow cytometry and slide-based enumeration. [Abstract]. *Toxicol Sci* 90(1-S):112.
- Reid E. 2003. Science in motion: Common molecular pathological themes emerge in the hereditary spastic paraplegias. *J Med Genet* 40(2):81-86.
- Reid E, Kloos M, Ashley-Koch A, et al. 2002. A kinesin heavy chain (KIF5A) mutation in hereditary spastic paraplegia (SPG10). *Am J Hum Genet* 71(5):1189-1194.

9. REFERENCES

- *Rice JM. 2005. The carcinogenicity of acrylamide. *Mutat Res* 580(1-2):3-20.
- +*Rohm and Haas Company. 1975. Acrylamide and methacrylamide. Subchronic percutaneous toxicity study in new-born rabbits. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS205982.
- *Russell LB, Hunsicker PR, Cacheiro NA, et al. 1991. Induction of specific-locus mutations in male germ cells of the mouse by acrylamide monomer. *Mutat Res* 262:101-107.
- *Russo A, Gabbani G, Simoncini B. 1994. Weak genotoxicity of acrylamide on premeiotic and somatic cells of the mouse. *Mutat Res* 309(2):263-272.
- Rydberg P, Eriksson S, Tareke E, et al. 2005. Factors that influence the acrylamide content of heated foods. *Adv Exp Med Biol* 561:317-328.
- Sabri MI, Spencer PS. 1990. Acrylamide impairs fast and slow axonal transport in rat optic system. *Neurochem Res* 15(6):603-608.
- Sabri MI, Dairman W, Fenton M, et al. 1989. Effect of exogenous pyruvate on acrylamide neuropathy in rats. *Brain Res* 483(1):1-11.
- Sahenk Z, Mendell JR. 1981. Acrylamide and 2,5-hexanedione neuropathies: Abnormal bidirectional transport rate in distal axons. *Brain Res* 219(2):397-405.
- +*Sakamoto J, Hashimoto K. 1986. Reproductive toxicity of acrylamide and related compounds in mice: Effects on fertility and sperm morphology. *Arch Toxicol* 59:201-205.
- +*Sakamoto J, Kurosaka Y, Hashimoto K. 1988. Histological changes of acrylamide-induced testicular lesions in mice. *Exp Mol Pathol* 48(3):324-334.
- Sakurai H. 2000. Recommendation of occupational exposure limits. *J Occup Health* 42(4):213-228.
- Saleh SI, El-Okazy AM. 2007. Assessment of the mean daily dietary intake of acrylamide in Alexandria. *J Egypt Public Health Assoc* 82(3-4):331-345.
- Sanchez J, Cabrer JM, Rossello CA, et al. 2008. Formation of hemoglobin adducts of acrylamide after its ingestion in rats is dependent on age and sex. *J Agric Food Chem* 56(13):5096-5101.
- Satchell PM. 1984. Circulatory control in canine acrylamide neuropathy. *J Auton Nerv Syst* 10(2):93-106.
- Satchell PM. 1990. Baroreceptor dysfunction in acrylamide axonal neuropathy. *Brain* 113 (Pt 1):167-176.
- +*Satchell PM, McLeod JG. 1981. Megaesophagus due to acrylamide neuropathy. *J Neurol Neurosurg Psychiatry* 44(10):906-913.
- Satchell PM, McLeod JG. 1984. Abnormalities of oesophageal mechanoreceptors in canine acrylamide neuropathy. *J Neurol Neurosurg Psychiatry* 47(7):692-698.

9. REFERENCES

- Schaumburg HH, Spencer PS, Arezzo JC. 1985. Studies of the primate somatosensory system in experimental acrylamide intoxication. *Neurotoxicology* 6(4):3-6.
- Schaumburg HH, Wisniewski HM, Spencer PS. 1974. Ultrastructural studies of the dying-back process: 1. Peripheral nerve terminal and axon degeneration in systemic acrylamide intoxication. *J Neuropathol Exp Neurol* 33(2):260-284.
- *Schettgen T, Kutting B, Hornig M, et al. 2004a. Trans-placental exposure of neonates to acrylamide-a pilot study. *Int Arch Occup Environ Health* 77(3):213-216.
- Schettgen T, Rossbach B, Kutting B, et al. 2004b. Determination of haemoglobin adducts of acrylamide and glycidamide in smoking and non-smoking persons of the general population. *Int J Hyg Environ Health* 207(6):531-539.
- Schettgen T, Weiss T, Drexler H, et al. 2003. A first approach to estimate the internal exposure to acrylamide in smoking and non-smoking adults from Germany. *Int J Hyg Environ Health* 206(1):9-14.
- Schmid TE, Xu W, Adler ID. 1999. Detection of aneuploidy by multicolor FISH in mouse sperm after *in vivo* treatment with acrylamide, colchicine, diazepam or thiabendazole. *Mutagenesis* 14(2):173-179.
- Schmidt RE, Plurad SB, Clark HB. 1987. Acrylamide-induced sympathetic autonomic neuropathy resulting in pineal denervation. *Lab Invest* 56(5):505-517.
- Schotman P, Gipon L, Jennekens FG, et al. 1977. Polyneuropathies and CNS protein metabolism: II. Changes in the incorporation rate of leucine during acrylamide intoxication. *Neuropathol Appl Neurobiol* 3(2):125-136.
- Schotman P, Gipon L, Jennekens FG, et al. 1978. Polyneuropathies and CNS protein metabolism: III. Changes in protein synthesis rate induced by acrylamide intoxication. *J Neuropathol Exp Neurol* 37(6):820-837.
- Schulz MR, Hertz-Picciotto I, van Wijngaarden E, et al. 2001. Dose-response relation between acrylamide and pancreatic cancer. *Occup Environ Med* 58(9):609.
- +*Schulze GE, Boysen BG. 1991. A neurotoxicity screening battery for use in safety evaluation: Effects of acrylamide and 3',3'-iminodipropionitrile. *Fundam Appl Toxicol* 16(3):602-615.
- *Schumacher JN, Green CR, Best FW, et al. 1977. Smoke composition. An extensive investigation of the water-soluble portion of cigarette smoke. *J Agric Food Chem* 25(2):310-320.
- *Sega GA, Generoso EE. 1990. Measurement of DNA breakage in specific germ-cell stages of male mice exposed to acrylamide, using an alkaline-elution procedure. *Mutat Res* 242:79-87.
- *Sega GA, Alcota RP, Tancongo CP, et al. 1989. Acrylamide binding to the DNA and protamine of spermiogenic stages in the mouse and its relationship to genetic damage. *Mutat Res* 216(4):221-230.
- *Sega GA, Generoso EE, Brimer PA. 1990. Acrylamide exposure induces a delayed unscheduled DNA synthesis in germ cells of male mice that is correlated with the temporal pattern of adduct formation in testis DNA. *Environ Mol Mutagen* 16(3):137-142.

9. REFERENCES

- *Segerback D, Calleman CJ, Schroeder JL, et al. 1995. Formation of N-7-(2-carbamoyl-2-hydroxyethyl)guanine in DNA of the mouse and the rat following intraperitoneal administration of [¹⁴C]acrylamide. *Carcinogenesis* 16(5):1161-1165.
- *Setchell BP, Waites GMH. 1975. The blood testis barrier. In: Creep RO, Astwood EB, Geiger SR, eds. *Handbook of physiology: Endocrinology V*. Washington, DC: American Physiological Society, 143-172.
- Settels E, Bernauer U, Palavinskas R, et al. 2008. Human CYP2E1 mediates the formation of glycidamide from acrylamide. *Arch Toxicol* 82(10):717-727.
- Shabana AH, Oboeuf M, Forest N. 1994. Cytoplasmic desmosomes and intermediate filament disturbance following acrylamide treatment in cultured rat keratinocytes. *Tissue Cell* 26(1):43-55.
- *Shelby MD, Cain KT, Cornett CV, et al. 1987. Acrylamide: Induction of heritable translocation in male mice. *Environ Mutagen* 9(4):363-368.
- Shelby MD, Cain KT, Hughes LA, et al. 1986. Dominant lethal effects of acrylamide in male mice. *Mutat Res* 173(1):35-40.
- Shell L, Rozum M, Jortner BS, et al. 1992. Neurotoxicity of acrylamide and 2,5-hexanedione in rats evaluated using a functional observational battery and pathological examination. *Neurotoxicol Teratol* 14(4):273-283.
- *Shipp A, Lawrence G, Gentry R, et al. 2006. Acrylamide: Review of toxicity data and dose-response analyses for cancer and noncancer effects. *Crit Rev Toxicol* 36(6-7):481-608.
- *Shiraishi Y. 1978. Chromosome aberrations induced by monomeric acrylamide in bone marrow and germ cells of mice. *Mutat Res* 57:313-324.
- Shivakumar BR, Ravindranath V. 1992. Selective modulation of glutathione in mouse brain regions and its effect on acrylamide-induced neurotoxicity. *Biochem Pharmacol* 43(2):263-270.
- Shukla PK, Khanna VK, Ali MM, et al. 2002. Protective effect of *Acorus calamus* against acrylamide induced neurotoxicity. *Phytother Res* 16(3):256-260.
- Sickles DW. 1987. Neural specificity of acrylamide action upon enzymes associated with oxidative energy-producing pathways: I. Histochemical analysis of NADH-tetrazolium reductase activity. *Neurotoxicology* 8(4):623-630.
- Sickles DW. 1989a. Toxic neurofilamentous axonopathies and fast anterograde axonal transport: I. The effects of single doses of acrylamide on the rate and capacity of transport. *Neurotoxicology* 10(1):91-102.
- Sickles DW. 1989b. Toxic neurofilamentous axonopathies and fast anterograde axonal transport. II. The effects of single doses of neurotoxic and non-neurotoxic diketones and B,B'-iminodipropionitrile (IDPN) on the rate and capacity of transport. *Neurotoxicology* 10:103-112.

9. REFERENCES

- *Sickles DW. 1991. Toxic neurofilamentous axonopathies and fast anterograde axonal transport. III. Recovery from single injections and multiple dosing effects of acrylamide and 2,5-hexanedione. *Toxicol Appl Pharmacol* 108(3):390-396.
- Sickles DW. 1992. Toxic neurofilamentous axonopathies and fast anterograde axonal transport. IV. *In vitro* analysis of transport following acrylamide and 2,5-hexanedione. *Toxicol Lett* 61(2-3):199-204.
- Sickles DW, Goldstein BD. 1985. Acrylamide alters oxidative enzyme activity in rat motoneurons. *Toxicol Lett* 26(2-3):111-118.
- Sickles DW, Goldstein BD. 1986. Acrylamide produces a direct, dose-dependent and specific inhibition of oxidative metabolism in motoneurons. *Neurotoxicology* 7(1):187-196.
- Sickles DW, Testino A. 2005. Effects of subacute and chronic acrylamide dosing on fast-transported proteins into rat neuromuscular junctions [Abstract]. *Toxicol Sci* 84(1-S):316-317.
- *Sickles DW, Brady ST, Testino A, et al. 1996. Direct effect of the neurotoxicant acrylamide on kinesin-based microtubule motility. *J Neurosci Res* 46(1):7-17.
- Sickles DW, Sperry AO, Testino A, et al. 2007. Acrylamide effects on kinesin-related proteins of the mitotic/meiotic spindle. *Toxicol Appl Pharmacol* 222(1):111-121.
- Sickles DW, Stone JD, Friedman MA. 2002. Fast axonal transport: A site of acrylamide neurotoxicity? *Neurotoxicology* 23(2):223-251.
- Sickles DW, Welter DA, Friedman MA. 1995. Acrylamide arrests mitosis and prevents chromosome migration in the absence of changes in spindle microtubules. *J Toxicol Environ Health* 44(1):73-86.
- Sidenius P, Jakobsen J. 1983. Anterograde axonal transport in rats during intoxication with acrylamide. *J Neurochem* 40(3):697-704.
- Simmons JE, Berman E, Jackson M, et al. 1987. In-vitro and in-vivo toxicity a comparison of acrylamide cyclophosphamide chlordecone and diethylstilbestrol. *J Environ Sci Health A Environ Sci Eng Toxic Hazard Sub* 22(7):639-664.
- Sisti R, Lorenzon G, Cicales R. 2001. Neurotoxic effects of acrylamide in Sprague-Dawley rats by oral administration. [Abstract]. *Front Fetal Health* 3(11-12):288.
- Smith EA, Oehme FW. 1991. Acrylamide and polyacrylamide a review of production use environmental fate and neurotoxicity. *Rev Environ Health* 9(4):215-225.
- Smith CJ, Perfetti TA, Rumble MA, et al. 2000. "IARC Group 2A Carcinogens" reported in cigarette mainstream smoke. *Food Chem Toxicol* 38(4):371-383.
- +*Smith MK, Zenick H, Preston RJ, et al. 1986. Dominant lethal effects of subchronic acrylamide administration in the male Long-Evans rat. *Mutat Res* 173(4):273-277.
- SNF S.A.. 2000. Toxicological review of acrylamide (CAS No. 79-06-01). Ruston, Louisiana. Environ International Corporation. SNF S. A. Saint-Etienne Cedex, France. November 2002.

9. REFERENCES

- +*Sobel W, Bond GG, Parsons TW, et al. 1986. Acrylamide cohort mortality study. *Br J Ind Med* 43(11):785-788.
- *Solomon JJ, Fedyk J, Mukai F, et al. 1985. Direct alkylation of 2'-deoxynucleosides and DNA following *in vitro* reaction with acrylamide. *Cancer Res* 45:3465-3470.
- *Sörgel F, Weissenbacher R, Kinzig-Schippers M, et al. 2002. Acrylamide: Increased concentrations in homemade food and first evidence of its variable absorption from food, variable metabolism and placental and breast milk transfer in humans. *Chemotherapy* 48(6):267-274.
- Souyri F, Chretien M, Droz B. 1981. Acrylamide-induced neuropathy and impairment of axonal transport of proteins: 1. Multifocal retention of fast transported proteins at the periphery of axons as revealed by light microscope radioautography. *Brain Res* 205(1):1-13.
- Spano M, Cordelli E, Leter G, et al. 1993. Genotoxic and cytotoxic effects of acrylamide on male germ cells. [Abstract]. *Eur J Histochem* 37(Suppl 2):84.
- *Spencer PS, Schaumburg HH. 1974. A review of acrylamide neurotoxicity. Part I. Properties, uses and human exposure. *Can J Neurol Sci* 1(2):143-150.
- *Spencer PS, Schaumburg HH. 1977. Ultrastructural studies of the dying back process. IV. Differential vulnerability of PNS and CNS fibers in experimental central-peripheral distal axonopathies. *J Neuropath Exp Neurol* 36:300-320.
- Spencer PS, Schaumburg HH. 1991. A review of acrylamide neurotoxicity. Part 2. Experimental animal neurotoxicity and pathologic mechanisms. *Can J Neurol Sci* 1(3):152-169.
- Spencer PS, Schaumburg HH. 1988. Analysis of neurotoxic disease: The acrylamide experience. In: Galli CL, Manzo L, Spencer PS, eds. *Recent advances in nervous system toxicology*. New York, NY: Plenum Press, 163-176.
- *Spencer PS, Sabri MI, Schaumburg HH, et al. 1979. Does a defect of energy metabolism in the nerve fiber underlie axonal degeneration in polyneuropathies? *Ann Neurol* 5:501-507.
- *SRI. 2008. Acrylamide. In: 2008 Directory of chemical producers. Menlo Park, CA: SRI Consulting, 410.
- Srivastava SP, Das M, Mukhtar H, et al. 1984. Effect of acrylamide on glutathione-s-transferase activity in different regions of rat brain. *Bull Environ Contam Toxicol* 32(2):166-170.
- Srivastava SP, Das M, Seth PK. 1983. Enhancement of lipid peroxidation in rat liver on acute exposure to styrene and acrylamide a consequence of glutathione depletion. *Chem Biol Interact* 45(3):373-380.
- Srivastava SP, Sabri MI, Agrawal AK, et al. 1986. Effect of single and repeated doses of acrylamide and bisacrylamide on glutathione-s-transferase and dopamine receptors in rat brain. *Brain Res* 371(2):319-323.
- Srivastava SP, Seth PK, Das M, et al. 1985. Effects of mixed-function oxidase modifiers on neurotoxicity of acrylamide in rats. *Biochem Pharmacol* 34(7):1099-1102.

9. REFERENCES

- Stadler RH. 2005. Acrylamide formation in different foods and potential strategies for reduction. *Adv Exp Med Biol* 561:157-169.
- Stadler RH, Anklam E. 2007. Update on the progress in acrylamide and furan research. *Food Addit Contam* 24(Suppl 1):1-2.
- *Stadler RH, Blank I, Varga N, et al. 2002. Acrylamide from Maillard reaction products. *Nature* 419(6906):449-450.
- *Stamler JS, Lamas S, Fang C. 2001. Nitrosylation: the prototypic redox-based signaling mechanism. *Cell* 106(6):675-683.
- Sterman AB. 1983. Altered sensory ganglia in acrylamide neuropathy: Quantitative evidence of neuronal reorganization. *J Neuropathol Exp Neurol* 42(2):166-176.
- Sterman AB, Panasci DJ, Persons W. 1983a. Does pyruvate prevent acrylamide neurotoxicity? Implications for disease pathogenesis. *Exp Neurol* 82(1):148-158.
- Sterman AB, Panasci DJ, Sheppard RC. 1983b. Autonomic-cardiovascular dysfunction accompanies sensory-motor impairment during acrylamide intoxication. *Neurotoxicology* 4(1):45-52.
- Stetkiewicz J, Wronska-Nofer T, Klimczak J, et al. 1988. Metabolic interaction and neurological effect of combined exposure to acrylamide and ethanol. *Pol J Occup Med* 1(2):127-136.
- Stobiecka A, Radecka H, Radecki J. 2007. Novel voltammetric biosensor for determining acrylamide in food samples. *Biosens Bioelectron* 22(9-10):2165-2170.
- +*Sublet VH, Zenick H, Smith MK. 1989. Factors associated with reduced fertility and implantation rates in females mated to acrylamide-treated rats. *Toxicology* 55:53-67.
- Sumizawa T, Igisu H. 2007. Apoptosis induced by acrylamide in SH-SY5Y cells. *Arch Toxicol* 81(4):279-282.
- Sumizawa T, Igisu H. 2008. Release of heat shock proteins from human neuroblastoma cells exposed to acrylamide. *J Toxicol Sci* 33(1):117-122.
- Sumner AJ, Asbury AK. 1975. Physiological studies of the dying-back phenomenon. Muscle stretch afferents in acrylamide neuropathy. *Brain* 98:91-100.
- Sumner A, Pleasure D, Ciesielka K. 1976. Slowing of fast axoplasmic transport in acrylamide neuropathy. *J Neuropathol Exp Neurol* 35(3):319.
- *Sumner SC, Fennell TR, Moore TA, et al. 1999. Role of cytochrome P450 2E1 in the metabolism of acrylamide and acrylonitrile in mice. *Chem Res Toxicol* 12(11):1110-1116.
- *Sumner SC, Macneela JP, Fennell TR. 1992. Characterization and quantitation of urinary metabolites of (1,2,3-¹³carbon)acrylamide in rats and mice using ¹³carbon nuclear magnetic resonance spectroscopy. *Chem Res Toxicol* 5(1):81-89.

9. REFERENCES

- *Sumner SC, Williams CC, Snyder RW, et al. 2003. Acrylamide: A comparison of metabolism and hemoglobin adducts in rodents following dermal, intraperitoneal, oral, or inhalation exposure. *Toxicol Sci* 75(2):260-270.
- *Suzuki K, Pfaff L. 1973. Acrylamide neuropathy in rats. *Acta Neuropathol (Berl)* 24(3):197-213.
- *Svensson K, Abramsson L, Becker W, et al. 2003. Dietary intake of acrylamide in Sweden. *Food Chem Toxicol* 41(11):1581-1586.
- *Swaen GM, Haidar S, Burns CJ, et al. 2007. Mortality study update of acrylamide workers. *Occup Environ Med* 64(6):396-401.
- *Takahashi M, Ohara T, Hashimoto K. 1971. Electrophysiological study of nerve injuries in workers handling acrylamide. *Int Arch Arbeitsmed* 28(1):1-11.
- Takahashi M, Shibutani M, Inoue K, et al. 2008. Pathological assessment of the nervous and male reproductive systems of rat offspring exposed maternally to acrylamide during the gestation and lactation periods-a preliminary study. *J Toxicol Sci* 33(1):11-24.
- *Takahashi M, Shibutani M, Nakahigashi J, et al. 2009. Limited lactational transfer of acrylamide to rat offspring on maternal oral administration during the gestation and lactation periods. *Arch Toxicol* [March 21; Epub ahead of print]
- Tandrup T, Braendgaard H. 1994. Number and volume of rat dorsal root ganglion cells in acrylamide intoxication. *J Neurocytol* 23(4):242-248.
- Tanii H, Hashimoto K. 1981. *In vitro* metabolism of acrylamide and related compounds. *Arch Toxicol* 48(2-3):157-166.
- +*Tanii H, Hashimoto K. 1983. Neurotoxicity of acrylamide and related compounds in rats: Effects on rotarod performance, morphology of nerves and neurotubulin. *Arch Toxicol* 54(3):203-214.
- Tanii H, Hashimoto K. 1985. Effect of acrylamide and related compounds on glycolytic enzymes of rat brain. *Toxicol Lett* 26(1):76-84.
- Tanii H, Hashimoto K. 1991. *In vitro* neurotoxicity study with dorsal root ganglia for acrylamide and its derivatives. *Toxicol Lett* 58(2):209-213.
- Tanii H, Hayashi M, Hashimoto K. 1988. Neurofilament degradation in the nervous system of rats intoxicated with acrylamide related compounds or 2,5-hexanedione. *Arch Toxicol* 62(1):70-75.
- Tanii H, Miki N, Hayashi M, et al. 1988. Cytotoxicity of acrylamide and related compounds to mouse neuroblastoma and rat Schwannoma cells. *Arch Toxicol* 61(4):298-305.
- Tareke E, Lyn-Cook B, Robinson B, et al. 2008. Acrylamide: a dietary carcinogen formed *in vivo*? *J Agric Food Chem* 56(15):6020-6023.
- *Tareke E, Rydberg P, Karlsson P, et al. 2002. Analysis of acrylamide, a carcinogen formed in heated foodstuffs. *J Agric Food Chem* 50(17):4998-5006.

9. REFERENCES

- Tareke E, Twaddle NC, McDaniel LP, et al. 2006. Relationships between biomarkers of exposure and toxicokinetics in Fischer 344 rats and B6C3F1 mice administered single doses of acrylamide and glycidamide and multiple doses of acrylamide. *Toxicol Appl Pharmacol* 217(1):63-75.
- Taubert D, Glockner R, Muller D, et al. 2006. The garlic ingredient diallyl sulfide inhibits cytochrome P450 2E1 dependent bioactivation of acrylamide to glycidamide. *Toxicol Lett* 164(1):1-5.
- Taylor DD, Jortner BS, Walton A, et al. 1995. An immunohistochemical study of the cytoskeleton of SH-SY5Y human neuroblastoma cells, and the effects of acrylamide. *In Vitro Toxicol* 8(3):323-333.
- Thomann P, Koella WP, Krinke G, et al. 1974. The assessment of peripheral neurotoxicity in dogs: Comparative studies with acrylamide and clioquinol. *Inflamm Res* 4(1):47-53.
- *Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. *Chemically induced alterations in sexual and functional development: The wildlife/human connection*. Princeton, NJ: Princeton Scientific Publishing, 365-394.
- +*Tilson HA, Cabe PA. 1979. The effects of acrylamide given acutely or in repeated doses on fore- and hindlimb function of rats. *Toxicol Appl Pharmacol* 47(2):253-260.
- Tilson HA, Cabe PA, Spencer PS. 1979. Acrylamide neurotoxicity in rats: A correlated neurobehavioral and pathological study. *Neurotoxicology* 1(1):89-104.
- *Tong GC, Cornwell WK, Means GE. 2004. Reactions of acrylamide with glutathione and serum albumin. *Toxicol Lett* 147:127-131.
- Tornqvist M, Paulsson B, Vikstrom AC, et al. 2008. Approach for cancer risk estimation of acrylamide in food on the basis of animal cancer tests and *in vivo* dosimetry. *J Agric Food Chem* 56(15):6004-6012.
- Totani N, Yawata M, Ojiri Y, et al. 2007. Effects of trace acrylamide intake in Wistar rats. *J Oleo Sci* 56(9):501-506.
- *TRI 07 2009. 2009. TRI Explorer. Providing access to EPA's toxics release inventory data. U. S. Environmental Protection Agency. <http://www.epa.gov/triexplorer>. March 18, 2009.
- *Tripathy NK, Patnaik KK, Nabi MJ. 1991. Acrylamide is genotoxic to the somatic and germ cells of *Drosophila melanogaster*. *Mutat Res* 259(1):21-27.
- *Tsuda H, Shimizu CS, Taketomi MK, et al. 1993. Acrylamide; induction of DNA damage, chromosomal aberrations and cell transformation without gene mutations. *Mutagenesis* 8(1):23-29.
- *Twaddle NC, McDaniel LP, Gamboa da Costa G, et al. 2004. Determination of acrylamide and glycidamide serum toxicokinetics in B6C3F₁ mice using LC-ES/MS/MS. *Cancer Lett* 207(1):9-17.
- *Tyl RW, Friedman MA. 2003. Effects of acrylamide on rodent reproductive performance. *Reprod Toxicol* 17(1):1-13.
- +*Tyl RW, Friedman MA, Losco PE, et al. 2000a. Rat two-generation reproduction and dominant lethal study of acrylamide in drinking water. *Reprod Toxicol* 14(5):385-401.

9. REFERENCES

- +*Tyl RW, Marr MC, Myers CB, et al. 2000b. Relationship between acrylamide reproductive and neurotoxicity in male rats. *Reprod Toxicol* 14(2):147-157.
- +*Union Carbide Corporation. 1947. Range finding tests on acrylamide with cover letter. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. EPA878212200. OTS206058.
- U.S. Army. 2007. A Population-based study of dietary acrylamide and prostate cancer risk. Fort Detrick, MD: U.S. Army Medical Research and Materiel Command. W81XWH-04-1-0288.
- Uphouse LL, Nemeroff CB, Mason G, et al. 1982. Interactions between "handling" and acrylamide on endocrine responses in rats. *Neurotoxicology* 3(1):121-125.
- Urban M, Kavvadias D, Riedel K, et al. 2006. Urinary mercapturic acids and a hemoglobin adduct for the dosimetry of acrylamide exposure in smokers and nonsmokers. *Inhal Toxicol* 18(10):831-839.
- Valdivia RP, Lafuente NM, Katoh M. 1989. Acrylamide-induced chromosome-type aberrations in spermiogenic stages evaluated in the first cleavage metaphases in the mouse. [Abstract]. *Environ Mol Mutagen* 14(Suppl 15):205.
- *van Dijk-Looijaard AM, van Genderen J. 2000. Levels of exposure from drinking water. *Food Chem Toxicol* 38(Suppl. 1):S37-S42.
- Vanhorick M, Moens W. 1983. Carcinogen-mediated induction of SV40 DNA amplification is enhanced by acrylamide in Chinese hamster CO60 cells. *Carcinogenesis* 4(11):1459-1463.
- Vesper HW, Bernert JT, Ospina M, et al. 2007. Assessment of the relation between biomarkers for smoking and biomarkers for acrylamide exposure in humans. *Cancer Epidemiol Biomarkers Prev* 16(11):2471-2478.
- Vesper HW, Slimani N, Hallmans G, et al. 2008. Cross-sectional study on acrylamide hemoglobin adducts in subpopulations from the European Prospective Investigation into Cancer and Nutrition (EPIC) Study. *J Agric Food Chem* 56(15):6046-6053.
- *Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238(2):476-483.
- Vikstrom AC, Eriksson S, Paulsson B, et al. 2008. Internal doses of acrylamide and glycidamide in mice fed diets with low acrylamide contents. *Mol Nutr Food Res* 52(8):974-980.
- *Von Tungeln LS, Churchwell MI, Doerge DR, et al. 2009. DNA adduct formation and induction of micronuclei and mutations in B6C3F1/Tk mice treated neonatally with acrylamide or glycidamide. *Int J Cancer* 124(9):2006-2015.
- Voogd CE, van der Stel JJ, Jacobs JJ. 1981. The mutagenic action of aliphatic epoxides. *Mutat Res* 89(4):269-282.
- Walden R, Schiller CM. 1981. Quantitative analysis of acrylamide in the milk of lactating rats following oral in-vivo exposure. [Abstract]. *Fed Proc* 40(3 Part 1):678.

9. REFERENCES

- +*Walden R, Squibb RE, Schiller CM. 1981. Effects of prenatal and lactational exposure to acrylamide on the development of intestinal enzymes in the rat. *Toxicol Appl Pharmacol* 58:363-369.
- *Walker K, Hattis D, Russ A, et al. 2007. Approaches to acrylamide physiologically based toxicokinetic modeling for exploring child-adult dosimetry differences. *J Toxicol Environ Health A* 70(24):2033-2055.
- Walker RA, O'Brien ET, Epstein DL, et al. 1997. N-ethylmaleimide and ethacrynic acid inhibit kinesin binding to microtubules in a motility assay. *Cell Motil Cytoskeleton* 37(4):289-299.
- Walum E, Flint OP. 1988. Sub-teratogenic neurotoxic effects of acrylamide. [Abstract]. *Teratology* 37(5):499.
- Walum E, Flint OP. 1993. Selective effects of acrylamide, methylene bisacrylamide, and haloperidol on neuronal development in rat embryo midbrain micromass cultures. *In Vitro Toxicol* 6(2):125-134.
- *Warr TJ, Parry JM, Callander RD, et al. 1990. Methyl vinyl sulphone: A new class of Michael-type genotoxin. *Mutat Res* 245:191-199.
- Weir RL, Glaubiger G, Chase TN. 1978. Inhibition of fast axoplasmic transport by acrylamide. *Environ Res* 17(2):251-255.
- Wenzl T, Anklam E. 2007. European Union database of acrylamide levels in food: Update and critical review of data collection. *Food Addit Contam* 24 (Suppl 1):5-12.
- *West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.
- *WHO. 2000. Air quality guidelines. 2nd edition. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/air/activities/20050223_4. May 11, 2009.
- *WHO. 2002. Health implications of acrylamide in food. World Health Organization. http://www.who.int/foodsafety/publications/chem/en/acrylamide_full.pdf. May 5, 2009.
- *WHO. 2003. Acrylamide in drinking-water. World Health Organization. WHO/SDE/WSH/03.04/71. http://www.who.int/water_sanitation_health/dwq/chemicals/acrylamide.pdf. May 5, 2009.
- *WHO. 2006. Guidelines for drinking-water quality. 3rd edition. Geneva, Switzerland: World Health Organization. http://www.who.int/water_sanitation_health/dwq/gdwq3/en/. May 11, 2009.
- *Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. *Mineral metabolism: An advance treatise. Volume II: The elements Part A*. New York, NY: Academic Press, 1-247.
- Wild HM, Kulikowski JJ. 1984. Neurotoxic effects of acrylamide on rat retinogeniculate fibres. *Behav Brain Res* 13(3):201-207.
- +*Wilson KM, Balter K, Adami HO, et al. 2009a. Acrylamide exposure measured by food frequency questionnaire and hemoglobin adduct levels and prostate cancer risk in the cancer of the prostate in Sweden study. *Int J Cancer* 124(10):2384-2390.

9. REFERENCES

- +*Wilson KM, Mucci LA, Cho E, et al. 2009b. Dietary acrylamide intake and risk of premenopausal breast cancer. *Am J Epidemiol* 169(8):954-961.
- Wilson KM, Vesper HW, Tocco P, et al. 2009c. Validation of a food frequency questionnaire measurement of dietary acrylamide intake using hemoglobin adducts of acrylamide and glycidamide. *Cancer Causes Control* 20(3):269-278.
- Wirfalt E, Paulsson B, Tornqvist M, et al. 2008. Associations between estimated acrylamide intakes, and hemoglobin AA adducts in a sample from the Malmö diet and cancer cohort. *Eur J Clin Nutr* 62(3):314-323.
- +*Wise LD, Gordon LR, Soper KA, et al. 1995. Developmental neurotoxicity evaluation of acrylamide in Sprague-Dawley rats. *Neurotoxicol Teratol* 17(2):189-198.
- Wispiyono B, Matsuoka M, Igisu H. 2004. Acrylamide does not cause apparent changes in genetic expression of creatine kinase in rat cerebellum. *Journal of UOEH* 26(1):51-57.
- Woo GH, Shibutani M, Kuroiwa K, et al. 2007. Lack of preventive effects of dietary fibers or chlorophyllin against acrylamide toxicity in rats. *Food Chem Toxicol* 45(8):1507-1515.
- +*Working PK, Bentley KS, Hurtt ME, et al. 1987. Dominant lethal assay of acrylonitrile and acrylamide in the male rat. [Abstract]. *Environ Mutagen* 9(Suppl 8):115.
- Wu Q, Sidoryk M, Mutkus L, et al. 2005. Acrylamide stimulates glutamine uptake in Fischer 344 rat astrocytes by a mechanism involving upregulation of the amino acid transport system N. *Ann N Y Acad Sci* 1053:435-443.
- *Xiao Y, Tates AD. 1994. Increased frequencies of micronuclei in early spermatids of rats following exposure of young primary spermatocytes to acrylamide. *Mutat Res* 309(2):245-253.
- Xie Q, Liu Y, Sun H, et al. 2008. Inhibition of acrylamide toxicity in mice by three dietary constituents. *J Agric Food Chem* 56(15):6054-6060.
- Xie Q, Sun H, Liu Y, et al. 2006. Adduction of biomacromolecules with acrylamide (AA) in mice at environmental dose levels studied by accelerator mass spectrometry. *Toxicol Lett* 163(2):101-108.
- Yang HJ, Lee SH, Jin Y, et al. 2005a. Genotoxicity and toxicological effects of acrylamide on reproductive system in male rats. *J Vet Sci* 6(2):103-109.
- Yang HJ, Lee SH, Jin Y, et al. 2005b. Toxicological effects of acrylamide on rat testicular gene expression profile. *Reprod Toxicol* 19(4):527-534.
- Yaylayan VA, Stadler RH. 2005. Acrylamide formation in food: A mechanistic perspective. *J AOAC Int* 88(1):262-267, 253.
- Yi C, Xie K, Song F, et al. 2006. The changes of cytoskeletal proteins in plasma of acrylamide-induced rats. *Neurochem Res* 31(6):751-757.
- *Young JF, Luecke RH, Doerge DR. 2007. Physiologically based pharmacokinetic/pharmacodynamic model for acrylamide and its metabolites in mice, rats, and humans. *Chem Res Toxicol* 20(3):388-399.

9. REFERENCES

- Yousef MI, El-Demerdash FM. 2006. Acrylamide-induced oxidative stress and biochemical perturbations in rats. *Toxicology* 219(1-3):133-141.
- Youssef AF, Santi BW. 1997. Simple neurobehavioral functional observational battery and objective gait analysis validation by the use of acrylamide and methanol with a built-in recovery period. *Environ Res* 73(1-2):52-62.
- Youssif MI, Ismail AA, Fayed MA, et al. 1995. Effect of acrylamide on the central nervous system of rats histochemical and histopathological studies. *J Med Res Inst* 16(4):1-17.
- Yu S, Son F, Yu J, et al. 2006. Acrylamide alters cytoskeletal protein level in rat sciatic nerves. *Neurochem Res* 31(10):1197-1204.
- Yuan Y, Chen F, Zhao GH, et al. 2007. A comparative study of acrylamide formation induced by microwave and conventional heating methods. *J Food Sci* 72(4):C212-C216.
- Zaidi SI, Raisuddin S, Singh KP, et al. 1994. Acrylamide induced immunosuppression in rats and its modulation by 6-MFA, an interferon inducer. *Immunopharmacol Immunotoxicol* 16(2):247-260.
- *Zeiger, E, Anderson, B, Haworth, S, et al. 1987. *Salmonella* mutagenicity tests. III. Results from the testing of 255 chemicals. *Environ Mutagen* 9(Suppl 9):1-110.
- *Zeiger E, Recio L, Fennell TR, et al. 2009. Investigation of the low-dose response in the *in vivo* induction of micronuclei and adducts by acrylamide. *Toxicol Sci* 107(1):247-257.
- +*Zenick H, Hope E, Smith MK. 1986. Reproductive toxicity associated with acrylamide treatment in male and female rats. *J Toxicol Environ Health* 17(4):457-472.
- Zhang X, Jiang L, Geng C, et al. 2008. Inhibition of acrylamide genotoxicity in human liver-derived HepG2 cells by the antioxidant hydroxytyrosol. *Chem Biol Interact* 176(2-3):173-178.
- Zhang X, Zhao C, Jie B. 2009. Various dietary polyunsaturated fatty acids modulate acrylamide-induced preneoplastic urothelial proliferation and apoptosis in mice. *Exp Toxicol Pathol* [Jan 29; Epub ahead of print]
- *Zhu YJ, Zeng T, Zhu YB, et al. 2008. Effects of acrylamide on the nervous tissue antioxidant system and sciatic nerve electrophysiology in the rat. *Neurochem Res* 33(11):2310-2317.
- *Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. *Pediatr Res* 12(1):29-34.
- Zodl B, Schmid D, Wassler G, et al. 2007. Intestinal transport and metabolism of acrylamide. *Toxicology* 232(1-2):99-108.

9. REFERENCES

This page is intentionally blank.

10. GLOSSARY

Absorption—The taking up of liquids by solids, or of gases by solids or liquids.

Acute Exposure—Exposure to a chemical for a duration of 14 days or less, as specified in the Toxicological Profiles.

Adsorption—The adhesion in an extremely thin layer of molecules (as of gases, solutes, or liquids) to the surfaces of solid bodies or liquids with which they are in contact.

Adsorption Coefficient (K_{oc})—The ratio of the amount of a chemical adsorbed per unit weight of organic carbon in the soil or sediment to the concentration of the chemical in solution at equilibrium.

Adsorption Ratio (K_d)—The amount of a chemical adsorbed by sediment or soil (i.e., the solid phase) divided by the amount of chemical in the solution phase, which is in equilibrium with the solid phase, at a fixed solid/solution ratio. It is generally expressed in micrograms of chemical sorbed per gram of soil or sediment.

Benchmark Dose (BMD)—Usually defined as the lower confidence limit on the dose that produces a specified magnitude of changes in a specified adverse response. For example, a BMD_{10} would be the dose at the 95% lower confidence limit on a 10% response, and the benchmark response (BMR) would be 10%. The BMD is determined by modeling the dose response curve in the region of the dose response relationship where biologically observable data are feasible.

Benchmark Dose Model—A statistical dose-response model applied to either experimental toxicological or epidemiological data to calculate a BMD.

Bioconcentration Factor (BCF)—The quotient of the concentration of a chemical in aquatic organisms at a specific time or during a discrete time period of exposure divided by the concentration in the surrounding water at the same time or during the same period.

Biomarkers—Broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility.

Cancer Effect Level (CEL)—The lowest dose of chemical in a study, or group of studies, that produces significant increases in the incidence of cancer (or tumors) between the exposed population and its appropriate control.

Carcinogen—A chemical capable of inducing cancer.

Case-Control Study—A type of epidemiological study that examines the relationship between a particular outcome (disease or condition) and a variety of potential causative agents (such as toxic chemicals). In a case-controlled study, a group of people with a specified and well-defined outcome is identified and compared to a similar group of people without outcome.

Case Report—Describes a single individual with a particular disease or exposure. These may suggest some potential topics for scientific research, but are not actual research studies.

10. GLOSSARY

Case Series—Describes the experience of a small number of individuals with the same disease or exposure. These may suggest potential topics for scientific research, but are not actual research studies.

Ceiling Value—A concentration of a substance that should not be exceeded, even instantaneously.

Chronic Exposure—Exposure to a chemical for 365 days or more, as specified in the Toxicological Profiles.

Cohort Study—A type of epidemiological study of a specific group or groups of people who have had a common insult (e.g., exposure to an agent suspected of causing disease or a common disease) and are followed forward from exposure to outcome. At least one exposed group is compared to one unexposed group.

Cross-sectional Study—A type of epidemiological study of a group or groups of people that examines the relationship between exposure and outcome to a chemical or to chemicals at one point in time.

Data Needs—Substance-specific informational needs that if met would reduce the uncertainties of human health assessment.

Developmental Toxicity—The occurrence of adverse effects on the developing organism that may result from exposure to a chemical prior to conception (either parent), during prenatal development, or postnatally to the time of sexual maturation. Adverse developmental effects may be detected at any point in the life span of the organism.

Dose-Response Relationship—The quantitative relationship between the amount of exposure to a toxicant and the incidence of the adverse effects.

Embryotoxicity and Fetotoxicity—Any toxic effect on the conceptus as a result of prenatal exposure to a chemical; the distinguishing feature between the two terms is the stage of development during which the insult occurs. The terms, as used here, include malformations and variations, altered growth, and *in utero* death.

Environmental Protection Agency (EPA) Health Advisory—An estimate of acceptable drinking water levels for a chemical substance based on health effects information. A health advisory is not a legally enforceable federal standard, but serves as technical guidance to assist federal, state, and local officials.

Epidemiology—Refers to the investigation of factors that determine the frequency and distribution of disease or other health-related conditions within a defined human population during a specified period.

Genotoxicity—A specific adverse effect on the genome of living cells that, upon the duplication of affected cells, can be expressed as a mutagenic, clastogenic, or carcinogenic event because of specific alteration of the molecular structure of the genome.

Half-life—A measure of rate for the time required to eliminate one half of a quantity of a chemical from the body or environmental media.

Immediately Dangerous to Life or Health (IDLH)—The maximum environmental concentration of a contaminant from which one could escape within 30 minutes without any escape-impairing symptoms or irreversible health effects.

10. GLOSSARY

Immunologic Toxicity—The occurrence of adverse effects on the immune system that may result from exposure to environmental agents such as chemicals.

Immunological Effects—Functional changes in the immune response.

Incidence—The ratio of individuals in a population who develop a specified condition to the total number of individuals in that population who could have developed that condition in a specified time period.

Intermediate Exposure—Exposure to a chemical for a duration of 15–364 days, as specified in the Toxicological Profiles.

In Vitro—Isolated from the living organism and artificially maintained, as in a test tube.

In Vivo—Occurring within the living organism.

Lethal Concentration_(LO) (LC_{LO})—The lowest concentration of a chemical in air that has been reported to have caused death in humans or animals.

Lethal Concentration₍₅₀₎ (LC₅₀)—A calculated concentration of a chemical in air to which exposure for a specific length of time is expected to cause death in 50% of a defined experimental animal population.

Lethal Dose_(LO) (LD_{LO})—The lowest dose of a chemical introduced by a route other than inhalation that has been reported to have caused death in humans or animals.

Lethal Dose₍₅₀₎ (LD₅₀)—The dose of a chemical that has been calculated to cause death in 50% of a defined experimental animal population.

Lethal Time₍₅₀₎ (LT₅₀)—A calculated period of time within which a specific concentration of a chemical is expected to cause death in 50% of a defined experimental animal population.

Lowest-Observed-Adverse-Effect Level (LOAEL)—The lowest exposure level of chemical in a study, or group of studies, that produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.

Lymphoreticular Effects—Represent morphological effects involving lymphatic tissues such as the lymph nodes, spleen, and thymus.

Malformations—Permanent structural changes that may adversely affect survival, development, or function.

Minimal Risk Level (MRL)—An estimate of daily human exposure to a hazardous substance that is likely to be without an appreciable risk of adverse noncancer health effects over a specified route and duration of exposure.

Modifying Factor (MF)—A value (greater than zero) that is applied to the derivation of a Minimal Risk Level (MRL) to reflect additional concerns about the database that are not covered by the uncertainty factors. The default value for a MF is 1.

10. GLOSSARY

Morbidity—State of being diseased; morbidity rate is the incidence or prevalence of disease in a specific population.

Mortality—Death; mortality rate is a measure of the number of deaths in a population during a specified interval of time.

Mutagen—A substance that causes mutations. A mutation is a change in the DNA sequence of a cell's DNA. Mutations can lead to birth defects, miscarriages, or cancer.

Necropsy—The gross examination of the organs and tissues of a dead body to determine the cause of death or pathological conditions.

Neurotoxicity—The occurrence of adverse effects on the nervous system following exposure to a chemical.

No-Observed-Adverse-Effect Level (NOAEL)—The dose of a chemical at which there were no statistically or biologically significant increases in frequency or severity of adverse effects seen between the exposed population and its appropriate control. Effects may be produced at this dose, but they are not considered to be adverse.

Octanol-Water Partition Coefficient (K_{ow})—The equilibrium ratio of the concentrations of a chemical in *n*-octanol and water, in dilute solution.

Odds Ratio (OR)—A means of measuring the association between an exposure (such as toxic substances and a disease or condition) that represents the best estimate of relative risk (risk as a ratio of the incidence among subjects exposed to a particular risk factor divided by the incidence among subjects who were not exposed to the risk factor). An OR of greater than 1 is considered to indicate greater risk of disease in the exposed group compared to the unexposed group.

Organophosphate or Organophosphorus Compound—A phosphorus-containing organic compound and especially a pesticide that acts by inhibiting cholinesterase.

Permissible Exposure Limit (PEL)—An Occupational Safety and Health Administration (OSHA) allowable exposure level in workplace air averaged over an 8-hour shift of a 40-hour workweek.

Pesticide—General classification of chemicals specifically developed and produced for use in the control of agricultural and public health pests.

Pharmacokinetics—The dynamic behavior of a material in the body, used to predict the fate (disposition) of an exogenous substance in an organism. Utilizing computational techniques, it provides the means of studying the absorption, distribution, metabolism, and excretion of chemicals by the body.

Pharmacokinetic Model—A set of equations that can be used to describe the time course of a parent chemical or metabolite in an animal system. There are two types of pharmacokinetic models: data-based and physiologically-based. A data-based model divides the animal system into a series of compartments, which, in general, do not represent real, identifiable anatomic regions of the body, whereas the physiologically-based model compartments represent real anatomic regions of the body.

Physiologically Based Pharmacodynamic (PBPD) Model—A type of physiologically based dose-response model that quantitatively describes the relationship between target tissue dose and toxic end

10. GLOSSARY

points. These models advance the importance of physiologically based models in that they clearly describe the biological effect (response) produced by the system following exposure to an exogenous substance.

Physiologically Based Pharmacokinetic (PBPK) Model—Comprised of a series of compartments representing organs or tissue groups with realistic weights and blood flows. These models require a variety of physiological information: tissue volumes, blood flow rates to tissues, cardiac output, alveolar ventilation rates, and possibly membrane permeabilities. The models also utilize biochemical information, such as air/blood partition coefficients, and metabolic parameters. PBPK models are also called biologically based tissue dosimetry models.

Prevalence—The number of cases of a disease or condition in a population at one point in time.

Prospective Study—A type of cohort study in which the pertinent observations are made on events occurring after the start of the study. A group is followed over time.

q_1^* —The upper-bound estimate of the low-dose slope of the dose-response curve as determined by the multistage procedure. The q_1^* can be used to calculate an estimate of carcinogenic potency, the incremental excess cancer risk per unit of exposure (usually $\mu\text{g/L}$ for water, mg/kg/day for food, and $\mu\text{g/m}^3$ for air).

Recommended Exposure Limit (REL)—A National Institute for Occupational Safety and Health (NIOSH) time-weighted average (TWA) concentration for up to a 10-hour workday during a 40-hour workweek.

Reference Concentration (RfC)—An estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious noncancer health effects during a lifetime. The inhalation reference concentration is for continuous inhalation exposures and is appropriately expressed in units of mg/m^3 or ppm.

Reference Dose (RfD)—An estimate (with uncertainty spanning perhaps an order of magnitude) of the daily exposure of the human population to a potential hazard that is likely to be without risk of deleterious effects during a lifetime. The RfD is operationally derived from the no-observed-adverse-effect level (NOAEL, from animal and human studies) by a consistent application of uncertainty factors that reflect various types of data used to estimate RfDs and an additional modifying factor, which is based on a professional judgment of the entire database on the chemical. The RfDs are not applicable to nonthreshold effects such as cancer.

Reportable Quantity (RQ)—The quantity of a hazardous substance that is considered reportable under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA). Reportable quantities are (1) 1 pound or greater or (2) for selected substances, an amount established by regulation either under CERCLA or under Section 311 of the Clean Water Act. Quantities are measured over a 24-hour period.

Reproductive Toxicity—The occurrence of adverse effects on the reproductive system that may result from exposure to a chemical. The toxicity may be directed to the reproductive organs and/or the related endocrine system. The manifestation of such toxicity may be noted as alterations in sexual behavior, fertility, pregnancy outcomes, or modifications in other functions that are dependent on the integrity of this system.

10. GLOSSARY

Retrospective Study—A type of cohort study based on a group of persons known to have been exposed at some time in the past. Data are collected from routinely recorded events, up to the time the study is undertaken. Retrospective studies are limited to causal factors that can be ascertained from existing records and/or examining survivors of the cohort.

Risk—The possibility or chance that some adverse effect will result from a given exposure to a chemical.

Risk Factor—An aspect of personal behavior or lifestyle, an environmental exposure, or an inborn or inherited characteristic that is associated with an increased occurrence of disease or other health-related event or condition.

Risk Ratio—The ratio of the risk among persons with specific risk factors compared to the risk among persons without risk factors. A risk ratio greater than 1 indicates greater risk of disease in the exposed group compared to the unexposed group.

Short-Term Exposure Limit (STEL)—The American Conference of Governmental Industrial Hygienists (ACGIH) maximum concentration to which workers can be exposed for up to 15 minutes continually. No more than four excursions are allowed per day, and there must be at least 60 minutes between exposure periods. The daily Threshold Limit Value-Time Weighted Average (TLV-TWA) may not be exceeded.

Standardized Mortality Ratio (SMR)—A ratio of the observed number of deaths and the expected number of deaths in a specific standard population.

Target Organ Toxicity—This term covers a broad range of adverse effects on target organs or physiological systems (e.g., renal, cardiovascular) extending from those arising through a single limited exposure to those assumed over a lifetime of exposure to a chemical.

Teratogen—A chemical that causes structural defects that affect the development of an organism.

Threshold Limit Value (TLV)—An American Conference of Governmental Industrial Hygienists (ACGIH) concentration of a substance to which most workers can be exposed without adverse effect. The TLV may be expressed as a Time Weighted Average (TWA), as a Short-Term Exposure Limit (STEL), or as a ceiling limit (CL).

Time-Weighted Average (TWA)—An allowable exposure concentration averaged over a normal 8-hour workday or 40-hour workweek.

Toxic Dose₍₅₀₎ (TD₅₀)—A calculated dose of a chemical, introduced by a route other than inhalation, which is expected to cause a specific toxic effect in 50% of a defined experimental animal population.

Toxicokinetic—The absorption, distribution, and elimination of toxic compounds in the living organism.

10. GLOSSARY

Uncertainty Factor (UF)—A factor used in operationally deriving the Minimal Risk Level (MRL) or Reference Dose (RfD) or Reference Concentration (RfC) from experimental data. UFs are intended to account for (1) the variation in sensitivity among the members of the human population, (2) the uncertainty in extrapolating animal data to the case of human, (3) the uncertainty in extrapolating from data obtained in a study that is of less than lifetime exposure, and (4) the uncertainty in using lowest-observed-adverse-effect level (LOAEL) data rather than no-observed-adverse-effect level (NOAEL) data. A default for each individual UF is 10; if complete certainty in data exists, a value of 1 can be used; however, a reduced UF of 3 may be used on a case-by-case basis, 3 being the approximate logarithmic average of 10 and 1.

Xenobiotic—Any chemical that is foreign to the biological system.

10. GLOSSARY

This page is intentionally blank.

APPENDIX A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99–499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that

APPENDIX A

are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as 100-fold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology and Environmental Medicine, expert panel peer reviews, and agency-wide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology and Environmental Medicine, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road NE, Mailstop F-32, Atlanta, Georgia 30333.

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Acrylamide
CAS Numbers: 79-06-1
Date: August 27, 2009
Profile Status: Draft for Public Comment
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 33
Species: Rat

Minimal Risk Level: 0.02 mg/kg/day ppm

Reference: Sublet VH, Zenick H, Smith MK. 1989. Factors associated with reduced fertility and implantation rates in females mated to acrylamide-treated rats. *Toxicology* 55:53-67.

Experimental design: Groups of male Long-Evans hooded rats (15/group) were administered acrylamide (in distilled water) by oral gavage for 5 days at doses of 0, 30, 45, or 60 mg/kg/day (experiment 1) and 0, 5, or 15 mg/kg/day (experiment 2). Males were then mated on weeks 1, 2, 3, 4, 7, and 10 to proestrus females (weeks 1-4 only for experiment 2). Mating was confirmed by vaginal lavage. Females were sacrificed on gestation day 15 and numbers of corpora lutea, implantation sites, and fetuses were determined and estimates of pre- and postimplantation losses were calculated.

Effect noted in study and corresponding doses: Effects of acrylamide exposure on mating, fertility, and pre- and postimplantation losses are summarized in Table A-1. There were no significant treatment-related effects on mating index (number of sperm positive/number mated) at any dose level and no significant effects on any measures of reproductive success at the 5 mg/kg/day dose level. Assessments of reproductive indices at week 1 revealed significantly depressed fertility rates and elevated preimplantation losses at doses ≥ 15 mg/kg/day. Increased postimplantation loss was seen at weeks 2 and 3 at doses ≥ 15 mg/kg/day. In sperm samples collected from the 45 mg/kg/day group, the percentage of motile sperm was modestly (but significantly) decreased (58 vs. 73% in controls) at week 3, but not at weeks 2 or 4. This study identified a NOAEL of 5 mg/kg/day and a LOAEL of 15 mg/kg/day for significantly decreased fertility and increased preimplantation loss during week 1 posttreatment and significantly increased postimplantation loss during weeks 2 and 3 posttreatment.

APPENDIX A

Table A-1. Effects of Acrylamide on Selected Reproductive Indices Following Gavage Dosing of Male Long-Evans Rats for 5 Days

Treatment (mg/kg)	Post-treatment mating week	Mating index (percent) ^a	Fertility index (percent) ^b	Preimplantation loss index (percent) ^c	Postimplantation loss index (percent) ^d
Control ^e	1	25/30 (83)	22/25 (88)	19.57	4.20
	2	26/30 (87)	25/26 (96)	11.54	5.70
	3	29/30 (97)	28/29 (97)	11.46	6.80
	4	27/30 (90)	26/27 (96)	9.48	5.00
5	1	15/15 (100)	12/15 (80)	24.29	8.30
	2	14/15 (93)	11/14 (79)	25.40	10.80
	3	15/15 (100)	14/15 (93)	10.77	5.40
	4	13/15 (87)	11/13 (85)	17.08	7.10
15	1	13/15 (87)	6/13 (46) ^f	74.69 ^f	10.30
	2	11/15 (73)	11/11 (100)	8.5	15.40 ^f
	3	13/15 (87)	12/13 (92)	13.42	15.70 ^f
	4	13/15 (87)	12/13 (92)	10.36	5.50
30	1	12/15 (80)	2/12 (17) ^f	97.83 ^f	0.00
	2	15/15 (100)	15/15 (100)	15.90	25.40 ^f
	3	15/15 (100)	14/15 (93)	14.51	35.00 ^f
	4	15/15 (100)	15/15 (100)	4.90	17.50
45	1	13/15 (87)	2/13 (15) ^f	85.89 ^f	26.77
	2	15/15 (100)	12/15 (80)	33.04 ^f	60.77 ^f
	3	15/15 (100)	10/15 (67) ^f	59.13 ^f	55.50 ^f
	4	15/15 (100)	15/15 (100)	11.60	7.67
60	1	15/15 (100)	1/15 (7) ^f	98.33 ^f	33.33
	2	15/15 (100)	8/15 (53) ^f	73.39 ^f	77.87 ^f
	3	15/15 (100)	1/15 (7) ^f	98.80 ^f	50.00 ^f
	4	15/15 (100)	9/15 (60) ^f	70.84 [*]	17.11 [*]

^a Mating index = number sperm/positive females/number mated.

^b Fertility index = number pregnant/number sperm-positive females.

^c Preimplantation loss = (number of corpora lutea/female – number of implant sites/female)/(number of corpora lutea/female) x 100.

^d Postimplantation loss = (number of implant sites/female – number of fetuses/females)/(number of implant sites/female) x 100.

^e Control data from "high" and "low" dose dominant lethal studies are combined for tabular presentation. However statistical differences associated with acrylamide treatment are based upon comparison with appropriate control group.

^f p≤0.05

Source: Sublet et al. 1989

Dose and end point used for MRL derivation: 1.78 mg/kg/day (BMDL₁₀) for decreased fertility in rats (as assessed by number of nonpregnant rats/number of sperm-positive females) following oral administration of acrylamide to male Long-Evans hooded rats for 5 days and subsequent mating with untreated female rats (see Table A-2).

All dichotomous models in the EPA Benchmark Dose Software (Version 2.1) were fit to the fertility data shown in Table A-2. Although the magnitude of the fertility response represents a more serious effect,

APPENDIX A

the group sizes in this study (15 male rats/dose) did not support a benchmark response (BMR) lower than 10%; thus, an extra risk incidence of 10% above controls was selected as the BMR. Model results for the data on fertility are shown in Table A-3. All dichotomous models provided adequate fit to the data ($p > 0.1$). Comparing across models, a better fit is generally indicated by a lower Akaike's Information Criteria (AIC). The multistage (1-degree polynomial) and quantal linear models converged on the same model providing the best fit (as assessed by AIC) to the fertility data; these models both predicted a BMD_{10} of 2.44 mg/kg/day and a $BMDL_{10}$ of 1.78 mg/kg/day. Figure A-1 shows the fit of the multistage (1-degree polynomial) model to the fertility data.

Table A-2. Selected Results of Attempted Impregnation by Male Rats Administered Acrylamide by Gavage for 5 Days Prior to Mating with Untreated Female Rats During Week 1 Posttreatment

	Dose (mg/kg/day)					
	0	5	15	30	45	60
Number sperm-positive females	25	15	13	12	13	15
Number nonpregnant females	3 (12%)	3 (20%)	7 (54%)	10 (83%)	11 (85%)	14 (93%)

Source: Sublet et al. 1989

Table A-3. Model Predictions for the Fraction of Unsuccessful Impregnation (Number of Nonpregnant Females/Number of Sperm-Positive Females) Among Male Rats Administered Acrylamide by Gavage for 5 Days and Mated to Untreated Female Rats During Week 1 Posttreatment

Model	BMD_{10} (mg/kg/day)	$BMDL_{10}$ (mg/kg/day)	χ^2 p-value	AIC
Gamma ^a	3.51	1.80	0.84	87.48
Logistic	6.02	4.48	0.53	87.71
Log-logistic ^b	4.76	1.55	0.95	86.98
Log-probit ^b	4.79	3.01	0.94	87.01
Multi-stage^c	2.44	1.78	0.91	85.69
Probit	6.02	4.67	0.44	88.31
Weibull ^a	3.13	1.80	0.83	87.56
Quantal linear	2.44	1.78	0.91	85.69

^aPower restricted to ≥ 1 .


^bSlope restricted to ≥ 1 .

^cBetas restricted to ≥ 0 ; lowest degree polynomial with adequate fit is reported; degree of polynomial=1.

AIC = Akaike's Information Criteria; BMD = benchmark dose

APPENDIX A

Figure A-1. Predicted and Observed Fractions of Unsuccessful Impregnation (Number of Nonpregnant Females/Number of Sperm-Positive Females) by Male Rats Administered Acrylamide by Gavage for 5 Days and Mated to Untreated Female Rats During Week 1 Posttreatment *



*BMD and BMDL associated with a 10% extra risk increase over control are shown; doses given in units of mg/kg/day.

NOAEL LOAEL Benchmark

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose? No.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

Was a conversion used from intermittent to continuous exposure? No.

APPENDIX A

Other additional studies or pertinent information that lend support to this MRL: As demonstrated in Table 3-4 (Levels of Significant Exposure to Acrylamide – Oral), male-mediated implantation loss and decreased mean body weight gain represent the most sensitive effects of acute-duration oral exposure to acrylamide. The lowest dose level reported to induce significantly decreased male-mediated implantation loss was 15 mg/kg/day in male Long-Evans rats administered acrylamide by gavage for 5 days followed by mating sessions with unexposed female rats (Sublet et al. 1989); this study identified a NOAEL of 5 mg/kg/day. A similarly-designed study (Tyl et al. 2000b) confirmed the findings of acrylamide-induced increased implantation loss, although, based on pairwise comparisons with controls, statistical significance was achieved only at the highest dose level (60 mg/kg/day). Tyl et al. (2000b) also noted significantly decreased body weight gain (>40% lower than controls) during the 5 days of dosing, a finding not included in the study report of Sublet et al. (1989). Based on the reproducible results for male-mediated implantation loss in the studies of Sublet et al. (1989) and Tyl et al. (2000b) and the lack of supporting information regarding the body weight effects, the male-mediated implantation loss was selected as the critical effect for deriving an acute-duration oral MRL for acrylamide. The study of Sublet et al. (1989) identified the lowest LOAEL for the critical effect and was therefore selected as the principal study.

Results of several other studies corroborate the findings of male-mediated decreases in fertility and increases in implantation losses following oral exposure to acrylamide (Chapin et al. 1995; Sakamoto and Hashimoto 1986; Smith et al. 1986; Tyl et al. 2000a; Zenick et al. 1986).

Agency Contacts (Chemical Managers): Patricia Ruiz, Ph.D.; Obaid Faroon, Ph.D.; Moiz Mumtaz, Ph.D.

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Acrylamide
CAS Numbers: 79-06-1
Date: August 27, 2009
Profile Status: Draft for Public Comment
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 63
Species: Rat

Minimal Risk Level: 0.002 mg/kg/day ppm

Reference: Burek JD, Albee RR, Beyer JE, et al. 1980. Subchronic toxicity of acrylamide administered to rats in the drinking water followed by up to 144 days of recovery. J Environ Pathol Toxicol 4(5-6):157-182.

Experimental design: Groups of 6-week-old male (23–29/group) and female (10/group) F344 rats were administered acrylamide in the drinking water for up to 93 days at concentrations designed to result in acrylamide intakes of 0, 0.05, 0.2, 1, 5, or 20 mg/kg/day. Ten rats/sex/group were assigned to the basic 90-day study and were observed for body weight and water consumption (recorded weekly) throughout the treatment period. Following 7 and 33 days of treatment, three control and three high-dose male rats were sacrificed for interim electron microscopic examination of the sciatic nerve. Ten male (nine in the high-dose group, due to one death prior to treatment termination) and all female rats from each treatment group were subjected to gross and histopathologic examination of all major organs and tissues at the end of the treatment period, at which time, three other male rats from each group were processed for electron microscopic examination of the sciatic nerve. The remaining rats (all males) in each group were observed for signs of recovery from treatment-related effects for up to 144 days following cessation of treatment. Three rats/group were subjected to electron microscopic examination of the sciatic nerve on recovery days 25, 111, and 144. Body weights were recorded for two rats/dose level prior to sacrifice on recovery day 111. At the end of the 144-day recovery period, the remaining four rats of each dose level were weighed and sacrificed for gross and histopathologic examination of all major organs and tissues.

All rats were observed daily (during the 5-day work week) for general health and clinical signs. Hindlimb foot splay was measured weekly in four control and four high-dose (20 mg/kg/day) male and female rats until the onset of neuropathy was detected, after which neuropathy in the high-dose group was monitored by clinical signs. After neuropathy was detected in high-dose rats, male and female rats in the 5 mg/kg/day dose groups were also subjected to weekly testing of foot splay (rats in the lower treatment groups were not tested due to the lack of response at 5 mg/kg/day). Blood samples collected from seven rats/sex in the control and high-dose groups on treatment day 76 and from all rats alive on day 60 of the recovery period were examined for packed cell volume, total erythrocyte count, total and differential leukocyte counts, and hemoglobin concentration. The study design included urinary sampling from 10 control and 10 high-dose rats per sex on treatment day 76 and at the end of the treatment period. Blood serum was collected from the 10 rats/sex/dose that were sacrificed at the end of treatment and from the 4 male rats/group that were maintained throughout the 144-day recovery period. Blood urea nitrogen, alkaline phosphatase, serum glutamic pyruvic transaminase, and serum cholinesterase activity were determined.

Light microscopic examinations were performed on brain, spinal cord, and peripheral nerves (including brachial plexus, sciatic, and femoral nerves) from selected male and female rats of each dose group. Nervous tissues were fixed in glutaraldehyde-paraformaldehyde and stained with hematoxylin eosin.

APPENDIX A

Additional sections of brain, spinal cord, and peripheral nerves were subjected to the luxol fast blue-periodic acid Schiff (LFB/PAS) reaction for myelin staining and to Bodian's stain to elucidate more subtle axonal changes. Myelin and axonal degeneration was classified as severe (degeneration in approximately 50% of the observed fibers), moderate (degeneration in 20–50% of observed fibers), slight (degeneration in <20% of observed fibers), very slight (effects restricted to focal or multifocal changes in individual nerves), or equivocal (nerves could not be graded as clearly normal). Only sciatic nerve tissues from male rats were examined by electron microscopy. Three blocks of sciatic nerve fibers, two longitudinal and one transverse, were selected per rat for thin sectioning and ultrastructural analysis. Ultrastructural alterations were counted by examining a maximum of 50 fields per block, a field defined as a section through any Schwann cell. This resulted in an examined maximum of 150 fields/rat or 450 fields/treatment group of three rats.

Hematology, urinary and clinical chemistry parameters, body weights, organ-to-body weight ratio data, foot spread results, and water consumption were statistically analyzed by one-way analysis of variance followed by Dunnett's test. The level of significance chosen was $p < 0.05$. The study report did not, however, include individual or averaged incidences or extent of changes in these parameters, so an independent analysis of the results of body and organ weights, water consumption, foot splay, hematology, urinalysis, or serum chemistry was not possible.

Effect noted in study and corresponding doses: Significantly lower body weights were reported in the high-dose male and female rats relative to controls: 8% lower (males and females) on treatment days 13 and 20, and 21 and 24% lower (males and females, respectively) on treatment day 91. No significant body weight effect was seen in rats of lower dose groups. High-dose rats also exhibited treatment-related effects on organ weights including significantly decreased absolute liver, kidney, and thymus weights in males (also testicular) and females, significantly decreased absolute brain and heart weights in females (trend for decreased weights in males), increased relative brain, heart, liver, and kidney weights in males and females, and decreased relative thymus (females only) and testicular weight in males. Absolute and relative liver weight was increased in 5 mg/kg/day males. Marginally statistically significant increases in relative heart weight in 0.05 and 0.2 mg/kg/day females were not considered to be of toxicological significance due to the lack of a dose response. High-dose female rats exhibited significantly decreased water consumption (15–39% decreased) between treatment days 20 and 90. Although decreased water consumption was noted in high-dose males, the decrease reached the level of statistical significance in only 4 of the 13 intervals recorded. The few instances of significantly increased water consumption in low-dose rats did not follow a consistent pattern or trend, and may be of no toxicological significance. By day 144 of the posttreatment recovery period, the high-dose group had recovered with higher (but not statistically significant) body weights than controls, significantly higher absolute liver and kidney weights, and as significantly higher relative brain and liver weights.

Significantly increased instances of hindlimb foot splay were observed in high-dose male and female rats on treatment day 22 (incidences were not reported), which became more pronounced on treatment day 29. Foot splay testing was terminated with this treatment group (to prevent injury), but clinical signs of neuropathy (including curling of the toes, rear limb splay, incoordination, and posterior weakness) progressed in severity throughout the remainder of the treatment period. Beginning on treatment day 29, rats of the 5 mg/kg/day dose level were tested, but foot splay was not detected at this treatment level in either males or females. No other treatment-related clinical effects were observed in the 5 mg/kg/day males or females or any of the lower dose groups. By day 7 of the posttreatment recovery period, the high-dose groups showed clear signs of improvements continuing to day 111 with only slight posterior weakness and curling of the toes. By day 144, these high-dose rats appeared clinically similar to the controls.

APPENDIX A

At the end of the treatment period, serum cholinesterase activity was increased and alkaline phosphatase activity was statistically significantly increased in high-dose females. Significant decreases in packed cell volume, total erythrocyte count, and hemoglobin concentrations in high-dose males and females and 5 mg/kg/day females were noted. Results of urinalysis did not reveal any acrylamide-induced abnormalities. By day 144 posttreatment, the high-dose group (sex not specified) had statistically significant decreased serum cholinesterase levels and no significant differences in other clinical chemistry parameters.

Upon necropsy, gross observations of rats following the 92- or 93-day treatment period revealed treatment-related alterations only in the high-dose group, including perineal soiling, decreased adipose tissue, decreased liver size, darkened kidneys, foci or mottled appearance of lungs, decreased size or flaccid testicles, decreased size of male accessory genitalia, decreased uterus size, altered appearance of peripheral nerves, atrophy of skeletal muscle in the posterior portion of the body, bladder distention, and diffuse mural thickening of the stomach. The authors did not include incidence data regarding gross examination data, however. Histopathologic examination of high-dose rats revealed effects such as atrophy of skeletal muscle (2/10 males, 8/10 females), slightly increased hematogenous pigment in the spleen (4/9 males), ulcerative gastritis or hyperkeratosis in the nonglandular stomach (4/10 males), atrophy of mesenteric fat (8/10 females), vacuolization of the smooth muscle in the bladder wall (1/10 males, 2/9 females), inflammation in the lungs (3/10 males, 5/10 females), and testicular effects that included atrophy (10/10), mineralization in seminiferous tubules (5/10), and increased cellular debris and/or decreased spermatogenic segments in the tubular lamina of the epididymides (9/10). The statistical significance of these findings could not be assessed because incidence data for controls were not reported. By day 144 posttreatment, only the high-dose rats had persistent gross pathological effects, primarily dark testicles and slightly distended bladders. The testicular histological lesions consisted of focal or multifocal atrophy to individual seminiferous tubules, some with mineral and cellular debris, and indication of partial reversibility of the testicular atrophy. Light microscopic examination of the sciatic nerve sections (stained with hematoxylin and eosin) revealed severe degeneration in the high-dose groups that was characterized by demyelination (LFB/PAS-treated sections) and axonal degeneration (Bodian's-treated sections) in 10/10 females and similar but less severe effects in males (degeneration moderate in 5/10 and severe in the other 5). These lesions were also seen in other peripheral nerve sections (brachial plexus and femoral nerve) but varied in severity from equivocal to severe (incidences not reported). The authors noted equivocal to very slight degenerative changes in peripheral nerves of 5 mg/kg/day males (9/10) and females (6/10) but found no light microscopic evidence of peripheral nerve lesions in 0.05, 0.2, or 1 mg/kg/day treatment groups. Very slight to slight degenerative changes (demyelination, swollen astrocytes and axons) were seen in spinal cord sections of high-dose male (5/10) and female (9/10) rats. No treatment-related lesions were observed at any dose level within brain sections examined by light microscopy. After 144 days of posttreatment recovery, no nerve tissue alterations were observed in any of the 5 mg/kg/day or lower dose groups. In the high-dose group, alterations ranged from very slight to slight in the sciatic nerve and no alterations were noted in sections of the brachial nerve. The authors stated that if the recovery period had been extended beyond 144 days, the remaining tissue changes would likely have completely reversed.

Ultrastructural (electron microscope) examinations of sciatic nerve preparations from three male rats/group included the examination of fields (defined as a section through any Schwann cell) for signs of axolemma invaginations, axonal invaginations with cell organelles and/or dense bodies, and Schwann cells without axons and/or with degenerating myelin. After 7 days of treatment, no significant differences were seen between control and high-dose rats (other treatment groups were not subjected to 7-day interim sacrifice). After 33 days of treatment, high-dose male rats exhibited increased prevalence of fields showing axolemma invaginations with cell organelles and/or dense bodies and fields exhibiting Schwann cells without axons and/or with degenerating myelin (other groups were not subjected to 33-day interim sacrifice). Following 90 days of treatment, severe axonal degeneration and axonal loss were seen in high-

APPENDIX A

dose rats. Approximately 55% of the fields examined exhibited alterations in myelinated nerves or Schwann cells (compared with 12 and 21% after treatment days 7 and 33, respectively). Similar, but less severe, ultrastructural alterations in approximately 34% of the fields examined were seen in the 5 mg/kg/day dose group. At the 1 mg/kg/day dose level, approximately 24% of the fields examined showed axolemma invaginations with cell organelles and/or dense bodies, but not more severe signs of ultrastructural alterations. The alterations in the sciatic nerve fields examined in the control, 0.05, and 0.2 mg/kg-day groups were roughly comparable (15, 9, and 12%, respectively), suggesting that there were no adverse effects at the 0.05 and 0.2 mg/kg/day doses. Importantly, the increase in lesions observed via electron microscopy in the 1 and 5 mg/kg/day groups appeared to have completely reversed by days 25 and 111 posttreatment, respectively. The observed lesions in the high-dose group were partially or completely reversed by day 144 posttreatment.

In summary, the study of Burek et al. (1980) identified a NOAEL of 0.2 mg/kg/day and a LOAEL of 1 mg/kg/day, based on ultrastructural degeneration (axolemma invaginations with cell organelles and/or dense bodies) in the sciatic nerve of male rats (as detected by electron microscopic examinations, which were limited to males). The increased frequency was characterized by the study authors as "slight" for the LOAEL at 1 mg/kg/day, and the lesions were reversible (back to control levels) by day 25 posttreatment in all 1 mg/kg/day treated rats. At the resolution of the light microscope, the 5 mg/kg/day dose was the lowest dose resulting in degenerative effects in the sciatic nerve of male and female rats.

Dose and end point used for MRL derivation: 0.2 mg/kg/day

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose? No.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

Was a conversion used from intermittent to continuous exposure? No.

Other additional studies or pertinent information that lend support to this MRL: As demonstrated in Table 3-4 (Levels of Significant Exposure to Acrylamide – Oral), degenerative nerve changes and male-mediated implantation losses represent the most sensitive effects of intermediate-duration oral exposure to acrylamide. The lowest dose level reported to induce male-mediated implantation losses was 2.8 mg/kg/day in male Long-Evans rats receiving acrylamide from the drinking water for 80 days; this study identified a NOAEL of 1 mg/kg/day (Smith et al. 1986). Ultrastructural degenerative peripheral nerve changes were observed at a dose level as low as 1 mg/kg/day in male F344 rats receiving acrylamide from the drinking water for up to 93 days; this study identified a NOAEL of 0.2 mg/kg/day (Burek et al. 1980). Available data suggest that neurological effects represent a more sensitive point of departure than reproductive effects for deriving intermediate- and chronic-duration oral MRLs for acrylamide. Therefore, degenerative nerve change was selected as the critical effect for deriving an intermediate-duration oral MRL for acrylamide. The study of Burek et al. (1980) was selected as the principal study because it identified the lowest LOAEL for the critical effect. A NOAEL/LOAEL approach was selected because results of the ultrastructural evaluations included only 3 of 10 rats/group

APPENDIX A

and were reported only as the total numbers of fields (per group) with ultrastructural changes as axolemma invaginations or Schwann cells without axons and/or with degenerating myelin. This reporting of the electron microscopy data does not support a statistical comparison of the incidence of changes between the exposed and control groups because it is unknown within any exposure group how the numbers of changes were distributed among the three rats (i.e., whether the apparent increase in incidence of fields with changes was due to one, two, or all three rats in the 1, 5, and 20 mg/kg-day groups). Therefore, a BMD approach was not feasible.

Interim data from a chronic toxicity and carcinogenicity bioassay of male and female F344 rats receiving acrylamide from the drinking water for up to 2 years (Johnson et al. (1984, 1985, 1986) provide support to the findings of Burek et al. (1980). In the 2-year study, interim sacrifices were performed at 3, 6, 12, and 18 months on some rats from each exposure group. One aspect of interim sacrifices was to evaluate the condition of selected peripheral nerve sections using light and electron microscopy. The most significant noncancer effects were increased incidences of axolemma invaginations (observed by electron microscopy) in the tibial branch of the sciatic nerve of male rats following 3 and 6 months of treatment and increased prevalence of "moderate" to "severe" degeneration (observed by light microscopy) in both males and females following 2 years of treatment. This study identified a NOAEL of 0.5 mg/kg/day and a LOAEL of 2.0 mg/kg/day for light and electron microscope evidence of peripheral nerve fiber degeneration in the male rats. Electron microscope evaluations at interims ≥ 12 months were not considered meaningful due to age-related degenerative effects that could not be distinguished from acrylamide-induced effects.

Agency Contacts (Chemical Managers): Patricia Ruiz, Ph.D.; Obaid Faroon, Ph.D.; Moiz Mumtaz, Ph.D.

APPENDIX B. USER'S GUIDE

Chapter 1

Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public, especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

Chapter 2

Relevance to Public Health

This chapter provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions:

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The chapter covers end points in the same order that they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, and dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this chapter.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal Risk Levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Chapter 3 Data Needs section.

APPENDIX B

Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, ATSDR has derived MRLs for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not meant to support regulatory action, but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans.

MRLs should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2, "Relevance to Public Health," contains basic information known about the substance. Other sections such as Chapter 3 Section 3.9, "Interactions with Other Substances," and Section 3.10, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology that the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses (RfDs) for lifetime exposure.

To derive an MRL, ATSDR generally selects the most sensitive end point which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen end point are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest no-observed-adverse-effect level (NOAEL) that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the levels of significant exposure (LSE) tables.

Chapter 3**Health Effects****Tables and Figures for Levels of Significant Exposure (LSE)**

Tables and figures are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, MRLs to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in

APPENDIX B

conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of NOAELs, LOAELs, or Cancer Effect Levels (CELs).

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 3-1 and Figure 3-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

LEGEND**See Sample LSE Table 3-1 (page B-6)**

- (1) **Route of Exposure.** One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. Typically when sufficient data exist, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Tables 3-1, 3-2, and 3-3, respectively). LSE figures are limited to the inhalation (LSE Figure 3-1) and oral (LSE Figure 3-2) routes. Not all substances will have data on each route of exposure and will not, therefore, have all five of the tables and figures.
- (2) **Exposure Period.** Three exposure periods—acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more)—are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) **Health Effect.** The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) **Key to Figure.** Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the two "18r" data points in sample Figure 3-1).
- (5) **Species.** The test species, whether animal or human, are identified in this column. Chapter 2, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 3.4, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) **Exposure Frequency/Duration.** The duration of the study and the weekly and daily exposure regimens are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to "Chemical x" via inhalation for 6 hours/day, 5 days/week, for 13 weeks. For a more complete review of the dosing regimen, refer to the appropriate sections of the text or the original reference paper (i.e., Nitschke et al. 1981).

APPENDIX B

- (7) System. This column further defines the systemic effects. These systems include respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered in these systems. In the example of key number 18, one systemic effect (respiratory) was investigated.
- (8) NOAEL. A NOAEL is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system, which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").
- (9) LOAEL. A LOAEL is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific end point used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less Serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference. The complete reference citation is given in Chapter 9 of the profile.
- (11) CEL. A CEL is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.
- (12) Footnotes. Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates that the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

LEGEND

See Sample Figure 3-1 (page B-7)

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period. The same exposure periods appear as in the LSE table. In this example, health effects observed within the acute and intermediate exposure periods are illustrated.
- (14) Health Effect. These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure. Concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m³ or ppm and oral exposure is reported in mg/kg/day.

APPENDIX B

- (16) NOAEL. In this example, the open circle designated 18r identifies a NOAEL critical end point in the rat upon which an intermediate inhalation exposure MRL is based. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the extrapolation from the exposure level of 3 ppm (see entry 18 in the table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).
- (17) CEL. Key number 38m is one of three studies for which CELs were derived. The diamond symbol refers to a CEL for the test species-mouse. The number 38 corresponds to the entry in the LSE table.
- (18) Estimated Upper-Bound Human Cancer Risk Levels. This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels (q_1^*).
- (19) Key to LSE Figure. The Key explains the abbreviations and symbols used in the figure.

SAMPLE

1 →

Table 3-1. Levels of Significant Exposure to [Chemical x] – Inhalation


	Key to figure ^a	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference
						Less serious (ppm)	Serious (ppm)	
2 →	INTERMEDIATE EXPOSURE							
		5	6	7	8	9		10
3 →	Systemic	↓	↓	↓	↓	↓		↓
4 →	18	Rat	13 wk 5 d/wk 6 hr/d	Resp	3 ^b	10 (hyperplasia)		Nitschke et al. 1981
	CHRONIC EXPOSURE							
	Cancer						11	
							↓	
	38	Rat	18 mo 5 d/wk 7 hr/d			20	(CEL, multiple organs)	Wong et al. 1982
	39	Rat	89–104 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, nasal tumors)	NTP 1982
	40	Mouse	79–103 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, hemangiosarcomas)	NTP 1982

12 →

^a The number corresponds to entries in Figure 3-1.^b Used to derive an intermediate inhalation Minimal Risk Level (MRL) of 5×10^{-3} ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

SAMPLE

Figure 3-1. Levels of Significant Exposure to [Chemical X] - Inhalation



DRAFT FOR PUBLIC COMMENT

APPENDIX B

APPENDIX B

This page is intentionally blank.

APPENDIX C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACGIH	American Conference of Governmental Industrial Hygienists
ACOEM	American College of Occupational and Environmental Medicine
ADI	acceptable daily intake
ADME	absorption, distribution, metabolism, and excretion
AED	atomic emission detection
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
ALT	alanine aminotransferase
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
AOEC	Association of Occupational and Environmental Clinics
AP	alkaline phosphatase
APHA	American Public Health Association
AST	aspartate aminotransferase
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	best available technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BMD/C	benchmark dose or benchmark concentration
BMD _x	dose that produces a X% change in response rate of an adverse effect
BMDL _x	95% lower confidence limit on the BMD _x
BMDS	Benchmark Dose Software
BMR	benchmark response
BSC	Board of Scientific Counselors
C	centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	cancer effect level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CI	confidence interval
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy

APPENDIX C

DOL	Department of Labor
DOT	Department of Transportation
DOT/UN/ NA/IMDG	Department of Transportation/United Nations/ North America/Intergovernmental Maritime Dangerous Goods Code
DWEL	drinking water exposure level
ECD	electron capture detection
ECG/EKG	electrocardiogram
EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F ₁	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
FR	Federal Register
FSH	follicle stimulating hormone
g	gram
GC	gas chromatography
gd	gestational day
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IARC	International Agency for Research on Cancer
IDLH	immediately dangerous to life and health
ILO	International Labor Organization
IRIS	Integrated Risk Information System
K _d	adsorption ratio
kg	kilogram
kgg	metric ton
K _{oc}	organic carbon partition coefficient
K _{ow}	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC ₅₀	lethal concentration, 50% kill
LC _{Lo}	lethal concentration, low
LD ₅₀	lethal dose, 50% kill
LD _{Lo}	lethal dose, low
LDH	lactic dehydrogenase
LH	luteinizing hormone
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
LT ₅₀	lethal time, 50% kill
m	meter
MA	<i>trans,trans</i> -muconic acid
MAL	maximum allowable level

APPENDIX C

mCi	millicurie
MCL	maximum contaminant level
MCLG	maximum contaminant level goal
MF	modifying factor
MFO	mixed function oxidase
mg	milligram
mL	milliliter
mm	millimeter
mmHg	millimeters of mercury
mmol	millimole
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ND	not detected
NFPA	National Fire Protection Association
ng	nanogram
NHANES	National Health and Nutrition Examination Survey
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSH TIC	NIOSH's Computerized Information Retrieval System
NLM	National Library of Medicine
nm	nanometer
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OR	odds ratio
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA

APPENDIX C

OTS	Office of Toxic Substances
OW	Office of Water
OWRS	Office of Water Regulations and Standards, EPA
PAH	polycyclic aromatic hydrocarbon
PBPD	physiologically based pharmacodynamic
PBPK	physiologically based pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
pg	picogram
PHS	Public Health Service
PID	photo ionization detector
pmol	picomole
PMR	proportionate mortality ratio
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PSNS	pretreatment standards for new sources
RBC	red blood cell
REL	recommended exposure level/limit
RfC	reference concentration
RfD	reference dose
RNA	ribonucleic acid
RQ	reportable quantity
RTECS	Registry of Toxic Effects of Chemical Substances
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
SGOT	serum glutamic oxaloacetic transaminase
SGPT	serum glutamic pyruvic transaminase
SIC	standard industrial classification
SIM	selected ion monitoring
SMCL	secondary maximum contaminant level
SMR	standardized mortality ratio
SNARL	suggested no adverse response level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD ₅₀	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	total organic carbon
TPQ	threshold planning quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	time-weighted average
UF	uncertainty factor
U.S.	United States
USDA	United States Department of Agriculture
USGS	United States Geological Survey
VOC	volatile organic compound
WBC	white blood cell
WHO	World Health Organization
>	greater than

APPENDIX C

\geq	greater than or equal to
=	equal to
<	less than
\leq	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer
μg	microgram
q_1^*	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result

APPENDIX C

This page is intentionally blank.

APPENDIX D. INDEX

absorbed dose.....	104, 125
adduct.....	9, 24, 26, 66, 67, 80, 81, 82, 83, 84, 86, 88, 89, 90, 92, 96, 98, 100, 101, 102, 112, 114, 115, 116, 124, 125, 138, 170, 178, 180, 183
adenocarcinoma.....	68, 69
adsorption.....	173
aerobic.....	160
anaerobic.....	160
atrophy.....	23
axolemma.....	13, 61
bioaccumulation.....	173
bioavailability.....	93, 173
bioconcentration factor.....	160
biodegradation.....	7, 152, 160
biomarker.....	22, 67, 88, 124, 125, 133, 137, 138, 139, 140, 170, 178, 180, 184
body weight.....	3, 7, 9, 12, 17, 21, 58, 59, 64, 65, 71, 91, 101, 102, 112, 121, 122, 123, 131, 132, 136, 154, 171, 180
body weight effects.....	9, 12, 17, 58, 59, 71
breast milk.....	3, 123, 171, 172, 174, 178, 183
cancer.....	3, 8, 16, 28, 29, 65, 66, 67, 68, 70, 75, 118, 121, 122, 126, 134, 137, 169, 189, 190
carcinogen.....	9, 70, 154, 169, 186, 188, 190
carcinogenic.....	9, 10, 15, 16, 70, 118, 122, 186, 190
carcinogenicity.....	9, 13, 68, 70, 96, 119, 126, 127, 134, 137, 139, 190
carcinoma.....	68, 69, 70, 75, 134
cardiovascular.....	17, 56, 71
cardiovascular effects.....	56
chromosomal aberrations.....	82, 83
clastogenic.....	75, 82, 117, 118, 135
death.....	15, 16, 30, 59, 71
degenerative effects.....	8, 11, 61, 63, 115, 133, 134, 135
deoxyribonucleic acid (see also DNA).....	75, 81, 87
dermal effects.....	17, 58, 71, 133
developmental effects.....	3, 27, 63, 64, 65, 74, 137, 139
DNA (see also deoxyribonucleic acid).....	75, 80, 81, 82, 83, 84, 86, 87, 88, 92, 96, 98, 99, 112, 114, 117, 118, 122, 124, 125, 127, 135, 138, 180
DNA adducts.....	83, 86, 92, 96, 98, 99, 112, 118, 124, 138, 180
dominant lethality.....	8, 62, 74, 135
dopamine.....	64, 136
endocrine.....	17, 57, 58, 71, 119, 120
endocrine effects.....	57, 58, 71
erythema.....	22
fertility.....	12, 63, 118, 131, 186
fetal tissue.....	123
fetus.....	94, 121, 122, 127, 171
foot splay.....	60, 137
gait.....	22, 23, 25, 26, 60, 64
gastrointestinal effects.....	56
general population.....	2, 131, 170, 171, 172
genotoxic.....	15, 75, 107, 118, 134, 140
genotoxicity.....	9, 75, 118, 134, 139

APPENDIX D

groundwater	158, 159, 173
half-life	96, 99, 103, 124
hematological effects	17, 56
hemoglobin	17, 22, 24, 56, 66, 67, 88, 89, 90, 93, 94, 96, 97, 98, 100, 101, 107, 111, 112, 114, 115, 125, 133, 134, 138, 170, 178, 180
hemoglobin adduct	17, 22, 66, 67, 88, 89, 90, 93, 94, 98, 100, 101, 107, 112, 114, 115, 125, 133, 134, 138, 180
heritable translocations	75, 118, 135
hindlimb splay	11, 60, 61, 64, 123, 133
hydrolysis	7, 96, 99, 110, 138, 160, 161
hydroxyl radical	160, 173
immunological	15, 21, 60, 74, 136
implantation loss	12, 13, 62, 126, 131, 132
K _{ow}	143
LD ₅₀	30, 70, 71, 131
leukemia	83, 85
lymphatic	28
lymphopoietic	28
lymphoreticular	21, 60, 74
male-mediated	12, 13, 62, 126, 131, 135
mammary gland tumors	9, 68, 69
mass spectroscopy	100
mesotheliomas	9, 68, 126, 134
micronuclei	82, 83
milk	8, 66, 123, 154, 172, 178, 179
motor function	21, 132, 136
musculoskeletal effects	57
neonatal	96, 132
neurobehavioral	120
neurochemical	132
neurodevelopmental	62, 137
neurological effects	8, 11, 13, 21, 22, 27, 58, 60, 61, 62, 74, 115, 116, 126, 131, 133, 137
neurophysiological	24
neurotransmitter	117
nuclear	100, 163
ocular effects	17, 58, 71
odds ratio	66
partition coefficients	114
peripheral neuropathy	8, 11, 24, 26, 57, 60, 61, 116, 123, 129, 131
pharmacodynamic	104, 113
pharmacokinetic	104, 105, 106, 107, 113, 121, 138
placenta	8, 94, 123, 178
placental barrier	154, 169, 171
rate constant	90, 114, 160
renal effects	57
reproductive effects	8, 13, 27, 62, 63, 74, 117, 126, 135
respiratory effects	17, 56
retention	150, 182
rotarod	11, 60, 61, 122, 137
solubility	152, 155, 159, 162, 170, 173
sperm mobility	8, 62, 135

APPENDIX D

spermatogonia.....	77, 82
systemic effects.....	17, 30, 71
T3 (see also triiodothyronine).....	57, 119, 120, 132
T4 (see also thyroxine).....	57, 119, 120, 132
testicular atrophy.....	8, 63, 126, 135
thyroid tumors.....	118
thyroxine (see also T4).....	57
toxicokinetic.....	15, 91, 92, 114, 138
tremors.....	60, 137
triiodothyronine (see also T3).....	57
tumors.....	9, 68, 69, 70, 75, 118, 119, 126, 134
tunica vaginalis.....	9, 68, 119, 134
vapor phase.....	190
vapor pressure.....	152, 155, 162, 170, 190
volatilization.....	155, 159

