



Published in final edited form as:

Biometrics. 2012 June ; 68(2): 466–476. doi:10.1111/j.1541-0420.2011.01677.x.

A latent variable approach to study gene-environment interactions in the presence of multiple correlated exposures

Brisa N. Sánchez, Shan Kang, and Bhramar Mukherjee

Department of Biostatistics, University of Michigan, Ann Arbor, MI USA 48109

Brisa N. Sánchez: brisa@umich.edu

Summary

Many existing cohort studies initially designed to investigate disease risk as a function of environmental exposures have collected genomic data in recent years with the objective of testing for gene-environment interaction ($G \times E$) effects. In environmental epidemiology, interest in $G \times E$ arises primarily after a significant effect of the environmental exposure has been documented. Cohort studies often collect rich exposure data, as a result, assessing $G \times E$ effects in the presence of multiple exposure markers further increases the burden of multiple testing, an issue already present in both genetic and environment health studies. Latent variable (LV) models have been used in environmental epidemiology to reduce dimensionality of the exposure data, gain power by reducing multiplicity issues via condensing exposure data, and avoid collinearity problems due to presence of multiple correlated exposures. We extend the LV framework to characterize gene-environment interaction in presence of multiple correlated exposures and genotype categories. Further, similar to what has been done in case-control $G \times E$ studies, we use the assumption of gene-environment ($G-E$) independence to boost the power of tests for interaction. The consequences of making this assumption, or the issue of how to explicitly model $G-E$ association has not been previously investigated in LV models. We postulate a hierarchy of assumptions about the LV model regarding the different forms of $G-E$ dependence and show that making such assumptions may influence inferential results on the G , E , and $G \times E$ parameters. We implement a class of shrinkage estimators to data adaptively trade-off between the most restrictive to most flexible form of $G-E$ dependence assumption and note that such class of compromise estimators can serve as a benchmark of model adequacy in LV models. We demonstrate the methods with an example from the Early Life Exposures in Mexico City to Neuro-Toxicants (ELEMENT) study of lead exposure, iron metabolism genes, and birth weight.

Keywords

Gene-environment independence; principal components; shrinkage estimation; structural equation models

Correspondence to: Brisa N. Sánchez, brisa@umich.edu.

Supplementary Materials

Supplementary Materials referenced in Section 4 are available under the Paper Information link at the Biometrics website <http://www.tibs.org/biometrics>.

1. Introduction

It is now clear from many lines of evidence that pure genetics or pure environmental factors play only a partial role in the etiology of most complex diseases. Instead, it is now accepted that the majority of chronic diseases likely stem from interactions between genetic traits, “*G*”, and environmental factors, “*E*” –an exponentially growing area of study (Khoury and Wacholder, 2009). Characterizing gene-environment interactions, “*G* × *E* effects”, is critical in understanding the biological mechanisms of disease etiology and can impact preventive medicine and public health by informing the way clinicians advise their patients and the way public health practitioners assess risk and set policy. Statistical approaches that heighten our ability to understand *G* × *E* effects can accelerate mapping of the so far elusive environmental footprint of disease etiology.

Established environmental health cohorts that have demonstrated modest health effects of the environment are now collecting genomic data to test *G* × *E* effects. However, *G* × *E* interaction studies are statistically difficult problems because of exposure measurement error, multiple potential exposure markers, and prohibitive sample sizes required to reach adequate power. Statistical methods to boost efficiency for testing *G* × *E* effects have primarily been developed for case-control studies, where imposing the assumption of independence between environmental exposures and inherited genetic susceptibility factors, so called *G-E* independence, boosts efficiency of *G* × *E* effect estimates (Chatterjee and Carroll, 2005, and references therein). Hybrid approaches that protect against bias under departures from independence constraints have also been proposed (e.g., Mukherjee and Chatterjee, 2008; Li and Conti, 2009; Chen et al., 2009).

Latent variable (LV) models have been used in environmental health studies to extract features from a set of correlated biomarkers, thus reducing dimensionality of exposure data and multiple testing burden, and enhancing power (e.g., Budtz-Jørgensen et al., 2003b). These models have enormous potential for modeling gene-environment interaction between multiple genes and multiple environmental exposures, an area which is still in its formative phase. Chatterjee et al. (2006) motivate a one degree of freedom test for gene-gene interactions from a latent variable perspective in a case-control study. However the issue of modeling *G-E* dependence through the LV framework is not discussed. More general LV models, as proposed here, can incorporate and estimate *G-E* association structures, which are of interest to environmental health researchers, and also impose constraints such as independence. In our motivating example, the relationship between iron metabolism genes and lead exposure is of interest in itself (e.g., Hopkins et al., 2008). Very few attempts exist that pose a general LV model for studying *G* × *E* effects in cross-sectional or cohort studies (e.g., Dhungana et al., 2007; Rathouz et al., 2008; and Javaras et al., 2010), especially those investigating an array of *G-E* dependence structures.

The Early Life Exposures in MEXICO City to Neuro-Toxicants (ELEMENT) study motivates our work. ELEMENT consists of four longitudinal birth cohorts in Mexico City, constituting over 2,000 mother infant pairs with prospectively collected exposure data and several anthropometric, cardiovascular, neuro-development, and behavioral outcomes. Genotyping for these cohorts is underway, with genotyping for the first cohort completed on a set of

candidate genes (≈ 400 pairs). Figure 1 is a path diagram describing relationships between four biomarkers of prenatal lead exposure, their interaction with iron metabolism genes, and birth weight.

In Section 2, we describe a model structure to summarize a group of biomarkers into latent variables, and the spectrum of G - E dependence models we consider. In Section 3, we describe maximum likelihood estimation (MLE) and a general class of shrinkage estimators to data-adaptively compromise between the most stringent and most flexible models for G - E association. Small scale simulation studies (Section 4) bring out salient features of our methodology. Section 5 presents analyses of our motivating example. Section 6 discusses the use of LV models for $G \times E$ studies, and shrinkage estimators in general LV modeling beyond the $G \times E$ context.

2. A Latent Exposure Model for $G \times E$ Studies

2.1 Model Representation

For the i^{th} of N individuals, let Y_i represent a univariate health outcome (here birth weight), and \mathbf{U}_i be an $l \times 1$ vector of latent exposures measured indirectly by a set of p measurements \mathbf{E}_i . In the example, $l = 1$, $p = 4$, and U_i is prenatal lead exposure (Figure 1). Let \mathbf{Z}_i and \mathbf{W}_i represent $q \times 1$ and $r \times 1$ covariate vectors, respectively. Without any loss of generality, genotype classes are represented by a categorical variable G_i with classes $g = 0, \dots, G$. Genotype classes may arise from data on biallelic polymorphisms where a ‘risk’ allele ‘A’ may alter the exposure metabolism pathway or affect health (with the reference allele denoted by ‘a’) or from combinations of genetic markers measured at multiple loci (e.g., risk alleles A, B). In the lead example we consider two single nucleotide polymorphisms (SNPs) implicated in iron metabolism, but due to sparsity of data, we assume G_i can take two possible values: zero for wild-type on both SNPs (‘aa’ and ‘bb’), and 1 for at least one copy of either of the risk alleles (i.e., ‘Aa’, ‘AA’, ‘Bb’, or ‘BB’), consistent with dominant/recessive models for genetic susceptibility. Alternatively, genetic groups can arise as categorization of an underlying genetic risk score that combines several markers identified by existing genome wide association studies (Qi et al, 2011), or from infant-mother genotype combinations at a single locus in studies of pre-natal fetal exposure.

The latent variable model is then specified in two stages: a health outcome model and an exposure model. In the outcome model, the association between the outcome Y_i , exposure \mathbf{U}_i and genetic category $G_i = g$ conditional on covariates \mathbf{Z}_i is characterized by either

$$Y_i = \beta_{0,g} + \beta_{U,g}^T \mathbf{U}_i + \beta_{Z,g}^T \mathbf{Z}_i + \varepsilon_i, \quad (1)$$

$$\text{or } Y_i = \beta_0 + \beta_U^T \mathbf{U}_i + \sum_{g=1}^G \left\{ \beta_g I_{gi} + \beta_{g \times U}^T I_{gi} \cdot \mathbf{U}_i + \beta_Z^T \mathbf{Z}_i + \beta_{g \times Z}^T I_{gi} \cdot \mathbf{Z}_i \right\} + \varepsilon_i, \quad (2)$$

using genotype class indicators $I_{gi} = I(G_i = g)$. The mean-zero error ε_i has variance σ^2 . Equation (1) is written using the multiple group notation (Bollen 1989), which is useful in writing the likelihood (Section 3), and $\beta_{0,g}$, $\beta_{U,g}$, $\beta_{Z,g}$ are parameters specific to class $g = 0$,

\dots, G . Such notation is standard in widely used LV software. In (1), the gene-environment interaction test is specified as $H_0 : \beta_{U,0} = \dots = \beta_{U,g}$, i.e., homogeneity of the environment's effects across genetic groups, and is equivalent to testing the interaction parameters in (2), i.e., $H_0 : \beta_{g \times U} = \beta_{U,g} - \beta_{U,0} = 0$ for all $g = 1, \dots, G$. $G \times E$ interactions are of interest in both environmental and genetic epidemiology, but in environmental epidemiology the $G \times E$ question arises after showing main effects of exposure on outcome, and heterogeneity of effects across genetic subgroups as well as the exposure effect within class g , $\beta_{U,g}$, are of primary interest.

The exposure model consists of a model for the latent variable (3) dependent on covariates W_i , and a measurement model (4) relating the observed exposure measurements to the latent variable

$$U_i = \alpha_{0g} + \alpha_W W_i + \xi_i \quad (3)$$

$$E_i = \nu_g + \Lambda_g U_i + \delta_i \quad (4)$$

with (3) and (4) again written in the multiple group notation. Regression coefficients α_{0g} and α_W are $l \times 1$ and $l \times r$ matrices with $\gamma_g = \alpha_{0g} - \alpha_{00}$ being effect of genotype class on exposure U_i , and covariates W_i ($r \times 1$) may help predict exposure levels for a given subject (e.g., occupation). The zero-mean error terms, ξ_i , are assumed independent of ε_i , and have category-specific $l \times l$ covariance matrices Φ_g . Means vector ν_g and factor loading matrix Λ_g are $p \times 1$ and $p \times l$, respectively, and δ_i has zero mean and $p \times p$ covariance matrix Θ_g .

Although LV models are helpful in many respects, one well known problem is the potential for lack of identifiability. Standard identifiability constraints have been developed for linear latent variable models (Bollen, 1989), and identifiability of latent class models has also been investigated (Huang and Bandeen-Roche, 2004). Essentially, model parameters are constrained to ensure identifiability; for example, some entries of ν_g and Λ_g are fixed to 0 or 1, although sometimes algebraic proofs of identifiability are needed (Sánchez et al., 2005). In the lead example, the constraints $\nu_g = (0, \nu_{g,2}, \nu_{g,3}, \nu_{g,4})^\top$ and $\Lambda_g = (1, \lambda_{g,2}, \lambda_{g,3}, \lambda_{g,4})^\top$ fix the mean and scale of the latent exposure to those of patella lead. Parameters in Φ_g are typically unconstrained, while the off-diagonal elements of Θ_g are typically, although not necessarily, restricted to be zero denoting conditional independence between E_i 's given U_i . However, theoretical identifiability may not necessarily guarantee numerical stability of results, which also depends on sample sizes. Some investigators recommend at least 5 to 10 observations per parameter estimated (Westland, 2010). Hence, users need to be attentive as to what model the available sample size allows them to fit.

2.2 Modeling G-E Dependence

In many $G \times E$ studies, it may be natural to assume that an individual's environmental exposure is independent of genetic factors, but may not be realistic when the gene and the exposure share a metabolic pathway. For example, iron metabolism genes may increase lead absorption (Hopkins et al., 2008), and characterizing such dependence may shed insight into the mechanistic process.

Varying degrees of *G-E* dependence can be modeled through imposing further constraints on the exposure model parameters (Figure 2). The most restrictive assumption is that parameters are homogeneous across genotypes. We use ‘A0’ to denote this full *G-E* independence

$$A0: (\alpha_{0g}, \Phi_g, \nu_g, \Lambda_g, \Theta_g) = (\alpha_0, \Phi, \nu, \Lambda, \Theta).$$

With A0, the exposure model (3)–(4) has at least *l* parameters in each of α_0 and Φ , *p* – *l* factor loadings Λ and *p* – *l* intercepts ν , and *p* parameters Θ , for a total of at least 3*p* parameters. In the lead exposure model, A0 totals 13 exposure model parameters.

A first step at relaxing A0 is to allow the intercepts and variances in the latent variable equation (3) to differ by genotype. Letting $\gamma_g = \alpha_{0g} - \alpha_0$ be the gene effect on the latent exposure we have

$$A1: (\nu_g, \Lambda_g, \Theta_g) = (\nu, \Lambda, \Theta), \text{ but } \gamma_g \neq 0 \text{ or } \Phi_g^{-1} \Phi_0 \neq I \text{ for at least one } g.$$

Relaxing these constraints is very natural, since genotype status may increase absorption of pollutants from the environment as well as change exposure variability. Genotype status may change the distribution of the observed exposure measures, E_i , but modeling change in the distribution of the underlying exposure is a parsimonious way of modeling changes in the actual E_i . This assumption has at least 3*p* + 2*lG* exposure model parameters; 15 in the lead example. Alternatively, one could restrict the latent variable variances Φ_g to be equal across genotype subgroups; that is, a slightly modified assumption A1* : $(\Phi_g, \nu_g, \Lambda_g, \Theta_g) = (\Phi, \nu, \Lambda, \Theta)$.

Next, constraints of equal means ν and factor loadings Λ can be removed,

$$A2: \Theta_g = \Theta.$$

Biological mechanisms that modify transfer of pollutants from one compartment to another are consistent with this assumption. In the lead example, three of the observed prenatal lead exposure biomarkers are measured on the mother, while umbilical cord blood on the offspring. The transfer rates from maternal compartments (i.e., blood and bone) to offspring compartment may vary by child’s genotype, such that factor loading λ_4 may vary by genotype. A2 has at least 3*p* + 2*Gp* exposure model parameters; 21 in the lead example.

Lastly, all equality constraints on the parameters can be removed, namely,

$$A3: \text{All model parameters differ by genotype,}$$

yielding at least 4*p* + 2*Gp* exposure model parameters; 26 in the lead example. In classical multiple group analyses (Bollen, 1989), one might additionally posit that the structure of the whole model might differ by genotype; e.g., that the number of latent variables differs by

genotype. We restrict our attention to assumptions A0–A3, where the model structure is the same across genotypes.

Assumptions A0–A3 have different number of parameters; an increase from 13 to 26 parameters in the example. Constraints on exposure model parameters may increase efficiency and power for the main hypotheses (G , E or $G \times E$ tests), but could induce bias in parameters of interest if they are incorrectly assumed. Model evaluation strategies are available that may help select which assumption A0–A3 fits the data best (Bentler and Hu, 1995). However, this task may not be straightforward since many fit criteria exist for LV models. Furthermore, testing all parameter constraints may incur a high rate Type 1 error rate, since the study will unlikely be powered to detect significant differences across genotypes in all model parameters.

3. Parameter estimation

Although various estimation procedures have been proposed for LV models (Bollen, 1989), full MLE is the most common estimation procedure given its wide availability in software packages. We review MLE and describe the implementation of shrinkage estimators that combine MLE estimates. Using shrinkage estimators may be a more suitable approach for parameter estimation. The strategy would be to fit the most restrictive model A0 and the most flexible model A3 (or A2 depending on sample size), and then use shrinkage to derive the final composite estimates.

3.1 Maximum likelihood estimation

Let $\mathbf{Y}_i^* = (Y_i, \mathbf{E}_i^\top)^\top$ and $\boldsymbol{\theta}$ be the vector of all model parameters. Assuming that ε_i , ξ_i and δ_i are normally distributed, and integrating over the latent variable, the joint marginal distribution of the observed outcome and exposures, $f(\mathbf{Y}_i^* | G_i = g, \mathbf{Z}_i, \mathbf{W}_i; \boldsymbol{\theta})$, is a multivariate normal density, with moments given by

$$E(\mathbf{Y}_i^* | G_i = g, \mathbf{Z}_i, \mathbf{W}_i; \boldsymbol{\theta}) = \boldsymbol{\nu}_g^* + \boldsymbol{\Lambda}_g^* (\boldsymbol{\alpha}_{0,g} + \mathbf{W}_i \boldsymbol{\alpha}_w) + \boldsymbol{\beta}_{z,g}^* \mathbf{Z}_i \text{ and}$$

$$\text{Var}(\mathbf{Y}_i^* | G_i = g, \mathbf{Z}_i, \mathbf{W}_i; \boldsymbol{\theta}) = \boldsymbol{\Lambda}_g^* \boldsymbol{\Phi}_g(\boldsymbol{\Lambda}_g^*)^\top + \boldsymbol{\Theta}_g^* \text{ where}$$

$$\boldsymbol{\nu}_g^* = \begin{pmatrix} \beta_{0,g} \\ \boldsymbol{\nu}_g \end{pmatrix} \boldsymbol{\Lambda}_g^* = \begin{pmatrix} \boldsymbol{\beta}_{U,g} \\ \boldsymbol{\Lambda}_g \end{pmatrix} \boldsymbol{\beta}_{z,g}^* = \begin{pmatrix} \boldsymbol{\beta}_{z,g} \\ \mathbf{0}_{p \times q} \end{pmatrix} \boldsymbol{\Theta}_g^* = \begin{pmatrix} \sigma^2 & \mathbf{0}_{1 \times p} \\ \mathbf{0}_{p \times 1} & \boldsymbol{\Theta}_g \end{pmatrix}$$

The log likelihood of $\boldsymbol{\theta}$ is then, $\ell(\boldsymbol{\theta}) = \sum_{i=1}^N \sum_{g=0}^G \log f(\mathbf{Y}_i^* | G_i = g, \mathbf{Z}_i, \mathbf{W}_i; \boldsymbol{\theta})$. Parameter estimates are obtained by maximizing $\ell(\boldsymbol{\theta})$, or equivalently, solving the score equations

$S(\boldsymbol{\theta}) = \sum_{i=1}^N \sum_{g=0}^G S_i = 0$, where $S_i = \partial \log f(\mathbf{Y}_i^* | G_i = g, \mathbf{Z}_i, \mathbf{W}_i; \boldsymbol{\theta}) / \partial \boldsymbol{\theta}$ is the contribution of the i^{th} observation to the score. Variances for parameters can be obtained by inverting the information matrix, $\mathbf{I}_\theta = -E(\partial^2 \ell(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top)$, or by computing robust variances

$$\hat{\text{var}}_R(\hat{\boldsymbol{\theta}}) = \mathbf{B}^{-1} \mathbf{A} \mathbf{B}^{-\top}, \text{ where } \mathbf{A} = 1/N \sum_{i=1}^N \sum_{g=0}^G S_i S_i^\top \text{ and } \mathbf{B} = 1/N \sum_{i=1}^N \sum_{g=0}^G \partial S_i / \partial \boldsymbol{\theta}.$$

3.2 Shrinkage estimation

Shrinkage estimators have been used in outcome-dependent sampling based studies as a way to balance bias and efficiency gains from assuming G - E independence while using a retrospective likelihood formulation of the model. Shrinkage approaches will enhance efficiency only when the retrospective model improves efficiency. A typical formulation will factorize the retrospective likelihood $p(G, E, Z|Y; \theta_1, \theta_2, \theta_3)$ as $p(Y|G, E, Z; \theta_1)p(G|E, Z; \theta_2)p(E, Z; \theta_3)/p(Y; \theta_1, \theta_2, \theta_3)$ with θ_1 being the outcome model parameters, θ_2 and θ_3 describing the G - E dependence and exposure-covariate associations, respectively, and the G - E association reflected through the term $p(G|E, Z; \theta_2)$. Because $p(Y; \theta_1, \theta_2, \theta_3) = \int p(Y|G, E, Z; \theta_1)p(G|E, Z; \theta_2)p(E, Z; \theta_3) dG dE dZ$ (in the denominator) depends on the specification of $p(G|E, Z; \theta_2)$, the MLE of θ_1 depends on the assumed model for the G - E association conditional on covariates. It has been shown (Chatterjee and Carroll, 2005) that assuming conditional G - E independence, $p(G|E, Z; \theta_2) = p(G|Z; \theta_2)$, in case-control studies leads to large efficiency gain for estimating the $G \times E$ interaction parameter in the outcome model $p(Y|G, E, Z; \theta_1)$. However, under violation of the independence assumption, these estimators are biased. Shrinkage estimators then arise as a weighted average of two estimators: one obtained under dependence and the other obtained under independence; the weights are chosen in a data-adaptive fashion and reflect the uncertainty around the conditional G - E association (Mukherjee and Chatterjee, 2008). Chen et al. (2009) propose a class of shrinkage estimators applicable, in principle, to estimation problems beyond $G \times E$ effects.

In a cohort or cross-sectional study, maximization of the joint likelihood with respect to outcome model parameters, even from modeling the joint distribution $p(Y, G, E|Z; \theta_1, \theta_2, \theta_3)$, will be independent of the specification of $p(G, E|Z; \theta_2, \theta_3)$. Because of the lack of outcome dependent sampling, and thus absence of conditioning on Y , the log likelihood becomes a sum of two terms that can be maximized separately. However, the proposed LV framework allows us to incorporate constraints on the G - E association (Section 2.2), raising the possibility of gaining efficiency. In the LV framework, estimation of G - E model parameters cannot be completely disentangled from estimation of outcome model parameters due to integration of the likelihood over the latent variable. The extent of efficiency and power gains for testing of outcome parameters due to such constraints depends on design and effect size settings that we investigate in our simulation study. Although extensive work exists on using G - E independence assumptions in case-control studies, this paper is the first to propose and study the implications such assumptions in a LV setting.

We follow Chen et al. (2009) to describe how estimates obtained under assumptions A_0 and $A_1 - A_3$ can be combined. Denoting the two estimates by $\hat{\theta}_{A_0}$ and $\hat{\theta}_{A^*}$,

$$\hat{\theta}_{\text{shrink}} = \hat{\theta}_{A^*} + K_{\text{MV}}(\hat{\theta}_{A_0} - \hat{\theta}_{A^*}) \quad (5)$$

is the shrinkage estimator, where A^* is any of $A_1 - A_3$ and shrinkage weights are $K_{\text{MV}} = V(\hat{\psi} + \psi\hat{\psi}^{-1})^{-1}$, with $\hat{\psi} = \hat{\theta}_{A_0} - \hat{\theta}_{A^*}$ and V is the estimated asymptotic covariance matrix of $\hat{\psi}$. Alternatively, one may use a diagonal weight matrix K_{CW} where the k^{th} diagonal element is $\hat{V}_k / (\hat{V}_k + \hat{\psi}_k^2)$; \hat{V}_k being the k^{th} diagonal element of V and $\hat{\psi}_k$ the k^{th} component of $\hat{\psi}$.

Choosing K_{CW} leads to ‘component-wise’ shrinkage, since the weights used for a given component of $\hat{\theta}_{shrink}$ depend only on the variance and bias related to that component; we call these estimates EB_{CW} . In contrast, using K_{MV} leads to so-called multivariate shrinkage (Chen et al., 2009), which we refer to as EB_{MV} . In both cases we use superscripts to denote which estimates were combined, e.g., EB_{MV}^{03} denotes combination of estimates obtained under assumptions A0 and A3. Note that (5) is only defined for parameters that are common to both models, and we have slightly abused notation by using $\hat{\theta}_{A^*}$ in (5) to only represent the subset of parameters that are equivalent to those in $\hat{\theta}_{A0}$.

Additional considerations about shrinkage estimators are worth mentioning. First, CW shrinkage may be desirable in terms of efficiency gain, compared to multivariate shrinkage in small samples, because large sampling error in the off-diagonals of \hat{V} undermines the potential efficiency gain from multivariate shrinkage (Chen et al., 2009). Second, the form of the weights imply that EB_{MV} estimates will be more prone to favoring the more flexible models. To see this, note that the component-wise shrinkage weights (the k^{th} diagonal element of K_{CW}) can be re-written as $1/(1+\chi_k^2)$, where $\chi_k^2 = \hat{\psi}_k^2 / \hat{V}_k$ is the ratio of the squared difference in the k^{th} parameter between the two models divided by variance of the difference. The MV shrinkage weights can be similarly re-written: $1/(1 + \chi^2)$ where $\chi^2 = \hat{\psi}^T V^{-1} \hat{\psi} / \hat{\chi}_k^2$ and χ^2 can be interpreted as a bias-variance ratios; when they are smaller than one, the EB estimates will lean toward the simpler model. In contrast to CW shrinkage, the MV shrinkage weight is the same for all parameters, but the ratio χ^2 is a weighted sum of all the bias-variance ratios for all model parameters. Hence, in MV shrinkage, a given parameter might be shrunk given not only its own bias-variance ratio χ_k^2 but also given bias-variance ratios for other parameters. Furthermore, χ_k^2 has expectation (approx) 1 when independence holds, whereas χ^2 has expectation equal to the number of model parameters estimated with assumption A0. Hence, MV shrinkage weights $1/(1 + \chi^2)$ will almost always be small, leading to EB estimates closer to those from more flexible models. Third, the weights summarize information about model fit in the sense of comparing differences in estimated parameters. If the models fit equally well, then corresponding parameters would likely be similar. Large differences in corresponding parameters indicate a poorer fitting model (e.g., constrained model). Hence, shrinkage estimates can help assess model adequacy, with smaller weights for the constrained model implying the more flexible model is preferred. Finally, both CW and MV shrinkage estimators will asymptotically converge to those from the more flexible model.

Chen et al. (2009) provide formal arguments to derive the variance for $\hat{\theta}_{shrink}$. Heuristically, the variance can be obtained by treating $\hat{\theta}_{shrink}$ as a function of two random variables, $\hat{\theta}_{A0}$ and $\hat{\theta}_{A^*}$, with joint covariance matrix $\Sigma = var((\hat{\theta}_{A0}^T, \hat{\theta}_{A^*}^T)^T)$. Letting $h(\theta_{A0}, \theta_{A^*}) = \theta_{A^*} + V(V + \psi\psi^T)^{-1} \psi$, with $\psi = \theta_{A0} - \theta_{A^*}$, and employing the multivariate Delta theorem, then $Var(\hat{\theta}_{shrink}) \approx H^T \hat{\Sigma} H$, where $H = \partial h(\theta_{A0}, \theta_{A^*}) / \partial (\theta_{A0}^T, \theta_{A^*}^T) |_{\theta_{A0} = \hat{\theta}_{A0}, \theta_{A^*} = \hat{\theta}_{A^*}}$. Matrix $\hat{\Sigma} = D^{-1} CD^{-T}$ is constructed using the sandwich-variance formula where $C = 1/N \sum_{i=1}^N P_i P_i^T$ and

$D=1/N\sum_{i=1}^N\partial P_i/\partial\theta$, and $P_i=(S_{A0,i}^\top S_{A*,i}^\top)^\top$ is a stacked vector of likelihood score contributions from each model.

3.3 Simpler estimation strategies

Instead of positing a latent exposure model, (1)–(4), one may fit separate multiple linear regression models (MLR) on each exposure measure, or one regression on their first principal component (PCA), and its interactions with G . We include these simple approaches in Sections 4 and 5.

4. Simulation Studies

We conducted a small scale simulation study to examine the finite sample properties of estimators under various settings of the true data generating model using $l = 1$, $p = 4$, and two genetic classes. Genetic class was generated as a binary variable with prevalence 0.2, similar to our data example. Since there are only two gene classes and one latent exposure, in this section (and section 5) we denote the gene effect among unexposed, the exposure effect among wild types, and the interaction parameters as β_G , β_U , $\beta_{G\times U}$. We investigate the estimators' properties under two scenarios of the G - E association: independence (A0) and dependence (A3). We used either $\beta_U = \beta_G = \beta_{G\times U} = 0$ or $\beta_U = 1$, $\beta_G = \beta_{G\times U} = 2$ (i.e., standardized effects of 0.2, 0.4, 0.4, respectively, since outcome variance was $\sigma^2 = 5^2$). See Supplementary Materials for full design.

Type I error

When G and E are independent, all approaches retain rejection probabilities (P(R)) of approximately 0.05 for tests at the 0.05 significance level (Table 1, scenario A0). When the data are generated under G - E dependence (Table 1, scenario A3), MLE estimates derived assuming independence (A0) have inflated Type I error probabilities for $\hat{\beta}_U$ and $\hat{\beta}_G$. While EB_{CW} retain inflated Type I error rates, EB_{MV} estimates do not.

Efficiency/Power

When G - E independence holds (Table 2, scenario A0), gains in efficiency for $\hat{\beta}_G$ and $\hat{\beta}_{G\times U}$ estimated from A0 compared to A1–A3 are very clear: the variance ratio (Var.R) for $\hat{\beta}_{G\times U}$ estimated under A3 vs A0 is 1.90. Efficiency gains translate to large power gains: power for $\hat{\beta}_{G\times U}$ is 0.44 under A3 and 0.66 under A0. Compared to A3, EB_{CW}^{03} and EB_{MV}^{03} have lower variance ratios (1.46 and 1.74, respectively) and higher power (0.53 and 0.49, respectively). The PCA approach has power comparable to that from A0, despite the bias in $\hat{\beta}_U$ and $\hat{\beta}_{G\times U}$.

Bias

When G - E independence does not hold, all parameter estimates have large biases, except those from A3 and EB_{MV}^{03} (Table 2, scenario A3). Of the MLEs, bias is larger when A0 or A1 are assumed; further, A1 vs. A0 does not result in uniformly less (absolute) bias for both β_U and $\beta_{G\times U}$. Simply relaxing the assumption of different mean and variance for the latent variable, but not the measurement model, may not be sufficient to reduce bias, and could in fact increase it.

EB_{CW} estimates are approximately half way between $A0$ and the more flexible models, although the exact distance varies depending on the magnitude of the coefficient. As such, they retain some of the bias of $A0$ estimates when G - E dependence exists. For EB_{CW}^{01} , the bias persists and is larger for β_U and $\beta_{G \times U}$ than the bias in $A0$ for these parameters. In contrast, EB_{MV}^{02} and EB_{MV}^{03} are generally closer to the more flexible model (see Section 3.2), and mostly eliminate the bias in $A0$.

As would be expected from the measurement error literature, parameter estimates using only $E1$ or PCA as the predictor in multiple regression analysis are biased, i.e., $\hat{\beta}_U$ and $\hat{\beta}_{G \times U}$ are attenuated. However, note that $\hat{\beta}_G$ have large bias as well, due to the G - E dependence and the measurement error in $E1$ or the first PC in measuring exposure. Measurement error in one predictor (e.g., $E1$) can induce bias in regression coefficients of covariates measured without error (e.g., G) that are correlated with the error-prone predictor (Budtz-Jørgensen et al., 2003a). The bias in $\hat{\beta}_G$ can be in either positive or negative under the alternative hypothesis (Huang et al., 2005), although estimates will be unbiased when there is no exposure effect ($\beta_U = \beta_{G \times U} = 0$, Table 1).

MSE

EB_{MV}^{03} estimates eliminate the bias in $A0$, but are less efficient; EB_{CW}^{03} retain some bias, but achieve smaller mean squared error (MSE), hence a better bias-efficiency tradeoff. For example, although EB_{CW}^{03} incurs 15% bias, its MSE is 0.93, in contrast to an MSE of 1.31 for EB_{MV}^{03} . EB_{CW} estimators achieve a better bias-variance compromise in small samples compared to EB_{MV} .

Additional simulation results for the case of null main effects and small interaction parameter: $\beta_U = 0$, $\beta_G = 0$, and $\beta_{G \times U} = 0.1$ demonstrate that efficiency gains in $A0$ vs $A3$ are still observed (Var.R=1.5). Although to a smaller degree, bias in $\hat{\beta}_{G \times E}$ persisted when incorrectly assuming $A0$ (11% v.s. 3% bias in $A3$).

Recommendation

For hypothesis testing alone, PCA approaches may be just as good as using a full latent variable model because they maintain Type I error (Table 1) and have power comparable to $A0$ (Table 2). However, in terms of both bias and efficiency, using EB_{MV}^{03} is our recommended estimation strategy. Although EB_{MV}^{03} has slightly higher MSE than EB_{CW}^{03} , it yields unbiased estimates, provides better control of Type I error, and has higher power than EB_{CW}^{03} .

5. Modeling lead exposure, iron metabolism genes, and birth weight

We use data from the first ELEMENT cohort, where the following prenatal lead exposure biomarkers were collected on the mother and child: maternal blood lead levels at delivery and umbilical cord blood lead as well as maternal bone lead levels (patella and tibia) (Gonzalez-Cossio et al., 1997). Birth weight is the health outcome of interest in our analysis.

To be included in this analysis, children had to be genotyped, and have measured birth weight and least one of the four prenatal exposure biomarkers (N=406). Missing data on covariates was imputed five times (Raghunathan et al., 2002). Parameter estimates were obtained from the imputed data sets and combined across the imputed data sets according to standard formulae (Little and Rubin, 2002, pg. 86).

Deleterious effects of prenatal lead exposure on birth weight have been demonstrated (Gonzalez-Cossio et al., 1997), and the main effects of lead exposure in this sample using the LV model are significant ($\hat{\beta}_U = -54.12$, $se(\hat{\beta}_U) = 25.1$, $T_U = -2.16$, Supplementary Materials Table 2). However, individuals with at least one iron metabolism gene variant may be protected against reduced birth weight due to lead exposure (Cantonwine et al., 2010). However, since iron metabolism genes appear to up-regulate iron *and* lead absorption (Hopkins et al., 2008), there may be dependence between genotype status and lead exposure. We use two SNPs related to iron metabolism, variants of the hemochromatosis gene (C282Y and H63D), and dichotomize genotype into wild type for both ('aa' and 'bb') or variant for any ('Aa', 'AA', 'bB', or 'BB'); both SNPs were in Hardy-Weinberg equilibrium, and 83 participants (20%) were classified as variants.

Increasing lead exposure among wild types is associated with decreased birth weight (negative $\hat{\beta}_U$ in Table 3). The largest point estimate is obtained under assumption A1, whereas the lowest is obtained using MLR with the observed patella lead measure as the exposure marker. Such large attenuation in the MLR estimate is due to measurement error of patella lead in capturing prenatal exposure. Similarly, the MLR effect estimated from a PCA-derived exposure measure is attenuated, consistent with the simulation studies. Among the MLE and *EB* estimates, those obtained from assumptions A0 and A1* have the smallest standard errors, that increase with increasing flexibility of the model as expected. The PCA and MLR standard errors are much smaller, and therefore, even though the point estimates are also largely attenuated, the test statistics are similar to those for the MLE estimates. Component-wise shrinkage estimates are approximately half-way between those from A0 and those from the more flexible models A1*-A3. EB_{CW}^{01} is closer to the estimates obtained assuming A1, but EB_{CW}^{02} and EB_{CW}^{03} are closer to the estimates from A0 than from those obtained with A2 or A3. In contrast, EB_{MV}^{02} and EB_{MV}^{03} are closer to the estimates from A2 or A3 compared to those from A0. CW shrinkage favors simpler models since it only trades off bias-variance in one parameter at a time, whereas MV shrinkage favors the more flexible model because it simultaneously considers differences in all model parameters (Section 3.2).

Estimates and standard errors for $\hat{\beta}_G$ are fairly constant across *G-E* assumptions and *EB* estimates. However, $\hat{\beta}_G$ from MLR and PCA are much higher (more negative) than those from MLE. This can be due to bias arising due to exposure-gene correlation and exposure measurement error (Budtz-Jørgensen et al., 2003a; Huang et al., 2005).

Being variant for iron metabolism genes is protective against reduced the birth weight due to lead exposure ($\hat{\beta}_{G \times U}$ are positive), as hypothesized (Cantonwine et al., 2010). Whereas assumptions about *G-E* independence did not ultimately alter the conclusions for the main effect among wild types (e.g., all t-test statistics $T_U < -2.4$ in all assumptions), conclusions

about $\hat{\beta}_{G \times U}$ are impacted by the assumed G - E model. The most flexible model yields largest estimated effects (105.0g with A3 vs 92.4g with A0) and the standard error is lower (49.2 in A3 vs 63.5 in A0), resulting in t-statistics for the $G \times U$ effect as low as $T_{G \times U} = 1.46$ (A0) and as large as 2.13 (A3). This is likely due to a higher degree of overall model residual variance explained in A3 due to an increased number of parameters (i.e., lowest $-2 \log$ likelihood, Table 4).

Differences in MLE estimates for the outcome parameters can be largely explained by a few key differences exposure model parameters by genotype (Table 4). While there is little difference in average exposure levels between wild types and variants (small $\hat{\gamma}_g$), the variance of the latent variable is twice as high among variants ($\hat{\Phi}_{g=1} = 2.11$) than among wild types ($\hat{\Phi}_{g=0} = 1.05$). Residual variances for E_1 and E_2 , Θ_{11} and Θ_{22} , also appear to differ between genotypes (51% and 72% difference, respectively), as does λ_4 (17% difference). This deserves further study—e.g., differences in Θ_{11} and Θ_{22} might be due to maternal genotypes, which are inherently correlated to infant genotype. Such investigation is out of the scope of the current work, but this finding highlights the utility of LV models in elucidating potential biological pathways.

In this example, implementing multivariate shrinkage was possible only for combining estimates from A0 with those from A3 and A0 with A2 estimates. MV shrinkage using A0 and A1 (and A1*) estimates resulted in a numerically singular variance matrix

$\Sigma = \text{Var}((\hat{\theta}_{A0}^\top, \hat{\theta}_{A*}^\top)^\top)$, likely due to measurement model parameters being too similar (and correlated) when only making small changes in the latent variable model (3). Although in the example we implemented all approaches for exposition, and even though standard model fit criteria (Table 4) would point toward model A1 being a better model in this particular example, as a general strategy we prefer outcome model parameters estimated using the EB_{MV}^{03} approach. This approach avoids the potential for increased Type I errors due to fitting multiple models before arriving at a final model, and minimizes bias in outcome model parameters that may persist due to differences in exposure model parameters associated to genotype that may not be declared “significantly different” due to lack of power.

6. Discussion

The presence of multiple correlated measures of exposure exacerbates existing challenges in $G \times E$ studies. The current paper is the first step towards an integrated framework where LV models are used to reduce the dimensionality of the exposure measure, thereby limiting the number of tests made and boosting power. Due to the general model formulation, it is easy to accommodate measurement errors in predictors, a pervasive problem in environmental epidemiology, and reduce multicollinearity concerns. Furthermore, a major intuitive appeal of the LV approach is that it provides not only estimates of the disease model parameters, but also a clearer picture of the underlying G - E association and helps capture the essence of the scientific problem. For a genetic marker and exposure which may have a common metabolic pathway, this model is more meaningful to practitioners than a multiple regression model relating Y_i to G_i and E_i which is not informative about the association between G_i and E_i .

Because of the flexibility afforded by LV models, one challenge is the potential for model misspecification. In this particular application of LV models, we described various specifications of the G - E association, and discussed how restrictions in the G - E model boost efficiency of the $G \times E$ associations, but may incur bias when such restrictions are incorrectly made. We proposed a strategy where one would fit a restricted model and the most flexible model afforded by the data, and then combine estimates based on shrinkage ideas. The proposed approach yields estimates that data-adaptively compromise between bias and variance, and avoids having to fit and re-fit models until a best fitting model is found. Alternatively, estimation could proceed in two stages. First, the most flexible model could be estimated, and genotype differences in exposure model parameters tested. In the second stage, parameters that were found to not differ by genotype would be constrained to be equal across genotypes. However, such two-stage approach would also suffer from inflated Type I error (Mukherjee and Chatterjee, 2008). Yet another alternative, with a similar flavor to what we proposed here, is to average parameter estimates obtained under various G - E assumptions according to prior information of the G - E association (Li and Conti, 2009) or using model fit criteria as weights (Hjort and Claeskens, 2003). Further still, one could use LASSO or Ridge penalties to select which exposure model parameters vary by genotype (Leoutsakos et al., 2010). Lastly, extensions of the methods proposed could include using a continuous genetic risk score G , such that a larger number of genetic categories can be (indirectly) included without collapsing to a few categories due to limited sample size. Such extension may not be straightforward since the multiple group analysis used here would not apply. Compromise estimators like the ones presented have not been used in the LV modeling literature, but can be a tool to achieve improved modeling strategies and robustness in LV models in applications even beyond $G \times E$ studies.

It is possible that one may use the proposed approach for screening $G \times E$ effects in genome-wide interaction studies. In our simulation studies, the estimation procedure takes approximately 0.36 minutes per data set in a desktop computer with 3.2GHz Intel processor and 1GB RAM. In the advent of cluster and parallel computing the proposed approach is scalable to genome-wide studies. Nevertheless, if the intent is solely testing, and not estimation, the PCA approach may be suitable, since, as shown in the simulation studies, it had comparable power to the proposed shrinkage estimates, despite substantial bias. Employing dimension-reduction approaches to the environmental exposure data will reduce multiple testing problems since only one genome-wide scan would be needed, instead of one scan for each observed exposure. Our methods are particularly appealing to study $G \times E$ effects with a given environmental exposure and genetic subclasses defined through genes on a related metabolic pathway.

The availability of higher dimensional genomic data, and multiple continuous or categorical outcomes point to several extensions of our work. General LV models encompass latent class models (Skrondal and Rabe-Hesketh 2004), hence one could posit a latent class model for multiple genetic factors, G_i , which borrows strength from multiple loci and can minimize the chance of false positives (Schumacher and Kraft, 2007). Recent proposals (Chatterjee et al., 2006) pose gene-gene interaction models based on a latent variable approach, and can be extended to reduce the dimension of gene-gene-environment interaction models. Similar to

what we have done for the exposure model in the present paper, a latent outcome model to summarize correlated multivariate or longitudinal outcome data Y_i can be proposed. One would summarize multivariate correlated outcomes by latent traits, i.e., express Y_i in terms of latent outcomes f_i (e.g., Budtz-Jørgensen et al., 2003b), and estimate model parameters for a regression of f_i on U_i and G_i . When Y_i involves repeated measures over time (e.g., growth curves), the model for the observed multivariate vector Y_i for subject i , measured at multiple time points may contain a random slope and random intercept, which are inherently latent variables. The random effects can be modeled as dependent on U_i and G_i and other covariates, such that inferences on how exposure and genes modify growth rates can naturally be obtained. Moreover, multivariate observations reflecting latent variables repeated over time (Roy and Lin, 2000), and time-to-event data (Proust-Lima et al., 2009) can be incorporated. In summary, extensions of the present model can involve summarization of all three data components: Y , G , E .

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors thank ELEMENT investigators for providing data for the example, as well as the following NIEHS grants that supported data collection: K23ES000381; P01 ES012874; P42 ES05947; R01 ES013744; R01 ES014930; R01 ES007821. The authors also acknowledge salary support from grants NSF DMS 1007494, NIEHS R01 ES016932 and R01 ES017022, and NIEHS/EPA 1-P20-SE018171-01.

References

- Bentler, PM.; Hu, LT. Evaluating model fit. In: Hoyle, RH., editor. Structural Equation Modeling. London: Sage; 1995. p. 76-99.
- Bollen, KA. Structural Equations with Latent Variables. New York: John Wiley & Sons; 1989.
- Budtz-Jørgensen E, Keiding N, Grandjean P, Weihe P, White RF. Consequences of exposure measurement error for confounder identification in environmental epidemiology. *Statistics in Medicine*. 2003a; 22:3089–3100. [PubMed: 12973789]
- Budtz-Jørgensen E, Keiding N, Grandjean P, Weihe P, White RF. Statistical methods for the evaluation of health effects of prenatal mercury exposure. *Environmetrics*. 2003b; 14:105–120.
- Cantonwine D, Hu H, Tellez-Rojo MM, Sánchez BN, Lamadrid-Figueroa H, Ettinger AS, Mercado-García A, Hernandez-Avila M, Wright RO. HFE gene variants modify the association between maternal lead burden and infant birthweight: a prospective birth cohort study in Mexico City, Mexico. *Environmental Health*. 2010; 9:43. [PubMed: 20659343]
- Carroll, RJ.; Ruppert, D.; Stefanski, L.; Crainiceanu, C. Measurement Error in Nonlinear Models: A Modern Perspective. 2nd edition. Boca Raton, FL: Chapman & Hall; 2006. Monographs on Statistics and Applied Probability.
- Chatterjee N, Carroll RJ. Semiparametric maximum likelihood estimation exploiting gene-environment independence in case-control studies. *Biometrika*. 2005; 92:399–418.
- Chatterjee N, Kalaylioglu Z, Moslehi R, Peters U, Wacholder S. Powerful multilocus tests of genetic association in the presence of gene-gene and gene-environment interactions. *American Journal of Human Genetics*. 2006; 79:1002–1016. [PubMed: 17186459]
- Chen YH, Chatterjee N, Carroll RJ. Shrinkage estimators for robust and efficient inference in haplotype-based case-control studies. *Journal of the American Statistical Association*. 2009; 104:220–233. [PubMed: 19430598]

- Dhungana P, Eskridge KM, Baenziger PS, Campbell BT, Gill KS, Dweikat I. Analysis of genotype-by-environment interaction in wheat using a structural equation model and chromosome substitution lines. *Crop Science*. 2007; 47:477–484.
- Gonzalez-Cossio T, Peterson KE, Sanin LH, Fishbein E, Palazuelos E, Aro A, Hernandez-Avila M, Hu H. Decrease in birth weight in relation to maternal bone-lead burden. *Pediatrics*. 1997; 100:856–862. [PubMed: 9346987]
- Hjort NL, Claeskens G. Frequentist model average estimators. *Journal of the American Statistical Association*. 2003; 98:879–899.
- Hopkins MR, Ettinger AS, Hernandez-Avila M, Schwartz J, Tellez-Rojo MM, Lamadrid-Figueroa H, Bellinger D, Hu H, Wright RO. Variants in iron metabolism genes predict higher blood lead levels in young children. *Environmental Health Perspectives*. 2008; 116:1261–1266. [PubMed: 18795173]
- Huang GH, Bandeen-Roche K. Building an identifiable latent class model with covariate effects on underlying and measured variables. *Psychometrika*. 2004; 69:5–32.
- Huang LS, Wang HK, Cox C. Assessing interaction effects in linear measurement error models. *Journal of the Royal Statistical Society Series C-Applied Statistics*. 2005; 54:21–30.
- Javaras KN, Hudson JI, Laird NM. Fitting ACE structural equation models to case-control family data. *Genetic Epidemiology*. 2010; 34:238–245. [PubMed: 19918760]
- Khoury MJ, Wacholder S. Invited commentary: from genome-wide association studies to gene-environment-wide interaction studies—challenges and opportunities. *American Journal of Epidemiology*. 2009; 169:227–230. discussion 234–235.
- Li DL, Conti DV. Detecting gene-environment interactions using a combined case-only and case-control approach. *American Journal of Epidemiology*. 2009; 169:497–504. [PubMed: 19074774]
- Little, RJA.; Rubin, DB. *Statistical Analysis with Missing Data*. 2nd edition. New Jersey: John Wiley & Sons, Inc; 2002.
- Mukherjee B, Chatterjee N. Exploiting gene-environment independence for analysis of case-control studies: An empirical bayes-type shrinkage estimator to trade-off between bias and efficiency. *Biometrics*. 2008; 64:685–694. [PubMed: 18162111]
- Proust-Lima C, Joly P, Dartigues JF, Jacqmin-Gadda H. Joint modelling of multivariate longitudinal outcomes and a time-to-event: A nonlinear latent class approach. *Computational Statistics & Data Analysis*. 2009; 53:1142–1154.
- Qi L, Ma J, Qi Q, Hartiala J, Allayee H, Campos H. Genetic risk score and risk of myocardial infarction in Hispanics. *Circulation*. 2011; 123:374–380. [PubMed: 21242481]
- Raghunathan, TE.; Solenberger, P.; Van Hoewyk, J. *Survey Methodology Program*. Ann Arbor: University of Michigan; 2002. IVEware: Imputation and Variance Estimation Software User guide.
- Rathouz P, Van Hulle C, Rodgers J, Waldman I, Lahey B. Specification, testing, and interpretation of gene-by-measured-environment interaction models in the presence of gene-environment correlation. *Behavioral Genetics*. 2008; 38:301–315.
- Roy J, Lin XH. Latent variable models for longitudinal data with multiple continuous outcomes. *Biometrics*. 2000; 56:1047–1054. [PubMed: 11129460]
- Sánchez BN, Budtz-Jørgensen E, Ryan LM, Hu H. Structural equation models: A review with applications to environmental epidemiology. *Journal of the American Statistical Association*. 2005; 100:1443–1455.
- Schumacher FR, Kraft P. A bayesian latent class analysis for whole-genome association analyses: an illustration using the gaw 15 simulated rheumatoid arthritis dense scan data. *BMC Proc*. 2007; 1(Suppl 1):S112. [PubMed: 18466453]
- Skrondal, A.; Rabe-Hesketh, S. *Generalized Latent Variable Modeling: Multilevel, Longitudinal, and Structural Equation Models*. Boca Raton, FL: Chapman & Hall; 2004.
- Westland JC. Lower bounds on sample size in structural equation modeling. *Electronic Commerce Research and Applications*. 2010; 9:476–487.




Figure 1. Path diagram showing relationships between exposure biomarkers, latent prenatal lead exposure, iron metabolism genes, and birth weight.




Figure 2. Path diagrams showing gene-environment dependence assumptions.

Table 1

Bias, variance ratios (Var.R), mean squared error (MSE), and rejection probabilities (P(R)) for outcome model parameter estimates under two scenarios of true exposure model parameters. Outcome model parameters set at $\beta_U = \beta_G = \beta_{G \times U} = 0$, $\sigma^2 = 5$; sample size was $N = 350$, with 500^8 replicates.

Data Model	Est. Method ^a	$\hat{\beta}_U$ Bias	Var.R ^b	MSE	P(R)	$\hat{\beta}_G$ Bias	Var.R ^b	MSE	P(R)	$\hat{\beta}_{G \times U}$ Bias	Var.R ^b	MSE	P(R)
A0	A0	0.00	1(Ref)	0.15	0.063	-0.05	1(Ref)	0.46	0.048	-0.04	1(Ref)	0.65	0.036
	A1	0.00	(1.02)	0.16	0.059	-0.05	(1.05)	0.46	0.044	-0.04	(1.10)	0.69	0.038
	A2	0.00	(1.01)	0.15	0.059	-0.04	(1.11)	0.48	0.042	-0.05	(1.15)	0.72	0.036
	A3	0.00	(1.02)	0.15	0.061	-0.04	(1.12)	0.49	0.042	-0.04	(1.15)	0.74	0.044
EB _{CW} ⁰¹	EB _{CW} ⁰¹	0.00	(1.02)	0.15	0.061	-0.05	(0.99)	0.46	0.046	-0.04	(0.99)	0.66	0.050
	EB _{CW} ⁰²	0.00	(1.02)	0.15	0.069	-0.04	(1.02)	0.46	0.046	-0.04	(1.01)	0.66	0.050
	EB _{CW} ⁰³	0.00	(1.02)	0.15	0.069	-0.05	(1.03)	0.46	0.044	-0.04	(1.04)	0.67	0.052
EB _{MV} ⁰¹	EB _{MV} ⁰¹	0.00	(1.03)	0.16	0.063	-0.05	(1.02)	0.46	0.044	-0.04	(1.04)	0.69	0.046
	EB _{MV} ⁰²	0.00	(1.03)	0.15	0.063	-0.04	(1.09)	0.48	0.044	-0.05	(1.08)	0.71	0.044
	EB _{MV} ⁰³	0.00	(1.03)	0.15	0.063	-0.04	(1.09)	0.48	0.042	-0.04	(1.07)	0.73	0.055
PCA ^d	E1 ^c	0.00	(0.23)	0.03	0.044	-0.04	(1.38)	0.64	0.052	0.00	(0.24)	0.15	0.048
	PCA ^d	0.00	(0.23)	0.03	0.063	-0.04	(1.02)	0.46	0.048	0.01	(0.24)	0.15	0.042
	A0	0.03	1(Ref)	0.18	0.126	-0.07	1(Ref)	0.77	0.142	0.03	1(Ref)	0.52	0.043
EB _{CW} ⁰¹	A1	0.04	(2.59)	0.32	0.038	-0.10	(2.46)	1.15	0.063	0.02	(1.21)	0.66	0.050
	A2	0.03	(1.42)	0.18	0.036	-0.08	(2.07)	0.95	0.056	0.03	(1.08)	0.57	0.043
	A3	0.03	(1.27)	0.16	0.038	-0.09	(2.26)	1.04	0.047	0.04	(1.19)	0.63	0.050
EB _{CW} ⁰²	EB _{CW} ⁰¹	0.03	(1.38)	0.23	0.020	-0.07	(1.18)	0.85	0.101	0.02	(1.81)	0.56	0.007
	EB _{CW} ⁰²	0.03	(0.83)	0.18	0.171	-0.07	(1.10)	0.80	0.119	0.03	(1.65)	0.52	0.011

Data Model	Est. Method ^a	$\hat{\beta}_U$ Bias	Var.R ^b	MSE	P(R)	$\hat{\beta}_G$ Bias	Var.R ^b	MSE	P(R)	$\hat{\beta}_{G \times U}$ Bias	Var.R ^b	MSE	P(R)
	EB _{CW} ⁰³	0.03	(0.84)	0.17	0.153	-0.08	(1.16)	0.82	0.106	0.03	(1.61)	0.53	0.018
	EB _{MV} ⁰¹	0.04	(2.59)	0.32	0.036	-0.10	(2.37)	1.15	0.074	0.02	(1.16)	0.66	0.050
	EB _{MV} ⁰²	0.03	(1.39)	0.18	0.050	-0.08	(1.98)	0.95	0.056	0.03	(1.04)	0.57	0.054
	EB _{MV} ⁰³	0.03	(1.25)	0.16	0.054	-0.09	(2.14)	1.03	0.059	0.04	(1.13)	0.63	0.061
	E1 ^c	0.01	(0.28)	0.04	0.070	-0.00	(1.97)	0.87	0.052	-0.01	(0.23)	0.12	0.047
	PCA ^d	-0.01	(0.36)	0.04	0.038	-0.05	(1.57)	0.75	0.056	-0.01	(0.24)	0.12	0.061

^a A0, A1, A2, A3 denote MLE; A1* not included because combining A0 and A1* resulted in singular variance matrix Σ (see Section 5).

^b Ratios of empirical variances, comparing to variance of A0

^c Multiple regression using one exposure marker, E1

^d Multiple regression using 1st principal component of (E1, E2, E3, E4) as exposure marker

[§] 4.6% (A0) and 7.0% (A3) data sets excluded to lack of convergence or unstable results, see Supplementary Materials for details.

Table 2

Percent bias, variance ratios (Var.R), mean squared error (MSE), and rejection probabilities (P(R)) for outcome model parameter estimates under two scenarios of true exposure model parameters. Outcome model parameters set at $\beta_U = 1$, $\beta_G = \beta_{G \times U} = 2$, $\sigma^2 = 5$; sample size was $N = 350$, with 500^s replicates.

Data Model	Est. Method ^d	$\hat{\beta}_U$ Bias%	Var.R ^b	MSE	P(R)	$\hat{\beta}_G$ Bias%	Var.R ^b	MSE	P(R)	$\hat{\beta}_{G \times U}$ Bias%	Var.R ^b	MSE	P(R)
A0	A0	-1.3%	1(Ref)	0.17	0.74	3.4%	1(Ref)	0.47	0.85	-1.4%	1(Ref)	0.74	0.66
	A1	-1.0%	(1.02)	0.18	0.74	3.0%	(1.14)	0.54	0.78	1.3%	(1.18)	0.83	0.62
	A2	-1.0%	(1.03)	0.18	0.74	3.0%	(1.73)	0.74	0.63	3.1%	(1.74)	1.06	0.47
	A3	-0.7%	(1.04)	0.18	0.74	2.9%	(1.76)	0.75	0.63	3.1%	(1.90)	1.18	0.44
	EB ⁰¹ _{CW}	-1.2%	(1.01)	0.17	0.73	3.2%	(1.08)	0.50	0.82	-0.3%	(1.08)	0.76	0.64
	EB ⁰² _{CW}	-1.2%	(1.02)	0.17	0.74	3.4%	(1.45)	0.56	0.71	-0.5%	(1.36)	0.82	0.56
	EB ⁰³ _{CW}	-1.0%	(1.02)	0.17	0.75	3.3%	(1.47)	0.56	0.72	-1.1%	(1.46)	0.85	0.53
	EB ⁰¹ _{MV}	-1.0%	(1.02)	0.18	0.73	3.0%	(1.12)	0.54	0.79	1.3%	(1.13)	0.83	0.64
	EB ⁰² _{MV}	-1.0%	(1.03)	0.18	0.73	3.0%	(1.66)	0.73	0.64	3.1%	(1.64)	1.04	0.50
	EB ⁰³ _{MV}	-0.7%	(1.04)	0.18	0.73	2.9%	(1.67)	0.72	0.65	2.8%	(1.74)	1.14	0.49
	E1 ^c	-65.4%	(0.22)	0.46	0.46	3.2%	(1.33)	0.49	0.83	-63.6%	(0.23)	1.79	0.41
	PCA ^d	-53.0%	(0.22)	0.31	0.74	3.3%	(0.94)	0.47	0.86	-51.5%	(0.23)	1.24	0.65
A3	A0	12.1%	1(Ref)	0.20	0.86	35.7%	1(Ref)	1.24	0.94	-27.6%	1(Ref)	0.92	0.48
	A1	45.6%	(2.57)	0.55	0.78	-18.7%	(2.57)	1.37	0.35	-42.0%	(1.24)	1.50	0.27
	A2	8.9%	(1.43)	0.19	0.79	10.3%	(3.79)	1.96	0.51	-13.5%	(1.86)	1.21	0.40
	A3	3.7%	(1.26)	0.17	0.80	-1.4%	(5.18)	2.14	0.41	0.9%	(2.65)	1.34	0.39
	EB ⁰¹ _{CW}	32.8%	(2.51)	0.43	0.72	0.0%	(2.83)	1.23	0.44	-37.5%	(1.94)	1.30	0.19
	EB ⁰² _{CW}	10.4%	(0.94)	0.19	0.88	28.4%	(3.47)	1.58	0.67	-21.3%	(1.98)	0.94	0.31

Data Model	Est. Method ^a	$\hat{\beta}_U$ Bias%	Var.R ^b	MSE	P(R)	$\hat{\beta}_G$ Bias%	Var.R ^b	MSE	P(R)	$\hat{\beta}_{GXU}$ Bias%	Var.R ^b	MSE	P(R)
EB ⁰³ _{CW}		7.4%	(0.95)	0.18	0.86	21.8%	(4.45)	1.56	0.57	-14.6%	(2.48)	0.93	0.30
		45.5%	(2.54)	0.55	0.77	-18.6%	(2.55)	1.36	0.37	-42.0%	(1.22)	1.51	0.27
		9.0%	(1.38)	0.19	0.79	10.7%	(4.03)	1.93	0.50	-13.7%	(1.97)	1.20	0.39
EB ⁰³ _{MV}		3.8%	(1.23)	0.17	0.79	-0.7%	(4.89)	2.09	0.44	0.4%	(2.50)	1.31	0.41
	E1 ^c	-63.5%	(0.27)	0.44	0.51	110.7%	(1.97)	5.59	0.99	-78.0%	(0.23)	2.59	0.23
PCA ^d		-44.2%	(0.33)	0.23	0.78	37.3%	(1.52)	1.29	0.88	-63.7%	(0.23)	1.77	0.48

^a A0, A1, A2, A3 denote MLE; A1* not included because combining A0 and A1* resulted in singular variance matrix Σ (see Section 5).

^b Ratios of empirical variances, comparing to variance of A0

^c Multiple regression using one exposure marker, E1

^d Multiple regression using 1st principal component of (E1, E2, E3, E4) as exposure marker

^e 3.6% (A0) and 2.4% (A3) data sets excluded to lack of convergence or unstable results, see Supplementary Materials for details.

Outcome model parameter estimates, robust standard errors, and t-statistics obtained using MLE under assumptions A0–A3 and shrinkage-based estimates combining assumptions. Coefficients $\hat{\beta}_U$ and $\hat{\beta}_{G \times U}$ have been re-scaled to represent changes in birth weight(g) associated with an increase of 10 µg Pb/g in patella bone mass. Models are adjusted for maternal age, parity, education and marital status

Table 3

Est. Method ^{a,b}	$\hat{\beta}_U$	$se(\hat{\beta}_U)$	T_U	$\hat{\beta}_G$	$se(\hat{\beta}_G)$	T_G	$\hat{\beta}_{G \times U}$	$se(\hat{\beta}_{G \times U})$	$T_{G \times U}$
A0	-80.59	29.97	-2.69 [†]	-97.75	51.30	-1.91	92.42	63.50	1.46
A1*	-79.51	29.79	-2.67 [†]	-99.09	51.45	-1.93	91.92	62.56	1.47
A1	-98.44	39.54	-2.49 [†]	-98.47	51.30	-1.92	103.09	52.71	1.96 [†]
A2	-85.14	33.75	-2.52 [†]	-98.83	51.42	-1.92	90.66	53.57	1.69
A3	-86.62	35.01	-2.47 [†]	-100.74	51.41	-1.96 [†]	105.00	49.24	2.13 [†]
EB ^{01*} _{CW}	-80.05	29.76	-2.69 [†]	-98.02	51.31	-1.91	91.90	63.25	1.45
EB ⁰¹ _{CW}	-91.12	40.15	-2.27 [†]	-97.89	51.31	-1.91	101.95	61.06	1.67
EB ⁰² _{CW}	-81.73	32.13	-2.54 [†]	-97.90	51.24	-1.91	92.58	60.07	1.54
EB ⁰³ _{CW}	-82.40	33.04	-2.49 [†]	-98.93	51.37	-1.93	94.63	62.10	1.52
EB ⁰² _{MV}	-85.02	33.64	-2.53 [†]	-98.80	51.40	-1.92	90.70	53.76	1.69
EB ⁰³ _{MV}	-86.28	34.66	-2.49 [†]	-100.58	51.39	-1.96 [†]	104.30	49.81	2.09 [†]
E ₁ ^c	-41.58	15.06	-2.76 [†]	-191.50	71.07	-2.69 [†]	60.59	31.19	1.94
PCA ^d	-46.58	15.34	-3.04 [†]	-187.92	69.27	-2.71 [†]	58.52	30.04	1.95

^b Estimates from models A0, A1, and A1*, were not sufficiently distinguishable from each other, hence EB⁰¹_{MV} and EB^{01*}_{MV} could not be obtained (Section 5).

^c Multiple regression using patella lead, E₁

^d Multiple regression using 1st principal component of E₁, E₂, E₃, E₄ as exposure marker

[†] p-value < 0.05

Exposure model parameter estimates and robust standard errors obtained under assumptions A0–A3 described by Figure 1.

Table 4

	A0	A1*	A1	A2	A3
<u>Model for latent variable</u>					
α_0	Est(SE) 1.536 (0.078)	Est(SE) 1.519 (0.086)	Est(SE) 1.526 (0.081)	Est(SE) 1.506 (0.087)	Est(SE) 1.506 (0.086)
γ_g		0.084 (0.178)	0.050 (0.182)	0.148 (0.207)	0.148 (0.208)
$\Phi_{g=0}$	1.230 (0.312)	1.253 (0.317)	0.835 (0.218)	1.090 (0.306)	1.056 (0.333)
$\Phi_{g=1}$			1.645 (0.489)	1.440 (0.460)	2.112 (0.916)
<u>Measurement Model</u>					
Estimates for wild types					
v_2	1.031 (0.053)	1.021 (0.057)	1.024 (0.056)	1.040 (0.057)	1.040 (0.056)
v_3	1.759 (0.024)	1.757 (0.025)	1.758 (0.025)	1.756 (0.027)	1.756 (0.027)
v_4	2.043 (0.023)	2.041 (0.023)	2.042 (0.023)	2.034 (0.026)	2.034 (0.026)
λ_2	0.553 (0.126)	0.542 (0.124)	0.680 (0.136)	0.527 (0.136)	0.558 (0.157)
λ_3	0.140 (0.036)	0.138 (0.035)	0.156 (0.036)	0.129 (0.041)	0.131 (0.043)
λ_4	0.125 (0.032)	0.124 (0.032)	0.140 (0.033)	0.114 (0.037)	0.116 (0.039)
Θ_{11}	1.172 (0.283)	1.148 (0.289)	1.401 (0.219)	1.236 (0.262)	1.258 (0.311)
Θ_{22}	0.710 (0.099)	0.717 (0.097)	0.624 (0.103)	0.671 (0.096)	0.627 (0.104)
Θ_{33}	0.218 (0.017)	0.218 (0.017)	0.218 (0.017)	0.218 (0.017)	0.217 (0.019)
Θ_{44}	0.194 (0.015)	0.194 (0.015)	0.194 (0.015)	0.195 (0.015)	0.197 (0.017)
Θ_{34}	0.168 (0.014)	0.168 (0.014)	0.168 (0.014)	0.168 (0.014)	0.168 (0.016)
Estimates for variants ^d					
v_2				0.880 (0.156)	0.920 (0.139)
v_3				1.746 (0.059)	1.751 (0.056)
v_4				2.053 (0.055)	2.056 (0.051)
λ_2				0.768 (0.162)	0.496 (0.205)
λ_3				0.180 (0.058)	0.142 (0.063)
λ_4				0.161 (0.053)	0.136 (0.058)
Θ_{11}					0.612 (0.809)

	A0	A1*	A1	A2	A3
Θ_{22}					1.077 (0.259)
Θ_{33}					0.226 (0.040)
Θ_{44}					0.183 (0.032)
Θ_{34}					0.171 (0.033)
Model fit criteria ^b , (criterion for good fit)					
Num Param ^c	23	24	25	31	36
-2LL (smaller is better)	12302.8	12302.6	12296.6	12291.4	<u>12287.0</u>
AIC (smaller is better)	12348.8	12350.6	<u>12346.5</u>	12353.4	12359.0
BIC (smaller is better)	<u>12440.9</u>	12446.7	12446.6	12477.6	12503.2
CFI ^d (>.95)	0.963	0.962	<u>0.970</u>	0.968	0.967
TLL ^d (>.95)	0.961	0.959	<u>0.967</u>	0.962	0.958
RMSEA ^d (<.05)	0.041	0.042	<u>0.038</u>	0.041	0.043

^a Parameters estimates for variants are the same for as for wild types unless shown here

^b For the exposure and outcome model combined

^c Including outcome model parameters

^d See Bentler and Hu (1995) for definitions