

## TECHNOLOGICAL VALUE OF SPRING WHEAT OF ZEBRA CULTIVAR AS RELATED TO THE WAY OF NITROGEN AND MAGNESIUM APPLICATION

### WARTOŚĆ TECHNOLOGICZNA PSZENICY JAREJ ODMIANY ZEBRA W ZALEŻNOŚCI OD SPOSOBU APLIKACJI AZOTU I MAGNEZU

RALCEWICZ M., KNAPOWSKI T., KOZERA W., BARCZAK B.

University of Technology and Life Sciences, Faculty of Agriculture,  
Chair of Agricultural Chemistry, 85-326 Bydgoszcz, 5 Seminaryjna St., Poland  
e-mail: ralce@utp.edu.pl, knap@utp.edu.pl, kozera@utp.edu.pl, barczak@utp.edu.pl

Manuscript received: August 28, 2009; Reviewed: September 17, 2009; Accepted for publication: September 18, 2009

#### ABSTRACT

A significant increase in grain yield was observed following soil fertilization with nitrogen up to the level of  $90 \text{ kg} \cdot \text{ha}^{-1}$ , independently of foliar fertilization. Among the examined variants of foliar fertilization on average, the significantly highest grain yield was obtained in the objects in which nitrogen and magnesium were administered in parallel. Within the entire range of nitrogen doses administered to soil, a moderately significant increase was noted in total protein content, to the level of  $90 \text{ kg N} \cdot \text{ha}^{-1}$  a moderately increased content of wet gluten, sedimentation ratio and bread volume were recorded and following application of  $60 \text{ kg N} \cdot \text{ha}^{-1}$  an increase in water absorbability of flour was observed. Independently of the soil fertilization with nitrogen, the most favourable of the studied variants of foliar fertilization for principal parameters of baking value proved to be a parallel application of nitrogen and magnesium or administration of nitrogen only.

Key words: spring wheat, nitrogen fertilization, foliar fertilization, grain yield, technological parameters

#### STRESZCZENIE

Istotny wzrost plonu ziarna stwierdzono po zastosowaniu doglebowo dawki azotu do poziomu  $90 \text{ kg} \cdot \text{ha}^{-1}$ , niezależnie od nawożenia dolistnego. Spośród badanych wariantów nawożenia dolistnego średnio istotnie najwyższy plon ziarna uzyskano na obiektach, gdzie zastosowano łącznie azot i magnez. Stwierdzono, w całym zakresie zastosowanych doglebowo dawek azotu, średnio istotny wzrost zawartości białka ogólnego, do poziomu  $90 \text{ kg N} \cdot \text{ha}^{-1}$  średnio istotny wzrost zawartości mokrego glutenu, wskaźnika sedymentacji i objętości pieczywa, a po zastosowaniu  $60 \text{ kg N} \cdot \text{ha}^{-1}$  wodochłonności mąki. Najkorzystniejszym spośród badanych wariantów nawożenia dolistnego, niezależnie od zastosowanego doglebowo nawożenia azotem, dla najważniejszych wyróżników wartości wypiekowej okazało się łączne zastosowanie azotu i magnezu lub tylko azotu.

Słowa kluczowe: pszenica jara, nawożenie azotem, nawożenie dolistne, plon ziarna, parametry technologiczne

## DETAILED ABSTRACT

Celem niniejszej pracy było określenie wysokości dawki azotu zastosowanej doglebowo oraz wskazanie wariantu nawożenia nalistnego, które pozwoliłyby na osiągnięcie wysokiego plonu ziarna pszenicy jarej odmiany Zebra o wymaganych parametrach wypiekowych. Badania prowadzono w latach 2005-2007 w RZD Minikowo na glebie płowej typowej, według międzynarodowej klasyfikacji FAO-UNESCO jest to Albic Luvisols, zaliczanej do kompleksu żytniego bardzo dobrego, w układzie dwuczynnikowym w trzech powtórzeniach. Pierwszym czynnikiem doświadczenia było doglebowe nawożenie azotem ( $n=4$ ; obiekt kontrolny, 60, 90 i 120 kg N·ha<sup>-1</sup>), które stosowano w formie saletry amonowej. Natomiast drugim czynnikiem ( $n=4$ ) było nawożenie nalistne stosowane w następujących formach: mocznik, siarczan magnezu, mocznik + siarczan magnezu i obiekt bez nawożenia. Istotny wzrost plonu ziarna stwierdzono po zastosowaniu doglebowo dawki azotu do poziomu 90 kg·ha<sup>-1</sup>, niezależnie od nawożenia dolistnego. Spośród badanych wariantów nawożenia dolistnego średnio istotnie najwyższy plon ziarna uzyskano na obiektach, gdzie zastosowano łącznie azot i magnez. Stwierdzono, w całym zakresie zastosowanych doglebowo dawek azotu, średnio istotny wzrost zawartości białka ogólnego, do poziomu 90 kg N·ha<sup>-1</sup> średnio istotny wzrost zawartości mokrego glutenu, wskaźnika sedymentacji i objętości pieczywa, a po zastosowaniu 60 kg N·ha<sup>-1</sup> wodochłonności mąki. Najkorzystniejszym spośród badanych wariantów nawożenia dolistnego, niezależnie od zastosowanego doglebowo nawożenia azotem, dla najważniejszych wyróżników wartości wypiekowej okazało się łączne zastosowanie azotu i magnezu lub tylko azotu. Wyniki badań wskazują również, że w przypadku odmiany Zebra, nawożenie azotem już na poziomie 60 kg·ha<sup>-1</sup> zabezpiecza uzyskanie minimalnych parametrów jakościowych, które determinują ziarno dla potrzeb piekarnictwa.

## INTRODUCTION

Technological values of spring wheat grain determine high demand for products made of the flour. Thus, grain of consumable wheat must manifest a high technological value, which assures good quality of baking products and stability of the technological process [15,16]. The properties are genetically conditioned and are dependent on the involved cultivar but they may vary depending upon natural environment and agrotechnical procedures, including fertilization [6,9,11,12,13,21,27,]. Literature data indicate that nitrogen fertilization represents not only the most important yield-stimulating procedure but

also the most important variable, which affects quality of the yield [5,7,8,13,17,26,29]. Progressive increase in doses of the component not always is equivalent to its increased quality. Therefore, determination of optimum nitrogen doses remains to represent a current problem [2,12,15,26]. Foliar application of aqueous solution of urea proved to be a rational and highly effective procedure in field cultivation and, according to the literature of the subject, parallel application of urea and magnesium sulphate is particularly advantageous [4,23]. The principal benefit of foliar fertilization involves its speed of action and the high utilisation of the introduced in this way alimentary components [14].

Present study aimed at determining the size of nitrogen dose introduced to the soil and selection of foliar fertilization variant, which would assure high yield of spring wheat grain of Zebra cultivar, manifesting the required baking parameters.

## MATERIALS AND METHODS

The field experiment was conducted in 2005-2007, in RZD Minikowo (Kujawy-Pomerania voivodship) in a typical lessive soil, in the international FAO-UNESCO classification the Albic Luvisols. The soil, classified as a rye complex of a very good quality, manifests a neutral pH and a high or average content of absorbable forms of phosphorus, potassium, and magnesium.

The bifactorial experiment was established using the method of the drawn at random sub-blocks with three repetitions. The first variable of the experiment involved soil fertilization with nitrogen ( $n=4$ ), in which Norway saltpetre was used at the following levels of fertilization: I. 60 kg N·ha<sup>-1</sup> (N<sub>60</sub>) in a single dose before sowing, II. 90 kg N·ha<sup>-1</sup> (N<sub>90</sub>) in a divided dose, including 2/3 of the dose applied before sowing and 1/3 of the dose applied at the full phase of shooting (phase 33 according to Zadocx et al.); III. 120 kg N·ha<sup>-1</sup> (N<sub>120</sub>) in a divided dose, including 1/2 of the dose applied before sowing, 1/4 kg applied at the full phase of shooting (phase 33 according to Zadocx et al.) and 1/4 of the dose applied at the beginning of heading (phase 53 according to Zadocx et al.); IV. control object with no nitrogen fertilization (N<sub>0</sub>). The other variable ( $n=4$ ) included foliar fertilization applied in the following forms: urea, magnesium sulphate, urea + magnesium sulphate, and the object with no fertilization. Variants of foliar fertilization were applied at the beginning of shooting phase (phase 31 according to Zadocx et al.) in the form of 10 % solution of urea and 5 % solution of magnesium sulphate and at the beginning of the heading phase (phase 53 according to Zadocx et al.) (5 % solution of urea, 5 % solution of magnesium

sulphate). The applied doses of urea solution introduced  $20.7 \text{ kg N}\cdot\text{ha}^{-1}$  while the applied doses of magnesium sulphate introduced  $4.8 \text{ kg MgO}\cdot\text{ha}^{-1}$ . The cultivated plant involved spring wheat, cultivar Zebra.

The experiment was conducted basing on a uniform level of fertilization with phosphorus and potassium. The forecrop for spring wheat involved oats, collected to obtain green forage. Agrotechnical procedures were conducted in line with recommendations for the species.

In the experiment the size of grain yield was determined and indices of baking value were established, including falling number (according to Hagberg, PN-ISO-3093), total protein content (%N-5.7, PN-75A-04018), gluten content (PN-A-74-043), sedimentation value (test according to Zeleny, PN-ISO-5529), water absorption by flour (PN-ISO 5530) and volume of baking products obtained from 100 g flour (PN-A-74108).

The obtained results of studies were subjected to statistical analysis, employing analysis of variance, and limiting differences were appraised employing the test of Tukey at the significance level of 0.05. In order to establish relationships and dependencies between soil fertilization with nitrogen and the obtained values of qualitative traits in spring wheat, the obtained results were subjected to analysis of direct correlations and of linear regression.

Weather conditions throughout the experiment (2005-2007 growing seasons) are given in Table 1. In the first year of the experiments both the mean air temperature and the total rainfall were lower, as compared with the 1995-2005 means, respectively by:  $0.8^\circ\text{C}$  and  $41.9 \text{ mm}$ , which accounted for 15.6%, while in 2006 and 2007 mean air temperature and total rainfall were higher than the multi-year means by a respective of:  $0.2^\circ\text{C}$  and  $0.3^\circ\text{C}$  and  $38.2 \text{ mm}$ , which accounted for 14.2% and 74.7 mm, namely 27.8%.

## RESULTS AND DISCUSSION

The mean grain yield for Zebra cultivar amounted to  $3.82 \text{ t}\cdot\text{ha}^{-1}$  (Table 2) and it was higher by  $0.5 \text{ t}\cdot\text{ha}^{-1}$ , as compared to the yield obtained for the same cultivar in the experiment of Biskupski et al. [1]. Most of studies documented a favourable effect of increasing doses of nitrogen on yield of spring wheat, although the recommended doses of the component were quite variable [5,10,16,25,26,29]. A moderately significant increase in grain yield was obtained using nitrogen doses up to  $90 \text{ kg}\cdot\text{ha}^{-1}$  and, as compared to  $N_{60}$  and  $N_0$  objects, it was higher by 11.3 and 45.9 %, respectively (Table 2). The relationship between grain yield and variable nitrogen fertilization was described by the equation of linear regression (Table 4). Independently of soil fertilization

with nitrogen, foliar fertilization in all its variants increased grain yield as compared to the control object (Table 2). Similarly, to experiments of Czuba et al. [4], parallel application of urea and of magnesium sulphate proved to be particularly advantageous. Application of the parallel fertilization with nitrogen and magnesium induced a moderately significant increase in wheat grain yield, as compared to objects fertilized exclusively with nitrogen or magnesium or to the control object, by 4.7%, 5.7% and 14.7 %, respectively (Table 2).

In determination of grain quality, attention is paid to its enzymatic properties and, in particular, to amylolytic activity, which is characterized by a falling number. Similarly as proved by Nowak et al. [15], in our experiment average values of falling number for the studied wheat pointed to low activity of alpha-amylase and they did not disqualify the studied grain from the point of view of its baking suitability (Table 5). According to Sulek et al. [20,22], value of the falling number depends more from genetic traits of the cultivar and weather conditions than from the applied in their experiment ways of nitrogen application. Independently of the foliar fertilization, the applied to soil nitrogen doses up to the level of  $90 \text{ kg}\cdot\text{ha}^{-1}$  have induced a significant increase in falling number, but only in comparison to the value obtained in the control object (Table 5). Application of the foliar fertilization has resulted in falling numbers at the level resembling those obtained in objects fertilized in parallel with nitrogen and magnesium or with magnesium alone. However, the trait value has proven to be significantly higher than the value obtained in the control object only in objects fertilized with magnesium (Table 5).

A number of studies indicate that an increased level of fertilization with nitrogen induces in the wheat grain an increased content of total protein [3,8,12,16,17,25,29], representing one of the most important factors determining its technological quality. Ralcewicz and Knapowski [18] noted a significant increase in content of the component following application of fertilization doses up to  $90 \text{ kg N}\cdot\text{ha}^{-1}$ , while Kłupczyński et al. [10] noted this also after application of  $120 \text{ kg N}\cdot\text{ha}^{-1}$ . Similarly to our experiment, a moderately significant increase in total protein content resulted from soil fertilization with any nitrogen doses up to the level of  $120 \text{ kg}\cdot\text{ha}^{-1}$  and, as compared to  $N_{90}$ ,  $N_{60}$  and  $N_0$  objects, it was higher by 12.2% , 19.1% and 25.2 %, respectively (Table 5). Results of studies of Wróbel [29] also have shown that content of the component is most favourably affected by the dose of  $120 \text{ kg N}\cdot\text{ha}^{-1}$  but applied on two occasions, i.e.  $60 \text{ kg}$  before sowing and  $60 \text{ kg}$  at the beginning of heading. The relationship between the applied soil fertilization with nitrogen and content of total protein

Table 1. Weather conditions in the vegetation seasons of 2005-2007  
Tabela 1. Warunki pogodowe w sezonach wegetacyjnych 2005-2007

Month Miesiąc	2005		2006		2007		Mean (średnia) 1995 -2005	
	Temper. (°C)	Rainfall Opady (mm)	Temper. (°C)	Rainfall Opady (mm)	Temper. (°C)	Rainfall Opady (mm)	Temper. (°C)	Rainfall Opady (mm)
IV	8.0	23.8	7.7	66.0	8.5	17.6	8.2	29.3
V	12.7	86.1	12.4	58.8	13.8	73.1	13.4	53.2
VI	14.8	30.2	17.1	22.7	18.2	105.5	16.2	48.4
VII	18.6	43.2	21.0	46.1	18.0	104.7	18.3	77.9
VIII	16.5	43.1	17.0	112.9	17.8	42.1	18.4	59.5
Mean; Sum Średnia; Suma	14.1	226.4	15.1	306.5	15.2	343.0	14.9	268.3

Table 2 Size of grain yield as related to the applied soil fertilization with nitrogen and variants of foliar fertilization [ $t \cdot ha^{-1}$ ]

Tabela 2. Wielkość plonu ziarna w zależności od zróżnicowanego nawożenia azotem i wariantów nawożenia nalistnego [ $t \cdot ha^{-1}$ ]

Nitrogen fertilization (I factor) Nawożenie azotem (I czynnik) [ $kg \cdot ha^{-1}$ ]	Foliar fertilization (II factor) Nawożenie dolistne (II czynnik)				mean średnia
	without fertilization bez nawożenia	urea mocznik	magnesium sulphate siarczan magnezu	urea+magnesium sulphate mocznik+ siarczan magnezu	
N <sub>0</sub>	2.58	2.86	2.83	3.33	2.90
N <sub>60</sub>	3.55	3.87	3.83	3.95	3.80
N <sub>90</sub>	3.93	4.35	4.27	4.38	4.23
N <sub>120</sub>	4.05	4.40	4.40	4.52	4.34
średnia	3.53	3.87	3.83	4.05	3.82
LSD - NIR <sub>(p=0,05)</sub> for-dla:	I 0.394	II 0.172	I x II n.s. - n.i.		

in wheat grain has been confirmed by the calculated coefficients of direct correlation (Table 3) and it has been described by the equation of linear regression (Table 4). Sztuder and Świerczewska [23] have provided the opinion that percentage content of total protein in spring wheat grain is determined not only by specific cultivar traits but also by the applied foliar fertilization. The most extensive effect of applied foliar fertilization on the protein content in wheat grain has been noted in the case of Sigma cultivar, for which protein content has ranged from  $106 g \cdot kg^{-1}$  in an object with no foliar fertilization to  $138 g \cdot kg^{-1}$  in the object treated three times with spray of urea plus magnesium sulphate and with Polvit Z/J. In studies of Sulek et al. [21] peak protein content has been noted in the grain obtained following application of

$90 kg N \cdot ha^{-1}$  in a dose divided in the following manner: 40 kg to the soil before sowing, 30 kg at the phase of shooting and 20 kg at the phase of heading, in the form of a solution with double spray aimed to prevent diseases. In present experiment, independently of soil fertilization, all the applied variants of foliar fertilization have induced a significant increase in total protein content, as compared to the control object (Table 5). The most favourable values of the discussed parameter have been obtained in the objects in which either nitrogen or magnesium have been applied in parallel or urea only has been applied and the content has been significantly higher than that in control object and the object fertilized with magnesium only, by 4.7% and 2.3 %, respectively (Table 5). Similarly to studies of other authors [18,28], a significant positive



Table 3. Significant indices of direct correlation between soil fertilization with nitrogen and studied traits of spring wheat Zebra cultivar

Tabela 3. Istotne współczynniki korelacji prostej pomiędzy zastosowanym doglebowo nawożeniem azotem a badanymi cechami pszenicy jarej odmiany Zebra

Parameter - Parametr	1	2	3	4	5	6	7
Fertilization–Nawożenie (1)	-	0.80	0.43	0.93	0.85	0.87	0.73
Grain yield–Plon ziarna (2)	0.80	-	0.36	0.85	0.91	0.82	0.43
Falling number–Liczba opadania (3)	0.43	0.35	-	0.51	0.42	0.38	n.s.n.i.
Protein content – Zawartość białka (4)	0.93	0.85	0.52	-	0.93	0.91	0.62
Gluten content–Zawartość glutenu (5)	0.85	0.91	0.43	0.93	-	0.91	0.56
Sedimentation value–Wskaźnik sedymentacji (6)	0.87	0.82	0.38	0.91	0.91	-	0.79
Water absorbtion – Wodochłonność mąki (7)	0.73	0.43	n.s.n.i.	0.62	0.56	0.79	-
Bread volume–Objętość pieczywa (8)	0.85	0.88	0.49	0.93	0.92	0.84	0.50

Tabela 4. Współczynniki regresji i determinacji (d) dla zależności między zastosowanym doglebowo nawożeniem azotem (x) a badanymi parametrami (y)

Table 4. Coefficients of regression and determination (d) as related to soil fertilization with nitrogen (x) and studied parameters (y)

Parametr Parameter	y = ax + b		
	a	b	d (%)
Grain yield - Plon ziarna	0.013	2.90	64.00
Falling number - Liczba opadania	0.185	357.02	18.49
Protein content - Zawartość białka	0.240	114.81	86.49
Gluten content - Zawartość glutenu	0.092	26.65	72.25
Sedimentation value -Wskaźnik sedymentacji	0.128	33.08	75.69
Water absorbtion - Wodochłonność mąki	0.045	54.19	53.29
Bread volume - Objętość pieczywa	0.595	438.32	72.25

correlation has been noted between total protein content in the grain, content of gluten and sedimentation value (Table 3); the relationships have been described by the calculated equations of linear regression (Fig.1).

The positive traits of baking products obtained from wheat flour are determined by the amount and quality of the gluten contained in the flour. According to several authors [3,8,16,17], an increase in gluten content results from intensified fertilization with nitrogen, although the procedure can also deteriorate quality of gluten [27]. In our experiment soil fertilization with nitrogen up to the dose of 90 kg·ha<sup>-1</sup> has induced a significant increase in content of the component, which has been higher than that obtained in N<sub>60</sub> and N<sub>0</sub> objects by 10% and 32.8 %, respectively (Table 5). Independently of the applied soil fertilization with nitrogen doses, a similar gluten content has been disclosed in objects fertilized in parallel with urea and magnesium sulphate or with urea alone and the content has proven to be significantly higher both as compared to the object fertilized with magnesium only (magnesium sulphate) and the control object. Evident increase in gluten content under effect of foliar fertilization of spring wheat was disclosed by

Sztuder and Świerczewska [23] and, as compared to the control, it was most pronounced (higher by 17.6 %) in the case of Sigma cultivar, following parallel application of solution containing urea, magnesium sulphate and Polvit Z/J. The proven in earlier studies [18,28] high correlation between gluten content and sedimentation value causes that gluten content used to be defined by the index. The relationship between sedimentation value and wet gluten content is illustrated in Fig.2. According to criteria used in international wheat trade [19], the raw material of a good baking value in the aspect of its usefulness for bakery is provided in grain of the studied spring wheat Zebra cultivar, obtained in objects in which no soil fertilization with nitrogen but foliar fertilization with urea alone or with urea plus magnesium sulphate was applied in parallel as well as individual variants of foliar and soil fertilization with nitrogen were applied at the level of 60 and 90 kg N·ha<sup>-1</sup>. On the other hand, a raw material of a very high baking quality involves grain obtained from objects of foliar fertilization with urea and magnesium sulphate in parallel following soil fertilization with the dose of 90 kg N·ha<sup>-1</sup> and objects in which all variants of foliar and soil fertilization

Table 5. Values of technological parameters for spring wheat Zebra cultivar as related to the applied soil fertilization with nitrogen and variants of foliar fertilization (average values for 2005-2007)  
 Tabela 5. Wartości parametrów technologicznych pszenicy jarej odmiany Zebra w zależności od nawożenia azotem i wariantów nawożenia dolistnego (średnie z lat 2005-2007)

Parameter Parametr	N fertilization (I factor) Nawożenie N (I czynnik) [kg·ha <sup>-1</sup> ]	Foliar fertilization (II factor) Nawożenie nalistne (II czynnik)				Mean Średnio	LSD <sub>p=0,05</sub> - NIR <sub>p=0,05</sub>			
		without fertilization bez nawożenia	urea mocznik	magnesium sulphate siarczan magnezu	urea+magnesium sulphate mocznik+ siarczan magnezu		I	II	I x II	II x I
Falling number Liczba opadania [s]	0 60 90 120	448 462 473 463	453 466 470 477	458 476 481 486	462 475 484 475	455 470 477 476	21.3	12.2	ns ni	ns ni
Mean - Średnio		462	467	475	474	469				
Protein content Zawart. białka [g·kg <sup>-1</sup> ]	0 60 90 120	112 123 134 140	118 132 138 146	112 128 137 144	116 132 140 145	115 129 137 144	5.6	2.6	ns ni	ns ni
Mean - Średnio		127	133	130	133	131				
Gluten content Zawart. glutenu [%]	0 60 90 120	24.4 28.6 33.6 36.0	28.2 34.0 35.8 38.7	25.0 30.7 34.3 37.1	28.5 34.5 36.9 38.7	26.5 32.0 35.2 37.6	2.83	1.19	ns ni	ns ni
Mean - Średnio		30.7	34.2	31.8	34.7	32.8				
Sediment. value W. sedymentacji [cm <sup>3</sup> ]	0 60 90 120	30.7 35.9 43.2 44.1	35.2 44.3 48.4 50.2	31.2 38.7 44.0 46.0	34.4 43.4 47.6 49.7	32.9 40.6 45.8 47.5	5.04	1.72	ns ni	ns ni
Mean - Średnio		38.5	44.5	40.0	43.8	41.7				
Water absorbtion Wodochł. mąki [%]	0 60 90 120	53.2 56.3 57.7 58.6	54.3 58.0 58.9 59.5	53.2 57.1 58.4 59.7	54.8 57.9 59.0 59.2	53.9 57.3 58.5 59.3	2.30	0.44	0.87	0.84
Mean - Średnio		56.5	57.7	57.1	57.7	57.2				
Bread volume Objęt. pieczywa [cm <sup>3</sup> ]	0 60 90 120	418 464 493 500	445 489 502 513	431 466 485 499	451 488 501 510	434 477 496 506	14.1	7.3	ns ni	ns ni
Mean - Średnio		469	487	470	488	479				

were applied at the level of 120 kg N·ha<sup>-1</sup> (Table 5). In our experiment, similarly to the earlier conducted experiments with spring wheat [10,17], soil fertilization with increasing doses of nitrogen have induced increase in the sedimentation value. Nevertheless, a significant increase in the qualitative index has been induced by application of nitrogen doses up to the level of 90 kg·ha<sup>-1</sup> and, as compared to N<sub>60</sub> and N<sub>0</sub> objects, it has amounted to 12.8% and 39.2 %, respectively (Table 5). Similarly, to gluten content, the value of sedimentation value has significantly positively correlated with the nitrogen dose applied to the soil (Table 3). The relationships have been described by linear regression equations (Table 4). In objects fertilized with nitrogen and magnesium in parallel and in the object fertilized with nitrogen alone similar values of sedimentation test have been noted and they, on the average, have proven to be moderately significantly higher than values of the index obtained in the object fertilized with magnesium only or the control

object. Sulek et al. [21] have indicated that improved value of the sedimentation value has resulted from application of nitrogen fertilizers in three doses, i.e., before sowing in a solid form and in phases of shooting and heading in a liquid form. The calculated coefficients of direct correlation have indicated a significant positive relationship between sedimentation value and contents of not only protein and gluten, as confirmed by the earlier publications [18,28], but also, i.e., water absorption of the flour (Table 3).

The amount and quality of gluten and the extent of damage to starch exert a significant effect of water absorption of flour [2, 6]. Similarly to studies of Podleśna and Cacak [16], in our experiment increase in nitrogen doses applied to the soil has resulted in augmented water absorption of the flour. Nevertheless, the statistically proven maximum increase in value of the qualitative trait has been noted after application of 60 kg·ha<sup>-1</sup>, and the absorption has




Fig. 1. Relationships between the total protein content and the content of wet gluten and the sedimentation value of spring wheat Zebra cultivar

Rys.1 Zależności pomiędzy zawartością białka ogólnego a zawartością mokrego glutenu, wskaźnika sedymentacji i objętością pieczywa pszenicy jarej odmiany Zebra




Fig. 2. Relationships between the content of wet gluten and the sedimentation value and the bread volume of spring wheat Zebra cultivar

Rys.2 Zależności pomiędzy zawartością mokrego glutenu a wskaźnika sedymentacji i objętością pieczywa pszenicy jarej odmiany Zebra




Fig. 3. Relationships between the bread volume and the total protein content and sedimentation value of spring wheat Zebra cultivar

Rys.3 Zależności pomiędzy zawartością białka ogólnego i wskaźnika sedymentacji a objętością pieczywa pszenicy jarej odmiany Zebra

been higher by 6.3% than that noted in the control object (Table 5). The applied foliar fertilization variants, independently from soil fertilization have induced a significant increase in water absorption by the flour, as compared to the control object (Table 5). Similarly, in studies of Żmijewski et al. [30] water absorption proved to be significantly correlated with, i.e., protein content (Table 3).

A direct qualitative index, which indicates baking value of wheat grain, involves volume of bread following test baking. Similarly to other studies on spring wheat [10,16,17,18,20], in our experiment the applied to soil increasing doses of nitrogen fertilization on the average have induced increase in bread volume obtained from flour of Zebra cultivar, and a significant increase in the trait has been noted following the dose of up to 90 kg N · ha<sup>-1</sup>. The bread volume obtained in the N<sub>90</sub> object has been higher than the values obtained in objects of N<sub>60</sub> and N<sub>0</sub> by 4% and 14.3 %, respectively (Table 5). According to Sulek et al. [21], a more extensive effect on bread volume in exerted by cultivar properties than by the way of nitrogen fertilization. In our experiment, independently of the applied soil fertilization with nitrogen, bread volumes obtained from objects fertilized with nitrogen and magnesium or with nitrogen only have been significantly higher both as compared to

the control, unfertilized object and as compared to the object fertilized with magnesium only. Bread volume has significantly and positively correlated with, i.e., value of sedimentation value (Table 3), which confirms the view that higher content of high molecular weight glutenin increases, i.e., the bread volume [24]. In turn, Żmijewski et al. [30] have confirmed the significant positive correlation between bread volume and protein content while for the relationship between the discussed trait and sedimentation value they have noted a negative correlation (Table 3). The relationship between bread volume on one hand and total protein content, gluten content and sedimentation value are described by the calculated equations of linear regression (Fig. 2 and 3).

## CONCLUSIONS

1. Soil fertilization with nitrogen up to the dose of 90 kg · ha<sup>-1</sup> was followed by a moderately significant increase in size of grain yield while among the variants of foliar fertilization the moderately significant peak grain yield was noted in objects in which nitrogen and magnesium were applied in parallel.
2. The applied soil fertilization with nitrogen, within the entire range of applied doses, induced a moderately significant increase in total protein content. The



moderately significant increase in total protein content, as compared to the control object, resulted also from any variant of foliar fertilization.

3. Independently of the applied variants of foliar fertilization, significant increases in wet gluten content, sedimentation value and bread volume were detected following soil fertilization with nitrogen doses up to 90 kg·ha<sup>-1</sup>, and of water absorption by the flour following fertilization up to 60 kg·ha<sup>-1</sup>.

4. Independently of the applied soil fertilization with nitrogen, among the studied variants of foliar fertilization the most favourable for principal indices of baking value, such as wet gluten content, sedimentation value and bread volume, proved to be parallel application of nitrogen and magnesium or of nitrogen only.

## REFERENCES

- [1] Biskupski A., Kaus A., Pabin J., Włodek St., The influence of differentiated with nitrogen on leaf area index (LAI), mean tip angle (MTA) and yield of crop in selected cultivars of spring wheat, *Annales UMCS* (2004) E, 59, 2: 649-654.
- [2] Biskupski A., Kaus A., Włodek St., Pabian J., Influence of different level of nitrogen fertilization on yielding and selected indices of canopy architecture in several spring wheat cultivars, *Pam. Puł.* (2006) 142: 31-41.
- [3] Borkowska H., Grundas S., Styk B., Wysokość i jakość plonów niektórych odmian pszenicy jarej w zależności od nawożenia azotowego, *Annales UMCS* (2002) E, 57: 99-103.
- [4] Czuba R., Sztuder H., Świerczewska M., The results of leaf nutrition of field drops. P. III. Response of plants to foliar application of magnesium or magnesium combined with nitrogen. *Roczn. Glebozn.* (1999) L, 1/2: 41-50.
- [5] Gašiorowska B., Makarewicz A., Nowosielska A., Rymuza K., Efficiency of nitrogen fertilization of different varieties of spring wheat, *Pam. Puł.* (2006) 142: 117-125.
- [6] Gooding M. J., Smith G.P., The potential to use climate, variety and nitrogen relationships to optimise wheat quality, In: *Short Communications, Fifth ESA Congress, 28 June - 2 July, (1998): 229-230.*
- [7] Guarda G., Padovano S., Delogu G., Grain field, nitrogen-use efficiency and baking quality of old and modern Italia bread-wheat cultivars grown at different nitrogen levels, *Eur. J. Agron.* (2004) 21: 181-192.
- [8] Johansson E., Prieto-Linde M.L., Jonsson J.O., Effects of wheat cultivar and nitrogen application on storage protein composition and breadmaking quality, *Cer. Chem.* (2001) 78: 19-25.
- [9] Johansson E., Svensson G., Influence of yearly variation and fertilizer rate on bread-making quality in Swedish grown wheats containing HMW glutenin subunits 2+12 or 5+10 cultivated during the period 1990-1996, *J. Agric. Sci. Camb.* (1999) 132: 13-22.
- [10] Klupczyński Z., Ralcewicz M., Knapowski T., Murawska B., The try of optimal nitrogen fertilization level determination on the backing value of bread varieties of spring wheat, *Mat. Konf. "Konkurencyjność rolnictwa z uwzględnieniem uwarunkowań regionalnych w aspekcie integracji z Unią Europejską", AR Krakow* (2000) 2: 553-558.
- [11] Knapowski T., Ralcewicz M., Evaluation of qualitative features of Mikon cultivar winter wheat grain and flour depending on selected agronomic factors, *EJPAU, Agronomy* (2004) 7: 1.
- [12] Knapowski T., Ralcewicz M., The effect of selected agrotechnical factors on the protein amino acid composition in the spring wheat grain, *Makro and Trace Elements, Mengen- und Spurenelemente* (2006) 23: 409-414.
- [13] Kocoń A., Fertilization of quality spring and winter wheat and its impact on yield and grain quality, *Pam. Puł.* (2005) 139: 55-63.
- [14] Michałojć Z., Szewczuk C., Teoretical aspects of foliar nutrition, *Acta Agrophysica* (2003) 85: 9-17.
- [15] Nowak W., Zbroszczyk T., Kotwicz L., Effect of management level on some quality traits of wheat cultivars, *Pam. Puł.* (2004) 135: 199-211.
- [16] Podleśna A., Cacak-Pietrzak G., Formation of spring wheat yield as well as its milling and baking parameters by nitrogen and sulphur fertilization, *Pam. Puł.* (2006) 142: 381-392.
- [17] Ralcewicz M., Knapowski T., Klupczyński Z., The baking value of the spring wheat studiem as function of diversified nitrogen fertilization, *Makro and Trace Elements, Mengen- und Spurenelemente* (2002) 21: 284-289.
- [18] Ralcewicz M., Knapowski T., The effect of diverse nitrogen fertilization on the height of grain yield and the technological value of spring wheat, *Annales UMCS* (2004) E, 59, 2: 969-978.
- [19] Rothkaehl J., Abramczyk D., Wartość technologiczna odmian pszenicy uprawianych w Polsce, *Przegl. Zboż.-Młyn.* (2007) 7, 2-4.
- [20] Sulek A., Cacak-Pietrzak G., Ceglińska A., Effectiveness of different methods of nitrogen application in spring wheat cultivated for baking purpose, *Pam. Puł.*

(2006) 142: 505-512.

[21] Sułek A., Cacak-Pietrzak G., Ceglińska A., Haber T., The effect of differentiated nitrogen fertilization on technological value of selected cultivars of spring wheat, *Pam. Puł.* (2002) 130: 709-718.

[22] Sułek A., Cacak-Pietrzak G., Ceglińska A., The effect of different methods of nitrogen application on yielding, yield components structure and some quality characters of spring wheat cultivars, *Annales UMCS* (2004) E, 59, 2: 543-551.

[23] Sztuder H., Świerczewska M., Wpływ nawozów dolistnych na cechy jakościowe ziarna niektórych odmian pszenicy ozimej i jarej, *Zesz. Prob. Post. Nauk Roln.* (2002) 484: 669-674.

[24] Verbruggen I.M., Veraverbeke W. S., Delcour J. A., Significance of LMW-GS and HMW-GS for dough extensibility: „addition“ versus „incorporation“ protocols, *J. Cer. Scien.* (2001) 33: 253-260.

[25] Waclawowicz R., Parylak D., Śniady R., Residual effect of organic and nitrogen fertilization on

the yields and selected quality features of spring wheat grain, *Pam. Puł.* (2005) 139: 277-288.

[26] Wojciechowski W., Nitrogen fertilization response of spring wheat Torca grown after ploughed down stubble crops, *Biul. IHAR* (2005) 237/238: 23-37.

[27] Wooding A.R., Kavale S., Macritchie F., Stoddard F.L., Wallace A., Effects of nitrogen and sulfur fertilizer on protein composition, mixing requirements, and dough strength of four wheat cultivars, *Cer. Chem.* (2000) 77: 798-807.

[28] Woźniak A., The influence of preceding crops on the quality of spring wheat grain, *Pam. Puł.* (2004) 135: 325-330.

[29] Wróbel E., Response of spring wheat on rate and date of nitrogen application, *Pam. Puł.* (1999) 118: 447-453.

[30] Żmijewski M., Subda H., Kowalska M., Korczak B., Czubaszek A., Karolini-Skaradzińska Z., Chemical composition and baking value of grain and flour of varieties of spring wheat, P. II. Baking value, *Biul. IHAR* (1999) 212: 71-79.