

## Original scientific article

# Relationship between adiponectin and testosterone in patients with type 2 diabetes

Jie Bai<sup>1</sup>, Yu Liu<sup>1</sup>, Gui-Fen Niu<sup>1</sup>, Li-Xin Bai<sup>1</sup>, Xiao-Yan Xu<sup>2</sup>, Guang-Zhen Zhang<sup>2</sup>, Le-Xin Wang<sup>3\*</sup>

<sup>1</sup> Department of Endocrinology, Liaocheng People's Hospital and Liaocheng Clinical School of Taishan Medical University, Shandong, PR China

<sup>2</sup> Department of Pharmacy, Liaocheng People's Hospital and Liaocheng Clinical School of Taishan Medical University, Shandong, PR China

<sup>3</sup> School of Biomedical Sciences, Charles Sturt University, Wagga Wagga, NSW, Australia

\*Corresponding author: [lwang@csu.edu.au](mailto:lwang@csu.edu.au)

### Abstract

**Introduction:** This study was designed to investigate the relationship between serum adiponectin and testosterone in patients with type 2 diabetes.

**Materials and methods:** Serum level of adiponectin and testosterone were prospectively measured in 65 patients with type 2 diabetes and in 20 healthy subjects. Testosterone was determined by the radio-immunoassay whereas adiponectin levels were determined by enzyme-linked immunosorbent assay (ELISA).

**Results:** The average serum testosterone did not differ between the diabetes and the control group, but the average adiponectin in the diabetes group was lower (14.6 (14.2-15.0) vs. 24.3 (24.05-24.55) ng/mL,  $P = 0.001$ ). In the diabetes group, the serum adiponectin level in patients with renal dysfunction (22.3 (21.5-23.1) ng/mL) was higher than in patients with no complications (12.1 (11.45-12.75) ng/mL) and than in patients with coronary artery disease (11.2 (10.25-12.15) ng/mL) ( $P = 0.009$ ). Univariate correlation analysis showed an inverse weak correlation between adiponectin and testosterone concentrations in male diabetic patients ( $r = -0.27$ ,  $P = 0.009$ ). There was no significant correlation between adiponectin and testosterone in female patients ( $r = -0.05$ ,  $P = 0.167$ ).

**Conclusions:** We conclude that patients with type 2 diabetes have lower serum adiponectin concentration than healthy individuals, and that there is a weak inverse correlation between adiponectin and testosterone serum concentrations in male diabetics.

**Key words:** type 2 diabetes; adiponectin; testosterone; renal function; coronary artery disease

Received: August 12, 2010

Accepted: December 6, 2010

### Introduction

Adiponectin is a protein exclusively secreted by mature adipocytes, which circulates in high concentrations in the blood (1,2). The circulating concentrations of adiponectin are inversely proportional to adiposity, and low adiponectin concentrations predict the development of type 2 diabetes and cardiovascular disease (2). Serum adiponectin levels are influenced by several factors, such as age, sex, body mass index and diabetes (2-5) and are decreased in patients with type 2 diabetes mellitus or metabolic syndrome (5,6). In obese men, serum and interstitial total adiponectin levels are lower than in lean subjects (7). After hyperinsuline-

mia, serum and interstitial total adiponectin is reduced in both obese and lean men. Current biochemical and physiological evidence suggest that low adiponectin levels may be a consequence of insulin resistance with compensatory hyperinsulinemia (1).

Testosterone is another factor that influences concentration of circulating adiponectin. In healthy subjects, adiponectin production is inhibited by testosterone (2). However, whether testosterone is also associated with circulating adiponectin in patients with diabetes is not very well understood.

In school-aged girls with type 1 diabetes, the decrease in adiponectin concentrations was correlated with increasing levels of testosterone (8). In middle-aged patients with metabolic syndrome, blood concentrations of testosterone were positively correlated with adiponectin (9). In addition, testosterone therapy in combination with diet and exercise in type 2 diabetic men resulted in an increase of adiponectin (10). To date, there is little information about the association between circulating testosterone and adiponectin in adult diabetic patients. Therefore, we performed this prospective study to investigate the relationship between plasma testosterone and adiponectin in a group of adult patients with type 2 diabetes.

## Materials and methods

### Patient selection

This study was approved by the institution review board of Liaocheng People's Hospital. Written informed consent was obtained from all participants. Between January and July 2008, 65 patients with type 2 diabetes were recruited to this study. There were 39 males and 26 females, with an average age of 58 years (range 37-84 years). Twenty one (32.3%) of these patients had concurrent coronary artery disease (CAD), and 22 (33.8%) had renal dysfunction. CAD was confirmed by coronary angiogram and electrocardiogram (ECG) as well as clinical symptoms. Renal dysfunction was established by assessing glomerular filtration rate (GFR < 90 mL/min), which was calculated from serum creatinine (Cr) concentration as follows:

$$\text{Cockcroft-Gault GFR} = (140 - \text{age}) \times (\text{Weight in kg}) \times (0.85 \text{ if female}) / (72 \times \text{Cr}).$$

Twenty healthy subjects (Table 1) were also recruited as the control group. These subjects were recruited from the health-check clinics of our hospital. They had no previous medical history of diabetes, cardiovascular or renal disease. Physical examination, chest X-ray, electrocardiogram and laboratory tests for blood glucose, lipids, renal and liver function revealed no abnormalities.

### Biochemical analyses

Five milliliters of blood was collected from the median cubital vein in a sitting position with a vacuum tube by the investigators in the mornings between 7-8 am, after overnight fasting. Serum was stored in a freezer at -20°C for up to four weeks until testosterone and adiponectin were analyzed. Biochemical data were assayed as follows: fasting blood glucose (glucose-oxidase electrode method); total cholesterol (cholesterol oxidase enzymatic assay method); triglyceride (enzymatic colorimetric method); serum creatinine (Jaffe reaction method). The above analyses were conducted by using a fully automated Olympus AU2700 chemistry analyzer (Olympus CO Ltd., Tokyo, Japan). Olympus Life and Material Science Europa GmbH reagents (Hamburg, Germany) were used.

Serum testosterone was determined by the radioimmunoassay (Northern Biotech Inc, Beijing, China). Our laboratory intra- and inter-assay coefficients of variation were less than 6.0% based on 10 measurements using control samples at level of 2.8 ng/L. Total serum adiponectin levels were determined by an enzyme-linked immunosorbent assay kit (ELISA) (Shiruike Inc, Shanghai, China) as according to the manufacturer's instructions. The sensitivity of the assay was calculated to be 1 ng/mL. Our laboratory intra- and inter-assay coefficients of variation were less than 8% based on 10 measurements using control samples at level of 24.5 ng/L.

### Statistical analysis

Data are presented as median and interquartile range (Q1-Q3), except age which was presented as median and range (maximum-minimum). Comparison of variables between groups was performed by Mann-Whitney test. Kruskal-Wallis one-way analysis of variance was used for three group comparisons. *Post hoc* analysis was performed by Dunn's post test. Correlation analysis was done by Spearman's rank correlation. SPSS software (version 11.5) was used for the statistical analysis. P values less than 0.05 were considered statistically significant.

## Results

As shown in Table 1, there were no significant differences in the age and sex between the study and control groups. The mean testosterone concentration did not differ significantly between two groups. However, patients with type 2 diabetes had significantly lower adiponectin concentration than patients in the control group (14.6 (14.2-15.0) vs. 24.3 (24.05-24.55) ng/mL, P = 0.001).

Levels of adiponectin and testosterone according to gender in diabetes and control patient groups are presented in table 2. In both, patient and control group, serum adiponectin concentration was lower in males (P = 0.019 and 0.026, respectively), whereas the average testosterone concentration was higher in males (P = 0.012 and 0.009, respectively) than in females. Adiponectin concentration

in males and females of the diabetes group was lower than in males and females of the control group (P = 0.008 and 0.009, respectively).

Table 3 shows adiponectin and testosterone levels in patients with and without CAD or renal dysfunction. There were no significant differences in age, gender, blood glucose, BMI or serum testosterone concentration across three groups. However, the serum adiponectin level in the renal dysfunction group differed from the other groups (P = 0.016). *Post hoc* analysis showed that the median level of adiponectin in the renal dysfunction group was higher than the CAD and non-complication group (P = 0.009).

Univariate correlation analysis showed an inverse weak correlation between adiponectin and testos-

**TABLE 1.** Baseline characteristics and serum levels of adiponectin and testosterone of the study and control group.

Indices	Diabetes group (N = 65)	Control group (N = 20)	P
Age (years)	56 (37-84)	55 (37-82)	0.946
Male (N, proportion)	39 (0.60)	11 (0.55)	0.720
BMI (kg/m <sup>2</sup> )	24.6 (24.0-25.2)	23.6 (22.95-24.25)	0.644
Fasting blood glucose (mmol/L)	12.7 (12.2-13.2)	5.0 (4.8-5.2)	<b>0.001</b>
Adiponectin (ng/mL)	14.6 (14.2-15.0)	24.3 (24.05-24.55)	<b>0.001</b>
Testosterone (ng/mL)	2.7 (2.4-3.0)	2.8 (2.6-3.0)	0.669

**TABLE 2.** Serum levels of adiponectin and testosterone of the diabetes and control group.

	Diabetes group	Control group	P*
<b>Adiponectin (ng/mL)</b>			
Male	N = 39 10.3 (9.75-10.85)	N = 11 19.6 (18.85-20.35)	<b>0.006</b>
Female	N = 26 16.1 (15.5-17.7)	N = 9 26.2 (25.3-27.1)	<b>0.009</b>
p**	<b>0.019</b>	<b>0.016</b>	
<b>Testosterone (ng/mL)</b>			
Male	N = 39 4.3 (3.7-4.9)	N = 11 4.6 (3.95-5.25)	<b>0.522</b>
Female	N = 26 0.32 (0.285-0.355)	N = 9 0.36 (0.315-0.405)	<b>0.788</b>
p**	<b>0.012</b>	<b>0.009</b>	

\* Compared with the diabetes group. \*\* Compared with male in the same group.

**TABLE 3.** Comparison of adiponectin levels between diabetic patients with and without cardiovascular complications.

	<b>CAD (N = 21)</b>	<b>Renal dysfunction (N = 22)</b>	<b>No complications (N = 22)</b>	<b>P</b>
<b>Age (years)</b>	53 (42-67)	52 (39-66)	54 (33-64)	0.868
<b>Male (N, proportion)</b>	12 (0.57)	13 (0.59)	14 (0.64)	0.462
<b>FBG (mmol/L)</b>	12.6 (11.9-13.3)	12.5 (11.65-13.35)	12.2 (11.45-12.95)	0.742
<b>HbA1c (%)</b>	9.1 (8.8-9.4)	9.3 (8.95-9.65)	9.0 (8.55-9.45)	0.866
<b>BMI (kg/m<sup>2</sup>)</b>	24.4 (23.5-25.3)	24.0 (23.05-24.95)	24.6 (23.8-25.4)	0.894
<b>Adiponectin (ng/mL)</b>	11.2 (10.25-12.15)	22.3 (21.5-23.1)	12.1 (11.45-12.75)	<b>0.010</b>
<b>Testosterone (ng/mL)</b>	3.2 (2.8-3.6)	2.6 (2.2-3.0)	2.8 (2.5-3.1)	0.098

CAD - coronary artery disease; FBG - fasting blood glucose

terone levels in male diabetic patients ( $r = -0.27$ ,  $P = 0.009$ ). There was no significant correlation between adiponectin and testosterone in female patients ( $r = -0.05$ ,  $P = 0.167$ ). Also, no significant correlation between serum adiponectin and testosterone was found in the healthy subjects, when male ( $r = -0.06$ ,  $P = 0.412$ ) and female ( $r = -0.09$ ,  $P = 0.225$ ) subjects were analyzed either separately or as a unique group ( $r = -0.06$ ,  $P = 0.165$ ). An inverse weak correlation between BMI and adiponectin was also found in the diabetes group ( $r = -0.33$ ,  $P = 0.02$ ).

## Discussion

The major findings of this study are: 1) serum adiponectin concentration in patients with type 2 diabetes was significantly higher than in controls; 2) serum level of adiponectin in patients with renal dysfunction was higher than in those with normal renal function; 3) there was an inverse weak correlation between serum adiponectin level and BMI or serum testosterone.

The results of our study were consistent with the previous studies where a lower serum adiponectin in diabetics than in healthy subjects was revealed (6,8). A recent population study showed that high serum adiponectin levels were associated with a lower incidence of type 2 diabetes (11). This relationship was more significant in female subjects than in males. The authors of this study concluded that the baseline adiponectin levels play a protec-

tive role in the pathogenesis of diabetes and are independent predictors for diabetes. The level of adiponectin seems to be closely associated with gender of the healthy subjects. The median serum levels of adiponectin in women was found to be 50% higher than in men (12). In addition, serum levels of adiponectin are also related to age. Older age is associated with a higher adiponectin level in both men and women (12). In our study we also found that in the healthy subjects the adiponectin level in women was higher than in men. Perhaps more importantly, the serum adiponectin levels in male diabetic patients were lower than in the female patients. These results suggest that adiponectin as a cardiovascular protector has diminished more in male diabetic patients.

The lower level of adiponectin in male diabetic patients may be due to a number of factors but a higher concentration of serum testosterone is likely one of them. An inverse weak correlation was found between adiponectin and testosterone concentration in the male patients. Iniguez and colleagues (8) recently measured the serum adiponectin and testosterone in 56 pubertal girls with type 1 diabetes. They found that decreasing adiponectin levels observed in these adolescents with type 1 diabetes correlate with increasing testosterone levels. However, the mechanisms by which testosterone affected the adiponectin levels were unclear (8).

From numerous studies it was suggested that adiponectin constitutes an important protective factor in the pathogenesis of the metabolic syndrome

and cardiovascular disease by exerting antidiabetic and antiatherogenic actions (13). Diminished serum adiponectin concentrations, on the other hand, are predictive for the development of diabetes and cardiovascular disease (14). Testosterone also plays a role in the pathogenesis of type 2 diabetes. Serum testosterone was reduced in male patients with type 2 diabetes, whereas it was increased in the female patients (15). It was believed that testosterone reduces the risk of type 2 diabetes in men but increases the risk in women (15). Given the antidiabetic and antiatherogenic actions of adiponectin and the potential antidiabetic actions of testosterone in men, reduced adiponectin and testosterone concentrations in males may at least in part account for the higher incidence of cardiovascular disease in men due to a diminished protective effect of these endogenous hormones.

In this study, the serum adiponectin level in patients with CAD did not differ from those without CAD (Table 3). This may be due to the fact that the pathogenesis of CAD in our patients may be multifactorial, involving other risk factors, such as blood plasma lipoproteins, high blood pressure etc., which were not explored in detail in this study. The adiponectin level in patients with renal dysfunction, however, was significantly higher than in those without renal dysfunction. Previous studies have suggested that a decrease in adiponectin

clearance in the kidney may increase levels of adiponectin, and adiponectin seems to be influenced more strongly by renal function than by sex hormones (12). In diabetic patients, the adiponectin levels were increased and were closely associated with the glomerular filtration rate, or with the stages of renal insufficiency (16-18). Although correlation analysis between adiponectin and renal function was not performed in our study, the reduced renal clearance was probably the main cause of increased serum adiponectin.

A potential limitation of this study is that the number of patients (N = 65) is relatively small. Whether a larger patient population would yield a stronger correlation between plasma testosterone and adiponectin in male and female patients remains to be seen. In addition, majority of studies have shown that testosterone is significantly lower in men with type 2 diabetes than in women (19). In our study, the mean value of plasma testosterone in men was higher than in women. Whether this is due to the small number of patients needs further investigation.

In summary, this study has confirmed that the levels of serum adiponectin in adult patients with type 2 diabetes are lower than in the healthy subjects. An inverse correlation between serum adiponectin and testosterone has also been demonstrated in male diabetic patients.

## References

1. Cook JR, Semple RK. Hypoadiponectinemia--cause or consequence of human "insulin resistance"? *J Clin Endocrinol Metab* 2010;95:1544-54.
2. Swarbrick MM, Havel PJ. Physiological, pharmacological, and nutritional regulation of circulating adiponectin concentrations in humans. *Metab Syndrome Related Disorder* 2008;6:87-102.
3. Gavrilu A, Chan JL, Yiannakouris N, Kontogianni M, Miller LC, Orlova C, et al. Serum adiponectin levels are inversely associated with overall and central fat distribution but are not directly regulated by acute fasting or leptin administration in humans: cross-sectional and interventional studies. *J Clin Endocrinol Metab* 2003;88:4823-31.
4. Bottner A, Kratzsch J, Muller G, Kapellen TM, Bluher S, Keller E, et al. Gender differences of adiponectin levels develop during the progression of puberty and are related to serum androgen levels. *J Clin Endocrinol Metab* 2004;89:4053-61.
5. Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, et al. Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 2000;20:1595-9.
6. Kapoor D, Clarke S, Stanworth R, Channer KS, Jones TH. The effect of testosterone replacement therapy on adipocytokines and C-reactive protein in hypogonadal men with type 2 diabetes. *Eur J Endocrinol* 2007;156:595-602.



7. Murdolo G, Hammarstedt A, Schmelz M, Jansson PA, Smith U. Acute hyperinsulinemia differentially regulates interstitial and circulating adiponectin oligomeric pattern in lean and insulin-resistant, obese individuals. *J Clin Endocrinol Metab* 2009;94:4508-16.
8. Iniguez G, Torrealba MI, Avila A, Cassorla F, Codner E. Adiponectin serum levels and their relationships to androgen concentrations and ovarian volume during puberty in girls with type 1 diabetes mellitus. *Hormone Res* 2008;70:112-7.
9. Akishita M, Fukai S, Hashimoto M, Kameyama Y, Nomura K, Nakamura T, et al. Association of low testosterone with metabolic syndrome and its components in middle-aged Japanese men. *Hypertension Res* 2010;33:587-91.
10. Heufelder AE, Saad F, Bunck MC, Gooren L. Fifty-two-week treatment with diet and exercise plus transdermal testosterone reverses the metabolic syndrome and improves glycaemic control in men with newly diagnosed type 2 diabetes and subnormal plasma testosterone. *J Androl* 2009;30:726-33.
11. Snijder MB, Heine RJ, Seidell JC, Bouter LM, Stehouwer CD, Nijpels G, et al. Associations of adiponectin levels with incident impaired glucose metabolism and type 2 diabetes in older men and women: the Hoorn study. *Diabetes Care* 2006;29:2498-503.
12. Isobe T, Saitoh S, Takagis, Takeuchi H, Chiba Y, Katoh N, Shimamoto K. Influence of gender, age and renal function on plasma adiponectin level: the Tanno and Sobetsu Study. *Eur J Endocrinol* 2005;153:91-8.
13. Matsuzawa Y, Funahashi T, Kihara S, Shimomura I. Adiponectin and metabolic syndrome. *Arterioscler Thromb Vasc Biol* 2004;24:29-33.
14. Spranger J, Kroke A, Mohlig M, Bergmann MM, Ristow M, Boeing H, et al. Adiponectin and protection against type 2 diabetes mellitus. *Lancet* 2003;361:226-8.
15. Ding EL, Song Y, Malik VS, Liu S. Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2006;295:1288-99.
16. Schalkwijk CG, Chaturvedi N, Schram MT, Fuller JH, Stehouwer CD. EURODIAB Prospective Complications Study Group. Adiponectin is inversely associated with renal function in type 1 diabetic patients. *J Clin Endocrinol Metab* 2006;91:129-35.
17. Komaba H, Igaki N, Goto S, Yokota K, Doi H, Takemoto T, et al. Increased serum high-molecular-weight complex of adiponectin in type 2 diabetic patients with impaired renal function. *Am J Nephrol* 2006;26:476-82.
18. Saraheimo M, Forsblom C, Fagerudd J, Teppo AM, Pettersson-Fernholm K, Frystyk J, et al. Serum adiponectin is increased in type 1 diabetic patients with nephropathy: FinDiane study group. *Diabetes Care* 2005;28:1410-4.
19. Maric C, Sex, diabetes and the kidney. *Am J Physiol- Renal Physiol* 2009;296:F680-8.

## Odnos između koncentracija adiponektina i testosterona kod bolesnika sa šećernom bolesti tipa 2

### Sažetak

**Uvod:** Ovo je istraživanje načinjeno kako bi se istražio odnos između koncentracija adiponektina i testosterona u serumu kod bolesnika kod bolesnika sa šećernom bolesti tipa 2.

**Materijali i metode:** U ovom prospektivnom istraživanju mjerene su koncentracije adiponektina i testosterona kod 65 bolesnika oboljelih od šećerne bolesti tipa 2 i kod 20 zdravih ispitanika. Koncentracija testosterona određena je radiološkim imunotestom, dok je koncentracija adiponektina određena enzimskim imunotestom (engl. *enzyme-linked immunosorbent assay*, ELISA).

**Rezultati:** Koncentracija testosterona u serumu nije se razlikovala između skupina bolesnika s šećernom bolesti tipa 2 i zdravih ispitanika, no prosječna koncentracija adiponektina je u skupini dijabetičara bila niža (14,6 (14,2-15,0) vs. 24,3 (24,05-24,55) ng/mL,  $P = 0,001$ ) nego kod zdravih ispitanika. U skupini dijabetičara koncentracija adiponektina u serumu bila viša kod bolesnika s poremećajem bubrežne funkcije (22,3 (21,5-23,1) ng/mL) nego kod bolesnika koji nisu imali komplikacije (12,1 (11,45-12,75) ng/mL) i bolesnika s koronarnom arterijskom bolešću (11,2 (10,25-12,15) ng/mL) ( $P = 0,009$ ). Univarijantnom analizom korelacije utvrđena je slaba obrnuta korelacija između koncentracije adiponektina i testosterona kod muškaraca dijabetičara ( $r = -0,27$ ,  $P = 0,009$ ). Nije bilo značajne korelacije ( $r = -0,05$ ,  $P = 0,167$ ) između koncentracije testosterona i adiponektina kod žena sa šećernom bolesti.

**Zaključak:** Zaključujemo da su kod skupine bolesnika sa šećernom bolesti tipa 2 koncentracije adiponektina i testosterona u serumu niže nego kod kontrolne skupine te da postoji slaba obrnuta korelacija između koncentracija adiponektina i testosterona kod muškaraca sa šećernom bolesti.

**Cljučne riječi:** šećerna bolest; adiponektin; testosteron; bubrežna funkcija; koronarna arterijska bolest