

Temperature-dose relationships with aflatoxin M₁ in milk on the brine shrimp (*Artemia salina*) larvae

Lejla Duraković^{1*}, Mihaela Blažinkov², Andrea Skelin²,
Sanja Sikora², Frane Delaš¹, Mirna Mrkonjić-Fuka²,
Katarina Huić-Babić², Sulejman Redžepović²

¹Faculty of Food Technology and Biotechnology, Department of Biochemical Engineering, Laboratory for General and Food Microbiology, University of Zagreb, Pierottijeva 6, Zagreb, Croatia

²Faculty of Agriculture, Department of Microbiology, University of Zagreb, Svetošimunska 25, Zagreb, Croatia

Received - Prispjelo: 21.02.2011.
Accepted - Prihvaćeno: 04.05.2011.

Summary

Temperature-dose relationships with aflatoxin M₁ (AFM₁) were studied using the brine shrimp *Artemia salina* larvae as a biological indicator in the temperature range from 20 °C to 40 °C. Increase in the incubation temperature resulted in sensitivity increase by the brine shrimp to AFM₁. Optimum sensitivity occurred at 30 °C. Positive results were obtained at 0.18 µg AFM₁ x L⁻¹ of whole pasteurized milk with a mortality of over 15%. Greater than 90 % mortality occurred at dose levels of 0.9 µg AFM₁ x L⁻¹ and above. The test can be conducted during 30-60 hours.

Key words: aflatoxin M₁, bioassay, *Artemia salina*, LC₅₀ and T₅₀ doses

Introduction

It is known that some moulds produce various toxic metabolites under appropriate temperature and moisture conditions. These metabolites, called mycotoxins, may be hazardous for human health (van Egmond, 1989, 1991; Wood, 1991; Duraković and Duraković, 2003).

The discovery of aflatoxin in 1960 dramatically stimulated interest in mycotoxins and mycotoxicoses and clearly revealed the real and potential danger of toxic fungal metabolites to men and animals (van Egmond, 1989; Wood, 1991; IARC, 1993a; Bhat and Vasanthi, 1999).

Exposure to mycotoxins through food is widely recognized as a human health hazard (Bhat and Vasanthi, 1999; Duraković and Duraković, 1999, 2003; Duraković et al., 2008). Of all mycotoxins, aflatoxin B₁ (AFB₁) is considered to be the most tox-

ic/carcinogenic compound (Škrinjar et al., 1992; IARC, 1993b; Škrinjar et al., 1995; Duraković et al., 2008). It is biotransformed by hepatic microsomal cytochrome P₄₅₀ to AFM₁.

Milk is usually contaminated with small amounts of AFM₁ as a consequence of the AFB₁ metabolism by animals that are fed with AFB₁ contaminated feed (van Egmond, 1991; Škrinjar et al., 1992, 1995; Taveira and Midio, 2001; Tratnik et al., 2001). The forming of AFM₁ occurs in liver and is secreted into milk in mammary gland (Stoloff, 1980; van Egmond, 1989). Like AFB₁, AFM₁ is toxic and carcinogenic, although toxicity of AFM₁ is slightly lower than of AFB₁ (Trucksess, 1999). The toxicological concern with AFM₁ arises in principle from its close structure similarity to AFB₁ (Figure 1), which has been shown to be of the most potent carcinogens (van Egmond and Wagstaffe, 1987; Duraković, 2007).

*Corresponding author/Dopisni autor: Phone/Tel.: +385 1 4605 045; E-mail: Lejla.Durakovic@pbf.hr

Table 1. Current and proposed legal limits for aflatoxin M₁ in milk and milk products in some European countries* (van Egmond and Wagstaffe, 1987; Creppy, 2002)

Country	Milk ($\mu\text{g} \times \text{L}^{-1}$)	Other milk products ($\mu\text{g} \times \text{L}^{-1}/\text{kg}^{-1}$)
Belgium	0.05 (V)	0.10 (P) milk for infant foods
Bulgaria	0.50 (V)	0.10 (P) milk powder for infant foods
Croatia	0.05 (V)	0.025 (P) milk for infant foods
Czech Republic	0.50 (V)	0.10 (P) milk powder for infant foods
France	0.05 (P)	0.03 (P) milk powder for infant foods
Germany	0.05 (P)	0.10 (P) milk powder for infant foods
The Netherlands	0.05 (P)	0.04 (P) milk powder for infant foods 0.20 (P) cheese
Russia	0.50 (V)	0.10 (P) milk powder for infant foods
Sweden	0.05 (P)	0.10 (P) milk powder for infant foods
Switzerland	0.05 (C)	0.01 (C) milk powder for infant foods

* (C) = current limit ; (P) = proposed limit ; (V) = voluntary limit




Figure 1. Chemical structures of aflatoxins B₁ and M₁ (van Egmond and Wagstaffe, 1987; Duraković, 2007)

The direct evidence for the carcinogenicity of AFM₁ is slender, primarily because insufficient quantities of the pure compound are available for toxicological studies, and is based largely on limited experiments with rainbow trout (Abedi and Scott, 1969; Hsieh and Ruebner, 1984) and Fisher rats (Abbas et al., 1984). The toxicological risks, however, are sufficiently high to justify legal controls. Current and proposed limits for AFM₁ in European countries are given in Table 1 (van Egmond and Wagstaffe, 1987; Creppy, 2002).

Evidence of hazardous human exposure to AFM₁ through milk and dairy products has been shown by several investigators (Lukač et al., 1991; Galvano et al., 1998; Trucksess, 1999; Samaržija et al., 2003; Galvano et al., 2009). AFM₁ is of great concern because of high human, namely children consumption of milk, dairy products and protein concentrates based on milk such as whey protein concentrates (Tratnik, 1998; Brnčić et al., 2008a; Brnčić et al., 2008b; Brnčić et al., 2009). There are also some innovative techniques for milk processing like ultrasound, pulsed electric fields and cold plasma that could potentially solve problems like this one (Bosiljkov et al., 2011).

Regulatory limits throughout the world are influenced by economic consideration and vary from one country to another (Stoloff et al., 1991). The European Community and Codex Alimentarius prescribe that the maximum level of the AFM₁ in liquid milk and dried or processed milk products should not exceed 50 ng x kg/L⁻¹ (Škrinjar et al., 1995; Trucksess, 1999; Codex Alimentarius Commission, 2001; Kamkar, 2006). However, according to US regulation by FDA, the level of AFM₁ in milk should not be higher than 500 ng x kg/L⁻¹ (Stoloff et al., 1991; Kamkar, 2006).

Therefore, contamination of milk and dairy products with AFM₁ has been recognized as a significant human health hazard.

The aim of this study was to determine the contents of AFM₁ in milk and to estimate LC₅₀ and T₅₀ values to *Artemia salina* larvae in experiments with different temperatures of incubation.

There are several studies on the effects of aflatoxins on the brine shrimp *Artemia salina* eggs and larvae (Harwing and Scott, 1971; Duraković et al., 1986, 1987, 1989; Schmidt, 1989; Logrieco et al., 1996; Hartl and Humpf, 2000; Duraković et al., 2002; Moretti et al., 2007).

Thus, Duraković and co-workers (1989, 2002) reported that concentration resulting in 50 % mortality of AFB₁ exposed to temperature of 20 °C for 24 hours was 3.0 µg x L⁻¹. Investigations by the same authors at temperature of 30 °C during the 24 hours resulted in 50 % mortality of investigated larvae by 0.9 µg x L⁻¹ of this toxin. Research by Schmidt (1989) and Logrieco et al. (1996) on *Fusarium* and *Aspergillus* mycotoxins shows that *Artemia salina* are especially attractive for use in short-term bioassay for these toxins.

According to previous findings (Brown, 1969; Hartl and Humpf, 2000; Favilla et al., 2006) the *Artemia salina* larvae appear to be as susceptible as biological indicator of toxicity of some mycotoxins in foods and feeds.

Materials and methods

AFM₁ standard

AFM₁ standard was obtained from Immunolab GmbH, Kassel (Germany). Toxin was dissolved in methanol to obtain concentrations 5.0, 10.0 and 50.0 µg x mL⁻¹.

Milk

Whole pasteurized milk, from retail shop was used. The milk showed no detectable levels of AFM₁ (results not presented) (Antunac et al., 2002) and the toxin was added in concentration that varied from 0.1 to 5.0 µg x L⁻¹ of milk. The samples were stored frozen in the dark until used.

Larvae

Brine shrimp dry eggs were obtained from Hans Brustman Co., Düsseldorf (Germany). Hatching procedure followed the one described in ARC test, standardized short term toxicity test with *Artemia nauplii* (Vanhaecke and Persoone, 1981a; Vanhaecke et al., 1981b). The hatching medium was artificial seawater of normal seawater salinity (35 g x L⁻¹). For each experiment, 100-200 mg of brine shrimp was placed in each 100 mL of hatching medium contained in a 500 mL Erlenmeyer flasks, and these were shaken as described by Favilla et al., (2006). Hatching can occur in less than 24 hours at 27 °C. Throughout hatching period the same conditions of light sensitivity and temperature were maintained.

Assay method

Disk screening method

Blank paper discs (8 mm diameter) were saturated with a selected solution of AFM₁ (about 20 µL/disc), and each disc was placed directly in the well (Duraković et al., 1987, 1989). Toxicity of each solution was evaluated in triplicate. Two drops (about 0.1 mL) of a larvae suspension (containing 20 to 40 larvae) were added to each well. Trays were incubated at 20 °C, 25 °C, 30 °C, 35 °C and 40 °C, for about 48-96 hours. Mortality was determined by counting the immobile (dead) larvae under a stereoscopic microscope, after killing the living larvae with heat or formalin and then counting the total number. Mortality in controls was determined simultaneously with each screening test. Natural mortality associated with discs saturated in noninoculated media or water average 3 and 1 % respectively (Duraković et al., 1987, 1989; Ben Naceur et al., 2009; Tsolaki et al., 2010).

Results and discussion

Tests were conducted at 20 °C, 25 °C, 30 °C, 35 °C, and 40 °C. The sensitivity of the brine shrimp increased with increase of incubation temperature to about 40 °C; optimum sensitivity to AFM₁ occurred at 30 °C. Tests at 35 °C and 40 °C produced significant mortality among controls. The mortality

Table 2. Percent mortality of *A. salina* larvae at 30 °C after 48 hours at various dose levels of aflatoxin M₁

Aflatoxin M ₁ ($\mu\text{g} \times \text{L}^{-1}$)	Average number of brine shrimp/Test	% Mortality	Number of tests
Control	46	11	10
0.15	42	25	8
0.30	56	50	12
0.40	35	65	10
0.50	38	75	9
0.90	48	95	8

response of the larvae varied according to dosage and it was possible to develop a standard curves in an observation period of 48 and 96 hours, respectively (Figures 2a, 2b, 3a and 3b).

In experiments at 20 °C the larvae started to die in 30 hours and were all killed within 96 hours at a highest dose level ($4.8 \mu\text{g} \times \text{L}^{-1}$ of AFM₁). Data derived from conducting tests at this temperature are

presented in Figures 2a and 2b. At least eight tests were conducted at all dose levels. The tests give positive results at $0.5 \mu\text{g} \times \text{L}^{-1}$ of AFM₁ with a mortality of over 5 %. The dose of $2.5 \mu\text{g} \times \text{L}^{-1}$ of this toxin in 60 hours produced mortality of 50 % of investigated larvae. Greater than 90 % mortality was achieved after 96 hours at dose levels of $4.5 \mu\text{g} \times \text{L}^{-1}$ and above; $0.5 \mu\text{g} \times \text{L}^{-1}$ was the lowest level of AFM₁ tested.




Figure 2a. Dosage-mortality response of *A. salina* larvae to AFM₁ with observation period of 96 hours at 20 °C




Figure 2b. Time-mortality response of *A. salina* larvae to lethal concentration of AFM₁ at 20 °C

Total mortality (100 %) was reached with 4.8 $\mu\text{g} \times \text{L}^{-1}$ of this toxin (Figures 2a and 2b). The values are in good accordance with the findings of Duraković and co-workers (1987, 1989), Schmidt (1989), Duraković and co-workers (2002) and Duraković (2007) who have stated 0.5 and 1.0 $\text{ng} \times \text{mL}^{-1}$ as the minimal concentrations of aflatoxins that produce LC₅₀ value in experiments with *A. salina* larvae.

Figures 3a and 3b and Table 2 represent data from the tests conducted at 30 °C. AFM₁ at dose levels of 0.4 $\mu\text{g} \times \text{L}^{-1}$ and above produced greater than 60 % mortality after 48 hours (Table 2). The larvae started to die in 10 hours and were all killed within 55 hours; 30 hours was the time for a 50 % kill. Greater than 90 % mortality was achieved after 48 hours at dose levels of 0.9 $\mu\text{g} \times \text{L}^{-1}$ and above. At this incubation temperature and toxin concentration of 1.2 $\mu\text{g} \times \text{L}^{-1}$ all the larvae were killed within 55 hours. At a lower temperature of incubation and lower concentrations of AFM₁ this period was prolonged by a few hours up to several days (Figures 2a, 2b, 3a and 3b).

Dosage mortality (LC₅₀ - 48/96 hours) and time mortality (T₅₀ - 48/96 hours) were calculated by graphical interpolation.

The percentage mortality between 5 % and 95 % were calculated from the average number of dead larvae per concentration, and plotted on log-probability paper. A straight line is drawn at sight through the points. The intersection of this line with the 50 % mortality horizontal line determines LC₅₀.

The test aims at the determination of the LC₅₀ and T₅₀ in the 48/96 hours on the basis of the critical range concentrations obtained in the preliminary test. Concentrations and dilutions are chosen from a logarithmic scale (Duraković et al., 1986, 1987). In principle five concentrations should be sufficient. For satisfactory LC₅₀, however, at least three data must be situated in the mortality range 5-95 %. If this not the case, the test should be repeated with additional concentrations from dilution scale. For each concentration, including the control, three replications should be used.




Figure 3a. Dosage-mortality response of *A. salina* larvae to AFM₁ with observation period of 48 hours at 30 °C




Figure 3b. Time-mortality response of *A. salina* larvae to lethal concentration of AFM₁ at 30 °C

Since AFM₁ is soluble in artificial seawater to a concentration of about 20 µg x mL⁻¹, the sensitivity of this test is below the upper solubility limits. The test volume, 1 mL can maintain up to 50 brine shrimp larvae without affecting the results. Living culture of test organism needs to be maintained; enough larvae to conduct a test can be made available by placing eggs in artificial seawater at 27 °C from 18-24 hours before they are needed (Duraković et al., 1989; Dabrowski, 1991; Coldwell et al., 2003; Moretti et al., 2007; Munoz et al., 2008). The brine shrimp test was also used effectively with partially purified other mycotoxins and chloroform extracts of a toxigenous moulds cultures (Brown, 1969; Duraković et al., 1987; Logrieco et al., 1996; Sarabia et al., 2002; Favilla et al., 2006).

For various reasons the brine shrimp *Artemia salina* is a uniquely suitable test species for laboratory experiments. In cysts they can be stored for years under dry conditions without losing their viability. The advantages of *A. salina* as the best "first choice" for toxicity studies in milk can be summarized as follows:

- the cysts are commercially and readily available so that the tests can be carried out worldwide with the same original material and without any problem of provisioning; moreover, the quantity of cysts required per test is very small so the price of the biological material is negligible,
- the necessity of year-round maintenance of stock cultures, with all the biological and technical difficulties and the considerable economic repercussions, is completely eliminated,
- large number of test organisms of exactly the same age and physiological condition can be easily obtained to start the tests.

Conclusions

Over the past several years, the toxicity of mycotoxins to the brine shrimp *Artemia salina* has been studied. Most investigations were conducted over 16-24 hours period and showed LC₅₀ values as low as the 500 ng x kg/L⁻¹ level (Stoloff, 1980; van Egmond and Paulsch, 1986; Moretti et al., 2007; Galvano et al., 2009).

Although these studies promise in using *A. salina*

as a bioassay for mycotoxins, several inadequacies existed in the data. The conditions for the studies were not the same; therefore, comparisons of data from different researchers are difficult. Also, in consideration of current concern over the possible use of mycotoxins as biochemical warfare agents, the detection limit needed to be extended to the low µg x L⁻¹ level (van Egmond and Wagstaffe, 1987; van Egmond, 1989; Creppy, 2002).

A. salina are especially attractive for the use in a short-term bioassay for aflatoxins. The dry eggs can withstand adverse environmental conditions and still hatch out in 24-48 hours in the presence of artificial seawater. Continuous maintenance of stock culture is not necessary. Test populations can be as large as desired, and the fresh larvae can live for 48 hours, during which time most bioassay can be conducted. A continuous system of hatching can be used to provide fresh larvae daily.

Compared to other organisms, the larvae are free from the problem of sterilization, culture media, techniques of toxic administration, etc., and are economical, time saving and readily available.

Utjecaj inkubacijske temperature i koncentracije aflatoksina M₁ u mlijeku na larve račića Artemia salina

Sažetak

Proučavani su utjecaji inkubacijske temperature i koncentracije aflatoksina M₁ (AFM₁) na larve račića *Artemia salina* kao biologijskog indikatora u temperaturnom rasponu od 20 °C do 40 °C. Povišenje temperature inkubacije uzrokovalo je povećanu osjetljivost račića na AFM₁. Optimalna osjetljivost očitovala se pri 30 °C. Pozitivni rezultati dobiveni su pri koncentraciji AFM₁ 0,18 µg x L⁻¹ pasteriziranog mlijeka uz izraženu smrtnost višu od 15 %. Smrtnost viša od 90 % očitovala se pri koncentracijama AFM₁ 0,9 µg x L⁻¹ i višim. Test se može provesti u tijeku 30-60 sati.

Ključne riječi: aflatoxin M₁, biotest, *Artemia salina*, vrijednosti LC₅₀ i T₅₀

Acknowledgements

We are gratefully expressing our acknowledgements to gentlewoman Brigitta Duraković for language correcting the manuscript.

The investigations were supported by a grant No. 058-0582184-0432 from Croatian Ministry of Science and Technology.

References

1. Abbas, H.K., Mirocha, C.J., Shier, W.T. (1984): Mycotoxins produced from fungi isolated from foodstuff and soil: comparison of toxicity in fibroblasts and rats feeding tests. *Applied and Environmental Microbiology* 48, 654-661.
2. Abedi, Z.H., Scott, P.M. (1969): Detection of Toxicity of Aflatoxin, Sterigmatocystin, and other Fungal Toxins by Lethal Action on Zebra Fish Larvae. *Journal of the Association of Official Analytical Chemistry* 52, 963-969.
3. Antunac, N., Lukač-Havranek, J., Samaržija, D. (2001): Effect of breed on chemical composition of goat milk. *Czech Journal of Animal Science* 46 (6), 268-274.
4. Ben Naceur, H., Jenhani, A.B.R., Romdhane, M.S. (2009): Ecobiological survey of the brine shrimp *Artemia salina* from Sabkhet El Adhibet (south-east Tunisia). *Journal of the Marine Biological Association of the United Kingdom* 89 (6), 1109-1116.
5. Bhat, R.V., Vasanthy, S. (1999): Mycotoxin contamination of foods and feeds, Overview: occurrence and impact on food availability trade exposure of farm animals and related economic losses. *Presented in the 3rd Joint FAO / WHO / UNEP International Conference on Mycotoxins*, Tunis, Tunisia, 3-6 March, 1999.
6. Bosiljkov, T., Tripalo, B., Brnčić, M., Ježek, D., Karlović, S., Jaguš, I. (2011): Influence of high intensity ultrasound with different probe diameter on the degree of homogenization (variance) and physical properties of cow milk. *African Journal of Biotechnology* 10 (1), 34-41.
7. Brnčić, M., Ježek, D., Rimac-Brnčić, S., Bosiljkov, T., Tripalo, B. (2008a): Influence of whey protein concentrate addition on textural properties of corn flour extrudates. *Mljekarstvo* 58 (2), 131-149.
8. Brnčić, M., Karlović, S., Bosiljkov, T., Tripalo, B., Ježek, D., Cugelj, I., Obradović, V. (2008b): Obogaćivanje ekstrudiranih proizvoda proteinima sirutke. *Mljekarstvo* 58 (3), 275-295.
9. Brnčić, M., Bosiljkov, T., Ukrainczyk, M., Tripalo, B., Rimac-Brnčić, S., Karlović, S., Karlović, D., Ježek, D., Vikić-Topić, D. (2009): Influence of Whey Protein Addition and Feed Moisture Content on Chosen Physicochemical Properties of Directly Expanded Corn Extrudates. *Food and Bioprocess Technology: an International Journal*, DOI10.1007/s11947-009-0273-0.
10. Brown, R.F. (1969): The effect of some mycotoxins on the Brine Shrimp *Artemia salina*. *Journal of the American Oil Chemists' Society* 46, 119-119.
11. Codex Alimentarius Commission (2001): Comments submitted on the draft Maximum level for Aflatoxin M₁ in milk. *Codex committee on food additives and contaminants*, 33rd session, Hogue, The Netherlands.
12. Coldwell, G.S., Bentley, M.G., Olive, P.J.W. (2003): The use of a brine shrimp (*Artemia salina*) bioassay to assess the toxicity of diatom extracts and short chain aldehydes. *Toxicon* 42, 301-306.
13. Creppy, E.E. (2002): Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicology Letters* 127, 19-28.
14. Dabrowski, K. (1991): Some aspects of ascorbate metabolism in developing embryos of the brine shrimp (*Artemia salina*). *Canadian Journal of Fisheries and Aquatic Sciences* 48, 1905-1908.
15. Duraković, S., Beritić, T., Duraković, Z., Pospišil, O., Delaš, F., Vorkapić-Furač, J. (1986): Primjena dehidriranih jaja *Artemia salina* u dokazivanju toksičnosti aflatoksinina. *Hrana i ishrana* 27 (4), 199-204.
16. Duraković, S., Duraković, Z., Beritić, T., Pospišil, O., Radić, B. (1987): Larve *Artemia salina* kao test organizam u istraživanju sinergizma mikotoksina. *Arhiv za higijenu rada i toksikologiju* 38 (4), 307-313.
17. Duraković, S., Galić, J., Rajnović, P., Pospišil, O., Štilinović, L. (1989): Larve *Artemia salina* kao jednostavan biološki reagens za istraživanje toksičnosti mikotoksina. *Acta Biologica Iugoslavica (B Mikrobiologija)* 26 (1), 15-22.
18. Duraković, S., Duraković, L. (1999): *Specijalna mikrobiologija*. Popović N., ed. Kugler, Zagreb.
19. Duraković, S., Delaš, F., Duraković, L. (2002): Mikrobnii indikatori sigurnosti i kakvoće namirnica. In: *Moderna mikrobiologija namirnica-knjiga druga*, Duraković S., ed. Kugler, Zagreb, pp: 249-282.
20. Duraković, S., Duraković, L. (2003): *Mikologija u biotehnologiji*. Duraković S., ed. Kugler, Zagreb.
21. Duraković, L. (2007): *The influence of chosen parameters on the growth of the mold Aspergillus flavus ATCC 26949 in mixed culture and biosynthesis of aflatoxins B₁ and G₁*. PhD Thesis. Faculty of Food Technology and Biotechnology, University of Zagreb, Croatia.
22. Duraković, S., Duraković, L., Vahčić, N., Skelin, A., Duraković, Z. (2008): Affect of moulds growth in mixed cultures on production of aflatoxins on maize hybrid. *Cereal Research Communications* 36 (Part 3, Supplement 5), 1615-1618.
23. Favilla, M., Macchia, L., Gallo, A., Altomare, C. (2006): Toxicity assessment of metabolites of fungal biocontrol agents using two different (*Artemia salina* and *Daphnia magna*) invertebrate bioassays. *Food and Chemical Toxicology* 44, 1922-1931.
24. Galvano, F., Galofaro, V., De Angelis, A., Galvano, M., Bognanno, M., Galvano, G. (1998): Survey of occurrence of Aflatoxin M₁ in dairy products marketed in Italy. *Journal of Food Protection* 61, 738-741.

25. Galvano, F., Galofaro, V., Galvano, G. (2009): Occurrence and stability of aflatoxin M₁ in milk and milk products - a worldwide review. *Journal of Food Protection* 59 (10), 1079-1090.
26. Hartl, M., Humpf, H.U. (2000): Toxicity assessment of fumonisins using the brine shrimp (*Artemia salina*) bioassay. *Food and Chemical Toxicology* 38, 1097-1102.
27. Harwing, J., Scott, P.M. (1971): Brine shrimp (*Artemia salina* L.) Larvae as a Screening System for Fungal Toxins. *Applied Microbiology* 21 (6), 1011-1016.
28. Hsieh, D.P.H., Ruebner, B.H. (1984): An Assessment of Cancer Risk from aflatoxins B₁ and M₁. In: *Toxigenic fungi-their toxins and human health hazard*, Kurata H. and Ueno Y., eds. Elsevier, Amsterdam-New York-Oxford-Tokyo, pp: 332-338.
29. IARC (1993a): Aflatoxins: naturally occurring aflatoxins (Group 1). Aflatoxin M₁ (Group 2b). *International Agency for Research on Cancer*. Lyon, France, Vol. 56, pp: 245-252.
30. IARC (1993b): Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. *International Agency for Research on Cancer*. Lyon, France, Vol. 56, pp: 345-395.
31. Kamkar, A. (2006): A study of the occurrence of aflatoxin M₁ in Iranian Fetta cheese. *Food Control* 17, 768-775.
32. Logrieco, A., Moretti, A., Fornelli, F., Fogliano, V., Ritieni, A., Caraffa, M.F., Randazzo, G., Botalico, A., Macchia, L. (1996): Fusaproliferin production by *Fusarium subglutinans* and its toxicity to *Artemia salina* SF-9 insect cells, and IARC / LCL 171 human B lymphocytes. *Applied and Environmental Microbiology* 62, 3378-3384.
33. Lukač, J., Samaržija, D., Antunac, N. (1991): Kvaliteta mlijeka na području Hrvatske. *Mljekarstvo* 41 (3), 65-70.
34. Moretti, A., Mule, G., Ritieni, A., Logrieco, A. (2007): Further data on the production of beauvericin, enniatins and fusaproliferin and toxicity to *Artemia salina* by *Fusarium* species *Gibberella fujikuroi* species complex. *International Journal of Food Microbiology* 118 (2), 158-163.
35. Munoz, J., Gomez, A., Green, A.J. (2008): Phylogeography and local endemism of the native Mediterranean brine shrimp *Artemia salina* (Branchiopoda : Anostraca). *Molecular Ecology* 17 (13), 3160-3177.
36. Sarabia, R., del Ramo Varo, I., Diaz Mayans, J., Torrelblanca, A. (2002): Comparing the acute response to cadmium toxicity of nauplii from different populations of *Artemia*. *Environmental Toxicology and Chemistry* 21 (2), 437-444.
37. Samaržija, D., Antunac, N., Pećina, M., Havranek, J. (2003): Quality of artisanal hard cheeses produced in the Mediterranean area of Croatia. *Milchwissenschaft* 58 (1-2), 43-46.
38. Schmidt, R. (1989): The application of *Artemia salina* L. bioassay for screening of aflatoxins. In: *Aspergillus: Mycotoxins Taxonomy and Pathogenicity*, Chelkowski J., ed. Elsevier, Amsterdam, The Netherlands, pp: 121-130.
39. Stoloff, L. (1980): Aflatoxin M₁ in perspective. *Journal of Food Protection* 43, 226-230.
40. Stoloff, L., van Egmond, H.P., Park, D.L. (1991): Rationales for establishment of limits and regulations for mycotoxins. *Food Additives and Contaminants* 8, 213-222.
41. Škrinjar, M., Stubblefield, R.D., Vujičić, I.F., Stojanović, E. (1992): Distribution of aflatoxin-producing moulds and aflatoxins in Yugoslav dairy cattle feed and raw milk. *Acta Microbiologica Hungarica* 39 (2), 175-179.
42. Škrinjar, M., Danev, M., Dimić, G. (1995): Investigations on the presence of toxigenic fungi and aflatoxins in raw milk. *Acta Alimentaria* 24 (4), 395-402.
43. Taveira, J.A., Midio, A.F. (2001): Incidence of aflatoxin M₁ in milk marketed in Sao Paulo, Brazil. *Italian Journal of Food Science* 13 (4), 443-447.
44. Tratnik, Lj. (1998): *Mlijeko - tehnologija, biokemija i mikrobiologija*. Hrvatska mljekarska udruga, Zagreb.
45. Tratnik, Lj., Božanić, R., Mioković, G., Šubarić, D. (2001): Optimization of Manufacture and Quality of Cottage Cheese. *Food Technology and Biotechnology* 39 (1), 43-48.
46. Trucksess, M.W. (1999): Mycotoxins. *Journal of AOAC International* 82, 488-495.
47. Tsolaki, E., Pitta, P., Diamadopoulos, E. (2010): Electrochemical disinfection of stimulated ballast water using *Artemia salina* as indicator. *Chemical Engineering Journal* 156 (2), 305-312.
48. van Egmond, H.P., Paulsch, W.E. (1986): Mycotoxins in milk and milk products. *Netherlands Milk and Dairy Journal* 40, 175-188.
49. van Egmond, H.P., Wagstaffe, P.J. (1987): Development of Milk Powder Reference Materials Certified for Aflatoxin M₁ Content (Part I). *Journal of the Association of Official Analytical Chemists* 70 (4), 605-610.
50. van Egmond, H.P. (1989): Aflatoxin M₁: occurrence, toxicity, regulation. In: *Mycotoxins in Dairy Products*, Van Egmond H.P., ed. Elsevier Applied Science, London, pp: 11-55.
51. van Egmond, H.P. (1991): Mycotoxins. *Bulletin of the International Dairy Federation, Special Issue* 9 (101), 131-145.
52. Vanhaecke, P., Persoone, G. (1981a): Standardised short term toxicity test with *Artemia nauplii* (ARC test). *Institut National de la Santé et de la Recherche Médicale* 106, 370-376.
53. Vanhaecke, P., Persoone, G., Claus, C., Sorgelos, P. (1981b): Proposal for a Short-Term Toxicity Test with *Artemia nauplii*. *Ecotoxicology and Environmental Safety* 5, 382-387.
54. Wood, G.E. (1991): Aflatoxin M₁. In: *Mycotoxins and phytoalexins*, Sharma R.P. and Salunkhe D.K., eds. CRC Press, Boca Raton, pp: 145-164.