

The effect of different marinating baths on sensory properties and shelf life parameters of cold marinated anchovies (*Engraulis encrasicolus*, L.)

V. Šimat¹, T. Bogdanović², M. Bulić³

Znanstveni rad

Summary

Synergistic effect of acid and salt content plays the most important role in the marinating process. Selection of suitable procedures for preparing raw fish material for marination, the marinating baths compositions and storage conditions, are carried out in order to achieve a longer shelf life, to maintain quality and achieve good sensory characteristics of marinated products. In this paper 22 samples of cold marinated anchovies containing marinating baths of different composition (different proportions of vine vinegar, alcoholic vinegar, salt and water) were investigated, in order to determine changes in sensory characteristics during maturation, as well as the shelf life parameters. For this purpose, pH values, water activity, NaCl content and sensory properties of marinated anchovy fillets, the content of volatile amines and the index of fat oxidation were determined. The fillets marinated in the marination bath prepared from 45% of water, 30% of vine vinegar, 25% of alcoholic vinegar and 7% of salt obtained the best sensory scores, a pH value of 3.09, 3.60% NaCl and water activity of 0.84. A weak accumulation of volatile amines during storage (<20mg TVB-N / 100g; <1mgTMA/100g) and statistically significant increase of the TBA index (6.41 to 7.36 malondialdehyde / kg) were observed during ripening, but the rancidity was not described as an intensive sensory property of the marinated fillets.

Key words: *Engraulis encrasicolus*, cold marinade, shelf-life, TVB-N, TBAi

Introduction

Anchovy (*Engraulis encrasicolus*, L.) is a small pelagic fish and the only European representative of the *Engraulidae* family, thus an endemic species of the Mediterranean-Atlantic region. The anchovies are caught along the Mediterranean shores and in the Adriatic Sea all year round, but mainly within the spring-autumn period (Sinovčić, 2000). It represents a species of an important economic value in all parts of the Mediterranean. The average annual capture production of European anchovy from 2002 till 2008 in the Mediterranean and Black Sea was around 480000 tons (Anon., 2010). The Croatian catch of anchovies in the Adriatic Sea increased four times since 2003 and had reached 13200 tons in 2007 (Anon., 2009).

Most of the catch is processed by salting, marinating or freezing and utilized for human consumption. Healthy living and Mediterranean kitchen encourage the consumption of fish and seafood products; especially lightly preserved products, such as cold marinades or carpaccio-like products.

The basic principle of cold marinating preservation is associated with the synergic activity of organic acids and salt from marinating bath to the changes in fish fillets. These products are made out of fresh, frozen or salted fish. Organic acids and salt are added to the fish to retard the microbiological and enzymatic activity, but also to change the taste and textural properties of the fish, resulting

with a lightly preserved product with limited shelf life (Poligne and Collingan, 2000; Pons-Sanchez-Cascado et al., 2005; Yeannes and Casales, 2008; Olgunoğlu et al., 2009, Šimat, 2010). The microbiological activity of cold marinades is restricted by low pH values, low water activity and high NaCl content. However, in order to insure the continuity in production process and best sensory properties of the product acid type and concentration, and NaCl content must be carefully chosen. Numerous studies were conducted in search of right procedures and preparation of raw fish for marinating, compositions of marinating baths and cover brines of marinated products, adequate storage for these products, all with purpose to achieve a longer shelf life and preservation of

¹ dr.sc. **Vida Šimat**, assistant, Center of Marine studies, University of Split, Livanjska 5/III, 21000 Split vida@unist.hr

² **Tanja Bogdanović**, BSc. biotechnology, Regional Veterinary Laboratory, Croatian Veterinary Institute, Poljička cesta 33, 21000 Split

³ **Milko Bulić**, BSc. fishery science, Center of Marine studies, University of Split, Livanjska 5/III, 21000 Split

Table 1. Overview of water, vinegar and NaCl contents and ratios in production of different marinating baths used for anchovy's fillet marination.

Tablica 1. Pregled omjera i udjela vode, octa i soli u pripremi različitih naljeva korištenih za mariniranje fileta incuna.

Marination bath ingredients Sastojci naljeva za mariniranje	Sample number Redni broj uzorka																					
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17 ^A	18 ^A	19 ^A	20 ^A	21 ^A	22 ^B
Water (%) Voda (%)	45	40	35	45	40	35	60	50	40	60	50	40	45	45	50	45	45	45	40	50	40	45
Alcoholic vinegar (%) Alkoholni ocat (%)							40	50	60	40	50	60	25	25	20		55	30	60			
Vine vinegar (%) Vinski ocat (%)	55	60	65	55	60	65							30	30	30	55		25		50	60	55
Salt (%) Sol (%)	7	7	7	10	10	10	7	7	7	10	10	10	7	10	7	7						7

^A Before marination the fillets were placed in 24 % salt solution for 45 minutes

^A Prije mariniranja fileti su salamureni u 24 % salamuri 45 minuta

^B Vine vinegar used for marination bath of this product had 8% of acetic acid (vol/vol)

^B Vinski ocat korišten za mariniranje ovog proizvoda sadržavao je 8% octene kisline (vol/vol)

product quality, without the loss of their specific sensory properties (Huss et al., 2003; Kilnic and Cakli, 2004; Olgunoğlu et al., 2009, Šimat, 2010). The salt content of lightly preserved products is essential to the product safety. It is advisable to use a salt content >6% as a critical limit (Derrick, 2009). However, such quantity of salt might be unsuitable for consumers, so the reduced salt content in products could be compensated by adding preservatives, such as sodium benzoate, sorbates, nitrates, or others that also enhance flavour and aroma (spices, sugar, etc). Although many scientists have proved the effect of several preservatives to prevent the risk of pathogenic bacteria, their value and application are not standardized for all types of marinade or other lightly preserved fishery products (Lai and Roy, 2004; Zaika, 2007). According to Meyer (1965), the growth of microorganisms is decelerated in brines with salt content of 6.5% if the acetic acid is present, because its presence makes the bacteria more sensitive. Additionally, salt enhances the texture and taste, achieving good

sensory properties and safety of the cold marinated fish. The aim of this research was to investigate the right procedures and preparation of raw anchovies for marinating, the effect of different marinating baths on sensory properties and shelf life parameters, all with purpose to achieve a longer shelf life of products and preservation of product quality without the loss of their specific sensory properties.

Materials and methods


Marinating process

Anchovies (*Engraulis encrasicolus*) caught in January 2010 in Central Adriatic fishing region GSA 17 (GMU 37.2.1) were used for product preparation. The fish were placed in self-draining polystyrene boxes, packed in flake ice and delivered to the local factory where they were frozen using combined brine/air IQF freezing process (Tacore, Spain) and stored at -24(±1) for 6 months. Approximately 100 kg of anchovies were thawed and used in cold marinating process (Figure 1.). The size category of the fish

was 42 individuals per kilogram. The process consisted of thawing and filleting of the anchovies, followed by rinsing of the fillets in rinsing tank filled with mild salt solution, and the preparation of the products. Fillets were placed in the plastic containers, marination baths made out of different types of vinegar and salt contents (Table 1.). The ratio between fish and marination bath was 1:1.5. During the entire production process as well as storage, fish fillets were completely immersed in the marination bath. Products numbered 18 through 22 from the table were brined in 24% salt solution before marination, so no additional salt was added into the marination bath. After 48 hours 11 products with best sensory properties were selected, and placed in the second marinating bath, consisted of the same ratio of acid and salt as the first one. These 11 products were stored at 4(±2) °C for 25 days.

Sensory assessment

Sensory assessment of the marinated anchovy fillets was carried out by a sensory panel of six trained and



experienced members. The intensity of sensory properties was described through a descriptive scale with point range from 1 to 8, specially adopted for marinated anchovies (modified method by Sallam, 2006). The scale was used to describe fillet's overall acceptability with regard to appearance, colour, odour, flavour (acidity and salinity), rancidity and aftertaste intensity, texture and juiciness. One being extremely good/ white/ pleasant/ intensive/ tender/ juicy in sensory properties, and eight extremely bad/ brown/ unpleasant/ tasteless/ filamentous/ dry. The rejection point was set at 5.5. The fillets were placed in a plastic dish, held at room temperature for 30 minutes before served to the panellists. The panel evaluated the samples, coded with random three-digit numbers without any information on storage time. They were instructed to eat crackers and drink water between the product assessments.

Proximate analysis

For proximate composition analy-

sis 20 individual fish of both raw and thawed anchovies were sampled, gutted, filleted and homogenised using a laboratory blender (KINEMATICA Mikrotron MB 550, Switzerland). The proximate composition was determined from homogenates as water content (drying the samples at 105°C to constant weight), crude protein (Kjeldhal method, $N \times 6.25$), crude fat (acid hydrolysis method) and crude ash (calcinations at temperatures $\leq 500^\circ\text{C}$) (AOAC, 2000). Proximate composition of marinated fillets was determined for the three products with best sensory score after 25 days of storage. Fillets were drained and homogenized for analysis. All analyses were done in triplicate and presented as percentage wet weight of anchovy fillet.

pH, NaCl and water activity analysis

The pH, NaCl and water activity were determined for raw material (fresh, thawed, rinsed, brined anchovies) and 11 marinated anchovy products after 2 and 25 days of storage.

The pH was measured using digital pH meter (702 SET/MET Titrino, Metrohm), equipped with glass electrode, calibrated at 4 and 7. The electrode was dipped into the mixture of fish and distilled water (1:1), at ambient temperature (Kyrana and Lougouvios 2002).

A method by Morh was used to determine salt content in the fillets.

Water activity was determined on sample homogenate using a_w meter Rotonc AG, HygroPalm AW1-set40, equipped with a calibrated digital probe with measurement accuracy ± 0.005 .

Determination of thiobarbituric acid index and volatile amine content

Thiobarbituric acid index and volatile amine content were determined

for raw material (fresh, thawed, rinsed and brined) and marinated anchovy fillets from three products with best sensory scores at the end of marination process.

Thiobarbituric acid index (TBAi) was determined as previously described by Vyncke (1970) and Lemon (1975), using spectrophotometer (PRIM Advanced, Secomam, France). After extraction with trichloroacetic acid (100 g/L), antioxidant propyl gallate (1 g/L) and chelating agent ethylenediaminetetraacetic acid (1 g/L) were added into solution to lower the possibility of erroneously formed malondialdehyde or other TBA reactive substances during blending and filtering of the sample. Calibration curves were done using 1,1,3,3-tetraethoxy-propane (TEP). Results were expressed as mg malondialdehyde/kg muscle (mg MA/kg).

Total volatile base nitrogen (TVB-N) and trimethylamine nitrogen (TMA-N) were determined by direct distillation of fish extracts as previously described in Šimat et al. (2009) and expressed as mg TVB-N/ TMA-N per 100g of fish muscle.

Statistical analysis

Graphic figures and statistical evaluations (analysis of variance, least significance difference, correlations) of obtained data were done using Microsoft Office Excel 2007 package and software Statgraphics® Plus v. 5.1 Professional (Manugistics, Inc., Rockville, MD, USA).

Results and discussion

The limited shelf-life of the cold marinades and other lightly preserved products is changeable due to high post mortal pH of fish muscle, high content of free amino acids, the presence of trimethylamine oxide (TMAO), the presence of specific spoilage bacteria and the differences in chemical composition and condition of the specimens of wild fish populations (Dalgaard, 2000). The

Table 2. Water activity (a_w), pH values and NaCl content of raw anchovy fillet used for the preparation of the products.Tablica 2. Vrijednosti aktiviteta vode (a_w), pH i udjela soli u svježem inčunu i sirovini korištenoj za izradu proizvoda.

Analyzed anchovy sample Analizirani uzorak inčuna	Parameters determined Određeni parametri						Proximate analysis (%) Kemijski sastav (%)			
	a_w	pH	NaCl (%)	TBAi (mg MA/kg)	TVB-N (mg/100g)	TMA-N (mg/100g)	Water content Udio vode	Protein content Udio bjelančevina	Fat content Udio masti	Ash content Udio pepela
Fresh Svježi	0.994 ^a (±0.001)	6.16 ^a (±0.01)	0.47 ^a (±0.11)	1.16 ^a (±0.19)	10.21 ^a (±0.26)	0.42 ^a (±0.09)	76.23 ^a (±0.24)	19.31 ^a (±0.04)	1.47 ^a (±0.08)	1.72 ^a (±0.05)
Thawed Odmrznuti	0.987 ^b (±0.002)	6.30 ^b (±0.02)	2.57 ^b (±0.24)	2.04 ^b (±0.14)	13.05 ^b (±0.32)	0.69 ^b (±0.12)	71.88 ^b (±0.26)	20.11 ^b (±0.11)	1.79 ^b (±0.11)	2.31 ^b (±0.11)
Rinsed Ispirani	0.844 ^c (±0.001)	5.86 ^c (±0.01)	1.30 ^c (±0.14)	1.89 ^c (±0.16)	12.56 ^b (±0.28)	0.62 ^b (±0.10)	ND	ND	ND	ND
Brined Salamureni	0.766 ^d (±0.002)	5.77 ^d (±0.01)	10.66 ^d (±0.62)	2.61 ^d (±0.18)	14.34 ^c (±0.34)	0.71 ^b (±0.14)	70.05 ^c (±0.14)	20.89 ^c (±0.11)	1.32 ^a (±0.11)	4.62 ^c (±0.12)

^{a-c} values in the same column labelled with different letters are significantly different ($p < 0.05$).^{a-c} vrijednosti u istoj koloni označene različitim slovom se statistički značajno razlikuju ($p < 0.05$).

knowledge of proximate composition of fish species is important in the application of different technological processes. It affects the quality aspect of raw material, sensory attributes and storage stability (Huss, 1988; Sikorski et al., 1990). In small pelagic fish seasonal changes in fat content might affect the flavour and sensory characteristics of seafood products. Relatively low fat content of the anchovy fillet (Table 2.) makes it a good choice of raw material for marinades. Principal factors in controlling and preventing the growth and activity of microorganisms are water activity, salt content and pH value of the product. The pH values <4.5, water activity values <0.85 and salt content >6.0% might inhibit the growth of most pathogens (Meyer, 1965; Poligne and Colligan, 2000; Derrick, 2009; Šimat, 2010).

The quality and freshness of the final product directly depend on the raw material used for its preparation, therefore water activity (a_w), the salt content and pH value, as well as TBA index, volatile amine content and proximate composition of fresh and thawed anchovies were determined (Table 2.). Same parameters were analysed after the rinsing and brining processes to determine differences

that might influence further development of the product.

The typical pH of the live fish is around 7, so the low initial pH of the anchovy muscle could reflect a good nutritional state of the fish or that the fish underwent some stress before the capture (Kristoffersen et al., 2006). Statistically significant differences ($p < 0.05$) were determined between the analysed quality parameters of fresh and thawed anchovies. Freezing process resulted in higher salt and ash content of thawed samples and long storage time at $-24(\pm 1)^\circ\text{C}$ in higher TBA index and TVB-N content, plus lower water content and pH value. The pre-treatments (rinsing and brining of the fillets) are necessary in production processes to ensure the removal of scales, blood or intestine parts from the fillets. Compared to raw anchovies, both pre-treatments resulted in lower pH of the muscle and statistically significant difference in NaCl content in the fillets.

The total volatile base nitrogen (TVB-N) and individual methylamine, especially trimethylamine (TMA) have a long history of use as a measure of deterioration of fish and fishery products, and as a criterion of freshness and degree of deterioration

in monitoring quality and ensuring safety of these products (Howgate, 2010). The concentration of TVB-N in freshly caught fish range from 5 to 20 mg/100g, while levels of 30-35 mg TVB-N/100g are considered the limit of acceptability for most marine fish species (Connell, 1995, Ababouch et al., 1996). Levels <15 mg TVB-N/100g were found in all the analyzed samples. Similar to our results, other authors reported TVB-N contents of fresh anchovies in range from 7.32 to 12.77 mg/100g (Pons-Sanchez-Cascado et al., 2005; Olgunoğlu et al., 2009; Özogul et al., 2010). The TMA content of anchovies was <1 mg/100g, indicating a very good quality of the raw material. Thiobarbituric test was used to determine the secondary oxidation of unsaturated fatty acids, malondialdehyde (MA) content, of the raw materials and marinated anchovies. Good quality fish is considered to be those whose values of MA do not exceed 3 mg/kg, while the quality of the fish with the amount of 8 mg (MA)/kg is considered deteriorated (Schormüller, 1969). The TBA indexes of raw material ranged from 1.16 to 2.61 mg MA/kg, indicating good quality fish.

The marinated products were prepared using the rinsed and brined fillets with the addition of differ-

Table 3. Water activity (a_w), pH value and NaCl content of selected marinated anchovy fillets after 2 and 25 days of storage.Tablica 3. Aktivitet vode (a_w), pH vrijednost i udio NaCl u mariniranim filetima inćuna nakon 2. i 25. dana skladištenja.

Sample number* Redni broj uzorka	Determined parameters Određivani parametri					
	a_w		pH		NaCl (%)	
	Number of days in marinating bath Broj dana u kupelji za mariniranje					
	2	25	2	25	2	25
2	0.853	0.805	3.53	3.15	4.39	3.51
3	0.854	0.803	3.60	3.15	4.76	3.84
7	0.856	0.806	3.57	3.14	4.91	4.51
9	0.828	0.809	3.44	3.01	5.60	3.54
11	0.801	0.811	3.49	3.00	7.19	5.08
13	0.846	0.840	3.47	3.09	4.92	3.60
14	0.811	0.843	3.49	3.04	6.10	4.79
16	0.802	0.847	3.88	3.31	4.18	4.65
17	0.812	0.856	3.45	3.17	3.70	3.23
18	0.851	0.849	3.48	3.11	4.52	2.18
20	0.818	0.848	3.45	3.23	4.29	2.43

* Samples descriptions are given in Table 1.

* Uzorci su opisani u Tablici 1.

ent marinating baths as described in Table 1. For products numbered from 1-16 rinsed anchovy fillets were used, while products from 17 to 22 were made with brined fillets. Two days after marination, the first sensory assessment was carried out and products with best sensory properties were selected for further investigation. Different composition of the marinating baths significantly affected the sensory properties of the products. Among the 22 products, 11 were selected for best sensory properties. Eight products (2, 3, 7, 9, 11, 13, 14, 16) made with rinsed anchovies and three products (17, 18, and 20) from brined anchovy fillets were chosen. All were put in a new marinating bath and stored at $4(\pm 2)$ °C for 25 days. After 2 and 25 days of marination water activity, pH value of the muscle and NaCl content were determined (Table 3.), as well as sensory assessment of these 11 products (Figures 2. and 3.). The marinating baths of the remaining products (1, 4, 5, 6, 8, 10, 12, 15, 19, 21, 22) were found not suitable for cold marination of anchovies.

The lowest water activity was recorded in samples with the highest NaCl content (11 and 14). Only these two samples had 10% of salt in the marinating baths. The lowest salt content was found in samples prepared with brined fillets (17, 18, 20) that had no additional salt in the marinating baths. It decreased even further with storage time. Generally, fillets in the products with a higher proportion of vinegar (3 and 9) absorbed more salt after marinating process. However, the NaCl content in all products decreased with storage, with the exception of the product 16 (55% of vine vinegar and 7% of NaCl). Using marinating baths without the addition of NaCl is not suitable for marination, regardless of pre-treatment of the raw material. Salt content of 4% is sufficient to insure marinated fish from spoiling (Fuselli et al. 1994). According to this criterion, the product 17, with salt content <4% during storage, was not satisfactory (Table 3.).

Although the pH value of all marinating baths used was 3.4-3.5, the ef-

fect of different sources of acid acted differently on the pH of fillets. The effectiveness of organic acids to achieve an antimicrobial effect in food varies depending on the degree of acid dissociation (Ray and Bhunia, 2008). With respect to the achieved pH of the fillets in all products, both types of vinegar (vine and alcohol) have shown to be a good choice of acid for fish marinade. After 25 days of storage at $4(\pm 2)$ °C, pH value decreased in all samples and ranged from 3.0 to 3.3 (Table 3.). Samples marinated with alcohol vinegar had lower pH than others, which affected the sensory evaluation, and the acidity of these products was rated as more intensive sensory characteristics (Figure 2.). Varlik et al. (2000) found that the value of pH in the marinated products should range between 4.1 and 4.5. However, lower pH values of marinated fish were recorded in scientific literature. In the study by Poligne and Collignan (2000) the pH value of the marinated anchovies increased from 3.9 to 4.2 with storage time, but after 20 days of storage it remains constant until the end of storage. According to Dokuzlu (1997), shelf-life is ensured if the pH value of anchovy fillets after marinating remains around 3.85. Similar results were reported for fish marination processes by other authors (Aksu et al., 1997; Kilinc and Cakli, 2004; Šimat, 2010).

The addition of vinegar affected the sensory perception of salt content in the fillets. Although the salt content in the marinated fillets ranged from 3.7 to 7.2%, due to the acidity of certain products salinity was not described as an expressive organoleptic parameter. Such influence of acid in product suppressing the salinity was recorded in the literature, but also a reverse effect, depending on the concentration of salt and acid in the product (Poligne and Collignan, 2000; Aubourg and Ugliano, 2002). Breslin (1996) concluded that regardless of NaCl concentration in the solu-

Table 4. Volatile amine content, TBA index and proximate analysis of three marinated samples with best sensory scores after 25 days.

Tablica 4. Sadržaj hlapljivih amina, TBA indeks i kemijski sastav tri marinirana proizvoda s najboljim senzorskim ocjenama nakon 25 dana.

Analyzed samples Analizirani uzorci	Parameters determined Određeni parametri			Proximate analysis (%) Kemijski sastav (%)			
	TBAi (mg MA/kg)	TVB-N (mg/100g)	TMA-N (mg/100g)	Water content Udio vode	Protein content Udio bjelančevina	Fat content Udio masti	Ash content Udio pepela
13	7.36 (±0.24)	18.21 (±0.36)	0.74 (±0.21)	70.52 (±0.18)	21.82 (±0.14)	2.42 (±0.11)	2.68 (±0.09)
14	7.84 (±0.22)	18.75 (±0.42)	0.82 (±0.12)	70.68 (±0.21)	22.14 (±0.16)	2.96 (±0.16)	2.74 (±0.12)
16	6.41 (±0.28)	19.09 (±0.21)	0.79 (±0.18)	70.87 (±0.22)	22.99 (±0.12)	3.79 (±0.14)	2.93 (±0.09)
13 (45% water/voda, 30% vine vinegar/vinski ocat, 25% alcoholic vinegar/alkoholni ocat and 7% NaCl)							
14 (45% water/voda, 30% vine vinegar/vinski ocat, 25% alcoholic vinegar/alkoholni ocat and 10% NaCl)							
16 (45% water/voda, 55% vine vinegar/vinski ocat and 7% NaCl)							

tion, the presence of acid reduces the effect of salt.

The overall sensory acceptability of the products with regard to appearance, colour, odour, flavour (acidity and salinity), rancidity and aftertaste intensity, texture and juiciness was determined on days 2 and 25 of the marination (Figures 2. and 3.). The general appearance and colour of the fillets after marination were unsatisfactory in majority of the samples: the redness of the fillet along the bone line, brownish colouring in the belly area, and slimy yellowish colour of meat were observed. The reasons for this can be different: poor handling of fish prior to processing, a long period of storage of frozen anchovies and improper defrosting. The described marinating processes could not bleach these spots on the fillets. The lowest rated products were 7, 17 and 20. Sample 7 had the lowest proportion of vinegar (40%), while samples 17 and 20 were previously brined, and had no additional salt in the marinating bath. The best general appearance and colour were observed for sample 13.

The lowest scores for texture were given to products 17, 18, 20 (Figure 2.). The fillets were too soft and pliant due to the lack of salt in the marinating bath. The highest points

for texture were given to sample 2. The odour was rated as very good to excellent in all the samples (a slight smell of vinegar was noted) and rancidity was not present in any of the products. Samples 18 and 20 received the lowest scores for juiciness (3, neither good nor bad), while products 11 and 14, whose marinated baths contained 10% of NaCl were described as good. The best sensory scores on the 2nd day of marinating were given to two samples: 2 (60% of vinegar, 40% of water, 7% of NaCl) and 13 (30% of vinegar, 25% of vinegar, 45% of water, 7% of NaCl).

After 25 days in marinating baths, all the evaluated sensory characteristics of products have improved in relation to the analysis of the second day of marinating, except for the samples 17, 18 and 20 that completely disintegrated under the influence of acid (Figure 4.). Given the fact that salt provides the necessary firmness, the main reason for the poor condition of these three samples might be found in the usage of the marinating bath without salt, which resulted in the extraction of salt that was in the fillet (Table 3.). The colour and general appearance of samples 2, 3, 7, 9, 11, 13, 14 have improved, and the fillets were more compact, whiter, and less slimy after 25 days of storage. The remaining of the dark areas on the

belly and the central part of the fillets faded in most samples. The highest scores for the appearance of the fillets were given to samples 11 and 14 marinated in baths containing 10% of NaCl. This improved appearance can be correlated with the salt's ability to draw water and blood from the flesh. The brining of fish fillets or the addition of salt into the marination bath can enhance the sensory properties of marinated fish (Badii and Howell, 2002; Esaiassen et al., 2004; Šimat, 2010). After 25 days the texture was better evaluated, rancidity was not observed in any of the samples, and the smell of vinegar was milder and finer than on the second day. Juiciness and acidity were best scored in the samples with a mixture of vine and alcoholic vinegar (13 and 14). Off flavours were not observed in any of the samples, so the lowest scores for the aftertaste (product 9 - 60% of vinegar) was associated with the acidity of the product.

After 25 days of marination the best sensory score was given to samples 13 and 14. The marination baths of these two products contained 30% of vine vinegar, 25% of alcoholic vinegar, 45% of water and they differed in the NaCl content (13-7%, 14-10%). The maturity of the fish was identified by taste, consistency and colour, and by shiny, glassy appearance of ma-




Figure 2. Scores of sensory assessment of 11 selected products (see product description in Table 1.) two days after marination. The horizontal line set at 5.5 represents the rejection point.

Slika 2. Rezultati senzorske ocjene 11 odabranih proizvoda (opis proizvoda dan je u Tablici 1) dva dana nakon mariniranja. Vodoravna linija (5.5) predstavlja senzorsku granicu prihvatljivosti.




Figure 3. Scores of sensory assessment of 11 selected products (see product description in Table 1.) 25 days after marination. The horizontal line set at 5.5 represents the rejection point.

Slika 3. Rezultati senzorske ocjene 11 odabranih proizvoda (opis proizvoda dan je u Tablici 1) 25 dana nakon mariniranja. Vodoravna linija (5.5) predstavlja senzorsku granicu prihvatljivosti.

ture fish meat (Šoša, 1989). The ripening process has shown a great importance in cold marinade production. After 25 days of slow ripening at $4(\pm 2)$ °C the sensory properties of marinated products were improved and the maturation was finalized. These semi-products were then packed in oil with spices, and stored at refrigeration temperatures (from 0 to 8 °C) to provide high quality life of the marinades for several weeks (Shenderyuk and Bykowski, 1990).

Volatile amine content, TBA index and proximate analysis of three marinated samples with best sensory scores were determined after 25 days of marination (Table 4.). Nutritive quality of the anchovy fillets was preserved by cold marinating process. The total volatile accumulation was weak; however the increase in TVB-N content during storage indicates the existence of an enzymatic maturation of the fillets. Both TVB-N and TMA content were under the limit of

acceptance for seafood of 25- 35 mg TVB-N/100g and 5- 10mg TMA/100g (Anon., 2008).

Statistically significant increase of TBA index was recorded during maturation, but not over the limit of acceptance (Table 4.). Olgunoğlu et al. (2009) found the amount of 4.24 ± 0.01 mg malondialdehyde / kg in the anchovy fillets after 7 months of storage of marinated anchovies (4.5% of vinegar, 10% of salt and 0.2% of citric acid) at $1 (\pm 1)$ °C. Kilinc and Cakli (2004) have not recorded a significant increase in the amount of MA during the 22 days of marinating fillets of sardine (7% of acetic acid and 10% of NaCl). However, further storage of marinated sardine fillets in a mild solution (2% of acetic acid and 4% of NaCl) through 6 months at +4 °C showed an increase from 4.33 to 9.25 mg MA/kg of pasteurized, and from 4.47 to 9.49 mg MA/ kg in non-pasteurized samples. Despite the increasing amount during storage, sensory evaluation of marinated anchovies did not indicate rancidity as a negative attribute.

Before packing in oil and spices, marinated fillets are centrifuged, thus some water from the fillet gets removed. This slight increase of the pH (3.86) and the salt content (4.52) were observed after centrifugation, but no changes in water activity of the fillets.

Conclusions

The quality of raw material directly determines the quality of the final product and the technological process of cold marinating of the highest quality raw materials needed, given that it is a process without thermal treatment, dependent on the ripening process. Salt and acid ratios have an important role in determining the shelf-life of the product, as well as in defining the final sensory properties. The products received better sensory scores after 25 days of marination. Marinating time (maturation) has

also a key role in defining the sensory properties of cold marinated anchovies. The samples containing vine vinegar had better sensory scores. However, the best sensory scores were given to sample 13, marinated in a bath with 45% of water, 30% of vine vinegar, 25% of alcoholic vinegar and 7% of NaCl, which corresponds to the NaCl content of 3.60% in the fillet, water activity of 0.840 and a pH value of the fillet 3.09 after marination process.

References

- Aksu, H., N. Erkan, H. Colak, C. Varlik, N. Gökoğlu, M. Ugur** (1997): Some changes in anchovy marinades during production in different acid – salt concentration and determination of shelf life. *Int. J. Anim. Vet. Adv.* 8 (1-2), 86–90.
- Anonymous** (2008): Pravilnik o provedbenim mjerama za određene proizvode na koje se primjenjuju propisi o hrani (2008), Narodne novine broj 154, (N.N. 154/08).
- Anonymous** (2009): Fishing towards compliance with EU regulations, *EUROFISH Magazine*, 2, 28-49.
- Anonymous** (2010): The State of World Fisheries and Aquaculture, Food and Agriculture Organization (FAO), Rome, Italy (<http://www.fao.org/docrep/013/i1820e/i1820e00.htm>), pristupljeno 02.02.2011.
- A.O.A.C.** (2000): Official methods of analysis of AOAC International, 17th ed., Gaithersburg, Maryland, USA.
- Aubourg, S. P., M. Ugliano** (2002): Effect of brine pre-treatment on lipid stability of frozen horse mackerel (*Trachurus trachurus*). *Eur. Food Res. Tech.* 215, 91-95.
- Badii, F., N.K. Howell** (2002): Changes in the texture and structure of cod and haddock fillets during frozen storage. *Food Hydrocolloids*, 16(4), 313-319.
- Breslin, P.A.S.** (1996): Interactions among salty, sour and bitter compounds. *Trends. Food Sci. Tech.* 12(7), 390-399.
- Dalgaard, P.** (2000). Fresh and lightly preserved seafood. In: Shelf life Evaluation of Foods 2nd edition, Man C.M.D., A.A. Jones (Eds.), Aspen Publishing Inc., Maryland, USA, pp. 110-139.
- Derrick, S.** (2009): Cured, smoked and dried fish. In: Microbiology handbook-fish and seafood. Fernandes, R. (Ed.), Leatherhead Food International Ltd, Cambridge, UK. pp. 93-121.
- Dokuzlu, C.** (1997): The Effects of Acid-Salt Ratio Used in the Production of Marinated Anchovy to the Microbiologic, Organoleptic Quality and Determination of Shelf-Life of the Product. *Journal of Pendik Veterinary Microbiology*, 28, 81–90.
- Esaiassen, M., J. Østli, E.O. Elvevoll, S. Joensen, K. Prytz, R. Richardsen** (2004): Brining of cod fillets: influence on sensory properties and consumer liking. *Food Quality and Preference*, 15, 421-428.
- Fuselli, S.R., M.R. Casales, R. Fritz, M.I. Yeannes** (1994): Microbiology of the Marination Process used in Anchovy (*Engraulis anchoita*) Production. *LWT – Food Science and Technology*, 27(3), 214-218.
- Huss, H.** (1988): Fresh fish - quality and quality changes, Training Programme on Fish Technology and Quality Control, FAO Fisheries Series, FAO, Rome, pp. 132.
- Huss, H. H., L. Ababuo, L. Gram** (2003): Assessment and management of seafood safety and quality, FAO Fisheries Technical Paper No. 444. Rome, pp. 230.
- Kilinc, B., S. Cakli** (2004): Chemical, microbiological and sensory changes in thawed frozen fillets of sardine (*Sardina pilchardus*) during marination. *Food Chem.* 88, 275–280.
- Kristoffersen, S., T. Tobiassen, V. Stein-sund, R.L. Olsen** (2006): Slaughter stress, post-mortem muscle pH and rigor development in farmed Atlantic cod (*Gadus morhua* L.). *Int. J. Food Sci. Technol.* 41, 861–864.
- Kyryna, V.R., V.P. Lougovious** (2002): Sensory, chemical and microbiological assessment of farm raised European sea bass (*Dicentrarchus labrax*) stored in melting ice. *Int. J. Food Sci. Technol.* 37, 319-328.
- Lai, P.K., J. Roy** (2004): Antimicrobial and chemopreventive properties of herbs and spices. *Curr. Med. Chem.* 11, 1451-1460.
- Lemon, D.W.** (1975): An improved TBA test for rancidity. New Series Circular No. 51. Environment Canada Fisheries and Marine Service. Halifax Laboratory. Canada. pp. 1-4.
- Meyer, V.** (1965): Marinades. In: Fish as food, Vol. 3, Borgstrom, G. (Ed.), Academic Press Inc., New York, pp. 165–193.
- Olgunoğlu, I. A., F. Özogul, Y. Özogul, E. Kuley** (2009): Chemical, sensory and microbiological assessment of marinated anchovy (*Engraulis encrasicolus* L., 1758) fillets stored at 1±1°C. *Advances in Food Sciences*, 31(2), 1-8.
- Özogul, Y., F. Özogul, E. Kuley** (2010): Effects of combining of smoking and marinating on the shelf life of anchovy stored at 4°C. *J. Food Sci. Biotechnol.* 19(1), 69-75.
- Poligne, I., A. Colignan** (2001): Quick Marination of Anchovies (*Engraulis encrasicolus*) Using Acetic and Gluconic Acids. Quality and Stability of the End Product. *LWT – Food Science and Technology*, 33, 202–209.
- Pons-Sanchez-Cascado, S., M. C. Vidal-Carou, A. Mariné-Font, M.T. Veciana-Nogues** (2005): Influence of the freshness grade of raw fish on the formation of volatile and biogenic amines during the manufacture and storage of vinegar-marinated anchovies. *J. Agr. Food Chem.* 53, 8586-8592.
- Ray, B., A. Bhunia** (2008): Fundamental food microbiology, 4. Edition, CRC Press, Boca Raton, Florida, SAD, pp. 492.
- Sallam, K.I.** (2006): Chemical, sensory and shelflife evaluation of sliced salmon treated with salts of organic acids. *Food Chem.* 101, 592-600.
- Schormüller, J.** (1969): Handbuch der Lebensmittel Chemie. Band 4. Fette und Lipide, Lipids. Springer, Berlin, 872-878.
- Sen, M.K.C., S. Temelli** (2003): Microbiological and Chemical Qualities of Marinated Anchovy Prepared with Different Vegetable Additives and Sauce. *Revue. Méd. Vét.* 154, 703–707.
- Shenderyuk, V. I., P.J. Bykowski** (1990): Salting and marinating of fish. In: Seafood: resources, nutritional composition and preservation, Z. E. Sikorski (Ed.), CRC Press Inc., Boca Raton, Florida, pp. 58–63.
- Sikorski, Z. E., A. Kolakiwska, J.R. Burt** (1990): Postharvest biochemical and microbial changes. In: Seafood, Resources, nutritional composition and preservation. Sikorski Z. E. (Ed.), CRC Press. Inc. Boca Raton, Florida. pp. 256.
- Sinovčić, G.** (2000): Anchovy, *Engraulis encrasicolus* (LINNAEUS, 1758) biology, population dynamics and fisheries case study. *Acta Adriatica*, 41(1), 3-53.
- Šimat, V., J. Maršić Lučić, M. Tudor, I. Mladineo** (2009): Long-term storage influence on volatile amines (TVB-N and TMA-N) in sardine and herring utilized as food for tune fattening. *J. Appl. Ichthyol.* 25, 766-770.
- Šimat, V.** (2010): Promjene parametara kvalitete u filetu hladno mariniranog inćuna (*Engraulis encrasicolus*). Prehrambeno-

Učinak naljeva različitih koncentracija octa i soli na senzorska svojstva i parametare kvalitete hladno mariniranih inćuna (*Engraulis encrasicolus* L.)

Sažetak

Sinerijski učinak kiseline i soli ima ključnu ulogu u procesu mariniranja. Odabir pogodnih postupaka pripreme sirove ribe za mariniranje, sastava kupelji za mariniranje, sastava naljeva mariniranog proizvoda i uvjeta skladištenja, provode se u svrhu postizanja duže trajnosti, očuvanja kvalitete i postizanja dobrih senzorskih svojstava mariniranih proizvoda. U radu je provedeno istraživanje na 22 uzorka hladno mariniranih inćuna koji su sadržavali naljeve različitog sastava (različiti omjeri vinskog octa, alkoholnog octa, soli i vode), s kako bi se, tijekom zrenja, istražile promjene senzorskih svojstava i čimbenika koji određuju rok trajnosti. U tu svrhu određivana je pH vrijednost, aktivitet vode, udio NaCl i senzorska svojstva mariniranih fileta inćuna, kao i sadržaj hlapljivih amina i indeks oksidacije masti. Dobiveni rezultati ukazali su da je za postizanje najboljih senzorskih svojstava najprikladnija kupelj pripravljena od 45% vode, 30% vinskog octa, 25% alkoholnog octa i 7% soli, čime se postiže pH vrijednost mesa od 3.09, udio NaCl od 3.60% i aktivitet vode od 0.84. Zabilježena je slaba akumulacija hlapljivih amina tijekom skladištenja mariniranih proizvoda (<20mg TVB-N/100g; <1mgTMA/100g) i statistički značajan porast TBA indeksa (6.41-7.36 malondialdehida/kg) tijekom zrenja, ali užeglost fileta nije bila opisana kao intenzivno svojstvo.

Ključne riječi: *Engraulis encrasicolus*, hladna marinada, rok trajanja, TVB-N, TBAi

Wirkung des Aufgusses von Essig- und Salzkonzentration auf sensorische Eigenschaften und Qualitätsparameter kaltmarinierter Anchovis (*Engraulis encrasicolus* L.)

Zusammenfassung

Die synergische Wirkung von Säure und Essig hat die Schlüsselrolle im Marinadeprozess. Die Auswahl der richtigen Verfahrensvorgänge bei Vorberetung des rohen Fisches für das Marinieren, die Zusammensetzung des Marinadebades, die Zusammensetzung des Aufgusses des marinierten Erzeugnisses und die Lagerungsbedingungen werden durchgeführt, um eine verlängerte Dauerhaftigkeit, Qualitätsschutz und Erreichung der guten sensorischen Eigenschaften der Marinadeerzeugnisse zu erzielen. In der Arbeit wurde die Forschung auf 22 Mustern der kalt marinierten Anchovis durchgeführt, die Aufgüsse verschiedener Zusammensetzung enthielten (verschiedene Verhältnisse von Weinessig, Alkoholesig, Salz und Wasser), um Veränderungen der sensorischen Eigenschaften und Faktoren, die die Dauerhaftigkeit bestimmen, während der Reifezeit zu untersuchen. Zu diesem Zweck wurden pH-Wert, Wasseraktivität, Anteil von NaCl und sensorische Eigenschaften der marinierten Anchovisfilets, Inhalt der verdunstbaren Amine und Index von Oxydationsfetten bestimmt. Die bekommenen Resultate haben gezeigt, dass für die Erzielung der besten sensorischen Eigenschaften das Bad zusammengesetzt aus 45 % Wasser, 30 % Weinessig, 25 % Alkoholesig und 7 % Salz am geeignetsten ist, wobei der pH Wert des Fischfleisches von 3.09, Anteil des NaCl von 3.60 % und Wasseraktivität von 0.84 erreicht werden. Es wurde eine schwache Akkumulation von verdunstbaren Aminen während der Lagerungszeit der marinierten Erzeugnisse beobachtet (<20 mg TVB-N/100 g; < 1 mg TMA/100 g) und der bedeutende Aufstieg des TBA Indexes (6.41-7.36 Malondialdehyd/kg) während der Reifezeit, aber die Ranzigkeit der Filets wurde nicht als intensive Eigenschaft beschrieben.

Schlüsselwörter: *Engraulis encrasicolus*, kalte Marinade, Dauerhaftigkeit, TVB-N, TBAi

Effetto di aggiunta di differenti concentrazioni d'aceto e sale sulle caratteristiche e i parametri di qualità di acciuge marinate a freddo (*Engraulis encrasicolus* L.)

Sommario

L'effetto sinergico di acido e di sale fa un ruolo chiave nel processo di marinata. La scelta del procedimento di preparazione del pesce crudo destinato alla marinata, la composizione di marinata e delle condizioni di immagazzinamento si esibiscono con lo scopo di conseguire una durata più lunga, una qualità mantenuta e delle buone caratteristiche sensoriche di prodotti marinati. Quest'articolo presenta anche una ricerca fatta su 22 campioni di acciuge marinate a freddo, le cui marinata avevano contenuti diversi (percentuali differenti di aceto di vino, aceto alcolico, sale e acqua), per esaminare i cambiamenti di caratteristiche sensoriche e dei fattori che determinano la data di scadenza, durante la maturazione. Perciò si dovevano determinare il valore pH, attività dell'acqua, la percentuale di NaCl e le caratteristiche sensoriche di filetti di acciuge marinate, e anche il contenuto delle amine evaporanti e l'indice d'ossidazioni di grassi. I risultati ottenuti dichiarano che per ottenere le migliori caratteristiche sensoriche bisogna preparare la marinata che contiene il 45% d'acqua, il 30% di aceto di vino, il 25% di aceto alcolico e il 7% di sale. Così il valore della carne diventa il 3.09, la percentuale di NaCl il 3.60% e l'attività dell'acqua lo 0.84. Si nota una debole accumulazione delle amine evaporanti durante l'immagazzinamento dei prodotti marinati (< 20 mg TVB-N/100 g; < 1 mg TMA/100 g) e uno staticamente notevole aumento dell'indice TVB (6.41-7.36 di malondialdehide/kg) durante la maturazione, però il danno ossidativo di filetti non era descritto come una caratteristica intensa.

Parole chiave: *Engraulis encrasicolus*, marinata a freddo, data di scadenza, TVB-N, TBAi

biotehnoški fakultet, Sveučilišta u Zagrebu, Zagreb, pp.193.

Šoša, B. (1989): Higijena i tehnologija prerađivanja morske ribe. Školska knjiga, Zagreb, pp. 183.

Varlik, C., N. Erkan, S. Metin, T. Bayhar, Ö.

Özden (2000): Determination of the Shelf-life of Marinated Fish Balls. Turk. J. Vet. Anim. Sci. 24, 593-597.

Vyncke, W. (1970): Direct determination of the thiobarbituric acid value in trichloroacetic acid extracts of fish as a measure of oxidative rancidity. Fette-Seifen Anstrichmittel, 72(12), 1084-1087.

Zaika, L.L. (2007): Spices and herbs: their antimicrobial activity and its determination. J. Food Qual. 9(2), 97-118.

Yeannes, M.I., M.R. Casales (2008): Modifications in chemical compounds and sensorial attributes of *Engraulis anchoita* fillet during marination process. Cienc. Technol. Aliment. 28(4), 798-803.

Received: March, 7, 2011

Accepted: March, 25, 2011 