

Valparaiso University ValpoScholar

Physics and Astronomy Faculty Presentations

Department of Physics and Astronomy

9-1-2006

Comparison of Tropospheric Ozone Columns Calculated from MLS, OMI, and Ozonesonde Data

Gary A. Morris

Valparaiso University, gary.morris@valpo.edu

Bojan Bojkov

GEST- University of Maryland Baltimore County

Mark R. Schoeberl

NASA GSFC

Amy Wozniak

SAIC, NASA GSFC

Jerry Ziemke

GEST - University of Maryland Baltimore County

See next page for additional authors

Follow this and additional works at: http://scholar.valpo.edu/phys_astro_fac_presentations



Part of the [Astrophysics and Astronomy Commons](#), [Earth Sciences Commons](#), and the [Physics Commons](#)

Recommended Citation

Morris, Gary A.; Bojkov, Bojan; Schoeberl, Mark R.; Wozniak, Amy; Ziemke, Jerry; Chandra, Sushil; Fishman, Jack; and Stajner, Ivanka, "Comparison of Tropospheric Ozone Columns Calculated from MLS, OMI, and Ozonesonde Data" (2006). *Physics and Astronomy Faculty Presentations*. Paper 5.

http://scholar.valpo.edu/phys_astro_fac_presentations/5

This is brought to you for free and open access by the Department of Physics and Astronomy at ValpoScholar. It has been accepted for inclusion in Physics and Astronomy Faculty Presentations by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Authors

Gary A. Morris, Bojan Bojkov, Mark R. Schoeberl, Amy Wozniak, Jerry Ziemke, Sushil Chandra, Jack Fishman, and Ivanka Stajner

Abstract

This poster shows a comparison of three derived tropospheric ozone residual (TOR) products with integrated tropospheric ozone columns from ozonesonde profile: (1) the method of Ziemke *et al.* (2006), (2) a modified version of Fishman *et al.* (2003), and (3) a trajectory mapping approach. In each case, MLS ozone profiles are integrated to the tropopause and subtracted from OMI (TOMS retrieval) total column ozone. The effectiveness of each of these techniques is examined as a function of latitude, time, and geographic region. In general, we find good agreement between the derived products and the ozonesondes, with the Fishman *et al.* TOR (labeled "Amy") generally high and the Schoeberl trajectory mapping (labeled "Mark") product generally low as compared to the integrated ozonesonde profiles (labeled "Sonde") as computed using the WMO tropopause definition. Differences in TOR results are due, at least in part, to non-uniform tropopause height definitions between the three approaches.

The Three TOR Products

We compare three approaches to computing TOR. All three approaches subtract stratospheric column ozone (SCO), as computed by integrating stratospheric ozone profiles (SOP), from a column ozone measurement by OMI. The TOMS retrieval is used with the OMI data for total column ozone. We analyze the period August 2004 through July 2006

"Amy"

The first approach is based on the TOR of Fishman *et al.* [1990] and Fishman *et al.* [2003]. A model that assimilates SBUV SOP provides daily, gridded SCO to be subtracted from the level 3 OMI data. An ozone climatology fills in ozone values between lowest reliable SBUV observation and the tropopause. A standard WMO tropopause definition of 2K/km is used.

"Jerry"

The second approach is described in Ziemke *et al.* [2006]. TOR is determined using the residual technique of Fishman *et al.* [1990] by subtracting MLS stratospheric column ozone (SCO) from OMI total column ozone. An adjustment for inter-calibration differences of the two instruments (~ +3 DU), computed using the convective-cloud differential method of Ziemke *et al.* [1998], is included in the TOR. Gridded global maps of SCO from MLS at 0.25° [lat] 0.25° and 1° [lon] 1.25° resolution are produced daily using a 2D interpolation scheme.

These SCO fields are then subtracted from the total column ozone fields. The WMO tropopause is used with a maximum pressure of 316 hPa. The derived TOR product is filtered for cloud errors by rejecting scenes where OMI reflectivity is greater than 0.3.

"Mark"

The third approach creates a high resolution trajectory maps [Morris *et al.*, 1995] of total ozone residual using forward trajectory projections of the previous six days of MLS ozone data. The integrated, trajectory-mapped MLS ozone data are then subtracted from the level-3 OMI data in the manner of Morris *et al.* [1997]. In addition to the WMO tropopause definition, a 3.5 PVU criteria is applied in the extra tropics, usually resulting in a lower tropopause height and smaller TOR values.




Figure 1. This cartoon illustrates the tropospheric ozone residual calculation made by each of the three approaches presented in this poster. Note that currently, Amy uses an assimilated SBUV product rather than MLS.




Figure 2. Monthly mean tropospheric ozone residual as computed using the method of Ziemke *et al.* [2006, JGR in press]




Figure 3. MLS SCO produced with 6-day trajectories. Notice the improved resolution over the one day MLS data (black dots)


The Ozonesonde Stations

Ozonesonde data from the Aura Validation Data Center (AVDC) include most available soundings. Thirteen stations provided 551 profiles used in the analysis presented. The Table summarizes the stations and indicates the latitude groupings used in this study. We thank all of the teams involved in gathering and processing the ozonesonde data.

Station Name	Principal Investigator	Latitude	Longitude	# Profiles
Alert	D. Tarasick	82.50	-62.33	21
Ascension Island	A. Thompson	7.98	-14.42	69
Boulder	S. Oltmans	40.00	-105.25	91
Cotonou	A. Thompson	6.21	2.23	29
Egbert	D. Tarasick	44.23	-79.78	25
Hilo	S. Oltmans	19.43	-155.04	50
Houston (Rice U)	G. Morris	29.70	-95.40	77
Huntsville	M. Newchurch	35.28	-86.59	85
Irene	A. Thompson	-25.90	28.22	35
LaReunion	A. Thompson	-21.06	55.48	16
Sepang	A. Thompson	2.73	101.70	26
Summit	S. Oltmans	72.60	-38.50	11
WatuKosek	A. Thompson	-7.50	112.60	16

Time Series of TOR Products vs Ozonesondes

Figure 4. Below time series for 6 of the 13 stations showing the derived TOR products and the integrated ozonesonde profiles. All three TOR products generally follow the seasonal cycles well. Amy tends to be high, while Mark tends to be low. Significantly tighter distributions are found in the tropics where day-to-day variability is smaller and the tropopause definitions used by the three approaches tend to be the same.



TOR Products vs Integrated Ozonesondes

Figure 5. Scatter plots illustrate the relationships between the TOR products and the integrated ozonesonde profiles as a function of latitude. Dots are colored by the latitude groupings of the sonde stations (see Table at left). A regression fit and correlation coefficient accompany each plot. Jerry's approach produces the highest correlation coefficient and a regression slope nearest to the expected value of 1.0. All three approaches perform better in the tropics.





Figure 6. A probability distribution function for co-located TOR and ozonesonde data. Mean, standard deviation, and mode statistics are provided with each data set. All TOR products produce far more small values (< 20 DU) than are seen in the sonde data. Jerry and Mark match the mode of the sonde data well, while Amy does a better job on the high side of the distribution.



Conclusions and Future Work

This poster has presented a preliminary evaluation of three TOR products derived from SBUV, Aura MLS, and OMI measurements. In general, all three reproduce well the tropospheric columns computed from ozonesonde profiles and the observed seasonal cycles. More variability is seen in the TOR products at midlatitudes than in the tropics and than seen in the ozonesonde data. If OMI cannot see the surface, ozone pollution (e.g. Houston) will result in a low bias of the TOR data as compared to the sondes.

Future work will integrate all MLS profiles to 200 hPa in the tropics and 100 hPa elsewhere to eliminate problems associated with using various tropopause definitions. Amy's TOR will be updated with assimilated MLS replacing SBUV in SCO calculations.

References

- Fishman, J., et al., Distribution of tropospheric ozone determined from satellite data, *J. Geophys. Res.*, 95(D4), 3599-3617, 1990.
- Fishman, J., A.E. Wozniak, and J.K. Creilson, Global distribution of tropospheric ozone from satellite measurements using the empirically corrected tropospheric ozone residual technique: Identification of the regional aspects of air pollution, *Atmos. Chem. Phys.*, 3, 893 - 907, 2003.
- Morris, G.A., J. Gleason, S. Hollandworth, J. Ziemke, and M.R. Schoeberl, Constructing synoptic maps of stratospheric column ozone from HALOE and SAGE using trajectory mapping, *EOS Trans. AGU*, 78, 17, Spring Meeting Suppl., S90, 1997.
- Morris, G.A., M.R. Schoeberl, L. Sparling, P.A. Newman, L.R. Lait, L. Elson, J. Water, A.E. Roche, J. Kumer, and J.M. Russell, III, Trajectory mapping and applications to data from the Upper Atmosphere Research Satellite, *J. Geophys. Res.*, 100, 16,491 - 16,505, 1995.
- Thompson, A.M., et al. Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998 - 2000 tropical ozone climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and ground-based measurements, *J. Geophys. Res.*, 108 (D2), 8238, doi: 10.1029/2001JD000967, 30 January 2003.
- Von den Oord, G.H.J., et al. The OMI Monitoring Instrument, *IEEE Trans. Geos. Rem. Sens.*, 44, 1093 - 1101, 2006.
- Waters, et al., The Earth Observing System Microwave Limb Sounder (EOS MLS) on the Aura Satellite, *IEEE Trans. Geos. Rem. Sens.*, 44, 1075 - 1092, 2006.
- Ziemke, J. R., S. Chandra, and P. K. Bhartia, Two new methods for deriving tropospheric column ozone from TOMS measurements: The assimilated UARS MLS/HALOE and convective-cloud differential techniques, *J. Geophys. Res.*, 103, 22,115-22,127, 1998.
- Ziemke, J. R., et al., Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's Chemical Transport Model, *J. Geophys. Res.*, in press, 2006.

Funding for this research was provided in part by Goddard Earth Science Technology (GEST) grant NCC5-494 and a separate grant to Valparaiso University, both under the Aura validation and atmospheric composition program of NASA's Earth-Sun System Division. Thanks to the MLS, OMI, SBUV, and ozonesonde teams!