Valparaiso University ValpoScholar

Symposium on Undergraduate Research and Creative Expression (SOURCE)

Office of Sponsored and Undergraduate Research

5-3-2014

Development of a Solar Rotary-Kiln Reactor for the Reduction of Metal Oxide Particles

Ali AlNuaimi Valparaiso University

Adam Berry Valparaiso University

Courtney Brandt Valparaiso University

Jesse Fosheim Valparaiso University, jesse.fosheim@valpo.edu

Eric Loria Valparaiso University

See next page for additional authors

Follow this and additional works at: https://scholar.valpo.edu/cus

Recommended Citation

AlNuaimi, Ali; Berry, Adam; Brandt, Courtney; Fosheim, Jesse; Loria, Eric; Ogland-Hand, Jonathan; and Schrader, Andrew, "Development of a Solar Rotary-Kiln Reactor for the Reduction of Metal Oxide Particles" (2014). *Symposium on Undergraduate Research and Creative Expression (SOURCE)*. 305. https://scholar.valpo.edu/cus/305

This Poster Presentation is brought to you for free and open access by the Office of Sponsored and Undergraduate Research at ValpoScholar. It has been accepted for inclusion in Symposium on Undergraduate Research and Creative Expression (SOURCE) by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Authors

Ali AlNuaimi, Adam Berry, Courtney Brandt, Jesse Fosheim, Eric Loria, Jonathan Ogland-Hand, and Andrew Schrader

Development of a Solar Rotary-Kiln Reactor for the Reduction of Metal Oxide Particles

Ali AlNuaimi, Adam Berry, Courtney Brandt, Jesse Fosheim, Eric Loria, Jonathan Ogland-Hand, Andrew Schrader

Departmental Affiliation: Mechanical and Electrical Engineering College of Engineering

A solar rotary-kiln reactor has been fabricated for the reduction of metal oxide particles at ~1650 K as part of a solar thermal decoupled water electrolysis process for the production of hydrogen. Particle motion is controlled through the reactor's angular speed of rotation. At rotational speeds greater than 65 rpm, the internal walls of the reactor are fully covered with particles. Simultaneously, mixing elements generate a particle cloud in the reactor in order to increase the absorption of incident solar radiation. A model of the reactor that solves the energy conservation equation and includes the kinetics of the metal oxide reduction suggests that peak thermal efficiencies of 47 percent are possible for the reduction of hematite to magnetite. In parallel, the solid state kinetics for the reduction of cobalt oxide (a promising alternative to iron oxide) in a low oxygen partial pressure atmosphere has been determined. Reduction follows the shrinking core model and is initially limited by the rate of oxygen diffusion in the gas phase and later limited by the chemical kinetics at the shrinking reactive interface. Regression of the model to isothermal and non-isothermal thermogravimetric analyzer data yielded the temperature-dependent reaction rate parameters.

Information about the Authors: No information provided.

Faculty Sponsor: Dr. Luke Venstrom

Student Contact: jesse.fosheim@valpo.edu