

**Valparaiso University**  
**ValpoScholar**

---

Symposium on Undergraduate Research and  
Creative Expression (SOURCE)

Office of Sponsored and Undergraduate Research

---

5-3-2014

# Using $\pi^0$ 's to Understand Proton Spin Structure through Polarized p+p Collisions at $1 < \eta < 2$ and $\sqrt{s} = 200$ GeV at STAR

Stephen Place

Valparaiso University, [stephen.place@valpo.edu](mailto:stephen.place@valpo.edu)

Follow this and additional works at: <https://scholar.valpo.edu/cus>

---

## Recommended Citation

Place, Stephen, "Using  $\pi^0$ 's to Understand Proton Spin Structure through Polarized p+p Collisions at  $1 < \eta < 2$  and  $\sqrt{s} = 200$  GeV at STAR" (2014). *Symposium on Undergraduate Research and Creative Expression (SOURCE)*. 348.  
<https://scholar.valpo.edu/cus/348>

This Poster Presentation is brought to you for free and open access by the Office of Sponsored and Undergraduate Research at ValpoScholar. It has been accepted for inclusion in Symposium on Undergraduate Research and Creative Expression (SOURCE) by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

## Using $\pi^0$ 's to Understand Proton Spin Structure through Polarized p+p Collisions at $1 < \eta < 2$ and $\sqrt{s} = 200$ GeV at STAR

Stephen Place

*Departmental Affiliation:* Physics and Astronomy  
College of Arts and Sciences

Measurements of spin dependent observables at the Relativistic Heavy Ion Collider at Brookhaven National Laboratory provide unique insight into the contribution of a proton's constituents to its spin. The Solenoidal Tracker at RHIC (STAR) can be used to measure effects of the quark and gluon spins in the proton by observations of neutral pions that result from longitudinally and transversely polarized proton-proton collisions in STAR. The neutral pions ( $\pi^0$ 's) decay into two photons that, for pseudorapidity ( $\eta$ ) 1 to 2, can be observed in the endcap electromagnetic calorimeter in STAR. These are used to reconstruct the kinematic properties of the  $\pi^0$ 's and we can then look for spin asymmetries in  $\pi^0$  production. Measurements of both the longitudinal and transverse spin asymmetries in the production of  $\pi^0$ 's from data taken in 2006 have made some contributions to our understanding of the structure of the proton. New data taken in 2009, in a longitudinal spin run with greater luminosity, will provide greater precision in the final results. Results from 2006 (STAR's most recent spin paper, submitted to Phys. Rev. D: arXiv:1309.1800) and preliminary work on 2009 data are shown.

*Information about the Author:*

Stephen Place is a sophomore physics major.

*Faculty Sponsor:* Dr. Adam Gibson-Even

*Student Contact:* [stephen.place@valpo.edu](mailto:stephen.place@valpo.edu)