

# The Great Lakes Entomologist

---

Volume 48  
Numbers 3/4 -- Fall/Winter 2015 *Numbers 3/4 --  
Fall/Winter 2015*

---

Article 12

October 2015

## Macromia alleghaniensis (Odonata: Macromiidae): New For Michigan, with Clarifications of Northern Records

Julie A. Craves

University of Michigan, Dearborn, [jcraves@umich.edu](mailto:jcraves@umich.edu)

Darrin S. O'Brien

Prairie Oaks Ecological Station

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Craves, Julie A. and O'Brien, Darrin S. 2015. "Macromia alleghaniensis (Odonata: Macromiidae): New For Michigan, with Clarifications of Northern Records," *The Great Lakes Entomologist*, vol 48 (3)  
Available at: <https://scholar.valpo.edu/tgle/vol48/iss3/12>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

***Macromia alleghaniensis* (Odonata: Macromiidae):  
New For Michigan, with Clarifications of Northern Records**

Julie A. Craves<sup>1,2\*</sup> and Darrin S. O'Brien<sup>2</sup>

**Abstract**

An Alleghany River Cruiser, *Macromia alleghaniensis* Williamson (Odonata: Macromiidae), collected in Cass County, Michigan on 18 June 2014, represents the first record of the species for the state, as well as the northernmost unequivocal record in North America. Other records north of 40° latitude are clarified and discussed.

---

*Macromia* Rambur is a genus of medium to large (61–91 mm) dragonflies of 84 species worldwide (Garrison et al. 2006). The 7 species in North America are dark brown or black with yellow on the thorax and abdomen in patterns that vary, often subtly, between and among species, with multiple subspecies and hybrids recognized in the genus (Donnelly and Tennessen 1994, Cannings et al. 2006, Paulson 2011).

*Macromia alleghaniensis* Williamson (Odonata: Macromiidae), the Alleghany River Cruiser, is found most frequently in the central southeastern United States. Records extend west to southeastern Missouri, western Arkansas and nearby Oklahoma and Texas, and south to the Gulf coast along the Mississippi-Alabama border. Scattered records exist in Georgia, Florida, and South Carolina.

*Macromia alleghaniensis* is uncommon or rare in the northern part of its range. It is considered rare and a species of special concern in New Jersey (Barlow et al. 2009, NJDEP 2012), critically imperiled in Pennsylvania (PHNP 2014), uncommon and imperiled in Maryland (MNHP 2010, MAIFS 2014), rare in West Virginia (WVDNR 2012), and uncommon and a species of greatest conservation need in Ohio (Glotzhober and McShaffrey 2002, ODNR 2005). This species is listed as critically imperiled in Illinois by NatureServe (2015), but it is not listed in the most current state wildlife action plan or endangered and threatened species lists (IDNR 2005, Mankowski 2011).

*Macromia alleghaniensis* has not been recorded in New York, Delaware, Indiana, Wisconsin, or Ontario (Donnelly 1992, Curry 2001, White et al. 2010, Pratt 2012, NHIC 2014, WOS 2014, Abbott 2006–2015). Prior to our record (described below), it had not been recorded in Michigan (Byers 1927, Kormondy 1958, Donnelly 2004, MOS 2008–2014).

*Macromia alleghaniensis* is typically found on small to medium, slow flowing streams and rivers (Dunkle 2000, Garrison et al. 2006, Beaton 2007, Paulson 2011). The flight season in the northern portion of its core range (~ 38–39° latitude) is June to August (Paulson 2011).

On 18 June 2014, DO captured a male *Macromia* along Wood Creek in the Three Rivers State Game Area, Cass County, Michigan (41.861°, -85.763°).

---

<sup>1</sup>Rouge River Bird Observatory, University of Michigan-Dearborn, Dearborn, MI 48128.

<sup>2</sup>Prairie Oaks Ecological Station, 2200 Centennial Lane, Ann Arbor, MI 48103.

\*Corresponding author: (e-mail: jcraves@umich.edu).

The creek at this location is 2 to 4 m wide and 0.25 to 0.5 m deep, clear, with a sandy/gravelly substrate and silty edges near the bank undercuts. Habitat along this ~ 120 m stretch of creek was fen or marsh with forbs, sedge hummocks, and areas of cattails interspersed with occasional shrubs. The stream corridor was 60 to 80 m wide and bordered by forest.

The voucher was deposited in the University of Michigan Museum of Zoology, Insect Division (cataloged by the Michigan Odonata Survey as MOS0035050) and a photograph was submitted to OdonataCentral (OC#423318, Abbott 2006–2015). It was initially identified as *Macromia illinoiensis* Walsh, the common *Macromia* species in the state. It was subsequently correctly identified as *M. alleghaniensis* by the abdominal markings (the combination of a nearly complete yellow ring on abdominal segment 7 and a yellow ring on abdominal segment 2 interrupted only dorsally) and the short keel on the middle tibia, which distinguish it from males of other species of *Macromia* (Williamson 1909, Donnelly and Tennessen 1994, Paulson 2011, Needham et al. 2014). This represents Michigan's first record of this species, and the northernmost unequivocal record in North America.

### Discussion

The majority of published records of *Macromia alleghaniensis* are from south of 40°. Table 1 lists records north of that latitude that are equivocal, invalid, or unconfirmed. Other than our Michigan record, the only records confirmed by specimens north of 40° are from 3 counties in Pennsylvania—Huntingdon, Bucks, and Chester—with the Huntingdon County location being the farthest north at 40.580° (Shiffer and White 2014). Our Michigan location is roughly 140 km north of that latitude.

Because their habitats tend to be less ecologically stable or persistent, lentic species have a higher capacity for dispersal and are more prone to move long distances, especially due to climate-related events such as drought (Arribas et al. 2012, Hof et al. 2012, Grewe et al. 2013). This may explain the occurrence of several southern lentic dragonfly species in northern states in the past decade (Craves and O'Brien 2007, Mauffray 2008, Craves and O'Brien 2011).

Conversely, lotic species like most *Macromia* are less apt to disperse. At the Michigan collection site, we saw several other *Macromia*, possibly also *M. alleghaniensis*. This possibility, combined with the scarcity of records north of 40°, even in areas well-covered by dragonfly enthusiasts, suggests that the Michigan location perhaps represents an established disjunct population.

Accurate demarcation of species distributions is an important component in understanding how best to conserve biodiversity, especially in the face of climate change (Bush et al. 2014, Preuss et al. 2014). Historical records and their locations should be verified to provide accurate baseline data for future biogeographic comparisons. Care should be taken to record precise locations for new records, as the county-level data that is common in invertebrate databases tends to over-estimate geographic ranges (Collins 2014).

The lower dispersal rates of lotic species also promote genetic differentiation between populations, which in turn may advance adaptation to local conditions or foster speciation (Hof et al. 2006, Marten et al. 2006). Securing vouchers of out-of-range individuals and from disjunct populations, especially of taxa most reliably identified by structural characters or in which hybridization is common such as in *Macromia*, will not only confirm identification and validate records but can further our understanding of gene flow, adaptation, and evolution in these populations.

Table 1. Equivocal, unconfirmed, and invalid records of *Macromia alleghaniensis* north of 40° N latitude.

Latitude	State	Location	Date	Notes and Sources
42.32° (county center)	Illinois	McHenry County	Unknown	OdonataCentral record OC#229282 (Abbott 2006-2015). This and the next record are county-level records (Donnelly 2004). There are no associated vouchers or published data on these records, and details are unknown to state experts (ISM 2006, T. Cashatt and T. Vogt, pers. comm.).
42.06° (county center)	Illinois	Carroll County	Unknown	OdonataCentral record OC#229280 (Abbott 2006-2015). See above.
41.658°	Ohio	Ottawa County, Put-in-Bay	1934	Two males taken by C. H. Kennedy in 1934 (not 1937, as stated in Glotzhober and McShaffrey 2002) at the "Put-in-Bay Lake Laboratory" were discovered at the University of Michigan Museum of Zoology and designated as the first Ohio records (Kormondy 1956). Originally located on South Bass Island, this Lab (operated by the Ohio State University) was relocated to Gibraltar Island in 1925 and renamed the Stone Laboratory (Anon. 1925). Kennedy taught at each location from 1920-1938; it is unclear why he would use the incorrect name and not provide complete dates on the specimens, as he did with most of his material. These inconsistencies and the lack of stream or river habitats on these Put-in-Bay islands make the provenance of these specimens uncertain. It has been speculated they were raised in the lab (Glotzhober and McShaffrey 2002) or brought to Kennedy from elsewhere and mixed in with his own collections (C. Trisler, pers. comm.).
40.41° (county center)	Illinois	Hancock County	Unknown	County-level record shown online (Konratieff 2000) is apparently a transcription error; as the data reference is the Illinois State Museum database (ISM 2006) which does not have a record for Hancock Co., but does have one for Hardin Co., which was omitted on the map.
40.119°	Ohio	Franklin County Olentangy River, 0.5 mi N of I-270	23 May 1991	This very early instar nymph is noted as "ID ?" (Glotzhober et al. 1995, OHC 2015). Most Odonata larval keys are for use with mature nymphs, and the <i>Macromia</i> key typically used at the time this nymph was collected (Needham and Westfall 1955) was later found to be unreliable (Donnelly and Tennesen 1994).

### Acknowledgments

The authors thank Steve Hummel and Brenda D. Smith-Patten for correcting our misidentification; John Beetham, Bob Dubois, Tim Cashatt, Nick Donnelly, Bob Glotzhober, Boris Kondratieff, Bill Mauffray, Mark O'Brien, Dennis Paulson, Tim Vogt, and Bill Smith for help tracking down records and other data; and Carmen Trisler for important context and insight on Kennedy's records. Additional thanks to Mark O'Brien for his always helpful comments, encouragement, and general bonhomie.

### Literature Cited

- Abbott, J. C. 2006–2015.** OdonataCentral: an online resource for the distribution and identification of Odonata. Available from <http://www.odonatacentral.org> (accessed 4 February 2015).
- Anonymous. 1925.** The Franz Theodore Stone Laboratory. Ohio State University Bulletin 30(5): 1–18.
- Arribas, P., J. Velasco, P. Abellán, D. Sánchez-Fernández, C. Andújar, P. Calosi, A. Millán, I. Ribera, and D. T. Bilton. 2012.** Dispersal ability rather than ecological tolerance drives differences in range size between lentic and lotic water beetles (Coleoptera: Hydrophilidae). *Journal of Biogeography* 39: 984–994.
- Barlow, A. E., D. M. Golden, and J. Bangma. 2009.** Field Guide to the Dragonflies of New Jersey. New Jersey Department of Environmental Protection, Flemington, NJ.
- Beaton, G. 2007.** Dragonflies and Damselflies of Georgia and the Southeast. University of Georgia Press, Athens, Georgia.
- Bush, A., V. Hermoso, S. Linke, D. Nipperess, E. Turak, and L. Hughes. 2014.** Freshwater conservation planning under climate change: demonstrating proactive approaches for Australian Odonata. *Journal of Applied Ecology* 51: 1273–1281.
- Byers, F. C. 1927.** An annotated list of the Odonata of Michigan. Occasional Papers of the Museum of Zoology, University of Michigan 183: 1–15.
- Cannings, R. A., P. M. Catling, and P. M. Brunelle. 2006.** New subspecific status for *Macromia rickeri* Walker. *Argia* 17(4): 23.
- Collins, S. D. 2014.** Fine-scale modeling of riverine Odonata distributions in the northeastern United States. Ph.D. dissertation, Department of Biology, Texas Tech University, Lubbock.
- Craves, J. A., and D. S. O'Brien. 2007.** *Erythrodiplax umbrata* (Odonata: Libellulidae): new for Michigan. *The Great Lakes Entomologist* 40: 95–97.
- Craves, J. A., and D. S. O'Brien. 2011.** *Tramea calverti* (Odonata: Libellulidae): new for Michigan with notes on other new reports from the Great Lakes region. *The Great Lakes Entomologist* 44: 78–82.
- Curry, J. R. 2001.** Dragonflies of Indiana. Indiana Academy of Science, Indianapolis.
- Donnelly, T. W. 1992.** The Odonata of New York State. *Bulletin of American Odonatology* 1: 1–27.
- Donnelly, T. W. 2004.** Distribution of North American Odonata. Part II: Macromiidae, Corduliidae, and Libellulidae. *Bulletin of American Odonatology* 8: 1–32.
- Donnelly, T. W., and K. J. Tennessen. 1994.** *Macromia illinoiensis* and *georgina*: a study of their variation and apparent subspecific relationship (Odonata: Corduliidae). *Bulletin of American Odonatology* 2: 27–61.
- Dunkle, S. W. 2000.** Dragonflies Through Binoculars. Oxford University Press, New York.

- Garrison, R. W., N. Von Ellenrieder, and J. A. Louton. 2006.** Dragonfly Genera of the New World, An Illustrated and Annotated Key to the Anisoptera. John Hopkins University Press, Baltimore.
- Glotzhober, R. C., and D. McShaffrey. 2002.** Dragonflies and Damselflies of Ohio. Ohio Biological Survey, Columbus.
- Glotzhober, R. C., R. A. Restifo, T. E. Perry, and R. W. Alrutz. 1995.** New dragonfly (Odonata) species in Ohio, and additions to county records. Ohio Journal of Science 95: 233–239.
- Grewe, Y., C. Hof, D. M. Dehling, R. Brandl, and M. Brändle. 2013.** Recent range shifts of European dragonflies provide support for an inverse relationship between habitat predictability and dispersal. Global Ecology and Biogeography 22: 403–409.
- Hof, C., M. Brändle, and R. Brandl. 2006.** Lentic odonates have larger and more northern ranges than lotic species. Journal of Biogeography 33: 63–70.
- Hof, C., M. Brändle, D. M. Dehling, M. Munguía, R. Brandl, M. B. Araújo, and C. Rahbek. 2012.** Habitat stability affects dispersal and the ability to track climate change. Biology Letters 8: 639–643.
- (IDNR) Illinois Department of Natural Resources. 2005.** The Illinois Comprehensive Wildlife Conservation Plan & Strategy. Version 1.0. Springfield, Illinois. 353 pp.
- (ISM) Illinois State Museum. 2006.** ISM Online Illinois Dragonfly and Damselfly Collection. Available from <http://www.museum.state.il.us/ismdepts/zoology/odonata/display.html?Genus=Macromia&Species=alleghaniensis> (accessed 14 February 2015).
- Kondratieff, B. C. 2000.** Dragonflies and Damselflies (Odonata) of the United States. Northern Prairie Wildlife Research Center, Jamestown, North Dakota. Version 12 Dec 2003. Available from <http://www.npwrc.usgs.gov/resource/distr/insects/dfly/il/278.htm> (accessed 15 February 2015).
- Kormondy, E. J. 1956.** *Macromia alleghaniensis* Williamson in Ohio with new records of other Odonata in Ohio. Ohio Journal of Science 56: 203–204.
- Kormondy, E. J. 1958.** A catalogue of the Odonata of Michigan. Miscellaneous Papers of the Museum of Zoology, University of Michigan 104: 1–43.
- (MAIFS) Mid-Atlantic Invertebrate Field Studies. 2014.** Odonata of Maryland and Washington D.C. Version 10 Nov 2014. Available from <http://www.marylandinsects.com/DragonfliesDamselflies.html> (accessed 18 February 2015).
- Mankowski, A. 2011.** Endangered and Threatened Species of Illinois: Status and Distribution, Volume 4 — 2009 and 2010 Changes to the Illinois List of Endangered and Threatened Species. Illinois Endangered Species Protection Board, Springfield, Illinois. iii + 38 pp.
- Marten, A., M. Brändle, and R. Brandl. 2006.** Habitat type predicts genetic population differentiation in freshwater invertebrates. Molecular Ecology 15: 2643–2651.
- Mauffray, B. 2008.** *Dythemix velox* (Swift Setwing) in Indiana. Argia 20(3): 26.
- (MNH) Maryland Natural Heritage Program. 2010.** Rare, Threatened and Endangered Animals of Maryland. April 2010 edition. Maryland Department of Natural Resources, Wildlife and Heritage Service, Annapolis, Maryland. 25 pp.
- (MOS) Michigan Odonata Survey. 2008–2014.** Michigan Odonata Survey online database. University of Michigan Museum of Zoology, Insect Division, Ann Arbor. Available from <http://insectsdatabaser.ummz.lsa.umich.edu/mos/home.php> (accessed 10 February 2015).
- NatureServe. 2015.** NatureServe Explorer: an online encyclopedia of life. Version 7.1. NatureServe, Arlington, Virginia. Available from <http://explorer.natureserve.org> (accessed 19 February 2015).

- Needham, J. G., and M. J. Westfall, Jr. 1955.** A manual of the dragonflies of North America (Anisoptera). University of California Press, Berkeley, California.
- Needham, J. G., M. J. Westfall, Jr., and M. L. May. 2014.** Dragonflies of North America, 3<sup>rd</sup> edition. Scientific Publishers, Gainesville, Florida.
- (NHIC) Natural Heritage Information Centre. 2014.** Dragonflies and damselflies [of Ontario]. Version 30 July 2014. Available from <https://www.ontario.ca/environment-and-energy/get-natural-heritage-information> (accessed 18 February 2015).
- (NJDEP) New Jersey Department of Environmental Protection. 2012.** NJ Endangered and Nongame Species Program Special Concern — Species Status Listing. Version 21 February 2012. Available from [http://www.conservewildlifenj.org/downloads/cwnj\\_481.pdf](http://www.conservewildlifenj.org/downloads/cwnj_481.pdf) (accessed 13 February 2015).
- (ODNR) Ohio Department of Natural Resources. 2005.** Ohio's State Wildlife Action Plan. Odonata Species of Greatest Conservation Need. Available from <http://wildlife.ohiodnr.gov/portals/wildlife/pdfs/species%20and%20habitats/odonata%20table.pdf> (accessed 13 February 2015).
- (OHC) Ohio History Connection. 2015.** Online collections catalog: Museum Collections. Available from <http://www.ohiohistory.org/collections--archives/online-collections-catalog> (accessed 9 February 2015).
- Paulson, D. 2011.** Dragonflies and Damselflies of the East. Princeton University Press, Princeton, New Jersey.
- (PNHP) Pennsylvania Natural Heritage Program. 2014.** PNHP Species Lists. Version November 2014. Available from <http://www.naturalheritage.state.pa.us/species.aspx> (accessed 13 February 2015).
- Pratt, P. D. 2012.** Regional lists of Ontario Odonata. Available from <http://home.primus.ca/~naturalist/odonata.html> (accessed 18 February 2015).
- Preuss, S., M. Low, A. Cassel-Lundhagen, and A. Berggren. 2014.** Evaluating range-expansion models for calculating nonnative species' expansion rate. *Ecology and Evolution* 4: 2812–2822.
- Shiffer, C. N., and H. B. White. 2014.** Pennsylvania Odonata Records. Available from <http://dx.doi.org/10.6084/m9.figshare.1056508> (accessed 6 February 2015).
- White, E. L., J. D. Corser, and M. D. Schlesinger. 2010.** The New York Dragonfly and Damselfly Survey 2005–2009, Distribution and Status of the Odonates of New York. New York Natural Heritage Program, Albany, New York. 324 pp.
- Williamson, E. B. 1909.** The North American dragonflies (Odonata) of the genus *Macromia*. *Proceedings of the U.S. National Museum* 37: 369–398.
- (WOS) Wisconsin Odonata Survey. 2014.** Wisconsin Dragonfly and Damselfly Species. Version 21 January 2014. Available from <http://wiatri.net/inventory/odonata/SpeciesAccounts/> (accessed 6 February 2015).
- (WVDNR) West Virginia Department of Natural Resources. 2012.** Rare, Threatened And Endangered Species In West Virginia. Version 14 June 2012. Available from [http://www.wvdnr.gov/Wildlife/PDFFiles/RTE\\_Animals\\_2012.pdf](http://www.wvdnr.gov/Wildlife/PDFFiles/RTE_Animals_2012.pdf) (accessed 6 February 2015).