

Journal of Mind and Medical Sciences

Volume 4 | Issue 1

Article 10

2017

Melatonin modulates inflammatory response and suppresses burn-induced apoptotic injury

Ganka Bekyarova

Medical University-Varna, Department of Pathophysiology, Bulgaria, ganka.bekyarova@gmail.com

Milena Atanasova

Medical University- Pleven, Department of Biology, Bulgaria

Maria Tzaneva

Medical University-Varna, Department of General and Clinical Pathology, Bulgaria

Anelia Dimitrova

Medical University- Pleven, Department of Biology, Bulgaria

Follow this and additional works at: <http://scholar.valpo.edu/jmms>

 Part of the [Medical Physiology Commons](#)

Recommended Citation

Bekyarova, Ganka; Atanasova, Milena; Tzaneva, Maria; and Dimitrova, Anelia (2017) "Melatonin modulates inflammatory response and suppresses burn-induced apoptotic injury," *Journal of Mind and Medical Sciences*: Vol. 4 : Iss. 1 , Article 10.

DOI: 10.22543/7674.41.P5966

Available at: <http://scholar.valpo.edu/jmms/vol4/iss1/10>

This Research Article is brought to you for free and open access by ValpoScholar. It has been accepted for inclusion in Journal of Mind and Medical Sciences by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Research Article

Melatonin modulates inflammatory response and suppresses burn-induced apoptotic injury

Ganka Bekyarova¹, Milena Atanasova², Maria Tzaneva³, Anelia Dimitrova²

¹Medical University-Varna, Department of Pathophysiology, Bulgaria

²Medical University- Pleven, Department of Biology, Bulgaria

³Medical University-Varna, Department of General and Clinical Pathology, Bulgaria

Abstract

Introduction: Melatonin, the principal secretory product of the pineal gland, has antioxidant functions as a potent antioxidant and free radical scavenger. **Objectives** of the present study were to investigate the effect of melatonin against inflammatory response, burn-induced oxidative damage and apoptotic changes of rat liver. **Methods:** Melatonin (10 mg /kg, i.p.) was applied immediately after 30% of total body surface area (TBSA) burns on male Wistar rats. The level of malondialdehyde (MDA) as a marker of an oxidative stress was quantified by thiobarbituric method. Hepatic TNF α and IL-10 as inflammatory markers were assayed by ELISA. Using light immunohistochemistry the expression Ki67 proliferative marker was investigated. **Results:** Hepatic MDA and TNF- α levels increased significantly following burns without any change in IL-10 level. Intracellular vacuolization, hepatic cell degeneration and apoptosis occurred in rats after burns. The number of apoptotic cells was increased whereas no significant increase in Ki67 proliferative marker. Melatonin decreased the MDA and TNF- α content and increased the IL-10 level. It also limited the degenerative changes and formation of apoptotic cells in rat liver but did not increase expression of the marker of proliferation. **In conclusion**, our data show that melatonin relieves burn-induced hepatic damage associated with modulation of the proinflammatory/anti-inflammatory balance, mitigation of lipid peroxidation and hepatic apoptosis.

Keywords: melatonin, cytokines, lipid peroxidation, apoptosis, liver, burn



Correspondence should be addressed to: Ganka Bekyarova; e-mail: ganka.bekyarova@gmail.com

Introduction

Severe burn is still a serious clinical problem in emergency medicine causing damages to organs distant from the burn wound (1-3). Pathophysiology of burn-induced liver injury includes many mechanisms and is not entirely defined yet. Inflammation is proposed to be the main event in the pathogenesis of thermal trauma (4-6). The inflammatory cascade has counter balancing factors that maintain a delicate balance of pro- and anti-inflammatory mediators that regulate cellular homeostasis (7-8). Many pro-inflammatory and anti-inflammatory cytokines are involved in the pathogenesis of thermal trauma having overlapping, antagonistic and synergistic effects (9). Tumor necrosis factor-alpha (TNF- α), interleukin-10 (IL-10) and C-reactive protein are thought to play an essential role in the modulation of immunological and inflammatory processes.

Hepatic apoptosis has been recognized as part of the pathological changes after burn injury (10). High apoptotic cell number after burn injury implicates hepatic damage and impairment to its function. Cytokines and oxidative stress are well known to induce hepatocytes apoptosis (11). Activation of NF- κ B by hypoxia induced endothelial cell death and apoptosis in Bcl-2-dependent manner (12).

Melatonin (N-acetyl-5-methoxytryptamine) is an indoleamine endogenously produced in the pineal gland and extrapineal tissues, including liver (13). Melatonin possesses a wide variety of biologic effects. Particularly, it has sedative, anxiolytic, and anti-nociceptive effects. Melatonin and its metabolites have potent antioxidant /anti-inflammatory properties and have been highly effective in various diseases linked to inflammation and oxidative stress (14). Melatonin not only neutralizes reactive oxygen species (ROS) and reactive nitrogen species (RNS) but acts through stimulation of antioxidant systems and stabilizes cell membranes. It modulates the expression of several protective enzymes

and reduces apoptosis and lipid peroxidation (15). There are no data about protection of melatonin against burn-induced apoptotic injury after burns.

The aim of the present study was to investigate the effect of melatonin on inflammatory response and burn-induced oxidative damage and apoptotic changes of rat liver.

Materials and Methods

- *Animals*

The experimental procedure was approved by the Home Office for Care and Use of Laboratory Animals, and experiments were performed in accordance with the European Communities Council Directives 86/609/EEC. Age-matched male rats weighing between 220 and 250 g fasted for 12 h were allowed free access to water before injury. Animals were housed in 20°C and offered rat food and water ad libitum. They were kept in dark-light cycles (DL = 12:12 hours) in individual wire-bottomed cages. Thus, lights were turned off at 8:00 p.m. and turned on at 8:00 a.m. for achieving satisfactory photoperiod.

- *Thermal injury and melatonin treatment*

After light ether inhalation, general anesthesia was performed using thiopental (30 mg/kg i.p.). In order to accomplish 30% of third degree burn hot boiling water (98°C) was applied on the back of the animals during a period of 10 sec. For those rats which were subjected to burn injury 4 ml of physiological saline was applied i.p. for immediate resuscitation following burn injury. No animals died within the first 24 hour post-burn period. Twenty-four male rats were randomly assigned to three groups of 8 animals each: control - non-burned, non-treated (C), vehicle-treated burned group (B), and melatonin-treated burned group (B+Mel).

Either melatonin (N-acetyl-5-methoxytryptamine, Merck, Germany) in a dose of 10 ml/kg body weight (b.w.) solved in vehicle, or vehicle (ethyl alcohol 2% in physiological saline in dose 5 ml/kg) was administered,

respectively. Melatonin and vehicle were applied immediately i.p. after burns in the morning between 8:00 a.m. and 9:00 a.m. and 12 hours after burns. All animals were given buprenorphine (0,3 mg /kg i.p.b.w. twice daily) for pain control post burn. They were re-anesthetized with thiopental and sacrificed 24 hours after burns as liver was sampled.

- *Biochemical analysis*

Blood was taken from the jugular vein and heparinized. Plasma was separated by centrifugation at 800 x g rpm for 10 min and aliquots were stored at -80oC until analysis. Liver was gently separated from the underlying tissue and homogenized in 1:5 w/v 50 mM phosphate buffer (pH 7,4) containing 0,1 mM EDTA, at 4000 rpm for 10 min. The homogenate was centrifuged at 800 x g rpm/15 min to discard the sediment and supernatant was frozen until analysis. All manipulations were performed at 4-8oC. Analysis was carried out immediately after thawing of the samples.

Membrane lipid peroxidation was assayed by MDA measured by its thiobarbituric acid (TBA) reactivity of hepatic homogenate using the method of Porter et al (16). Results were expressed as nmol MDA/g protein. They were determined using the extinction coefficient of MDA-TBA complex at 532 nm =1, 56 x 10⁻⁵ cm⁻¹ M⁻¹ solution.

- *Immunohistochemistry*

Rat liver specimens were fixed in 10% neutral buffered formalin and embedded in paraffin. The deparaffinized and rehydrated sections (5 µm thick) were treated with 1% hydrogen peroxide for peroxidase activity inhibition for 5 min. Then they were rinsed in 0.1 M phosphate buffered saline (PBS) (pH 7.4) and treated in normal goat serum for 20 min. Subsequently, the sections were incubated with polyclonal primary antibody for 24 hours at room temperature. Monoclonal Mouse Anti-Rat Ki-67 Antigen (DAKO, USA) was

used. After rinsing with PBS the sections were incubated for 20 min in goat anti-rabbit immunoglobulins at room temperature. Then they were rinsed in PBS again and treated with rabbit peroxidase-anti-peroxidase complex for 20 min at room temperature and then rinsed in PBS. Finally, peroxidase activity was estimated by the diaminobenzidine-tetrachloride H₂O₂-method.

Negative controls. Controls were incubated with non-immune sera instead of primary antibody.

- *Histopathological Examinations*

Tissue specimens were fixed in 10% buffered formalin (pH 7.2), dehydrated in ascending series of ethyl alcohol (70% - 100%), cleared in methyl benzoate and embedded in paraffin wax. Tissue sections of 5 µm were stained with hematoxylin and eosin (H&E) and examined using light microscope (Olympus BH-2, Tokyo, Japan). Apoptotic SECs were examined by light microscopy at a magnification of 400× (high power field). Approximately 500 SECs were examined in each high power field. The apoptotic index was defined as the number of stained cells per high power field.

- *Statistics*

Liver enzyme data were log-transformed to satisfy the assumptions required to perform parametric tests and, therefore, presented as geometric mean and 95% confidence intervals of the mean. Orthogonal contrasts in ANOVA were used to statistically analyze the difference between any two specified groups.

Results

Changes in the Hepatic Malondialdehyde and Hepatic TNF-α and IL-10 Levels

Hepatic TNF-α content was significantly increased by 112% (p<0.05) at 24 h in the burned group. TNF-α content was increased in the burned group treated with melatonin g, but the increase was significantly lower than in the burned group (Fig. 1).




Fig 1. Effect of melatonin on hepatic TNF level in the early post burn period. C-control, non-burned; B- burned group , and B+Mel- melatonin-treated burned group. Results are given as mean±SEM; * $p < 0.05$ C vs B; † $p < 0.05$ B+mel vs B.

The hepatic IL-10 content tended to increase in both groups following burn, but was significantly higher by 44% ($p < 0.01$) in the burned group treated with melatonin than in untreated burned group (Fig. 2). Hepatic MDA content persistently increased 166% ($p < 0.01$) at 24 h and was still significantly higher than the control level. However, MDA content was significantly lower in the burned group treated with melatonin (Fig.3).




Fig 3. Effect of melatonin on hepatic MDA level in the early post burn period

*** $P < 0.01$ B vs C; † $p < 0.05$ B+Mel vs C; †† $p < 0.05$ B+Mel vs B

A positive correlation was established between hepatic MDA and hepatic TNF α levels ($r = 0.50$, $p < 0.05$) (Fig. 4).




Fig 2. Effect of melatonin on hepatic IL-10 level in the early post burn period ** $P < 0.01$; B vs B+mel; † $p < 0.05$ B+Mel vs C

Histological (H&E) changes in liver

In the control group, hepatocytes had a normal shape and size with vesicular nuclei in (Fig. 5). In the burned group, hepatocytes showed an abnormal structure, acidophilic degeneration and vacuolization. In addition, an increased number of apoptotic cells were detected. In the burned group treated with melatonin, a reduced number of apoptotic cells was established.




Fig 4. Correlation between hepatic MDA and hepatic TNF- α levels after burn

Changes in the Expression on Ki-67-proliferative marker. The data showed single Ki67 positive cells in hepatocytes around the central vein of the control group. K67 proliferative marker was not changed in the untreated burned group and the burned group treated with melatonin.




Fig 5. Effect of melatonin on histological changes in liver in the early post burn period. Original magnification, 400. H & E staining, original magnification. Arrows indicate apoptotic cells

Discussion

Data from the present study show activation of inflammatory response, lipid peroxidation, degenerative damage and apoptosis in liver after thermal trauma. Melatonin inhibits inflammatory response and protects against hepatic oxidative damage and apoptosis.

Levels of TNF- α (a proinflammatory mediator) and IL-10 (an antiinflammatory mediator) in liver tissue increased significantly following severe burn (17). Our data demonstrate that TNF- α plays an important role in the occurrence and development of postburn liver damage (18). In vitro experiments demonstrated that TNF- α may directly act on rat liver cells. The application of TNF- α inhibitor prevents hepatic dysfunction in shock after burn in a guinea pig model (19).

IL-10 may inhibit inflammatory mediators such as TNF- α , upregulates humoral immune reaction and inhibits cellular immune reaction such as antigen presentation by macrophages (20). The imbalance between TNF- α and IL-10 expression may be one of the initiating factors for the development of systemic inflammatory response syndrome and multiple organ failure in the early after burn period (21).

High TNF α level activates cell-death pathways and promotes the production of reactive oxygen species and lipid peroxidation and thus hepatic injury (22). Furthermore, hepatocytes are much more sensitive to TNF- α and free radicals damaging effect when antioxidant capacity is low.

Liver tissue is subjected to severe peroxidation damage due to ischaemia/ hypoxia following severe burn (23). In the present study, levels of MDA in liver tissue were significantly increased, demonstrating severe lipid peroxidation in liver tissue following severe burn. MDA content was significantly decreased in the burned group treated with melatonin. However, this suggests that melatonin attenuates free radical activated burn-induced lipid peroxidative injury of hepatic tissue.

It appears that oxidative stress plays an important role in the primary cell and tissue destruction, and thus the secondary inflammatory reaction. ROS/RNS may activate the transcription factor NF- κ B and increase the level of TNF α , which has a cytotoxic activity and may cause hepatic injury. On the other hand, increased production of TNF α and free radicals increases the expression of transcription factor NF- κ B, likely forming a vicious cycle between oxidative stress and inflammation by the activation of NF- κ B. In support of this idea is the correlation between levels of TNF α and MDA in the liver after thermal skin injury ($r = 0.50$, $p < 0.05$) (Fig.4). Such a positive relationship between markers of lipid peroxidation and inflammation in liver was found in other experimental models of thermal trauma (24-26).

In the present study melatonin decreased the hepatic tissue TNF- α and MDA levels. In addition, it increased liver tissue IL-10. It may be assumed that the protective effect of melatonin against burn-induced oxidative injury

is associated with modulation of the pro-inflammatory/anti-inflammatory balance.

Melatonin significantly decreases the TNF- α and increases the IL-10 content in liver tissue in other pathological conditions (27-28) and this effect may be due to its anti-inflammatory properties. Melatonin has the capability of scavenging both oxygen and nitrogen-based reactants including ONOO- and blocking transcriptional NF-kB factor which induces pro-inflammatory cytokines (29). Thus melatonin affects the pro- and anti-inflammatory balance and restricts hepatic damage from uncontrollable inflammation, and this may be the main mechanism for the protective effect of melatonin against burn-induced oxidative damage.

Our data demonstrate that thermal injury induces degenerative changes in liver and increases number of apoptotic cells in this organ. Oxidative damage of mitochondria causes an elevation of pro-apoptotic Bax protein and of anti-apoptotic Bcl-2 protein in various diseases related with oxidative stress (30). Hence high temperature causes higher Bax/Bcl-2 ratio, which was similar to previous findings, and also elevated BAX/BCL-2 ratio suggesting the susceptibility of these cells to apoptosis (31). Proinflammatory cytokine TNF α triggers extrinsic apoptotic pathway through the binding of TNF α with cell-death receptors. The increased Bax proapoptotic protein expression may associate with translocation of proapoptotic Bid protein to mitochondria by the Fas/TNF-R1 pathway (29)

The present study shows that melatonin restricts burn-induced degenerative changes modulation of the pro-inflammatory/anti-inflammatory balance and decreased the number of apoptotic cells in liver. Melatonin attenuates the oxidative stress, activates antiapoptotic-regulating proteins and protects against apoptosis (15). Treatment with melatonin increases the glutathione synthesis and antioxidant defense and

modulates important steps in both mitochondrial and the death receptor apoptotic pathway.

The current study shows that the proliferative marker K67 was not changed in the thermal injured rats compared to the controls. Previous studies showed that the proliferation of hepatocytes increased on the second day after experimental thermal trauma (32). Probably, differences in experimental models of thermal injury can explain the difference in the results. Our data show that melatonin inhibits apoptosis of hepatocytes, but it does not stimulate their proliferation in the acute period after burns.

Conclusions

In conclusion, our data show that thermal trauma activates proinflammatory processes, causes oxidative damage and apoptosis in liver. The protection afforded by melatonin against burn-induced hepatic apoptotic injury probably occurs through modulation of the balance pro/anti-inflammatory cytokines and mitigation of lipid peroxidation.

References

1. Bertin-Maghit M, Goudable J, Dalmas E, Steghens JP, Bouchard C, Gueugniaud PY, Petit, P, Delafosse B. Time course of oxidative stress after major burns. *Intensive Care Med.* 2000; 26(6): 800-3.
2. Lausevic Z, Lausevic M, Trbojevic-Stankovic J, Krstic S, Stojimirovic B. Predicting multiple organ failure in patients with severe trauma. *Can J Surg.* 2008; 51(2): 97-102.
3. Jeschke MG. Hepatic response to thermal injury: Is the liver important for postburn outcomes. *Mol Med.* 2009; 15(9-10): 337-51.
4. Agay D, Andriollo-Sanchez M, Claeysen R, Touvard L, Denis J, Roussel AM, Chancerelle Y. Interleukin-6, TNF-alpha and interleukin-1 beta

- levels in blood and tissue in severely burned rats. *Eur Cytokine Netw.* 2008; 19(1): 1-7.
5. Bankey PE, Williams JG, Guice KS, Taylor SN. Interleukin-6 production after thermal injury: evidence for nonmacrophage sources in the lung and liver. *Surgery.* 1995; 118(2): 431-8.
 6. Finnerty CC, Herndon DN, Przkora R, Pereira CT, Oliveira HM, Queiroz DM, Rocha AM, Jeschke MG. Cytokine expression profile over time in severely burned pediatric patients. *Shock.* 2006; 26(1): 13-9.
 7. Despond O, Proulx F, Carcillo JA, Lacroix J. Pediatric sepsis and multiple organ dysfunction syndrome. *Curr Opin Pediatr.* 2001; 13(3): 247-53.
 8. Schneider CP, Schwacha MG, Chaudry IH. The role of interleukin-10 in the regulation of the systemic inflammatory response following trauma-hemorrhage. *Biochim Biophys Acta.* 2004; 1689(1): 22-32.
 9. Jeschke MG, Einspanier R, Klein D, Jauch KW. Insulin attenuates the systemic inflammatory response to thermal trauma. *Mol Med.* 2002; 8(8): 443-50.
 10. Jeschke MG, Kraft R, Song J, Gauglitz GG, Cox RA, Brooks NC, Finnerty CC, Kulp GA, Herndon DN, Boehning D: Insulin protects against hepatic damage postburn. *Mol Med.* 2011; 17(5-6), 516-522.
 11. Wang K. Molecular mechanisms of hepatic apoptosis regulated by nuclear factors. *Cell Signal.* 2015; 27(4):729-38.
 12. Matsushita H, Morishita R, Nata T, Aoki M, Nakagami H, Taniyama Y, Yamamoto K, Higaki J, Yasufumi K, Ogihara T: Hypoxia-induced endothelial apoptosis through nuclear factor-kappaB (NF-kappaB)-mediated Bcl-2 suppression: in vivo evidence of the importance of NF-kappaB in endothelial cell regulation. *Circ Res.* 2000; 86(9): 974-81.
 13. Reiter RJ, Tan DX, Maldonado MD. Melatonin as an antioxidant: physiology versus pharmacology. *J Pineal Res.* 2005; 39(2): 215-6.
 14. Jung KH, Hong SW, Zheng HM, Lee DH, Hong SS. Melatonin downregulates nuclear erythroid 2-related factor 2 and nuclear factor-kappaB during prevention of oxidative liver injury in a dimethylnitrosamine model. *J Pineal Res.* 2009; 47(2):173-83.
 15. Radogna F, Cristofanon S, Paternoster L, D'Alessio M, De Nicola M, Cerella C, Dicato M, Diederich M, Ghibelli L. Melatonin antagonizes the intrinsic pathway of apoptosis via mitochondrial targeting of Bcl-2. *J Pineal Res.* 2008; 44(3): 316-25.
 16. Porter NA, Nixon J, Isaac J. Cyclic peroxidase and thiobarbituric assay. *Biochim Biophys Acta.* 1976; 441: 596-599.
 17. Gauglitz GG, Song J, Herndon DN, Finnerty CC, Boehning D, Barral JM, Jeschke MG. Characterization of the inflammatory response during acute and post-acute phases after severe burn. *Shock.* 2008; 30(5): 503-7.
 18. Bekyarova G, Apostolova M, Kotzev I. Melatonin protection against burn-induced hepatic injury by down-regulation of nuclear factor kappa B activation. *Int J Immunopathol Pharmacol.* 2012; 25(3): 591-6
 19. Avlan D, Taşkinlar H, Tamer L, Camdeviren H, Ozturhan H, Oztürk C, Aksöyek S. Protective effect of trapidil against oxidative organ damage in burn injury. *Burns.* 2005; 31(7): 859-65.
 20. Sabat R. IL-10 family of cytokines. *Cytokine Growth Factor Rev.* 2010; 21(5): 315-24.
 21. Jaffer U, Wade RG, Gourlay T. Cytokines in the systemic inflammatory response syndrome: a

- review. *HSR Proc. Intensive Care Cardiovasc Anesth.* 2010; 2(3): 161-75.
22. Sun BW, Sun Y, Sun ZW, Chen X. CO liberated from CORM-2 modulates the inflammatory response in the liver of thermally injured mice. *World J Gastroenterol.* 2008; 14(4): 547-53.
23. Bekyarova G, Galunska B, Ivanova D, Yankova T. Effect of melatonin on burn-induced gastric mucosal injury in rats. *Burns.* 2009; 35(6): 863-8.
24. Sener G, Sehirlı AO, Gedik N, Dülger GA. Rosiglitazone, a PPAR-gamma ligand, protects against burn-induced oxidative injury of remote organs. *Burns.* 2007; 33(5): 587-93.
25. Chen XL, Dodd G, Thomas S, Zhang XL, Wasserman MA, Rovin BH, Kunsch C. Activation of Nrf2/ARE pathway protects endothelial cells from oxidant injury and inhibits inflammatory gene expression. *Am J Physiol Heart Circ Physiol.* 2006; 290(5): H1862-70.
26. Wu WS, Chou MT, Chao CM, Chang CK, Lin MT, Chang CP. Melatonin reduces acute lung inflammation, edema, and hemorrhage in heatstroke rats. *Acta Pharmacol. Sin.* 2012; 33(6): 775-82.
27. Oliveira LG, Kuehn CC, dos Santos CD, Miranda MA, da Costa CM, Mendonça, VJ, do Prado Júnior JC. Protective actions of melatonin against heart damage during chronic Chagas disease. *Acta Trop.* 2013; 128(3): 652-658.
28. Mauriz JL, Collado PS, Veneroso C, Reiter RJ, González-Gallego J. A review of the molecular aspects of melatonin's anti-inflammatory actions: recent insights and new perspectives. *J Pineal Res.* 2013; 54(1): 1-14.
29. Acuña Castroviejo D, López LC, Escames G, López A, García JA, Reiter RJ. Melatonin-mitochondria interplay in health and disease. *Curr Top Me. Chem.* 2011; 11(2): 221-240.
30. Somal A, Aggarwal A, Upadhyay RC. Effect of thermal stress on expression profile of apoptosis related genes in peripheral blood mononuclear cells of transition Sahiwal cow. *Iran J Vet Res.* 2015; 16(2): 137-43.
31. Yin XM. Bid, a critical mediator for apoptosis induced by the activation of Fas/TNF-R1 death receptors in hepatocytes. *J Mol Med (Berl).* 2000; 78(4): 203-11.
32. Jeschke MG, Low JF, Spies M, Vita R, Hawkins HK, Herndon DN, Barrow RE. Cell proliferation, apoptosis, NF-kappaB expression, enzyme, protein, and weight changes in livers of burned rats. *Am J Physiol Gastrointest Liver Physiol.* 2001; 280(6): G1314-20.