

December 1977

An Annotated List of Phytophagous Insects Collected on Immature Black Walnut Trees in Southern Illinois

P. L. Nixon
Southern Illinois University

J. E. McPherson
Southern Illinois University

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Nixon, P. L. and McPherson, J. E. 1977. "An Annotated List of Phytophagous Insects Collected on Immature Black Walnut Trees in Southern Illinois," *The Great Lakes Entomologist*, vol 10 (4)
Available at: <https://scholar.valpo.edu/tgle/vol10/iss4/8>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

AN ANNOTATED LIST OF PHYTOPHAGOUS INSECTS COLLECTED ON IMMATURE BLACK WALNUT TREES IN SOUTHERN ILLINOIS¹

P. L. Nixon and J. E. McPherson²

ABSTRACT

An annotated list of phytophagous insects on immature black walnut in southern Illinois was compiled between 26 April, 1974, and 9 October, 1975. Approximately 300 species, in 10 orders, were collected by hand-picking and sweeping. Notes taken on the various species included types of feeding damage, instars present, predators and parasites, and distribution in southern Illinois. Lepidoptera (about 80 species collected) were responsible for the majority of damage observed.

This study was conducted to compile an annotated list of the phytophagous insects found on immature black walnut trees (*Juglans nigra* L.) in southern Illinois. To our knowledge, the only other general list of black walnut insects is that by Barrett (1932) which was compiled from the literature. Notes were gathered on types of feeding damage, instars present, predators and parasites, and distribution in southern Illinois.

Black walnut grows for about 10 years before it is ready for earliest commercial seed production. Insect damage during this time is particularly important because the tree produces the first 9 to 17 feet of trunk height. This trunk must grow at a constant rate and be straight for veneer log production, the highest quality wood (Carmean, 1970).

In 1968, the north central region of the United States (i.e., Indiana, Illinois, Iowa, Missouri, Kansas, Nebraska, Michigan, Wisconsin, Minnesota, North Dakota, and South Dakota) produced most of the black walnut veneer logs used in the United States (Blyth, 1973). From 1963 to 1968, Illinois production of these logs increased from 2.3 to 3.0 million board feet (Blyth, 1973). In 1973, U.S. consumption and exports totaled 28.3 million board feet at a value of \$1414/1000 board feet (Anonymous, 1974).

Insects have been shown to decrease black walnut production. A weevil [*Conotrachelus retentus* (Say)] and the husk fly [*Rhagoletis suavis* (Lowe)] cause considerable damage to developing nuts (Miller, 1973). Several insects, including a shoot moth (*Gwendolina concitricana* Heinrich), case moths [*Acrobasis caryivorella* Ragonot and *A. juglandis* (LeBaron)], weevils (*C. retentus* and *C. juglandis* LeConte), and periodical cicadas (*Magicicada* spp.) cause stem deformation through terminal bud or elongating shoot injury (Miller, 1973). Several species, including walnut caterpillar (*Datana integerrima* Grote & Robinson) and walnut aphids (e.g., *Monellia* spp.), cause defoliation or sap removal, thus reducing biomass additions to stem and nut crops (Miller, 1973). Others, including the flatheaded apple tree borer [*Chrysobothris femorata* (Olivier)] and white oak borer [*Goes tigrinus* (DeGeer)], bore into the wood creating holes that may result in large wounds (Miller, 1973).

METHODS AND MATERIALS

Southern Illinois was arbitrarily defined as that part of the state south of an east-west line passing through St. Louis, Missouri, and Vincennes, Indiana. All plantations surveyed

¹Part of a research report submitted to Southern Illinois University at Carbondale by the senior author in partial fulfillment of the requirements of the M.S. degree in zoology.

²Graduate student and Associate Professor, respectively, Department of Zoology, Southern Illinois University at Carbondale, Carbondale, Illinois 62901.




Fig. 1. Location of black walnut plantations surveyed.

(27) had been planted since 1962 by private individuals under the direction of the Illinois State Department of Conservation, or by the U.S. Forest Service, and were located in 17 of the 28 counties. Each plantation was assigned a number for easy reference when referring to the various insect species collected (Table 1). Location of each plantation is shown in Figure 1, and plantation location, planting dates, acreage, and other tree species present are given in Table 1.

Insects were collected from 26 April to 1 October, 1974, and from 8 April to 9 October, 1975. The April and October dates were before leafing out and after leaf drop, respectively, of black walnut in southern Illinois during these two years. Although collecting trips were made to plantations prior to the April and after the October dates, no specimens were found.

This study consisted of two parts. During 1974 and most of 1975, sampling was confined to three plantations (Nos. 12, 13, 14) within 10 miles of the Southern Illinois University-Carbondale campus; trips were made to these plantations 1-2 times/week. Of the remaining 24 plantations, two were visited in 1974 (21 August, plantation 15; 27 August, plantation 11), the other 22 in 1975. Each of the 24 was visited only once.

Specimens were collected by hand-picking or sweeping the foliage and branches of various black walnut trees in each plantation. Notes taken on the various species included types of feeding damage (e.g., leaflet consumption, sap removal, bud boring), instars present, predators and parasites, and distribution in southern Illinois. A field feeding record by a chewing insect was defined as the observation of mandibles removing pieces of leaflet,

Table 1. Descriptions of black walnut plantations surveyed.

Plant. no.	County	Location	Year planted	Acres	Other tree species present
1	Monroe	T3S R11W S4	1970	1	none
2	St. Clair	T1N R7W S31	volunteer	3	mixed deciduous
3	Washington	T1S R5W S13	1969	2	none
4	Clinton	T3N R4W S30	1969	1	mixed deciduous
5	Marion	T4N R1E S16	1968-1969	2	none
6	Wayne	T2S R7E S12	1962	2	none
7	Hamilton	T4S R6E S15	1972	2	none
8	Franklin	T6S R2E S17	1973	8	mixed deciduous
9	Randolph	T7S R5W S21	1972-1973	2	none
10	Jackson	T8S R4W S12	1972	4	none
11	Jackson	T8S R4W S22	1973	3	none
12	Jackson	T9S R2W S25	1965	1	none
13	Jackson	T9S R1W S31	1970	2	none
14	Jackson	T10S R1W S25	1965	2	none
15	Union	T11S R2W S8	1969	12	none
16	Union	T12S R2W S8	1970	2	none
17	Alexander	T14S R2W S28	1967-1970	6	none
18	Alexander	T15S R2W S5	1969	7	autumn olive, alder & black locust
19	Massac	T14S R4E S34	1967	1	mixed deciduous
20	Johnson	T12S R4E S3&10	1965	5	mixed deciduous
21	Pope	T12S R7E S6	1972	4	none
22	Pope	T12S R7E S8	1970	11	mixed deciduous
23	Hardin	T12S R9E S17	1966-1967	7	none
24	Hardin	T12S R8E S2&3	1969	7	autumn olive, alder & black locust
25	Hardin	T12S R8E S4	1965	1	none
26	Hardin	T12S R8E S3	1963, 1966	2.5	none
27	Gallatin	T10S R8E S32	1973	1	none

and by a sucking insect, mouthparts inserted in the plant tissue for at least 30 consecutive seconds. Field feeding records were not possible with the sweeping technique.

Most specimens not observed feeding in the field were returned to the laboratory for further observation. A laboratory feeding record by a chewing insect was defined as either the observation of feeding, or the disappearance of pieces of leaflets from the insect's container. For a sucking insect, it was defined the same as a field observation.

Many insects collected as immatures were laboratory reared to adult to facilitate identification. Specimens returned to the laboratory for observation and rearing were placed in petri dishes (about 9 cm diam, 2 cm depth) with moistened filter paper on the bottom. Black walnut leaflets for chewing insects, and leaflets with a section of rachis for sucking

insects, were added for food. The dishes were kept in an incubator maintained at $23.9 \pm 1.1^\circ\text{C}$ and constant light of about 130 ft-c, and checked at least once/day to record possible feeding. Food was changed and water added about 1 and 2 times/week, respectively, for active specimens, water added less often for pupae. Filter paper was changed as needed, usually once every two weeks.

Pupae of several large Lepidoptera (i.e., Danaidae, Notodontidae, and Saturniidae) were placed in large, clear, plastic containers (29 X 18 X 13 cm) covered with cheesecloth to provide adequate room for wing expansion when the adults emerged. The containers were kept in the laboratory outside (room temperature, $26.7 \pm 1.1^\circ\text{C}$) or inside the incubator.

Several Lepidoptera reared from eggs or larvae apparently entered diapause upon pupation. Since diapause can be broken by a cold period (Beck, 1968), these pupae were placed in a refrigerator at $7.2 \pm 0.6^\circ\text{C}$. Those collected in 1974 were kept in the refrigerator from 27 November, 1974, to 27 February, 1975, those collected in 1975, from 6 November, 1975, to 6 March, 1976. A longer cold period was used in 1975-1976 in an attempt to increase the percentage of those breaking diapause and thus reaching adult.

It should be emphasized that those insects that fed on black walnut in the laboratory had no other food source available to them. Thus, these feeding records demonstrated only that these insects have the potential to feed on black walnut in the field, not that they actually do so.

Parasites that emerged from black walnut insects returned to the laboratory were preserved for later identification.

Certain species proved to be very common during this study, others very rare. The relative abundance of each (inclusive of all stages) is indicated as follows: rare, less than 5; uncommon, 5-15; common, 16-30; very common, more than 30.

All specimens, except the few retained by the U.S. National Museum, are housed in the Entomology Collection, Southern Illinois University at Carbondale Zoological Research Museum.

RESULTS AND DISCUSSION

Insects primarily caused chewing and sucking damage to leaves of immature black walnut trees in southern Illinois, although some ovipositional [e.g., *Anormenis septentrionalis* (Spinola)] and trunk boring damage were observed. Chewing species bored in buds and stems, and removed leaflets. Sucking species removed photosynthates from branches, twigs, rachises, and leaflets; the latter often resulting in spotting. Ovipositional damage presumably resulted in death of twigs. Boring beneath bark occurred in a dying tree.

Approximately 300 species, in 10 orders, were collected during the two years (1974-75) of this study (Table 2). The order Lepidoptera was best represented, with about 80 species collected, and was responsible for the majority of damage observed. Larvae of *Acrobasis* spp. (Pyralidae) killed buds and stems by their boring activity. Larvae of *Hyphantria cunea* (Drury) (Arctiidae), *Morrisonia confusa* (Hübner) (Noctuidae), *Datana integerrima* Grote & Robinson (Notodontidae), *Gretchena bolliana* (Slingerland) (Olethreutidae), and *Gracillaria blandella* Clemens (Gracillariidae) were most important in leaflet removal (partial defoliation).

Table 2. Phytophagous insects from immature black walnut in southern Illinois.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
COLLEMBOLA					
Poduridae					
<i>Hypogastrura packardii</i> (Folsom)	14	U	A&/or J (HP)	---	Ba
Isotomidae					
<i>Isotomurus palustris balteatus</i> (Reuter)	17	R	A&/or J (S)	---	---
Entomobryidae					
<i>Entomobrya atrocineta</i> Schott	14	R	A&/or J (S)	---	---

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
ORTHOPTERA					
Acrididae					
<i>Melanoplus differentialis</i> (Thomas)	11,15	R	A (HP)	---	Le
<i>M. femurrubrum</i> (DeGeer)	13	R	A (HP)	F*	Le
<i>M. punctulatus griseus</i> (Thomas)	22	R	A (HP)	F*	Le
<i>M. rusticus obovativipennis</i> (Blatchley)	13,22	R	A (HP)	F*	Le
<i>M. sanguinipes</i> (Fabricius)	13	R	A (HP)	F*	Le
<i>M. s. scudderii</i> (Uhler)	13,27	R	A (HP)	F*	Le
Tettigoniidae					
<i>Conocephalus brevipedis</i> (Scudder)	13	R	A (HP)	F*	Le
<i>C. fasciatus</i> (DeGeer)	16	R	A (HP)	F*	Le
<i>C. strictus</i> (Scudder)	21	R	N (HP)	F*	Le
<i>Microcentrum rettnerve</i> (Burmeister)	2,13	R	N (HP, S)	F*	Le
<i>M. rhombifolium</i> (Saussure)	13	R	A (S)	---	---
<i>Orechelimum</i> sp.	12,13,16,27	U	N (S)	---	---
<i>Pterophylla camellifolia</i> (Fabricius)	14	R	A (S)	---	---
<i>Scudderia curvicauda</i> (DeGeer)	16	R	A (HP)	F*	Le
<i>S. furoata</i> Brunner	8,13,15	U	A, N (HP)	F**	Le
Gryllidae					
<i>Rapithus agitator</i> Uhler	8,13,26	R	A, N (HP)	F*	Le
<i>Oecanthus argentinus</i> Saussure	12,13	C	A, N (HP, S)	F*	Le
<i>O. latipennis</i> Riley	4,13,14,16,20,27	U	A, N (HP, S)	F*	Le
<i>O. nigricornis</i> F. Walker	13	R	A, N (HP, S)	F*	Le
<i>O. niveus</i> (DeGeer)	3,9,13,14,16,20,27	U	A, N (HP)	F*	Le
<i>Orocharis saltator</i> Uhler	13,19,25	R	N (HP, S)	F*	Le
<i>Phyllopalpus pulchellus</i> Uhler	13,16	R	A, N (S)	---	---
Phasmatidae					
<i>Diaperomera femorata</i> (Say)	15	R	N (HP)	---	Br
PSOCOPTERA					
Psocidae					
<i>Metylophorus pumilus</i> (Walsh)	14	R	A (HP)	---	Br
<i>Psocus leidyi</i> Aaron	14	R	A (S)	---	---
THYSANOPTERA					
Thripidae					
<i>Anaphothrips obscurus</i> (Mueller)	17	R	A (S)	---	---
<i>Frankliniella fusca</i> (Hinds)	17	R	A (S)	---	---
<i>F. triticeae</i> (Fitch)	12,17,22	U	A (HP)	---	Le
HEMIPTERA					
Miridae					
<i>Ceratocapsus</i> sp. near <i>digitulus</i> Knight	2,12,13,14,16,25,26	VC	A, N (HP)	F*	Le
<i>Diaphnidea pellucida</i> Uhler	14	U	A (S)	---	---
<i>Hyaliodus vitripennis</i> (Say)	12,13,14	R	A (S)	F*	Le
<i>Lopidea confluenta</i> (Say)	13,14,17	VC	A (HP)	F**	Ra
<i>Lygus lineolaris</i> (Palisot de Beauvois)	9,12,13,14,16,17	VC	A (HP)	F*	Le
<i>Orthotylus</i> sp.	13,14,16,17	C	A (S)	F*	Le
<i>Plagiognathus flavicornis</i> Knight	12,13,14,17	U	A (HP, S)	F**	Le
<i>F. politus</i> Uhler	13	R	A (S)	---	---
<i>F. repletus</i> Knight	12,14,17	U	A, N (HP)	F**	Le
<i>Stenotus binotatus</i> (Fabricius)	17	R	A (S)	---	---
Tingidae					
<i>Corythucha juglandis</i> (Fitch)	12,13,14,16,17,24,26	VC	A, N (HP)	F**	Le
Lygaeidae					
<i>Lygaeus kalmii</i> (Stål)	12,13	U	A (HP, S)	---	Le
<i>Oncopeltus fasciatus</i> (Dallas)	12,13	C	A, N (HP)	F*	Le
Coreidae					
<i>Acanthocephala terminalis</i> (Dallas)	5,14,22	U	A, N (HP, S)	F*	Le
<i>Euthochtha galeator</i> (Fabricius)	27	R	A, N (HP)	F*	Ra
<i>Leptoglossus oppositus</i> (Say)	SIU campus	R	A (S)	---	---

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
Rhopalidae					
<i>Istorhyssus hyalinus</i> (Fabricius)	13	R	A (HP)	F**	Le
Alydidae					
<i>Alydus eurinus</i> (Say)	13,27	U	A (HP)	---	Le
<i>A. pilosulus</i> (Herrich-Schaeffer)	13	R	A (HP)	---	Le
Pentatomidae					
<i>Acrosternum hilare</i> (Say)	12,13,14	R	A,N,E (HP,S)	---	Le
<i>Brochymena arborea</i> (Say)	14	R	A,N,E (HP)	---	Br,Ra,Tw
<i>B. quadripustulata</i> (Fabricius)	12,13,14,17,18,22,23,27	VC	A,N (HP)	F**	Ra
<i>Euschistus servus</i> (Say)	12,13,14,24,27	VC	A,N,E (HP,S)	F**	Le,Ra
<i>E. t. tristigmus</i> (Say)	14	R	A (HP)	---	Le
<i>Mormidea lugens</i> (Fabricius)	14	R	A (S)	---	---
Cydnidae					
<i>Sehirus c. cinctus</i> (Palisot de Beauvois)	4	R	A (HP)	F*	Le
Corimelaenidae					
<i>Corimelaena agnella</i> McAtee	14	R	A (S)	---	---
<i>C. l. lateralis</i> (Fabricius)	13	R	A (S)	---	---
<i>C. pulicaria</i> (Germar)	12,13	VC	A (HP)	F**	Le
<i>Galgupha atra</i> Amyot & Serville	11	R	A (HP)	---	Le
HOMOPTERA					
Cicadidae					
<i>Magisicada tredeceassini</i> Alexander & Moore					
	14	R	A (HP)	F*	Ra
<i>Tibicen lyriceus</i> (DeGeer)					
	14	R	A (S)	---	---
Membracidae					
<i>Enchenopa binotata</i> (Say)	1,4,12,13,14,25	VC	A,N (HP)	F**	Ra
<i>Entylia bactriana</i> Germar	8,22	R	A (HP)	F*	Le,Ra
<i>Micrutalis calva</i> (Say)	17,18,27	U	A (HP)	F*	Ra
<i>Spissistilus borealis</i> (Fairmaire)	13,14	R	A (HP)	F**	Ra
<i>Stictoccephala hubalus</i> (Fabricius)	9,12,13,20	U	A (HP)	F**	Ra
<i>S. taurina</i> (Fitch)	13,26	R	A (HP,S)	F**	Ra
Cicadellidae					
<i>Agallia constricta</i> Van Duzee	12,14,16	C	A (HP)	F**	Le,Ra
<i>A. quadripunctata</i> (Provancher)	20	R	A (HP)	F*	Le
<i>Agallitopsis novella</i> (Say)	14,22	R	A (HP)	F**	Le
<i>Coelidia olitoria</i> (Say)	2,3,4,5,9,12,13,14,16,20,22,25,27	VC	A,N (HP)	F**	Le,Ra
<i>Colladonus olitellarius</i> (Say)	13	R	A (S)	F*	Le
<i>Draeculacephala antica</i> (Walker)	16,27	R	A,N (S)	---	---
<i>Empoasca fabae</i> (Harris)	17,18	C	A (S)	---	---
<i>Endria inimica</i> (Say)	18	R	A (S)	---	---
<i>Erythroneura</i> spp.	12,13,14,17,18,24,27	VC	A (S)	F*	Le
<i>Graphocephala coccinea</i> (Forster)	2,12,13,14,16,20	C	A (HP)	F**	Le,Ra
<i>G. versuta</i> (Say)	13,14,16,20,23,24,25	C	A (HP)	F**	Le
<i>Gyponana</i> sp.	2	R	N (HP,S)	F*	Le
<i>Latalus sayi</i> (Fitch)	13	R	A (S)	---	---
<i>Macrostelus fascifrons</i> (Stål)	17,18	R	A (S)	---	---
<i>Menosoma cincta</i> (Osborn & Ball)	12,13	U	A (HP)	F**	Le
<i>Nanopsis verticis</i> (Say)	1,2,3,4,5,8,12,13,14,16,17,20,22,25,26	VC	A,N (HP)	F**	Le,Ra
<i>Oncometopia orbona</i> (Fabricius)	18,24	R	A (HP)	F*	Ra
<i>Paraphlepsius irroratus</i> (Say)	1,12	R	A (HP,S)	F**	Le
<i>P. temessa</i> (DeLong)	14,16	R	A (HP)	F**	Ra
<i>Paraulacizea irrorata</i> (Fabricius)	13	U	A (HP)	---	Le
<i>Penthimia americana</i> Fitch	22	R	A (HP)	F*	Le
<i>Planicephalus flavocostatus</i> (Van Duzee)	12	R	A (S)	---	---
<i>Scaphoideus titanus</i> Ball	12	R	A (HP)	F**	Le
<i>Scaphytopius acutus</i> (Say)	12,13	R	A (S)	---	---
<i>S. frontalis</i> (Van Duzee)	12,13,18	R	A (HP)	F**	Le
<i>Stragania</i> sp.	14	R	N (S)	---	---
Cercopidae					
<i>Clastoptera achatina</i> Germar	13,25	R	A (S)	---	---
<i>C. laenata</i> Fowler	19	R	A (HP)	---	Ra

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
<i>C. obtusa</i> (Say)	14	R	A(S)	---	---
<i>Lepryonia quadrangularis</i> (Say)	27	R	A(HP)	F*	Ra
<i>Philaenus spumarius</i> (Linnaeus)	2, 3, 5, 8, 12, 13	C	A(HP)	F**	Le, Ra
Delphacidae					
<i>Liburniella ornata</i> (Stål)	8, 13	R	A(HP)	F**	Le
Derbidae					
<i>Anotia bonnetti</i> Kirby	14	R	A(S)	---	---
<i>Apache degeerii</i> (Kirby)	3	R	A(HP)	F*	Le
<i>Cedusa kedusa</i> McAtee	14	U	A(HP)	F**	Le
<i>Syntames uhleri</i> Ball	4, 5, 13, 14, 22, 27	U	A(HP)	F**	Le, Ra
Cixiidae					
<i>Cixius</i> sp.	14	R	A(HP)	---	Tr
<i>Haplaxius pictifrons</i> (Stål)	17	R	A(S)	---	---
<i>Oliarus</i> sp.	14	R	A(S)	---	---
Flatidae					
<i>Anomenis septentrionalis</i> (Spinola)	3, 5, 12, 13, 14, 17	VC	A, N, E(HP, S)	F**	Le, Ra
<i>Metacalfa pruinosa</i> (Say)	1, 3, 12, 13, 14, 16, 27	VC	A, N(HP, S)	F**	Br, Le, Ra
<i>Ormenoides verusta</i> (Melichar)	3, 14, 26	R	A(HP)	F**	Le, Ra
Acanaloniidae					
<i>Acanalonia bivittata</i> (Say)	3, 27	R	A(HP)	F**	Ra
<i>A. contea</i> (Say)	1, 3, 4, 8, 14, 19	U	A, N(HP)	F**	Ra
Issidae					
<i>Thionia bullata</i> (Say)	16	R	A(S)	---	---
<i>T. simplex</i> (Germar)	4	R	A(HP)	F**	Ra
Psylliidae					
<i>Livia vernalis</i> (Fitch)	13	R	A(S)	---	---
Aphididae					
<i>Acyrtosiphon pisum</i> (Harris)	14	R	A or N(S)	---	---
<i>Dactynotus</i> sp.	27	R	A(HP)	---	Le
<i>Hyalopterus pruni</i> (Geoffroy)	14	R	A(HP)	---	Le
<i>Monellia nigropunctata</i> Granovsky	9, 13, 20, 23	C	A(HP)	F*	Le
<i>Monellopsis caryae</i> (Monell)	12, 13, 14, 15	C	A or N(HP)	F*	Le
<i>Myzocallis</i> sp.	2	R	A(HP)	---	Le
<i>Nearctaphis bakeri</i> (Cowan)	13	R	A, A or N(HP, S)	---	Le
<i>Pemphigus</i> sp.	13	R	A or N(S)	---	---
<i>Rhopalosiphum maidis</i> (Fitch)	19	R	A(HP)	---	Le
<i>Schizaphis graminum</i> (Rondani)	14	R	A or N(S)	---	---
Coccidae					
<i>Lecanium corni</i> Bouche	14, 19	U	A or N(HP)	F**	Br, Tr, Tw
<i>Pulvinaria innumerabilis</i> (Rathvon)	14	R	A or N(HP)	F**	Br
COLEOPTERA					
Elateridae					
<i>Conoderus lividus</i> (DeGeer)	14, 25	U	A(HP)	---	Le
<i>Melanotus depressus</i> (Melshemer)	14, 17	R	A(HP)	F*	Le
Throscidae					
<i>Drapetes geminatus</i> Say	13	R	A(S)	---	---
Buprestidae					
<i>Agrilus arcuatus</i> (Say)	13	R	A(S)	---	---
<i>A. fallax</i> Say	13	R	A(S)	---	---
<i>A. transimpresus</i> Fall	13, 14, 17	U	A(HP)	F*	Le
<i>Dicaerpa lepida</i> LeConte	13	R	A(HP)	F*	Le
Ptilodactylidae					
<i>Ptilodactyla angustata</i> Horn	2	R	A(HP)	---	Le
<i>P. serricollis</i> (Say)	14	R	A(HP)	---	Le
Helodidae					
<i>Cyphon</i> sp.	12	R	A(S)	---	---
<i>Scolites orbiculatus</i> (Fabricius)	1	R	A(HP)	---	Le
Languriidae					
<i>Languria mozardi</i> Latreille	14	R	A(S)	---	---
Phalacridae					
<i>Phalacrus</i> sp.	13	R	A(HP, S)	---	Le
<i>Stilbus</i> sp.	13	R	A(S)	---	---
Pedilidae					
<i>Macratriva</i> sp.	14	R	A(HP, S)	---	Le

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
Mycetophagidae					
<i>Littargus quadrispilotus</i> LeConte	12	R	A(S)	---	---
Meloidae					
<i>Epicauta cinerea</i> (Forster)	15	R	A(S)	---	---
Mordellidae					
<i>Mordella</i> sp.	12	R	A(S)	---	---
<i>Mordellistena</i> sp.	12,14,17	R	A(S)	---	---
Scarabaeidae					
<i>Anomala flavipennis</i> Burmeister	14	R	A(HP)	F**	Le
<i>A. imnuba</i> (Fabricius)	14	R	A(S)	F*	Le
<i>A. marginata</i> (Fabricius)	14,17,18,25	U	A(HP)	F**	Le
<i>Ataenius strigatus</i> (Say)	12	R	A(HP)	---	Br
<i>Cotinis nitida</i> (Linnaeus)	24	R	A(HP)	F*	Le
<i>Euphoria herbacea</i> (Olivier)	14	R	A(S)	---	---
<i>Macrodactylus subspinosus</i> (Fabricius)	14	R	A(S)	F**	Le
<i>Valgus squamiger</i> (Beauvois)	13	R	A(HP)	F*	Le
Cerambycidae					
<i>Lepturges</i> sp. near <i>quercii</i> Fitch	14	R	L(HP)	F**	UBa
<i>L.</i> sp. near <i>symmetricus</i> (Haldeman)	14	R	L(HP)	F**	UBa
<i>Megacyllene robiniae</i> (Forster)	13	R	A(S)	F*	Le
<i>Tetraopes tetrophthalmus</i> (Forster)	13	R	A(HP,S)	F*	Le
Chrysomelidae					
<i>Anomoea flavokansiensis</i> Moldenke	12,13	U	A(HP)	F**	Le
<i>Babia quadriguttata</i> (Olivier)	14	R	A(S)	F*	Le
<i>Bassareus brunneipes</i> (Olivier)	14,20	R	A(HP)	F*	Le
<i>B. mammifer</i> (Newman)	13	R	A(S)	F*	Le
<i>Cerotoma trifurcata</i> (Forster)	16	R	A(HP)	F*	Le
<i>Chaetocnema confinis</i> Crotch	24	R	A(S)	---	---
<i>C. minuta</i> Melsheimer	5	R	A(HP)	---	Le
<i>Colaspis brunnea</i> (Fabricius)	12,14	R	A(S)	---	---
<i>Crepidodera nana</i> (Say)	9,13	R	A(HP)	---	Le
<i>Cryptocephalus guttulatus</i> Olivier	9,14,22,27	U	A(HP)	---	Le,Ra
<i>C. leucomelas</i> Suffrian	13,14	R	A(HP,S)	F*	Le
<i>C. mutabilis</i> Melsheimer	8,14	R	A(HP,S)	F*	Le
<i>C. nanus</i> Fabricius	2,13,14,17	U	A(HP,S)	F*	Le
<i>C. notatus</i> quadrimaculatus Say	13	R	A(S)	---	---
<i>C. quadruplea</i> Newman	13,14,17	U	A(HP)	F*	Le
<i>Diabrotica undecimpunctata howardi</i> Barber	9,12,13,14	C	A(HP)	F**	Le
<i>Ibidomera oliviacollis</i> (Kirby)	12,13	U	A,L(S)	---	---
<i>Longitarsus</i> sp.	21	R	A(HP)	---	Le
<i>Luperaltica nigripalpis</i> (LeConte)	14	R	A(S)	---	---
<i>Myochrous denticollis</i> (Say)	13	R	A(HP)	F*	Le
<i>Nodonota clypealis</i> Horn	14,17	U	A(HP,S)	---	Le
<i>N. tristis</i> (Olivier)	9,12,13,14,17,18	C	A(HP)	F*	Le
<i>Paohybrachys relictus</i> Fall	1	R	A(HP)	F*	Le
<i>P. subfasciatus</i> LeConte	14	R	A(S)	F*	Le
<i>Paria fragariae</i> Wilcox	14,17,26,27	U	A(HP)	F*	Le
<i>P. quadrinotata</i> (Say)	14	R	A(S)	---	---
<i>Phyllotreta zimmermanni</i> (Crotch)	17	R	A(S)	---	---
Bruchidae					
<i>Althaeus</i> sp.	12	R	A(S)	---	---
Curculionidae					
<i>Anthonomus</i> sp.	12	R	A(HP)	---	Le
<i>Apion</i> sp.	13	R	A(S)	---	---
<i>Conotrachelus elegans</i> (Say)	2	R	A(HP)	---	Le
<i>Cyrtopistomus castaneus</i> (Roelofs)	8,9,12,13,14,16,24,25,26,27	VC	A(HP,S)	F*	Le
<i>Eugnaptus</i> sp. near <i>collaris</i> (Fabricius)	14,17	R	A(S)	---	---
<i>Hypera punctata</i> (Fabricius)	20	R	A(HP)	---	Le
<i>Madarellus undulatus</i> (Say)	13	R	A(S)	F*	Le
<i>Magdalis pandura</i> Say	14,17	R	A(HP)	---	Le
<i>Odontopus calceatus</i> Say	13	R	A(HP)	---	Le
LEPIDOPTERA					
Danaidae					
<i>Danaus plexippus</i> (Linnaeus)	13	R	L(HP)	---	Le

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
Lycaenidae					
<i>Satyrium calamus</i> (Hübner)	12,13,14	C	A, L(HP,S)	F*	Le
Sphingidae					
<i>Cressonia juglandis</i> (J.E.Smith)	13	R	L(HP)	---	Le
Saturniidae					
<i>Actias luna</i> (Linnaeus)	14,15	R	L, E(HP)	F*	Le
<i>Antheraea polyphemus</i> (Cramer)	14	R	L(HP)	F*	Le
Citheroniidae					
<i>Citheronia regalis</i> (Fabricius)	15	R	L(HP)	F*	Le
Ctenuchidae					
<i>Cisseps fulvicollis</i> (Hübner)	9,12,13	U	A(HP)	---	Le
Arctiidae					
<i>Diacrisia virginica</i> (Fabricius)	25	R	L(HP)	F*	Le
<i>Eopantheria scribtonia</i> (Stoll)	14	R	L(HP)	F*	Le
<i>Halsidota tessellaris</i> (J.E.Smith)	14,26	R	L(HP,S)	F*	Le
<i>Haploa colona fulvicosta</i> (Clemens)	14	R	L(S)	F*	Le
<i>Hyphantria cunea</i> (Drury)	1,10,12,13,14	VC	L(HP)	F**	Le
Noctuidae					
<i>Acrionia impleta</i> (Walker)	12,13	R	L(HP)	F*	Le
<i>Baileya australis</i> (Grote)	12,13,14	R	L(HP,S)	F*	Le
<i>Bomolocha</i> sp. near <i>madefactalis</i> (Guenée)	1,13	R	A, L(HP,S)	F*	Le
<i>B. sp. near sordidula</i> Grote	14	R	L(S)	F*	Le
<i>Catocala</i> sp. near <i>neogama</i> J.E.Smith	20	R	L(HP)	F*	Le
<i>Eupellia</i> sp. near <i>sidus</i> Guenée	13	R	L(HP)	F*	Le
<i>Heliocontia apicella</i> (Grote)	13	R	A(S)	---	---
<i>Morrisonia confusa</i> (Hübner)	9,12,13,14,16,17,18	C	L(HP)	F**	Le
<i>Orthosia hibisci</i> (Guenée)	14	U	L(HP)	---	Le
<i>Peridroma saucia</i> (Hübner)	14	U	L(S)	---	---
<i>Spragueia leo</i> (Guenée)	13	R	A(S)	---	---
Notodontidae					
<i>Datana angustii</i> Grote & Robinson	14,27	U	L(HP)	F**	Le
<i>D. integerrima</i> Grote & Robinson	1,5,12,13,14,15	VC	L, E(HP)	F**	Le
<i>Heterocampa guttivitta</i> (Walker)	14	R	L(S)	F*	Le
<i>H. manteo</i> (Doubleday)	17	R	L(HP,S)	F*	Le
<i>Schizura ipomoeae</i> (Doubleday)	1	R	L(HP)	---	Le
<i>S. leptinoides</i> (Grote)	5,12,13,14	U	L(HP)	F*	Le
Lymantriidae					
<i>Dasychira</i> sp. near <i>basiflava</i> (Packard)	14	R	L(HP)	F*	Le
<i>Orgyia leucostigma</i> (J.E.Smith)	12,20	R	L(HP)	F*	Le
Lasiocampidae					
<i>Malacosoma americanum</i> (Fabricius)	14	R	L(S)	---	---
Geometridae					
<i>Abbotana clemataria</i> (J.E.Smith)	13,14	R	L(HP)	F*	Le
<i>Anacamptodes defectaria</i> (Guenée)	12	R	L(S)	F*	Le
<i>A. ephyraria</i> (Walker)	14	R	L(HP)	F*	Le
<i>Anavitrinella pampinaria</i> (Guenée)	13	R	L(S)	F*	Le
<i>Hypagyrtis esther</i> (Barnes)	13,14,20	U	L(HP)	F*	Le
<i>H. unipunctata</i> (Haworth)	12,22	R	L(HP)	F*	Le
<i>Lambdina fervidaria</i> (Hübner)	17	R	L(S)	F*	Le
<i>Melanolophia canadaria</i> (Guenée)	14	R	L(S)	F*	Le
<i>M. signataria</i> (Walker)	13,17	R	L(S)	F*	Le
<i>Synchlora aerata</i> (Fabricius)	12,13	R	A(S)	---	---
Limacodidae					
<i>Prolimacodes scapha</i> Harris	12	R	L(HP)	F*	Le
<i>Sib. e stimulea</i> (Clemens)	16	R	L(HP)	F**	Le
Zygaenidae					
<i>Acolloithus falsarius</i> Clemens	13,25	R	A(HP,S)	---	Le
Pyralidae					
<i>Aerobasis demotella</i> Grote	12 or 13,13	U	L(HP)	F**	IBu, ISt
<i>A. juglandis</i> (LeBaron)	12 or 13,13,14	U	L(HP)	F**	IRa, Le
<i>A. latifasciella</i> Dyar	14	U	L(HP,S)	F**	IBu, ISt
<i>Argyria nivalis</i> (Drury)	12	R	A(S)	---	---
<i>Crambus elegans</i> Clemens	14	U	A(S)	---	---
<i>C. sp. near teterellus</i> Zincken	14	R	A(S)	---	---
<i>Tetralopha asperatella</i> Clemens	13	R	L(HP)	F*	Le

Table 2. continued.

Taxon	Plantation numbers	Rel. freq. ^a	Stages collected ^b	Feeding record ^c	Assoc. plant parts ^{d,e}
<i>T. nephelotella</i> Hulst	14	R	L (HP)	F*	Le
<i>T. subcanalis</i> (Walker)	14	R	L (HP)	F*	Le
<i>Udea rubigalis</i> (Guenée)	14	R	A (S)	---	---
Olethreutidae					
<i>Episimus argutus</i> (Clemens) (?)	13	R	A (HP)	---	Le
<i>Grapholitha eclipsana</i> Zeller	13	R	A (S)	---	---
<i>G. interstinctana</i> (Clemens)	13	U	A (HP, S)	---	Le
<i>Gretchena amatana</i> Heinrich	13	R	L (S)	F*	Le
<i>G. bolliana</i> (Slingerland)	2, 3, 12, 13, 14, 17, 22, 23, 27	VC	A, L (HP, S)	F**	Le
<i>Gwendolina concitaticana</i> Heinrich	3 or 23	R	L (HP)	F*	Le
<i>Laspeyresia</i> sp. near <i>caryana</i> Fitch	13	R	A (HP)	---	Le
Tortricidae					
<i>Archips argyrospilus</i> (Walker)	13, 14	R	L (HP)	F*	Le
<i>Argyrotaenia juglandana</i> (Fernald)	12	R	P (HP)	---	Le
<i>A. velutinana</i> (Walker)	14	R	L (HP)	F*	Le
<i>Choristoneura rosaceana</i> (Harris)	12, 14, 23	U	P, L (HP)	F**	Le
<i>Pandemis limitata</i> (Robinson)	12, 13, 14, 16, 26	U	L (HP)	F*	Le
<i>Sparganothis distincta</i> (Walsingham)	11	R	A (HP)	---	Le
Gelechiidae					
<i>Arogalea cristifasciella</i> (Chambers)	12	R	A (HP)	---	Tr
<i>Chionodes</i> sp.	14	R	L (HP)	F*	Le
<i>Dichomeris ligulella</i> (Hübner)	14	R	L (HP)	F*	Le
<i>D. ventrella</i> (Fitch)	13, 14, 18	U	L (HP, S)	F*	Le
Stenomidae					
<i>Antaeotricha leucillana</i> (Zeller)	13	R	A (S)	---	---
Coleophoridae					
<i>Coleophora caryaefoliella</i> Clemens	18	R	L (HP)	---	Le
Gracillariidae					
<i>Gracillaria blandella</i> Clemens	1, 5, 9, 12, 13, 14, 24, 26	C	A, L (HP)	F**	Le
<i>Phyllonorycter juglandiella</i> (Chambers)	3, 4, 23	U	L (HP)	F**	Le
Psychidae					
<i>Thyridopteryx ephemeraeformis</i> (Haworth)	13, 14	R	L (HP)	F*	Le
DIPTERA					
Cecidomyiidae					
<i>Clinodiplosis</i> sp.	2, 22	C	L (HP)	---	Le
HYMENOPTERA					
Xyelidae					
<i>Megaxyela</i> sp.	6 & 12 or 13	R	L (HP)	---	Le
Tenthredinidae					
<i>Caulocampus acericaulis</i> (MacGillivray)	12	R	A (S)	---	---
<i>Dolerus aprilis</i> (Norton)	14	R	A (S)	---	---
<i>Ertocampa juglandis</i> (Fitch)	14, 17	U	L (HP)	F*	Le
<i>Macrophya flavicoxae</i> (Norton)	13	R	A (S)	---	---

^aC = common, R = rare, U = uncommon, VC = very common.
^bA = adult, E = egg, J = juvenile, L = larva, N = nymph, P = pupa, HP = hand-picking, S = sweeping.
^cF* = laboratory feeding record, F** = field feeding record.
^dRefers to plant part upon which insect was collected or, if noted feeding in the field or laboratory, the part upon which it was feeding.
^eBa = bark, Br = branch, IBu = in bud, IRa = in rachis, IST = in stem, Le = leaflet, Ra = rachis, Tr = trunk, Tw = twig, UBa = under bark.

Table 3. List of predators and parasites from black walnut insects.

Black Walnut Insects	Predator (Field) or Parasite (Laboratory)		Comments
	Family	Species	
Lygaeidae <i>Lygaeus kalmii</i> Stål	Tachinidae	<i>Leucostoma gravipes</i> Wulp	Emerged from adult
Pentatomidae <i>Euschistus servus</i> (Say)	Scelionidae	<i>Trissolus euschisti</i> (Ashmead)	Emerged from eggs
Cicadellidae <i>Agallia constricta</i> Van Duzee	Asilidae	<i>Cerotainia macrocera</i> (Say)	Feeding on adult in field
Chrysomelidae <i>Anomoea flavokansteniensis</i> Moldenke	Pentatomidae	<i>Podisus maculiventris</i> (Say)	Feeding on adult in field
Arctiidae <i>Halisidota tessellaris</i> (J.E.Smith)	Ichneumonidae	<i>Parania g. geniculata</i> (Holmgren)	Emerged from pupa
Noctuidae <i>Morrisonia confusa</i> (Hübner)	Braconidae	<i>Apanteles hyphantriae</i> Riley	Emerged from larva
Notodontidae <i>Datania integerrima</i> Grote & Robinson	Scelionidae	<i>Telenomus ichthyurae</i> Ashmead	Emerged from pupa
	Ichneumonidae	<i>Barylypa</i> sp.	Emerged from pupa
	Ichneumonidae	<i>Tanyptelma datanae</i> (Riley)	Emerged from pupa

Some of the more common sucking insects noted feeding in the field included *Lopidea confluenta* (Say) (Miridae), *Corythucha juglandis* (Fitch) (Tingidae), *Brochymena quadripustulata* (Fab.) (Pentatomidae), *Enchenopa binotata* (Say) (Membracidae), *Coelidia olitoria* (Say), *Nanopsis verticis* (Say) (Cicadellidae), *Anormenis septentrionalis* (Spinola), and *Metcalfa pruinosa* (Say) (Flatidae). However, the importance of sucking damage to the tree is difficult to determine.

The predators and parasites collected during this study, and their prey and hosts, respectively, are given in Table 3.

Additional biological information on southern Illinois black walnut phytophagous insects is provided by Nixon (1976).

ACKNOWLEDGMENTS

We wish to thank the following faculty and staff members of Southern Illinois University at Carbondale: W. G. Dyer and G. Garoian, Department of Zoology, for their critical reviews of parts of this paper; and K. A. Schmitt and A. L. Aamodt, Scientific Photography and Illustration Facility, Research and Projects, for the final production of Fig. 1. We also thank J. P. Cuda and M. L. Youther, former zoology graduate students, and S. W. Wilson, zoology graduate student, for assistance in field collections.

We acknowledge the following individuals for assistance in locating, and/or granting permission to survey, black walnut plantations: J. L. Clutts, G. H. Lyon, Jr., A. R. Mikich, J. Newcomb, J. R. Ward, USDA Forest Service, Illinois; G. Campbell, D. L. Cooper, D. A. Gillespie, R. C. Heavener, W. A. Keith, R. Moore, T. Seals, S. C. Springer, Department of Conservation, State of Illinois; and H. R. Andres, E. F. Bollman, J. L. Bremer, J. Eckert, A. Holden, P. Karhoff, and W. Schrage, private landowners.

We are grateful to the following specialists for identification of insects included in this study: J. A. Beatty, Department of Zoology, SIU-C; W. E. Miller, North Central Forest Experiment Station, St. Paul, Minnesota; D. R. Davis, W. D. Duckworth, W. D. Field, R. C. Froeschner, P. J. Spangler, U. S. National Museum, Washington, D.C.; D. M. Anderson, R. W. Carlson, D. C. Ferguson, R. J. Gagne, G. Gordh, R. D. Gordon, A. B. Gurney, J. L. Herring, R. W. Hodges, J. M. Kingsolver, L. V. Knutson, J. P. Kramer, P. M. Marsh, A. S.

Menke, D. R. Miller, E. L. Mockford, K. O'Neill, L. M. Russel, C. W. Sabrosky, D. R. Smith, T.J. Spilman, M. B. Stoetzel, E. L. Todd, R. E. Warner, D. M. Weisman, R. E. White, and D. L. Wray, Systematic Entomology Laboratory, Agric. Res. Serv., USDA, Beltsville, Maryland. We are also grateful to R. H. Foote and L. V. Knutson for expediting the identifications by personnel of the U.S. National Museum and USDA, Beltsville.

Our special thanks to D. T. Funk, R. E. Phares (now in Washington, D.C.), and B. C. Weber, Forestry Sciences Laboratory, North Central Forest Experiment Station, USDA, Carbondale, Illinois, for their willing assistance during this study.

This research was partially funded by the USDA Forest Service (Cooperative Research Agreement No. 13-364).

LITERATURE CITED

- Anonymous. 1974. U.S. Dep. Commerce News, Domestic & International Business Administration 74-39.
- Barrett, R. E. 1932. An annotated list of the insects and arachnids affecting the various species of walnuts or members of the genus *Juglans* Linn. Univ. Calif. Pub. Entomol. 5:275-309.
- Beck, S. D. 1968. Insect photoperiodism. Acad. Press, New York.
- Blyth, J. E. 1973. Timber demand and use, p. 7-9, in: Black walnut as a crop. USDA Forest Serv., Gen. Tech. Rep. NC-4:1-114.
- Carmean, W. H. 1970. Tree height-growth patterns in relation to soil and site, p. 499-512, in: C. T. Youngberg and C. B. Davey (eds.). Tree growth and forest soils. Third north Amer. Forest Soils Conf. Proc.
- Miller, W. E. 1973. Insects as related to wood and nut production, p. 91-96, in: Black walnut as a crop. USDA Forest Serv., Gen. Tech. Rep. NC-4:1-114.
- Nixon, P. L. 1976. Key and annotated list of phytophagous insects collected on immature black walnut trees in southern Illinois. M.S. Research Report, Southern Illinois University at Carbondale.