

# The Great Lakes Entomologist

---

Volume 14  
Number 3 - Fall 1981 *Number 3 - Fall 1981*

Article 2

---

October 1981

## Seasonal Flight Patterns of Hemiptera in a North Carolina Black Walnut Plantation. 6. Tingidae and Aradidae

J. E. McPherson  
*Southern Illinois University*

B. C. Weber  
*Southern Illinois University*

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

McPherson, J. E. and Weber, B. C. 1981. "Seasonal Flight Patterns of Hemiptera in a North Carolina Black Walnut Plantation. 6. Tingidae and Aradidae," *The Great Lakes Entomologist*, vol 14 (3)  
Available at: <https://scholar.valpo.edu/tgle/vol14/iss3/2>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**SEASONAL FLIGHT PATTERNS OF HEMIPTERA IN A  
NORTH CAROLINA BLACK WALNUT PLANTATION.  
6. TINGIDAE AND ARADIDAE**

J. E. McPherson<sup>1</sup> and B. C. Weber<sup>2</sup>

ABSTRACT

The seasonal flight patterns of 11 species of Tingidae and six species of Aradidae collected in window traps in a North Carolina black walnut plantation are described. Flying height distributions and seasonal flight activities of *Corythucha ciliata* (Say) and *Gargaphia solani* Heidemann are considered in detail.

---

This is the sixth in a series of papers on seasonal flight patterns of Hemiptera in a black walnut (*Juglans nigra* L.) plantation near Asheville, North Carolina, and deals with the families Tingidae and Aradidae; earlier papers dealt with the Pentatomoidea (1980), Coreoidea (1981a), Reduvoidea (1981b), Cimicoidea (1981c) and Lygaeoidea (1981d). The study was conducted from 24 March to 14 October in 1977, and from 24 March to 13 October in 1978. Specimens were collected weekly by window trapping; traps were suspended at 1, 2, 3, 4, 5, 6 and 7 m. The study site and trap construction were discussed in detail by McPherson and Weber (1980). All hemipteran specimens collected during this study are deposited in the Entomology Collection, Zoology Research Museum, Southern Illinois University, Carbondale.

RESULTS AND DISCUSSION

Eleven tingid and six aradid species were collected during the two years of this study; numbers of specimens collected for all taxa ranged from one to 91 (Table 1).

Most of the species were collected in numbers too low to permit conclusions about seasonal flight patterns. However, *Corythucha ciliata* (Say) and *Gargaphia solani* Heidemann were collected in sufficient numbers (Table 1) to allow a more detailed discussion of flying height distributions and seasonal flight activities.

*C. ciliata* occurs primarily on sycamore (e.g., Bailey 1951, Barber and Weiss 1922, Blatchley 1926, Drake 1919, Froeschner 1944, Horn et al. 1979, Morrill 1903, Wade 1917) but has also been collected from cypress (Froeschner 1944), ash, hickory, mulberry (Blatchley 1926, Drake 1919) and leather-leaf (Bailey 1951). It overwinters as adults (e.g., Bailey 1951, Barber and Weiss 1922, Froeschner 1944, Wade 1917); in central New England, overwintering adults may be found in abundance under loose tree bark in October and may not emerge until early June (Bailey 1951). Barber and Weiss (1922) felt this species probably has "two broods" (is bivoltine?) in New Jersey.

In the present study, *C. ciliata* adults were found from early April to early September (Table 1). They were collected at all seven flying heights with over 55% captured at 3-5 m (Fig. 1).

---

<sup>1</sup>Department of Zoology, Southern Illinois University, Carbondale, IL 62901.

<sup>2</sup>USDA Forest Service, North Central Forest Experiment Station, Forestry Sciences Laboratory, Carbondale, IL 62901.

Table 1. Seasonal flight activity of Tingidae and Aradidae during 1977-78 in a North Carolina black walnut plantation.

Taxon	No. Collected	Collection Height (m)		Range of Collection Dates
		$\bar{x} \pm SE$	Range	
TINGIDAE				
<i>Corythucha ciliata</i> (Say)	70	4.27 $\pm$ 0.21	1-7	1 April-2 Sept.
<i>Corythucha cydoniae</i> (Fitch)	1	3.00	—	23 June
<i>Corythucha juglandis</i> (Fitch)	26	4.62 $\pm$ 0.35	1-7	1 April-13 Oct.
<i>Corythucha marmorata</i> (Uhler)	3	3.67 $\pm$ 1.45	1-6	13 May-12 Aug.
<i>Corythucha pallida</i> Osborn & Drake	2	6.50 $\pm$ 0.50	6-7	22 April
<i>Gargaphia solani</i> Heidemann	91	1.85 $\pm$ 0.16	1-7	2 June-13 Oct.
<i>Gargaphia tiliae</i> (Walsh)	1	6.00	—	29 July
<i>Leptoypha costata</i> Parshley	8	4.75 $\pm$ 0.25	4-6	3 June-22 July
<i>Melanorhopala infuscata</i> Parshley	1	4.00	—	30 June
<i>Physatocheila brevisrostris</i> Osborn & Drake	2	2.50 $\pm$ 1.50	1-4	7 April-15 Sept.
<i>Physatocheila variegata</i> Parshley	1	6.00	—	6 May
ARADIDAE				
<i>Aradus cinnamomeus</i> Parshley	3	3.67 $\pm$ 0.67	3-5	22 Sept.-23 Sept.
<i>Aradus crenatus</i> Say	1	1.00	—	29 April
<i>Aradus niger</i> Stål	1	6.00	—	29 Sept.
<i>Mezira granulata</i> (Say)	30	4.07 $\pm$ 0.33	1-7	22 April-16 Sept.
<i>Mezira lobata</i> (Say)	1	4.00	—	20 May
<i>Neuroctenus</i> sp. <sup>a</sup>	2	3.50 $\pm$ 1.50	2-5	20 May-2 June


<sup>a</sup>Genus is currently being revised (Froeschner, pers. comm.).

This species overwintered as adults and was apparently bivoltine (Fig. 3) Adults began to emerge from overwintering sites in early April and were present to mid-May. Their adult offspring (summer generation) occurred from late June to late July or early August. Adults of the second (overwintering) generation were present in September (and possibly in August).


*G. solani* occurs primarily on eggplant (e.g., Bailey 1951, Fink 1915, Froeschner 1944, Horn et al. 1979) but has also been collected from horse-nettle (Blatchley 1926, Fink 1915, Froeschner 1944, Somes 1916), white horse-nettle (Froeschner 1944) and sunflower (Horn et al. 1979). It overwinters as adults (Bailey 1951, Fink 1915, Froeschner 1944). Fink (1915) stated that *G. solani* may have 7-8 generations per season in the vicinity of Norfolk, Virginia, with apparently six generations on eggplant and the remainder on horse-nettle.

In the present study, *G. solani* adults were found from early June to mid-October (Table 1). They were collected at all seven flying heights with almost 70% collected at 1 m (Fig. 2).

This species apparently overwintered as adults but the number of generations per year is unclear from the available data (Fig. 4). It appears that during spring (i.e., April and May), adults were either not flying or were not in the plantation. However, adults were quite active from early July to the end of the season.



Figs. 1-2 Flying height distributions of two tingid species during 1977-78 in a North Carolina black walnut plantation: (1) *Corythucha ciliata*, (2) *Gargaphia solani*.



Figs. 3-4. Seasonal flight activities of two tingid species during 1977-78 in a North Carolina black walnut plantation: (3) *Corythucha ciliata*, (4) *Gargaphia solani*.

#### ACKNOWLEDGMENTS

We wish to thank Dr. R. C. Froeschner, National Museum of Natural History, Washington, D.C., for confirming our identifications of these tingids and aradids. We also wish to acknowledge Mr. D. Brenneman and the staff of the North Carolina Division of Forestry, Morganton, for their help in collecting data and maintaining the window traps. This research was partially supported by the USDA Forest Service, North Central Forest Experiment Station.

#### LITERATURE CITED

Bailey, N. S. 1951. The Tingoidea of New England and their biology. *Entomol. Amer.* 31:1-140.  
 Barber, H. G. and H. B. Weiss. 1922. The lace bugs of New Jersey. *New Jersey Dept. Agric. Circ.* 54:1-24.  
 Blatchley, W. S. 1926. Heteroptera or true bugs of eastern North America with especial reference to the faunas of Indiana and Florida. *Nature Pub. Co., Indianapolis.* 1116 pp.  
 Drake, C. J. 1919. On some North American Tingidae (Hemip.). *Ohio J. Sci.* 19:417-421.  
 Fink, D. E. 1915. The eggplant lace-bug. *USDA Bull.* 239:1-7.

- Froeschner, R. C. 1944. Contributions to a synopsis of the Hemiptera of Missouri, Pt. III. Lygaeidae, Pyrrhocoridae, Piesmididae, Tingididae, Enicocephalidae, Phymatidae, Ploiaridae, Reduviidae, Nabidae. Amer. Midland Natur. 31:638-683.
- Horn, K. F., C. G. Wright, and M. H. Farrier. 1979. The lace bugs (Hemiptera:Tingidae) of North Carolina and their hosts. North Carolina Agric. Exp. Sta. Tech. Bull. 257:1-22.
- McPherson, J. E. and B. C. Weber. 1980. Seasonal flight patterns of Hemiptera in a North Carolina black walnut plantation. 1. Pentatomoidea. Great Lakes Entomol. 13:177-183.
- . 1981a. Seasonal flight patterns of Hemiptera in a North Carolina black walnut plantation. 2. Coreoidea. Great Lakes Entomol. 14:11-13.
- . 1981b. Seasonal flight patterns of Hemiptera in a North Carolina black walnut plantation. 3. Reduvidae. Great Lakes Entomol. 14:15-17.
- . 1981c. Seasonal flight patterns of Hemiptera in a North Carolina black walnut plantation. 4. Cimicoidea. Great Lakes Entomol. 14:19-22.
- . 1981d. Seasonal flight patterns of Hemiptera in a North Carolina black walnut plantation. 5. Lygaeoidea. Great Lakes Entomol. 14:133-136.
- Morrill, A. W. 1903. Notes on the immature stages of some tingitids of the genus *Corythuca* (sic). Psyche 10:127-134.
- Somes, M. P. 1916. Some insects of *Solanum carolinense* L., and their economic relations. J. Econ. Entomol. 9:39-44.
- Wade, O. 1917. The sycamore lace-bug (*Corythucha ciliata*, Say). Oklahoma Agric. Exp. Sta. Bull. 116:1-16.