

The Great Lakes Entomologist

Volume 18
Number 4 - Winter 1985 *Number 4 - Winter*
1985

Article 4

December 1985

Pammene Perstructana (Walker) (Lepidoptera: Tortricidae) Identified After More Than a Century

William E. Miller
University of Minnesota

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Miller, William E. 1985. "*Pammene Perstructana* (Walker) (Lepidoptera: Tortricidae) Identified After More Than a Century," *The Great Lakes Entomologist*, vol 18 (4)
Available at: <https://scholar.valpo.edu/tgle/vol18/iss4/4>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

**PAMMENE PERSTRUCTANA (WALKER) (LEPIDOPTERA:
TORTRICIDAE) IDENTIFIED AFTER MORE THAN A CENTURY**

William E. Miller¹

ABSTRACT

Sciaphila perstructana Walker has been known only from the female holotype, and *Pammene signifera* (Heinrich) only from males. Based on associated males and females from the Great Lakes area, the two names apply to one species. The resulting synonymy leads to a new combination, *Pammene perstructana*, and finally associates a species with this long-baffling specific name.

Sciaphila perstructana Walker is among myriads of British Museum (Natural History) insects laconically described by Francis Walker in the mid-19th century (Anon. 1874). It is also among 37 Walker names in Nearctic Olethreutinae (Powell 1983). Identities of many species represented by these names have been uncertain because Walker often worked from single specimens.


The Walker Nearctic olethreutine names have been resolved slowly. One reason is that Heinrich (1923, 1926), the first reviser to use genitalic characters, did not examine Walker syntypes. He relied instead on identifications made by Walsingham and C. H. Fernald, both of whom predated taxonomic use of genitalia. Use of nongenital characters led to many synonymical errors. Moreover, the Walker syntypes remained undissected until they were prepared by N. S. Obraztsov after World War II. Obraztsov also selected many lectotypes, but *Sciaphila perstructana*, like many Walker species, was represented only by a holotype. Photos of these types and their genitalia were deposited by Obraztsov in the American Museum of Natural History, New York (Diakonoff 1966). These have helped speed correction of errors, which has now progressed practically as far as specimen condition and present taxonomic methods permit.

Sciaphila perstructana has continued to be an enigma, however. It has been hesitantly referred to various genera (McDunnough 1959), most recently to *Cydia* (Powell 1983). The holotype is female, and males are often necessary for generic placement. Also, having been known only from the holotype, *Sciaphila perstructana* appeared to be an unusually rare moth. After assembling more than 20 specimens of a *Pammene* occurring in the Great Lakes area (Figs. 1, 3), I found that males matched the male holotype of *P. signifera* (Heinrich). The female of this species was hitherto unknown. That the female was the long-baffling *Sciaphila perstructana* became clear when I compared genitalia of *Pammene signifera* females with those of the *Sciaphila perstructana* holotype female (Fig. 2). The new findings are summarized below.

Pammene perstructana (Walker). New Combination
(Figs. 1-3)

Sciaphila perstructana Walker (1863: 343) (holotype: female, St. Martin's Falls, Albany River, Hudson Bay, no date, Barnston, genit. prep. 5319, forewing length 6.0 mm, in British Museum (Natural History), genitalia shown here in Figure 2).

¹Department of Entomology, University of Minnesota, St. Paul, MN 55108.



Figs. 1-3. *Pammene perstructana*: (1) wings of an example from Lake Katherine, Oneida Co., Wisconsin; (2) genitalia of holotype female; (3) genitalia of a male from Cass Lake, Cass Co., Minnesota.

Hemimene signifera Heinrich (1926: 22) (holotype: male, Lake of Bays, Ontario, 1-VII-20, McDunnough, forewing length 5.5 mm, in Canadian National Collection, genitalia illustrated by Heinrich 1926: Fig. 290). New Synonymy.

Pammene signifera; Powell (1983: 37).

I examined both of the above holotypes and the Obraztsov photos of *P. perstructana*. Nontype voucher specimens are in the University of Minnesota, St. Paul; Michigan State University Entomology Museum, East Lansing; University of Wisconsin Insectarium, Madison; University of California, Berkeley; and the Illinois Natural History Survey, Urbana.

There are five Nearctic species of *Pammene* (Powell 1983), and no larval hosts are known. The Palearctic species number nearly 60 (Danilevsky and Kuznetsov 1968). Their larvae feed in terminals, tied leaves, flowers, fruits, galls, and related tissues of *Quercus*, *Salix*, *Acer*, *Fagus*, and other perennials (Swatchek 1958). Larvae of the Palearctic *Pammene clanculana* (Tengström), a species which closely resembles *P. perstructana*, feed in seed-bearing catkins of *Betula* (Benander 1950). Should the latter two prove conspecific, the name *P. perstructana* would take precedence because of priority.

ACKNOWLEDGMENT

I thank curators of the above institutions for specimen loans and other help. Figure 2 is reproduced through the courtesy of the American Museum of Natural History.

LITERATURE CITED

- Anonymous. 1874. Francis Walker (Obituary). Entomol. Mo. Mag. 11:140-141.
 Benander, P. 1950. Andra Familjegruppen, Vecklarfjärilar, Tortricina, Part 10, No. 39, Svensk Insektfauna. Entomologiska Föreningen, Stockholm. 173 pp.

- Danilevsky, A. C. and V. I. Kuznetsov. 1968. Leafroller family Tortricidae, fruit moth tribe Laspeyresiini. Fauna USSR. lepidopterous insects, vol. 5, part 1. Leningrad. 635 pp. (in Russian)
- Diakonoff, A. 1966. Nicholas Sergeevich Obraztsov, 1906–1966. J. Lepidop. Soc. 20:255–265.
- Heinrich, C. 1923. Revision of the North American moths of the subfamily Eucosminae of the family Olethreutidae. U.S. Nat. Mus. Bull. 123. 298 pp.
- . 1926. Revision of the North American moths of the subfamilies Laspeyresiinae and Olethreutinae. U.S. Nat. Mus. Bull. 132. 216 pp.
- McDunnough, J. H. 1959. On some changes in nomenclature of microlepidoptera, with description of a new species. Amer. Mus. Novit. 1954, 9 pp.
- Powell, J. A. 1983. Tortricidae, pp. 31–41 in R. W. Hodges (ed.). Check list of the Lepidoptera of America north of Mexico. E. W. Classey Ltd. and the Wedge Entomological Research Foundation, London. 284 pp.
- Swatchek, B. 1958. Die Larvalsystematik der Wickler. Abh. Larvalsystem. Insekt. 3. Akademie-Verlag, Berlin. 269 pp.
- Walker, F. 1863. Tortricites and Tineites. pp. 287–561 in List of the specimens of lepidopterous insects in the collection of the British Museum. Part 28. British Museum, London.