The Great Lakes Entomologist

Volume 20 Number 2 - Summer 1987 Number 2 - Summer 1987

Article 1

June 1987

New Distribution Records of Scolytidae From Indiana and Florida

Mark Deyrup Archbold Biological Station

Thomas H. Atkinson University of Florida

Follow this and additional works at: https://scholar.valpo.edu/tgle



Part of the Entomology Commons

Recommended Citation

Deyrup, Mark and Atkinson, Thomas H. 1987. "New Distribution Records of Scolytidae From Indiana and Florida," The Great Lakes Entomologist, vol 20 (2)

Available at: https://scholar.valpo.edu/tgle/vol20/iss2/1

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

NEW DISTRIBUTION RECORDS OF SCOLYTIDAE FROM INDIANA AND FLORIDA

Mark Deyrup¹ and Thomas H. Atkinson²

ABSTRACT

Twenty-three new state records are provided for Florida and Indiana Scolytidae; 92 species are now known from Indiana, 112 from Florida.

While reviewing our collections of Scolytidae for a study reported in this journal, we found a number of records of species not previously reported from Indiana in a faunal review by Deyrup (1981) or from Florida in the comprehensive monograph by Wood (1982). We add 14 and subtract two species from the Indiana list, for a total of 92 species known from the state. We add nine species to the Florida fauna, for a total of 112 species. The specimens are in the collections of T. H. Atkinson (THA), Archbold Biological Station (ABS), Stephen L. Wood (SLW), and Florida State Collections of Arthropods (FSC). All identifications were made by the authors, with the exception of Indiana species of Hylocurus and Xyleborus, which were identified by Stephen L. Wood of Brigham Young University.

INDIANA

Crypturgus alutaceus Schwarz. Clark Co. State Forest, in gallery of Ips grandicollis (Eichhoff) in Pinus virginiana Mill.

Crypturgus borealis Swaine. Indiana Dunes State Park, Porter Co., in gallery of Pityogenes hopkinsi Swaine in Pinus strobus L.

Crypturgus pusillus (Gyllenhal). Deleted from Indiana list. Specimens listed (Deyrup 1981) are C. alutaceus and C. borealis.

Dendroctonus terebrans (Olivier): Clark Co. State Forest, in Pinus virginiana.

Dendroctonus valens LeConte. Pokagon State Park, Steuben Co., in Pinus sylvestris L. Hylastes porculus Erichson. Indiana Dunes State Park, Porter Co., in roots of Pinus strobus. Clark Co. State Forest, in roots of Pinus virginiana.

Hylastes tenuis Eichhoff. W. Lafayette, Tippecanoe Co., in flight. Clark Co. State Forest, in roots of *Pinus virginiana*.

Hylocurus bicornus Blackman. Delete from Indiana list. Specimens listed (Deyrup 1981) refer to next species.

Hylocurus binodatus Wood. W. Lafayette, Tippecanoe Co., in branches of Carya ovata (Mill.) K. Koch.

Hylocurus torosus Wood. Turkey Run State Park, Parke Co., in branches of Carya ovata. Shakamak State Park, Clay Co., in branches of Carya ovata.

Liparthrum squamosum (Blackman). Clark Co. State Forest, in twigs of Maclura pomifera (Raf.) Schneid.

¹Archbold Biological Station, P.O. Box 2057, Lake Placid, FL 33852.

²Department of Entomology and Nematology, University of Florida, Gainesville, FL 32611.

Pityogenes plagiatus (LeConte). Clark Co. State Forest, in branches of Pinus virginiana. Pityophthorus cariniceps LeConte. Clifty Falls State Park, Jefferson Co., in branches of Pinus strobus; Indiana Dunes State Park, Porter Co., in branches of P. banksiana Lamb.

Pityophthorus liquidambarus Blackman. Clark Co. State Forest, in branches of Liquidambar styraciflua L.

Pityophthorus pulchellus Ecihhoff. Hovey Lake, Posey Co.

Xyleborus planicollis Zimmermann. W. Lafayette, Tippecanoe Co., in light trap.

Specimens of all species are in the ABS collection except for X. planicollis (SLW) and P. pulchellus (FSC).

FLORIDA

Araptus sp. (accinctus Wood?). Monroe Station, Collier Co., in dead stem of Sarcostemma sp., one female specimen (ABS). First representative of genus known from U.S.

Chramesus hicoriae LeConte. Gainesville, Alachua Co., in fallen branch of Carya illinoensis (Wang.) K. Koch; in window trap in hardwood hammock (THA).

Corthylus punctatissimus (Zimmermann). Relay Management Tract. Flagler Co., in window trap with ethanol (THA).

Hylurgopinus rufipes (Eichhoff). Highlands Hammock State Park, Highlands Co., in bole of *Ulmus americana* L. (ABS).

Phloeosinus dentatus (Say). Gainesville, Alachua Co., in Juniperus silicicola (Small) Bailey (THA).

Phloeotribus frontalis (Olivier). Gainesville, Alachua Co., in Morus rubra L. (THA). Pityophthorus pullus (Zimmermann). Gainesville, Alachua Co., in Pinus elliotti Engelm., Tallahassee, Leon Co. (FSC).

Scolytus quadrispinosus Say. Gainesville, Alachua Co., in Carya sp. (THA). Xyleborus tachygraphus Zimmermann. Relay Management Tract, Flagler Co., in window trap with ethanol (THA).

Xylosandrus crassiusculus (Motschulsky). This species was recently reported from extreme northern Florida by Chapin and Oliver (1986). It also occurs on the Archbold Biological Station, Highlands Co., at Lake Worth, Palm Beach Co., and Collier-Seminole State Park, Collier Co. (ABS, FSC). This exotic species is therefore widely distributed in Florida, but unlike many exotic scolytids, seems to occur at very low population levels.

LITERATURE CITED

Chapin, J. B., and A. D. Oliver. 1986. New records for Xylosandrus and Xyleborus species (Coleoptera: Scolytidae). Proc. Entomol. Soc. Washington 88:680-683.

Devrup, M. 1981. Annotated list of Indiana Scolytidae. Great Lakes Entomol. 14:1-9.

Wood, S. L. 1982. The bark and ambrosia beetles of North and Central America (Coleoptera: Scolytidae), a taxonomic monograph. Great Basin Natur. Mem. 6:1-1359.