

June 1987

Distribution and Host Plants of *Corthylus Punctatissimus* (Coleoptera: Scolytidae) in the Lower Peninsula of Michigan

Richard A. Roeper
Alma College

Demetrios V. Zestos
Alma College

Brian J. Palik
Alma College

Larry R. Kirkendall
University of Oslo

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Roeper, Richard A.; Zestos, Demetrios V.; Palik, Brian J.; and Kirkendall, Larry R. 1987. "Distribution and Host Plants of *Corthylus Punctatissimus* (Coleoptera: Scolytidae) in the Lower Peninsula of Michigan," *The Great Lakes Entomologist*, vol 20 (2)

Available at: <https://scholar.valpo.edu/tgle/vol20/iss2/3>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

DISTRIBUTION AND HOST PLANTS OF *CORTHYLUS PUNCTATISSIMUS* (COLEOPTERA: SCOLYTIDAE) IN THE LOWER PENINSULA OF MICHIGAN

Richard A. Roeper¹, Demetrios V. Zestos¹, Brian J. Palik¹, and Larry R. Kirkendall²

The pitted ambrosia beetle, *Corthylyus punctatissimus* Zimmerman, infests woody saplings and shrubs 14 mm in diameter or less. The beetle bores an entrance hole into the main stem at soil level and constructs a main gallery tunnel which generally spirals downward in the stem. Egg-niche construction is followed by inoculation of symbiotic fungi and oviposition. The main stem of the host tree wilts as a result of the girdling activity of the beetle. Finnegan (1967) described the life history of *C. punctatissimus* infesting *Acer saccharum* Marshall in Ontario and Quebec.

The range of *C. punctatissimus* has been recorded from Colorado and Quebec to Arkansas and South Carolina by Finnegan (1967) and Wood (1982). Wood (1982) reported a specimen of *C. punctatissimus* from Midland County in Michigan. The woody host plants from which the beetle has been recorded include *Acer saccharinum* L. (silver maple); *A. saccharum* (sugar maple); *Berberis* sp. (barberry); *Carpinus caroliniana* Walter (American hornbeam); *Cornus* sp. (dogwood); *Corylus* sp. (hazel); *Gaylussacia* sp. (huckleberry); *Ostrya* sp. (hophornbeam); *Rhododendron* sp. (azaleas and rhododendron); *Sassafras* sp. (sassafras); *Ulmus* sp. (elm); *Vaccinium* sp. (blueberry) and *Viburnum* sp. (cranberry) (Finnegan 1967, Wood 1982).

In this study infested hosts were distinguished by their wilted condition. Infested plants wilt so quickly that leaves do not undergo abscission and can remain attached even after a year's time. The infested hosts were excavated and dissected to confirm the presence of the beetle. Collection sites of *C. punctatissimus* were recorded by county, hosts, number of collections (in parentheses), and range of collection dates.

COLLECTION RECORDS

Antrim Co. *Acer saccharum*, (3) 5–8 October.

Charlevoix Co. *A. saccharum*, (1) 5 October.

Genesee Co. *Cornus* sp., (1) 7 June.

Grand Traverse Co. *A. saccharum*, (1) 7 October.

Gratiot Co. *Acer negundo* L., (1) 11 October; *Acer rubrum* L., (214) 17 April–3 November; *A. saccharum*, (186) 15 March–15 November; *Quercus rubra* L., (12) 5 June–15 September; *Prunus serotina* Ehrhart, (15) 5 June–15 September; *Ulmus americana* L., (1) 7 October.

Jackson Co. *A. saccharum*, (1) 13 September.

Kalamazoo Co. *A. saccharum*, (1) 12 August.

Lapeer Co. *P. serotina*, (1) 7 June.

Livingston Co. *A. rubrum*, (3) 13 August–18 September; *A. saccharum*, (1) 12 August; *Carpinus caroliniana*, (1) 21 August; *Cornus racemosa* Larmarck, (1) 13 August; *Corylus americana* Walter, (1) 13 August; *Quercus velutina* Lamarck, (1) 13 August.

¹Biology Department, Alma College, Alma, MI 48801.

²Zoological Institute, University of Oslo, P.O. Box 1050, Blindern, Oslo 3, Norway.

Montcalm Co. *A. rubrum*, (1) 6 June; *A. saccharum*, (27) 5 May–20 October; *Carpinus caroliniana*, (1) 27 September; *Cornus racemosa*, (1) 27 September; *Ostrya virginiana* (Miller) K. Koch, (1) 7 October; *Sassafras albidum* (Nuttall) Ness, (1) 21 September; *Viburnum dentatum* L., (1) 1 October; *Viburnum dentatum* L., (1) 27 September.

Montmorency Co. *A. saccharum*, (1) 20 October; *Fagus grandifolia* Ehrhart, (1) 20 October; *Amelanchier* sp., (1) 20 October.

Oakland Co. *A. rubrum*, (1) 7 June.

Oceana Co. *S. albidum*, (1) 20 October.

Osceola Co. *F. grandifolia*, (1) 1 February.

St. Joseph Co. *A. saccharum*, (1) 12 August.

This study indicates *C. punctatissimus* is widely distributed throughout the Lower Peninsula of Michigan with all collections representing new county records. New hosts include *Acer rubrum* (red maple), *A. negundo* (boxelder), *Fagus grandifolia* (American beech), *Quercus rubra* (red oak), *Q. velutina* (black oak), *Amelanchier* sp. (serviceberry), *Viburnum dentatum* (arrowwood) and *V. trilobum* (highbush cranberry). The beetle probably can infest most deciduous woody tree saplings or shrubs within its range.

ACKNOWLEDGMENTS

The authors thank Dr. D. C. L. Gosling, C. Mathieson, C. Duke, P. Hesch, and D. Potter for assistance in collection of specimens, and Dr. R. O. Kapp for host plant determinations.

LITERATURE CITED

- Finnegan, R. J. 1967. Notes on the biology of the pitted ambrosia beetle, *Corthylus punctatissimus* (Coleoptera: Scolytidae), in Ontario and Quebec. Canadian Entomol. 99:49–54.
- Wood, S. L. 1982. The bark and ambrosia beetles of North and Central America (Coleoptera: Scolytidae), a taxonomic monograph. Great Basin Natur. Mem. 6:1–1359.