

The Great Lakes Entomologist

Volume 20
Number 4 - Winter 1987 *Number 4 - Winter*
1987

Article 2

December 1987

Influence of Temperature on Egg Production in Sunflower Stem Weevil *Cylindrocopturus Adspersus* (Coleoptera: Cucurlionidae)

John F. Barker
Metabolism and Radiation Research Laboratory

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Barker, John F. 1987. "Influence of Temperature on Egg Production in Sunflower Stem Weevil *Cylindrocopturus Adspersus* (Coleoptera: Cucurlionidae)," *The Great Lakes Entomologist*, vol 20 (4)
Available at: <https://scholar.valpo.edu/tgle/vol20/iss4/2>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in The Great Lakes Entomologist by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

**INFLUENCE OF TEMPERATURE ON EGG PRODUCTION IN
SUNFLOWER STEM WEEVIL *CYLINDROCOPTURUS ADSPERSUS*
(COLEOPTERA: CUCURLIONIDAE)**

John F. Barker¹

ABSTRACT

Controlled laboratory experiments were conducted to study the influence of temperature on production of eggs by a sunflower stem weevil *Cylindrocopturus adspersus*. Maximum egg production occurred when the temperature was $30 \pm 1^\circ\text{C}$. At 27°C the rate of egg production was lower than at 30°C but significant only at the $P < 0.1$ level. At 20, 23, and 35°C egg production was significantly lower than at 30°C ($P < 0.05$) indicating that temperature can be an important factor in regulating the fecundity of this species.

The known range of *Cylindrocopturus adspersus* (LeConte), a sunflower stem weevil, is from southern Canada throughout the Great Plains (Oseto 1977, Rogers and Jones 1979) into the Texas high plains (Rogers and Serda 1982). In the northern Great Plains, *C. adspersus* is reproductively active during late June, July, and August. Populations of this species are sporadic and cyclic which strongly suggests that various environmental factors influence their abundance. The influence of temperature on egg production was investigated to determine the influence of temperature on fecundity of this species.

MATERIALS AND METHODS

Larvae of *C. adspersus* were field collected from sunflower stalks and stored at 4°C . After chilling for at least five weeks the larvae were brought out to 27°C to pupate. Pupae were sexed as described by Reinecke (1981) and separated shortly after eclosion. Five to seven days after emergence as adults, the males were marked with a small spot of quick drying enamel paint and six males and six females were added to each of five hat boxes measuring $17 \times 26 \times 34$ mm ($n = 30$ females for each temperature tested). The box covers had ventilation holes cut into them, and the holes were covered with a screen to keep the weevils confined. Each cage contained a short section of sunflower stalk and a cut sunflower leaf. To maintain freshness the leaf and stem were inserted into a small bottle of water. The leaf and stem were changed every 48 h. All experiments were conducted in a controlled environment chamber. Temperature was held constant $\pm 1^\circ\text{C}$ with relative humidity maintained at 55%. Temperature effects were tested at 20, 23, 27, 30, and 35°C . Each test was made with five replicates. The photophase was held at 15:9 light:dark cycle. Fluorescent light intensity at the level of the holding shelf was 350 lux.

The eggs oviposited during each 48-h period were used to quantify the rate and total egg production. *C. adspersus* females oviposited on the cut stalk and leaf stem of the

¹Metabolism and Radiation Research Laboratory, USDA, ARS, Fargo, ND 58105.

Table 1. Effect of temperature on egg production in *C. adspersus*.

Temperature	Eggs/female/day	Total egg production ^a
35±1°C	3.2±0.3 ^b A	1483
30±1°C	4.7±0.4 B	5842
27±1°C	3.6±0.3 AB	4826
23±1°C	1.4±0.1 C	1945
20±1°C	0.4±0.0 D	711

^aUp to 75 days.

^bDifferent letters indicate the means are significantly different at the $P < 0.05$ level.

sunflower plant placed in the hat box. Eggs were removed from the section of stalk and cut leaf and counted. The mean numbers of eggs in different temperature groups were compared by a one-way analysis of variance and an *F* test (Guenther 1964).

RESULTS

The effect of temperature on total egg production and the number of eggs per female per day are summarized in Table 1. The total number of eggs oviposited and the number of eggs per female per day increased up to 30°C, while longevity decreased with increasing temperature. Each experiment at each temperature began with 30 females. The majority of females (22 of 30) held at 27 and 30°C lived at least 45 days while the majority (19 and 23 respectively of 30) held at 23 and 20°C were still alive at 75 days. In contrast, those held at 35°C were all dead by 30 days. The greatest number of eggs were produced when the temperature was 30±1°C. Oviposition at 35, 23, and 20°C was significantly lower than at 30°C. At 27°C egg production was lower than at 30°C but significant only at the $P < 0.1$ level.

DISCUSSION

The above results suggest temperature is important in the regulation of egg production in *C. adspersus*. Reproduction is inhibited at fairly warm temperatures of 20–23°C and is lower even at 27°C than at 30°C. It is probable that extended summertime temperatures below 27°C can contribute to down-cycles of wild populations of this species in the northern Great Plains. At 35°C the mortality was high with a subsequent reduction in total egg production. Temperatures above 30°C could reduce fecundity through increased stress and mortality. At temperatures of 20–23°C, the insects produce about 12–33% as many eggs, respectively, as at 30°C. Longevity increased almost two fold at these temperatures but the insects would have to live three times as long or longer to produce the equivalent number of eggs as at 30°C. It is unlikely that under natural conditions in the northern Great Plains, egg production at 20 or 23°C could catch up to production at 30°C due to the onset of fall conditions and the maturation and drying of the host plant occurring at the end of August.

ACKNOWLEDGMENTS

I thank Sharon Grugel and Kelly Jones for providing the insects for this study.

LITERATURE CITED

- Guenther, W. C. 1964. Analysis of variance. Prentice-Hall Inc., Englewood Cliffs, N. J.
- Oseto, C. Y. 1977. Biology and control of stem weevils, *Cylindrocopturus adspersus* and *Baris stremua*. Sunflower Forum 2:4. Sunflower Assoc. Am.
- Reinecke, J. P. 1981. Sexing living pupae and teneral adults of a sunflower stem weevil *Cylindrocopturus adspersus*. Southwestern Entomol. 6:201-204.
- Rogers, C. E., and O. R. Jones. 1979. Effects of planting date and soil water on infestation of sunflower by larvae of *Cylindrocopturus adspersus*. J. Econ. Entomol. 72:529-531.
- Rogers, C. E., and J. G. Serda. 1982. *Cylindrocopturus adspersus* in sunflower: Overwintering and emergence patterns on the Texas high plains. Environ. Entomol. 11:154-156.