

# The Great Lakes Entomologist

---

Volume 22  
Number 2 - Summer 1989 *Number 2 - Summer*  
1989

Article 8

---

June 1989

## Occurrence of *Pomerantzia Kethleyi* (Acari: Prostigmata: Pomerantziidae) in Illinois and Minnesota

J. Kethley  
*Field Museum of Natural History*

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Kethley, J. 1989. "Occurrence of *Pomerantzia Kethleyi* (Acari: Prostigmata: Pomerantziidae) in Illinois and Minnesota," *The Great Lakes Entomologist*, vol 22 (2)  
Available at: <https://scholar.valpo.edu/tgle/vol22/iss2/8>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**OCCURRENCE OF *POMERANTZIA KETHLEYI*  
(ACARI: PROSTIGMATA: POMERANTZIIDAE)  
IN ILLINOIS AND MINNESOTA**

J. Kethley<sup>1</sup>

ABSTRACT

The soil mite *Pomerantzia kethleyi*, is recorded from Illinois and Minnesota for the first time. The microhabitat of the Minnesota site is the first record of a pomerantziid from coarse sand-gravel near the surface.

---

Soil mites of the family Pomerantziidae are rarely collected because they tend to occur at depths that are difficult to sample. Two California species are deep-soil taxa having mean depths of 27.4 cm for *P. prolata* Price 1971, and 63.2 cm for *P. benhami* Price 1974 (Price and Benham, 1976).

*Pomerantzia kethleyi* Price, 1975, is known only from the type series, collected from clay prairie soil at a depth of 35–65 cm in Lake Co., Indiana. Two deep soil samples from Illinois and one surface soil sample from Minnesota have yielded additional material of *P. kethleyi*. ILLINOIS: Will Co., Goodenow Woods, 2 February 1974, W. Suter, one deutonymph from clay soil at a depth of 40–50 cm; Franklin Co., Fitzgerrel State Park, 25 October 1982, JK, one female from clay soil at a depth of 50–60 cm. MINNESOTA: Polk Co., 8.2 mi East of Crookston on U.S. route 2, 12 August 1986, JK, from coarse sand and gravel at a depth of 0–10 cm; the specific site is part of the eastern sand ridge formed by the Campbell Stage of glacial Lake Agassiz.

These new records significantly expand the distribution of *P. kethleyi*, and are suggestive of a considerable geographic range for this species. The specific microhabitat of the Minnesota site is the first indication that a species of *Pomerantzia* occurs in coarse sand-gravel near the surface.

LITERATURE CITED

- Price, D. W. 1975. Notes on the genus *Pomerantzia* Baker with a description of a new species from Indiana (Acarina: Pomerantziidae). Proc. Entomol. Soc. Wash. 77:487–490.  
Price, D. W. and G. S. Benham, Jr. 1976. Vertical distribution of pomerantziid mites (Acarina: Pomerantziidae). Proc. Entomol. Soc. Wash. 78:309–313.

---

<sup>1</sup>Division of Insects, Field Museum of Natural History, Chicago, Illinois 60605