

October 1990

Typocerus Deceptus in Southern Illinois (Coleoptera: Cerambycidae)

William T. McDowell
Southern Illinois University

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

McDowell, William T. 1990. "*Typocerus Deceptus* in Southern Illinois (Coleoptera: Cerambycidae)," *The Great Lakes Entomologist*, vol 23 (3)

Available at: <https://scholar.valpo.edu/tgle/vol23/iss3/10>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

**TYPOCERUS DECEPTUS IN SOUTHERN ILLINOIS
(COLEOPTERA: CERAMBYCIDAE)**William T. McDowell¹

ABSTRACT

The lepturine cerambycid *Typocerus deceptus* is reported from southern Illinois for the first time. It was collected in association with *T. v. velutinus* near, or in oak-hickory forest stands on *Hydrangea arborescens*, a new host plant record.

Typocerus deceptus was originally described by Knull (1929) from a small series of specimens collected in Pennsylvania. He collected it on *Rhus glabra* during July in Ohio but stated it was never common. Linsley and Chemsak (1976) added Indiana, Virginia, and South Carolina as new state records and found *T. deceptus* on *Rhus glabra* during June and July. Bouseman (1977) listed it from Kankakee, Mason, and Tazewell counties in north-central Illinois, and collected it on flowers of *Ceanothus americanus* and *Cornus* sp. during June and July, and felt that oak-hickory forest stands were used by larvae as host sites. Gosling (1981, 1983) collected it in St. Joseph County, Michigan on flowers of *Ceanothus americanus*, *Cephalanthus occidentalis*, *Rosa palustris* and *Daucus carota* during July and near oak-hickory forest stands. Gosling (1981) also found *T. deceptus* in copula and determined that *T. v. velutinus* (Olivier)(nearest sibling) was much more common (ratio of 11.7 *T. v. velutinus* to 1.0 *T. deceptus*).

MATERIALS AND METHODS

At irregular intervals, beginning in 1985, lepturine cerambycids occurring on *Hydrangea arborescens* were collected at Bell Smith Springs and Lusk Creek, Pope County. In addition, an extensive field study of lepturines occurring on *H. arborescens* and *Daucus carota* was conducted in the LaRue-Pine Hills Ecological Area, Union County, during 1987-1988. Both counties are located in southern Illinois. All specimens of *T. deceptus* are deposited in SIUC Entomology Collection.

RESULTS AND DISCUSSION

Thirty-one *Typocerus deceptus* were collected at the three localities. They were collected only on flowers of *H. arborescens* and only during June (N = 18) and July (N = 13), even though extensive collecting was conducted during the flowering period (June and July) of *D. carota* at LaRue-Pine Hills Ecological Area during 1987-88. At no time were any *T. deceptus* found in copula although *T. v. velutinus*,

¹Biological Sciences Program, Southern Illinois University, Carbondale, IL 62901

occurring with it on flowers, was commonly found mating. All specimens were found in or near oak-hickory forest stands and always in association with *T. v. velutinus*. At the LaRue-Pine Hills Ecological Area, *T. v. velutinus* was most abundant (N = 152), followed by *T. deceptus* (N = 31), *T. lugubris* (Say) (N = 9), and *T. acuticauda* Casey (N = 3). The *T. deceptus* sex ratio was 1.21 ♂:1 ♀.

Results of this study have shown *T. deceptus* present in southern Illinois and associated with *H. arborescens*; it was not found on *D. carota*. All host records were on plants associated with oak-hickory forest stands supporting results of Bouseman (1977) and Gosling (1981, 1983). At no time were any *T. deceptus* collected or sighted in copula as Gosling (1981) found.

ACKNOWLEDGMENTS

I thank Dr. Jeff Heuther (Pennwalt Corp.) for initially identifying the *T. deceptus* specimens and Dr. John K. Bouseman (Illinois Natural History Survey) for confirmations and sex identifications. For critically reviewing the manuscript, I thank Dr. J. A. Beatty and Dr. J. E. McPherson (Dept. of Zoology, Southern Illinois University) and two anonymous reviewers.

LITERATURE CITED

- Bouseman, J. R. 1977. *Typocerus deceptus* Knull in Illinois, with discussion of habitat (Coleoptera: Cerambycidae). Coleop. Bull. 31:179-180.
- Gosling, D. C. L. 1981. Ecology of the Cerambycidae in a southwest Michigan woodland. Ph. D. Thesis. Univ. Michigan, Ann Arbor.
- _____. 1983. New state records of Cerambycidae from Michigan (Coleoptera). Great Lakes Entomol. 16:187.
- Knull, J. N. 1929. Additions to the lists of Buprestidae and Cerambycidae of Pennsylvania (Coleop.). Entomol. News 40:144-145.
- _____. 1946. The longhorned beetles of Ohio (Coleoptera: Cerambycidae). Ohio Biol. Surv. Bull. 39:133-354.
- Linsley, E. G. and J. A. Chemsak, 1972. Cerambycidae of North America. Part VI, no. 2. Taxonomy and classification of the subfamily Lepturinae. Univ. California Publ. Ent. 80:1-186.