

October 1992

Ixodes Dentatus (Acari: Ixodidae) in Michigan: First State Records and Occurrence on a Human

Edward D. Walker
Michigan State University

Melvin L. Poplar
Michigan Department of Public Health

Howard L. Russell
Michigan State University

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

Recommended Citation

Walker, Edward D.; Poplar, Melvin L.; and Russell, Howard L. 1992. "*Ixodes Dentatus* (Acari: Ixodidae) in Michigan: First State Records and Occurrence on a Human," *The Great Lakes Entomologist*, vol 25 (4)
Available at: <https://scholar.valpo.edu/tgle/vol25/iss4/9>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

IXODES DENTATUS (ACARI: IXODIDAE) IN MICHIGAN: FIRST STATE RECORDS AND OCCURRENCE ON A HUMANEdward D. Walker¹, Melvin L. Poplar², and Howard L. Russell¹

ABSTRACT

An *Ixodes dentatus* adult female was taken from a cottontail rabbit in Kalamazoo County, and a nymph from a child in Berrien County, in 1992 in Michigan. These findings represent the first records of *I. dentatus* in the state, and document an unusual parasitization of a human being by this species of tick.

Ixodes dentatus Marx is distributed in the eastern U.S., mainly occurring in coastal states, but it has been reported from several inland states as well, including Indiana and Ohio (Cooley and Kohls 1945, Clifford et al. 1961, Keirans and Clifford 1978, Kollars 1992). There are no previous records of this tick from Michigan. *Ixodes dentatus* normally parasitizes cottontail rabbits (*Sylvilagus floridanus*). It has also been found on several species of passerine birds, woodchucks, a white-footed mouse, and Norway rats (*op. cit.*). In this note, we report the presence of *I. dentatus* in Michigan, and document parasitism of a human being.

On May 15, 1992, in Kalamazoo County, a pet dog flushed a young cottontail rabbit from a hole. The dog's owner, a county health department sanitarian, noticed a tick on the rabbit and submitted it to the Michigan Department of Public Health (MDPH) for identification. We identified the tick as an adult, female *I. dentatus*. On June 15, 1992, parents in Berrien County, Michigan noticed a tick embedded in their child's head, between the ear and hairline. They submitted the tick to the state health department, via their county health department, for identification. We identified the tick as an *I. dentatus* nymph. The tick was not bloodfed.

Ixodes dentatus has not previously been identified from among 4,462 tick identification records compiled by the MDPH since 1968. These records include tick submissions from the public for identification by the MDPH; the Parasitology Laboratory, Michigan State University Veterinary Clinic; and the insect diagnostic service at the Department of Entomology, Michigan State University. Thus, our observations may represent new state distribution records for *I. dentatus*. Indeed, these two records in the southwestern part of the state may indicate that *I. dentatus* has recently become established in Michigan. Alternatively, *I. dentatus* may be endemic in the state, but has not previously been documented simply owing to a lack of surveys.

Experimentally, *I. dentatus* is a competent vector of *Borrelia burgdorferi*,

¹Department of Entomology, Michigan State University, East Lansing, MI 48824.

²Michigan Department of Public Health, Bureau of Environmental and Occupational Health, Insect and Rodent Control Unit, 3500 N. Logan St., P.O. Box 30035, Lansing, MI 48909.

the spirochete that causes Lyme disease (Telford and Spielman 1989a). This spirochete has been isolated from *I. dentatus* in Connecticut, Massachusetts, and Virginia (Anderson et al. 1989, Telford and Spielman 1989b, Levine et al. 1991), suggesting that *B. burgdorferi* may be maintained in a zoonotic cycle involving *I. dentatus* and rabbits, separate from the well-documented cycle involving *Peromyscus* mice and *Ixodes dammini* Spielman, Clifford, Piesman, and Corwin (Lane et al. 1991). Although we do not suspect that *I. dentatus* currently is an enzootic vector of the Lyme disease spirochete in Michigan, its presence along with *I. dammini* and *B. burgdorferi* in the state (Strand et al. 1992; E.D. Walker and M.L. Poplar, unpublished data) intimates that it may have some future role. Whether *I. dentatus* could transmit *B. burgdorferi* to humans depends upon whether this tick ever bites people. The frequency of this event is apparently very low (Anderson et al. 1989; R.A. Restifo, Vector-Borne Disease Unit, Ohio Department of Health, personal communication). Our observation of *I. dentatus* parasitizing a human documents this rare event in Michigan.

ACKNOWLEDGMENTS

We are grateful to J.E. Keirans, curator of the National Tick Collection, Georgia Southern University, for confirming the *I. dentatus* identifications. The nymph was deposited with that collection, specimen number RML 120803.

LITERATURE CITED

- Anderson, J.F., L.A. Magnarelli, R.B. LeFebvre, T.G. Andreadis, J.B. McAninch, G.C. Perng, and R.C. Johnson. 1989. Antigenically variable *Borrelia burgdorferi* isolated from cottontail rabbits and *Ixodes dentatus* in rural and urban areas. *J. Clin. Microbiol.* 27:13-20.
- Clifford, C.M., G. Anastos, and A. Elbl. 1961. Larval Ixodid ticks of the eastern United States. *Misc. Publ. Entomol. Soc. Am.* 2:213-237.
- Cooley, R.A. and G.M. Kohls. 1945. The genus *Ixodes* in North America. *Natl. Inst. Health Bull.* 184, 246 pp.
- Keirans, J.E. and C.M. Clifford. 1978. The genus *Ixodes* in the United States: a scanning electron microscope study and key to the adults. *J. Med. Entomol.* 2:1-149 (suppl.).
- Kollars, T.M., Jr. 1992. New host of *Ixodes dentatus* (Acari: Ixodidae) and first record in Tennessee. *J. Med. Entomol.* 29:715-716.
- Lane, R.S., J. Piesman, and W. Burgdorfer. 1991. Lyme borreliosis: relation of its causative agent to its vectors and hosts in North America and Europe. *Ann. Rev. Entomol.* 36:587-609.
- Levine, J.F., D.E. Sonenshine, W.L. Nicholson, and R.T. Turner. 1991. *Borrelia burgdorferi* in ticks (Acari: Ixodidae) from coastal Virginia. *J. Med. Entomol.* 28:668-674.
- Strand, R.M., E.D. Walker, and R.W. Merritt. 1992. Field studies on *Ixodes dammini* in the upper peninsula of Michigan. *Vector Control Bull. North Cent. States* 1:111-118.
- Telford, S.R. and A. Spielman. 1989a. Competence of a rabbit-feeding *Ixodes* (Acari: Ixodidae) as a vector of the Lyme disease spirochete. *J. Med. Entomol.* 41:482-490.
- _____. 1989b. Enzootic transmission of the agent of Lyme disease in rabbits. *Am. J. Trop. Med. Hyg.* 41:482-490.