



Montowska, Magdalena and Alexander, Morgan R. and Tucker, Gregory A. and Barrett, David A. (2015) Authentication of processed meat products by peptidomic analysis using rapid ambient mass spectrometry. *Food Chemistry*, 187 . pp. 297-304. ISSN 0308-8146

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/48918/2/FoodChem_08-03-2015.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

1 **Authentication of processed meat products by peptidomic analysis**
2 **using rapid ambient mass spectrometry**

3

4 Magdalena Montowska ^{a,b,*} Morgan R. Alexander ^c, Gregory A. Tucker ^d,
5 David A. Barrett ^{a,*}

6

7 ^a Centre for Analytical Bioscience, School of Pharmacy, University of Nottingham, University Park,
8 Nottingham NG7 2RD, United Kingdom

9 ^b Institute of Meat Technology, Poznan University of Life Sciences, Wojska Polskiego 31, Poznan 60-
10 624, Poland

11 ^c Laboratory of Biophysics and Surface Analysis, School of Pharmacy, University of Nottingham,
12 University Park, Nottingham NG7 2RD, United Kingdom

13 ^d Division of Nutritional Sciences, School of Biosciences, University of Nottingham, Sutton Bonington
14 Campus, Leics LE12 5RD, United Kingdom

15

16 **Corresponding Authors**

17 *MM, Present address: Institute of Meat Technology, Poznan University of Life Sciences,
18 Wojska Polskiego 31, Poznan 60-624, Poland. Tel.: +48 61 8487251; fax: +48 61 8487254.
19 *E-mail address:* magdalena.montowska@gmail.com (M. Montowska).

20 *DB, Tel.: +44 115 95 15062. Fax: +44 115 95 15102.

21 *E-mail address:* david.barrett@nottingham.ac.uk (D.A. Barrett).

22

23

24

25

26 **ABSTRACT**

27 We present the application of a novel ambient LESA-MS method for the authentication of
28 processed meat products. A set of 25 species and protein-specific heat stable peptide markers
29 has been detected in processed samples manufactured from beef, pork, horse, chicken and
30 turkey meat. We demonstrate that several peptides derived from myofibrillar and sarcoplasmic
31 proteins are sufficiently resistant to processing to serve as specific markers of processed
32 products. The LESA-MS technique required minimal sample preparation without fractionation
33 and enabled the unambiguous and simultaneous identification of skeletal muscle proteins and
34 peptides as well as other components of animal origin, including the milk protein such as casein
35 alpha-S1, in whole meat product digests. We have identified, for the first time, six fast type II
36 and five slow/cardiac type I MHC peptide markers in various processed meat products. The
37 study demonstrates that complex mixtures of processed proteins/peptides can be examined
38 effectively using this approach.

39

40

41 ***Keywords:***

42 Food adulteration, Ambient Mass Spectrometry, Liquid Extraction Surface Analysis Mass
43 Spectrometry, Peptide markers, Fast and slow type MHC isoforms

44

45

46

47

48

49

50

51 **1. Introduction**

52 Despite the existence of extensive mandatory regulations in most countries, food
53 adulteration is still a global issue which attracts attention at international level and increases
54 public concern regarding food quality. In 2013, the horse meat scandal revealed the weaknesses
55 in the food safety system and contributed to a decrease of confidence in the food industry.
56 Fraudulent practices, i.e. the presence of undeclared horse DNA in food products labeled as
57 containing beef, were confirmed in 4.66% and 0.61% of controlled foods in 2013 and 2014,
58 respectively (European Commission, 2014) as a result of tests in the 28 EU countries. Recent
59 studies have revealed an even higher level of food mislabeling, for example 68% mislabeling
60 was found in sausages, burger patties and meats collected from butcheries and retail outlets in
61 South Africa (Cawthorn, Steinman, & Hoffman, 2013) and in seafood in the USA, a rate of
62 33% of investigated samples were mislabeled according to U.S. Food and Drug Administration
63 (FDA) guidelines (Kimberly, Walker, Lowell, & Hirshfield, 2013). Similarly, the results of
64 inspections carried out in Poland in 2011 by the Office of Competition and Consumer
65 Protection (UOKiK) and Department of Trade Inspection revealed that 24.7% of the examined
66 batches of luxury processed meat products, i.e. conventional, traditional and organic products
67 sold at high prices, were adulterated/labeled incorrectly (UOKiK, 2012). Continuous
68 monitoring of food quality and safety is now mandatory in the EU and other countries but the
69 increasing sophistication of adulteration means that analytical methods require continuous
70 improvement to ensure effective fraud detection. The rigorous analysis of complex and
71 processed products requires the development of novel analytical methodology which has
72 potential for high-throughput analysis and provides rapid, specific and reliable results.

73 At present, established methods for meat speciation are based on ELISA and PCR
74 techniques, which are robust when applied to raw or moderately processed samples (Chen &
75 Hsieh, 2000; Ballin, Vogensen, & Karlsson, 2009; Fajardo, González, Rojas, García, & Martín,

76 2010; Köppel, Eugster, Ruf, & Rentsch, 2012). The reported lower efficiency of these methods
77 in highly processed samples has been linked to processing conditions, thermal denaturation and
78 degradation of the markers compounds monitored (typically DNA or protein epitope) and
79 problems with cross-reactivity between species giving unreliable results (Arslan, Irfan-Ilhak,
80 & Calicioglu, 2006; Şakalar, Abasiyanik, Bektik, & Tayyrov, 2012; Musto, Faraone, Cellini,
81 & Musto, 2014). The difficulty with reliable multiplex detection in a single test and
82 contamination of DNA from other organisms also place severe limitations on analysis of
83 complex samples. However, some proteins are quite resistant to heating (Buckley, Collins,
84 Thomas-Oates, & Wilson, 2009; Montowska & Pospiech, 2012; Buckley, Melton, &
85 Montgomery, 2013) and hence peptidomic analysis techniques have potential advantages when
86 applied to authenticate processed (cooked) food.

87 Recently, considerable improvement in mass spectrometry (MS) instrumentation has
88 enabled the detection of peptide markers by liquid chromatography-MS techniques (LC-MS)
89 and this has enabled identification of specific proteins from soybean (Leitner, Castro-Rubio,
90 Marina, & Lindner, 2006), fish (Carrera et al., 2011) and meat species (Buckley et al., 2009;
91 Sentandreu, Fraser, Halket, Patel, & Bramley, 2010; Montowska & Pospiech, 2013; von
92 Bargaen, Brockmeyer, & Humpf, 2014). In our previous work, we evaluated ambient MS
93 techniques for standard protein identification in mixtures and for the analysis of meat digests
94 to discriminate between five meat species (Montowska, Rao, Alexander, Tucker, & Barrett,
95 2014a). Subsequently, we detected heat stable peptide markers derived from meat proteins after
96 thermal denaturation using our previously introduced ambient liquid extraction surface analysis
97 mass spectrometry (LESA-MS) methodology (Montowska, Alexander, Tucker, & Barrett,
98 2014b).

99 It is known that the primary structure of some meat proteins is relatively resistant to
100 processing and that certain skeletal muscle proteins are both species- and tissue-specific and

101 hence there is good potential for the use of specific muscle proteins and peptide markers for
102 meat authentication (Buckley et al., 2009, 2013; Sentandreu & Sentandreu, 2011; Montowska
103 & Pospiech, 2012). We consider that the ease of use and rapid nature of ambient MS has
104 advantages for high-throughput screening of processed food and we wish to explore the
105 potential application of our LESA-MS peptidomic approaches (Montowska et al., 2014a,b) for
106 this purpose. We suggest that the peptidomic analysis can serve as a tool not only for species
107 identification but also for the assessment of the quality of the product. In this study we define
108 ‘product quality’ as a general term linked with different authenticity issues, such as the
109 detriment of the quality of the product by illegal change of meat to less valuable components
110 of animal origin (e.g. meat of lower class, offal, connective tissue, blood plasma), undeclared
111 plant or milk additives as well as a change in proportion of ingredients. Analysis of myosin
112 isoforms due to their extensive diversity may help to trace some illegal practices in processed
113 meat products.

114 Unlike highly conserved actin, myosin exhibits extensive variations in vertebrate striated
115 muscles, which is translated into differences in fibre composition and shortening velocity. In
116 adult mammals, pure fibres (slow type I red, and fast type white IIA, IIX, IIB) are expressed
117 by a single myosin heavy chain (MHC) isoform (1, 2A, 2X, and 2B) whereas hybrid fibres may
118 contain several MHC isoforms (1/2A, 2A/1, 2AX, 2XA, 2XB, and 2BX) each encoded by a
119 separate gene (Pette & Staron, 2000). Therefore in this study, besides species identification,
120 we wish to identify heat stable peptides unique to fast and slow type MHC isoforms.
121 Peptidomic analysis may be a viable way to discriminate between the processed meat and non-
122 meat components to examine the quality of the processed meat products.

123 In this paper, we present the application of our previously established LESA-MS
124 methodology (Montowska et al., 2014a,b) for detection and identification of heat stable beef,
125 pork, horse and poultry peptide markers in various processed meat products. This rapid

126 peptidomic approach aims to identify heat stable peptides without the need for purification and
127 chromatographic separation. We also describe the application of in-solution tryptic digestion
128 of processed meat samples followed by deposition onto a polymer surface, desorption and
129 direct analysis by LESA-MS for protein/peptide composition of processed meat in order to
130 compare the identified MHC isoforms and select heat stable peptides unique to fast and slow
131 type MHCs.

132

133 **2. Materials and methods**

134 *2.1. Preparation of samples*

135 Meat products (n=18) were purchased at English and Polish supermarkets or
136 manufactured in our own pilot plant. Samples of raw sausages were cooked from chilled in an
137 oven at 190°C for 30 min according to the manufacturers' instructions. In-house processed
138 sausages (3 batches) were prepared in a pilot plant of the Institute of Meat Technology in
139 Poznan (Poland) exclusively from cured pork with the addition of spices and were coarsely
140 minced, smoked and cooked. All samples of about 5 cm length or 5 g were cut from fresh
141 products and kept at -80 °C until further MS analysis. Sample information and details about
142 processing methods and meat composition are given in Table 2.

143 Washing, digestion and mass spectrometry analysis were performed according to the
144 procedure described previously (Montowska et al., 2014b). Preparation for LESA-MS analysis
145 of samples of processed meat products involved washing procedures followed by digestion.
146 For this purpose, thin sections of sausages (slices of 0.5 g) or 1 g of meat spreads were
147 transferred to glass vials and washed to remove contaminants such as physiological salts, fat,
148 and other soluble low molecular weight compounds. Sample was rinsed twice for 30 s in
149 ethanol/water (70:30) followed by a 15 s wash in ethanol and then by a 30 s wash in
150 methanol/water (90:10). The sample then was rinsed for 2 x 30 s in deionized water, and finally

151 for 30 s in 100 mM of aqueous ammonium bicarbonate. Washed samples were placed to dry
152 for 30 min in a desiccator.

153

154 *2.2. In-solution digestion*

155 Dried samples (10 mg) were rehydrated in 100 μ L of water and subsequently digested in
156 a solution containing 0.083 μ g/ μ L of trypsin in ammonium bicarbonate at room temperature
157 over a period of 24 h. Digested solution was then centrifuged for 10 min at 13400 rpm, and the
158 supernatant was diluted 10-fold with deionized water. Samples of 1 μ L were spotted onto a
159 Permanox slide, 75 x 25 mm (Nunc, Thermo Fisher Scientific, Rochester, NY, USA) and
160 allowed to evaporate in air at room temperature prior to analysis.

161

162 *2.3. LESA Mass Spectrometry*

163 The LESA source was a TriVersa NanoMate (Advion, Ithaca, NY) coupled to a Thermo
164 Fisher LTQ Velos ion trap mass spectrometer (Thermo Fisher Scientific, San Jose, CA)
165 operated in positive-ion electrospray ionization mode. The NanoMate platform operated at
166 nanoESI tip voltage of 1.6 kV, with a gas pressure of 0.4 psi and a capillary temperature of
167 190°C. The same spray/extraction solvent acetonitrile/water/formic acid (50:50:1) was used in
168 all LESA experiments. Total solvent extraction volume was 5 μ L, dispensed and aspirated
169 volumes were 3.5 and 3.2 μ L, respectively. Each data set was collected from a single protein
170 spot. Data-dependent analysis (DDA) tandem MS/MS data were collected in full scan mode
171 with m/z range of 50-2000 divided into four segments (m/z 60-600, 550-1050, 1000-1550 and
172 1500-2000), 1 microscan, 100 ms max injection time, AGC mode on. DDA mode as well as
173 standard MS/MS experiments were used for the analysis of samples. Collision-induced
174 dissociation (CID) experiments were performed at a normalized collision energy of 38%. Data
175 were analyzed using Xcalibur software (Thermo Fisher Scientific). For protein and peptide

176 identification raw files were converted to MASCOT generic format using MSCONVERT
177 provided by the ProteoWizard project (<http://proteowizard.sourceforge.net/tools.shtml>). The
178 resulting files were searched via MS/MS ions search using MASCOT against the SwissProt
179 and the National Center for Biotechnology Information (NCBI) databases with the following
180 parameters: trypsin enzyme, taxonomy bone vertebrates, one missed cleavage, peptide mass
181 tolerance of 1.2 Da, MS/MS tolerance 0.6 Da, no modifications, peptide charge 1+, 2+ and 3+.
182 All samples were analyzed at least in three technical replicates. A decoy search was performed
183 automatically and the matches and MASCOT scores were evaluated at 1% of a false discovery
184 rate (FDR) for identity and homology threshold. Selected peptides in FASTA format were
185 searched against the NCBI database using the protein BLAST alignment research tool and
186 blastp algorithm for species and protein specificity.

187

188 **3. Results and discussion**

189 *3.1. Myofibrillar proteins as a source of heat stable peptide markers*

190 The workflow for the rapid identification of heat stable peptide markers in processed
191 meat products proposed in this study is presented in Figure 1. The methodology involves three
192 main steps: (a) washing thin slices of samples to remove contaminants which would interfere
193 with the electrospray signal, (b) in-solution tryptic digestion, and (c) data collection using
194 LESA-MS/MS with the acquisition time of 4 min. We focused on the identification of the more
195 abundant proteins using readily detectable peptides which were resistant to processing as well
196 as unique to both species and meat protein. Although, we observed that the sequence coverage
197 of proteins for processed meat products was lower compared to raw and cooked meats
198 (Montowska et al., 2014b) we found this methodology robust and specific and competitive to
199 LC-MS methods, especially when monitoring particular heat stable peptides (see following
200 sections). In our opinion three factors enhanced the efficiency of the analysis: (1) washing all

201 samples, (2) purification/centrifugation to remove undigested material and reduce the mixture
202 complexity as well as effect of ion suppression, and (3) achieving stable nanoelectrospray
203 during MS data acquisition.

204 We examined the applicability of the LESA-MS method using various commercial
205 processed meat products purchased from supermarkets as well as in-house processed sausages
206 as reference samples. Data sets were collected with data-dependent tandem LESA-MS for
207 screening of whole products digests and with standard MS/MS experiments using a list of 80
208 ions which was generated in our previously reported studies of cooked beef, pork, horse,
209 chicken and turkey meat (Montowska et al., 2014b). The dominant MHC isoforms found in
210 processed meat products and their MASCOT output scores are shown in Table 1. In the
211 processed samples we identified the same most abundant skeletal muscle and sarcoplasmic
212 proteins as previously detected for cooked meat, such as MHCs, MLCs, actin, tropomyosin,
213 myoglobin, GAPDH, beta-enolase (Montowska et al., 2014b), but the MASCOT scores and
214 sequence coverages were lower in the case of processed products analysed. In this study turkey
215 MHC was classified to the closely related chicken species since no full sequence of turkey
216 myosin has been published and only short fragments are available in the NCBI database.

217 A list of peptide markers identified in this study for the 18 different processed meat
218 products is presented in Table 2. Most of the observed peptides were identified as heat stable
219 markers belonging to MHCs, MLCs and myoglobin, and were unique to both species and single
220 muscle protein. Predominantly, MASCOT scores were above the homology or identity
221 threshold and all presented peptides were ranked first in the list of matched peptides of
222 MASCOT results. Figure 2 shows a typical MS/MS spectrum of pork fast type myosin-1 and
223 myosin-4 marker SALAHAVQSSR (563.67²⁺) obtained from frankfurters (sample 15).

224 The limit of detection is a critical step towards quantitative analysis using the peptidomic
225 LESA-MS approach. At present, sensitivity rises when chromatographic separation is

226 involved, for example pork or horse meat can be detected down to 0.24% in processed products
227 using HPLC-MS//MS method coupled with multiple reaction monitoring (Von Bargaen,
228 Brockmeyer, & Humpf, 2014). In the aforementioned article, five peptide markers for
229 processed pork and horse meat have been identified. These findings overlap with our studies,
230 since three of the peptides, i.e., TLAFLFAER (pork), SALAHAVQSSR (pork), and
231 LVNDLTGQR (horse) were identified in cooked meats using LESA-MS/MS as described
232 previously (Montowska et al., 2014b). It is likely, that lower sensitivity compared with LC-MS
233 methods due to dynamic range of protein concentration and the lack of fractionation stage, may
234 be enhanced by the use of the LESA instrument with high resolution/accurate mass/MSⁿ mass
235 spectrometer.

236

237 *3.2. Sarcoplasmic proteins as a source of heat stable peptide markers*

238 We also present the detection of heat-stable species-specific markers for sarcoplasmic
239 proteins (glyceraldehyde-3-phosphate dehydrogenase GAPDH, myoglobin, beta-enolase),
240 which can address the needs of species identification (Table 2). Myoglobin peptides were found
241 to be good markers for processed red meats, i.e., beef and horse meat, whereas pork GAPDH
242 peptide WGDAGATYVVESTGVFTTMEK (1125.32²⁺) is a good marker for the processed
243 products containing pork. This is consistent with previous studies, where enzymatic activity of
244 glycolytic enzymes, for instance GAPDH, was found to be higher in fast-twitch glycolytic
245 muscles (Takekura & Yoshioka, 1987; Okumura et al., 2005). In kabanos sausage (sample 10)
246 and frankfurters (sample 14) milk proteins were detected, and thus two peptides unique to
247 casein alpha-S1 (HQGLPQEVLNENLLR and EPMIGVNQELAYFYPELFR) were identified
248 with significant MASCOT output scores (Table 2). These results are consistent with the
249 product labels since the milk or cheese proteins were included in the list of ingredients. One
250 sample of pork sausages (sample 11) was declared to contain veal at 6%, another two samples

251 of cocktail sausage (sample 12) and frankfurters (sample 13) were labeled to contain turkey
252 meat at 6% and turkey MRM respectively, however we did not detect cattle and turkey markers
253 using our LESA methodology. We cannot exclude the possibility that the meat content was
254 below the limit of detection of the method. We have already shown in previous work that
255 ambient LESA-MS can detect 10% of cooked cattle, pork, horse, and turkey meat and 5% of
256 chicken meat in a beef matrix (Montowska et al., 2014b). Horse sausage (sample 9) turned out
257 to be made not only from horse meat but also from pork. However there was no list of
258 ingredients on the label, hence we cannot confirm the adulteration of this product.

259

260 *3.3. Discrimination between fast and slow type MHC isoforms in processed meat products*

261 Having identified markers for protein and species identification in processed meat
262 products our next stage was to discriminate between fast and slow type MHCs to identify heat
263 stable peptides unique to these isoforms. For this purpose, the same data sets as for meat
264 speciation obtained with DDA LESA-MS/MS were analysed individually. Similarly, each
265 potential marker was searched against the NCBI database with the BLAST tool for isoform
266 specificity. Table 3 shows six fast type II and five slow/cardiac type I MHC unique peptides
267 identified for beef, pork and horse meat using this approach. Examples of type I myosin-7
268 peptides obtained from cooked ham (sample 1) and fried horse sausage (sample 8) are shown
269 in Figure 3 as fragmented spectra for pig LLSNLFANYAGADTPVEK (962.10²⁺) and horse
270 MLSNLFANYLGADAPIEK (984.29²⁺). As far as we are aware, this is the first time that the
271 peptides specific to slow-twitch type 1 myosin-7 isoform were identified in processed meat
272 products.

273 Frequent detection of slow MHC isoform and peptides unique to slow type isoforms over
274 the fast MHC isoforms implies that processed pork and beef products investigated in this study
275 were manufactured mainly from smaller red or intermediate muscles. In our previous studies,

276 protein/peptide differentiation between samples of cattle, pig and horse *longissimus dorsi*
277 muscle was performed (Montowska et al., 2014a,b). Since this muscle is composed mostly of
278 fast, white fibres, peptides unique to fast myosin-1(2X) and myosin-4(2B) isoforms were
279 detected in those samples with the highest scores. In two samples of potted beef and corned
280 beef (samples 3 and 4) we were able to detect only slow myosin-7 isoform. The peptide
281 GQNVQQVVYAK, unique to beef myosin-7, was identified in both samples but with the
282 MASCOT scores below the identity and homology threshold (Table 2). This may indicate that:
283 (a) the amount of meat was below the declared content, (b) these products were manufactured
284 from the meat of lower class containing high amount of connective tissue and fat, and/or (c)
285 high amounts of non-meat components, such as collagen preparations, offal and fat were added
286 to the products. Only horse myoglobin was detected in horse sausage (sample 9), therefore the
287 ability/inability to detect markers of specific proteins and tissues may indicate a good/poor
288 quality ingredients in a given product or even an ingredient's substitution.

289 Because myosin is the most abundant muscle protein and its content corresponds to 40-
290 50% of the total muscle proteins, myosin peptides might be a good indicator of the meat content
291 in the processed products. They also might be used to assess the fibre-type composition of meat
292 components, and thus indirectly to assess the product quality. Although, markers obtained from
293 MHC isoforms, may be a robust tool to indicate the quality of meat ingredients, in our opinion
294 the authentication of processed meat product defined as the quality assessment needs to be
295 based on appropriate and reliable quantitative analysis of several peptide markers of meat and
296 non-meat origin in parallel.

297

298 **4. Conclusions**

299 We have examined the applicability of a novel and rapid LESA-MS method to identify
300 peptide markers in different types of processed meat products for authentication purposes. The

301 entire procedure is radically simplified over other peptidomic methodologies by excluding
302 fractionation steps before and after the protein digestion stage. Sample preparation is therefore
303 limited only to the processes of washing and digestion. By the application of data-dependent
304 LESA-MS/MS for fast screening of whole product digests, we were able to identify a set of 25
305 heat stable peptide markers derived from myofibrillar and sarcoplasmic proteins. We have also
306 proved that this novel method of peptidomic examination from processed meat products has
307 good specificity to readily identify peptide markers for fast and slow type MHC isoforms. We
308 have demonstrated that there is a set of specific peptides resistant to thermal treatment and
309 easily detectable in various industrially processed meat products which not only serve as
310 markers of meat speciation but also can help to track down other illegal practices linked with
311 the substitution of ingredients.

312 Since this work was focused on authenticity issues, only peptides from the most abundant
313 proteins were identified. However, the use of LESA interfaced with high resolution mass
314 spectrometry may enhance sensitivity sufficiently to enable analysis of less abundant proteins.
315 This easy to use and versatile ambient methodology has great potential to be implemented in
316 the routine, rapid high-throughput screening of processed products, and in addition displays
317 specificity sufficient to enable examination of other important issues in meat science, e.g.,
318 variations in muscle metabolism and meat quality.

319

320

321

322

323

324

325

326 **Acknowledgements**

327 The Postdoctoral Fellowship of Magdalena Montowska was supported by the European
328 Commission under the Marie Curie Intra-European Fellowship Programme (Call: FP7-PEOPLE-2011-
329 IEF).

330

331 *The contents reflect only the authors' views and not the views of the European Commission.*

332

333

334 **References**

335 Arslan, A., Ilhak, O.I., & Calicioglu, M. (2006). Effect of method of cooking on identification of heat
336 processed beef using polymerase chain reaction (PCR) technique. *Meat Science*, 72(2), 326–330.

337 Ballin, N.Z., Vogensen, F.K., & Karlsson, A.H. (2009). Species determination - Can we detect and
338 quantify meat adulteration? *Meat Science*, 83(2), 165–174.

339 Buckley, M., Collins, M., Thomas-Oates, J., & Wilson, J.C. (2009). Species identification by analysis
340 of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass
341 spectrometry. *Rapid Communications in Mass Spectrometry*, 23(23), 3843–3854.

342 Buckley, M., Melton, N.D., & Montgomery, J. (2013). Proteomics analysis of ancient food vessel
343 stitching reveals >4000-year-old milk protein. *Rapid Communications in Mass Spectrometry*,
344 27(4), 531–538.

345 Carrera, M., Cañas, B., López-Ferrer, D., Piñeiro, C., Vázquez, J., & Gallardo, J.M. (2011). Fast
346 monitoring of species-specific peptide biomarkers using high-intensity-focused-ultrasound-
347 assisted tryptic digestion and selected MS/MS ion monitoring. *Analytical Chemistry*, 83(14),
348 5688–5695.

349 Cawthorn, D.-M., Steinman, H.A., & Hoffman, L.C. (2013). A high incidence of species substitution
350 and mislabelling detected in meat products sold in South Africa. *Food Control*, 32(2), 440–449.

351 Chen, F.C., & Hsieh, Y.H. (2000). Detection of pork in heat-processed meat products by monoclonal
352 antibody-based ELISA. *Journal of AOAC International*, 83(1), 79–85.

353 European Commission (2014). Outcome of the coordinated control plan with a view to establish the
354 prevalence of fraudulent practices in the marketing of certain foods.
355 http://ec.europa.eu/food/food/horsemeat/tests_results_en.htm (accessed November 20, 2014).

356 Fajardo, V., González, I., Rojas, M., García, T., & Martín, R. (2010). A review of current PCR-based
357 methodologies for the authentication of meats from game animal species. *Trends in Food Science
358 & Technology*, 21(8), 408–421.

359 Kimberly, W., Walker, T., Lowell, B., & Hirshfield, M. (2013). Oceana study reveals seafood fraud
360 nationwide. *Oceana*, February, 1-69.
361 [http://oceana.org/sites/default/files/reports/National_Seafood_Fraud_Testing_Results_FINAL.p](http://oceana.org/sites/default/files/reports/National_Seafood_Fraud_Testing_Results_FINAL.pdf)
362 [df](http://oceana.org/sites/default/files/reports/National_Seafood_Fraud_Testing_Results_FINAL.pdf).

363 Köppel, R., Eugster, A., Ruf, J., & Rentsch, J. (2012). Quantification of meat proportions by measuring
364 DNA contents in raw and boiled sausages using matrix-adapted calibrators and multiplex real-
365 time PCR. *Journal of AOAC International*, 95(2), 494–499.

366 Leitner, A., Castro-Rubio, F., Marina, M.L., & Lindner, W. (2006). Identification of marker proteins
367 for the adulteration of meat products with soybean proteins by multidimensional liquid
368 chromatography-tandem mass spectrometry. *Journal of Proteome Research*, 5(9), 2424–2430.

369 Montowska, M., & Pospiech, E. (2012). Myosin light chain isoforms retain their species-specific
370 electrophoretic mobility after processing, which enables differentiation between six species: 2DE
371 analysis of minced meat and meat products made from beef, pork and poultry. *Proteomics*,
372 12(18), 2879–2889.

373 Montowska, M., & Pospiech, E. (2013). Species-specific expression of various proteins in meat tissue:
374 proteomic analysis of raw and cooked meat and meat products made from beef, pork and selected
375 poultry species. *Food Chemistry*, 136(3-4), 1461–1469.

376 Montowska, M., Rao, W., Alexander, M.R., Tucker, G.A., & Barrett, D.A. (2014a). Tryptic digestion
377 coupled with ambient desorption electrospray ionization and liquid extraction surface analysis
378 mass spectrometry enabling identification of skeletal muscle proteins in mixtures and
379 distinguishing between beef, pork, horse, chicken, and turkey meat. *Analytical Chemistry*, 86(9),
380 4479–4487.

381 Montowska, M., Alexander, M.R., Tucker, G.A., & Barrett, D.A. (2014b). Rapid detection of peptide
382 markers for authentication purposes in raw and cooked meat using ambient liquid extraction
383 surface analysis mass spectrometry. *Analytical Chemistry*, 86(20), 10257–10265.

384 Musto, M., Faraone, D., Cellini, F., & Musto, E. (2014). Changes of DNA quality and meat
385 physicochemical properties in bovine supraspinatus muscle during microwave heating. *Journal*
386 *of the Science of Food and Agriculture*, 94(4), 785–791.

387 Okumura, N., Hashida-Okumura, A., Kita, K., Matsubae, M., Matsubara, T., Takao, T., & Nagai, K.
388 (2005). Proteomic analysis of slow- and fast-twitch skeletal muscles. *Proteomics*, 5(11), 2896–
389 2906.

390 Pette, D., & Staron, R.S. (2000). Myosin isoforms, muscle fiber types, and transitions. *Microscopy*
391 *Research and Technique*, 50(6), 500–509.

392 Şakalar, E., Abasiyanik, M.F., Bektik, E., & Tayyrov, A. (2012). Effect of heat processing on DNA
393 quantification of meat species. *Journal of Food Science*, 77(9), N40–44.

- 394 Sentandreu, M.A., Fraser, P.D., Halket, J., Patel, R., & Bramley, P.M. (2010). A proteomic-based
395 approach for detection of chicken in meat mixes. *Journal of Proteome Research*, 9(7), 3374–
396 3383.
- 397 Sentandreu, M.A., & Sentandreu, E. (2011). Peptide biomarkers as a way to determine meat
398 authenticity. *Meat Science*, 89(3), 280–285.
- 399 Takekura, H., & Yoshioka, T. (1987). Determination of metabolic profiles on single muscle fibres of
400 different types. *Journal of Muscle Research and Cell Motility*, 8(4), 342–348.
- 401 Urząd Ochrony Konkurencji i Konsumentów (UOKiK - Office of Competition and Consumer
402 Protection) (2012). Raport: Produkty żywnościowe z segmentu luksusowych w świetle kontroli
403 przeprowadzonych przez Inspekcję Handlową. Warsaw, December 2012 [in Polish].
404 http://www.uokik.gov.pl/raporty_z_kontroli_inspekcji_handlowej.php (accessed November 20,
405 2014).
- 406 Von Barga, C., Brockmeyer, J., & Humpf, H.-U. (2014). Meat authentication: a new HPLC-MS/MS
407 based method for the fast and sensitive detection of horse and pork in highly processed food.
408 *Journal of Agricultural and Food Chemistry*, 62(39), 9428–9435.
409

Table 1. Dominant skeletal myosin heavy chain (MHC) isoforms identified in tryptic digests of processed meat products

No	Sample	Identified protein	NCBI accession number	Matched peptides ^a	Sequence coverage (%) ^b	MASCOT score ^c
1	Cooked ham	myosin-2 (<i>Sus scrofa</i>)	gi 55741490	55	20	388
		myosin-1 (<i>Sus scrofa</i>)	gi 157279731	51	19	388
		myosin-7 (<i>Sus scrofa</i>)	gi 55741486	31	12	226
2	Beef spread	myosin-2 (<i>Bos taurus</i>)	gi 75055812	54	21	429
3	Potted beef	myosin-7 (<i>Bos taurus</i>)	gi 41386711	25	14	162
4	Corned beef	myosin-7 (<i>Bos taurus</i>)	gi 41386711	54	21	519
5	Beef sausage	myosin-2 (<i>Bos taurus</i>)	gi 75055812	66	29	419
		myosin-1 (<i>Bos taurus</i>)	gi 41386691	66	28	434
6	Chorizo	myosin-1 (<i>Sus scrofa</i>)	gi 157279731	50	20	547
		myosin-7 (<i>Sus scrofa</i>)	gi 55741486	24	11	190
7	Hunters sausage	myosin-4 (<i>Sus scrofa</i>)	gi 178056718	57	24	392
8	Horse sausage	myosin-2 (<i>Equus caballus</i>)	gi 126352598	71	26	842
		myosin-7 (<i>Equus caballus</i>)	gi 126352320	48	19	702
9	Horse sausage (smoked)	myosin-7 (<i>Equus caballus</i>)	gi 126352320	30	14	155
		myosin-1 (<i>Sus scrofa</i>)	gi 157279731	20	9	86
10	Kabanos sausage with cheese	myosin (<i>Gallus gallus</i>)	gi 13432175	40	19	376
		myosin-4 (<i>Sus scrofa</i>)	gi 178056718	38	19	282
11	Pork sausage	myosin-4 (<i>Sus scrofa</i>)	gi 178056718	55	22	797
12	Cocktail sausage	myosin-2 (<i>Bos taurus</i>)	gi 75055812	54	22	448
13	Frankfurters poultry	myosin (<i>Gallus gallus</i>)	gi 13432175	54	21	520
14	Frankfurters classic	myosin-1 (<i>Sus scrofa</i>)	gi 157279731	46	16	425
15	Frankfurters	myosin-2 (<i>Bos taurus</i>)	gi 75055812	53	20	325
		myosin-1 (<i>Sus scrofa</i>)	gi 157279731	39	15	403
16	Hotdogs	myosin-1 (<i>Sus scrofa</i>)	gi 157279731	33	10	164
		myosin (<i>Gallus gallus</i>)	gi 13432175	23	11	220
18	In-house sausages	myosin-1 (<i>Sus scrofa</i>)	gi 157279731	44	17	274

^aNumber of matched peptides in the database search. ^bPercent of coverage of the entire amino acid sequence. ^cMASCOT score at FDR of 1%.

Table 2. Peptide markers for both species and protein identified in processed meat products; all presented peptides were ranked first in the list of matched peptides of MASCOT peptide view results

Sample	Processing method	Declared meat composition	Identified species	Protein	Peptide marker	NCBI accession number	MASCOT ion score ^a	Identity threshold ^b	Homology threshold ^c
1. Cooked ham	sliced, cured, cooked	pork 78%	pig	myosin-1	SALAHAVQSSR	gi 157279731	42	>59	>38
				myosin-2	TLAFLFSGAQTGEAEAGGTK	gi 55741490	41	>60	>57
				myosin-7	LLSNLFANYAGADTPVEK	gi 55741486	86	>64	>49
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	37	>55	>44
2. Beef spread	minced, canned	beef 66%	cattle	myosin-2	TLAFLFSGTPTGDSEASGGTK	gi 75055812	30	>62	>34
				MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	60	>62	>40
				myoglobin	HPSDFGADAQAAMSK	gi 27806939	56	>71	>67
3. Potted beef	minced, canned	beef 67%, beef heart	cattle	myosin-7	GQNVQQVVYAK	gi 41386711	28	>46	>44
				MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	88	>38	>26
4. Corned beef	cured, cooked, corned	beef	cattle	myosin-7	GQNVQQVVYAK	gi 41386711	32	>68	>50
				MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	54	>54	>22
5. Beef sausage	raw, oven cooked from chilled	beef 53%	cattle	myosin-2	TLAFLFSGTPTGDSEASGGTK	gi 75055812	51	>100	>63
				myosin-1	ALEDQLSELK	gi 41386691	38	>40	>39
				MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	87	>54	>28
				myoglobin	HPSDFGADAQAAMSK	gi 27806939	42	>60	>43
6. Chorizo style sausage	raw, oven cooked from chilled	pork 87%	pig	myosin-1	SALAHAVQSSR	gi 157279731	61	>64	>51
				myosin-7	LLSNLFANYAGADTPVEK	gi 55741486	62	>60	>33
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	98	>58	-
7. Hunters sausage	smoked, cooked, roasted	pork 70%, beef 20%	pig	myosin-4	SALAHAVQSSR	gi 178056718	60	>60	>50
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	112	>61	-
			cattle	MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	19	>35	>13
8. Horse sausage	raw, oven cooked from chilled	horse meat	horse	myosin-2	VVETMQTMLDAEIR	gi 126352598	87	>58	-
				myosin-7	MLSNLFANYLGADAPIEK	gi 126352320	83	>58	>40
				myoglobin	GLSDGEWQQVLNVWGK	gi 7546624	72	>56	>36
				myoglobin	VEADIAGHGQEVLR	gi 7546624	67	>54	-
				myoglobin	HGTVVLTALGGILK	gi 7546624	99	>59	-
9. Horse sausage	smoked, cooked, dried	no data	pig	myosin-1	SALAHAVQSSR	gi 157279731	16	>40	>24
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	63	>35	>17
			horse	myoglobin	GLSDGEWQQVLNVWGK	gi 7546624	59	>38	>22
				myoglobin	VEADIAGHGQEVLR	gi 7546624	80	>38	-
				myoglobin	HGTVVLTALGGILK	gi 7546624	83	>40	-

10. Kabanos sausage with cheese	smoked, cooked, dried	chicken meat 58%, pork 12%, cheese 7.5%	chicken	myosin	VAEQELLDATER	gi 13432175	81	>44	>43
				MLC1/3f	DQGTFFDFVEGLR	gi 212330	39	>59	>44
				MLC2f	GADPEDVIMGAFK	gi 223047	58	>60	-
			pig	myosin-4	SALAHAVQSSR	gi 178056718	43	>44	>43
			cattle	casein alpha-S1	HQGLPQEVLENLLR	gi 225632	43	>60	>36
				casein alpha-S1	EPMIGVNQELAYFYPELFR	gi 225632	48	>58	>26
11. Pork sausage	smoked, cooked	pork 92%, veal 6%	pig	myosin-4	SALAHAVQSSR	gi 178056718	64	>44	-
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	88	>45	>27
12. Cocktail sausage	cured, cooked	beef 60%, turkey meat 6%	cattle	myosin-2	TLAFLFSGTPTGDSEASGGTK	gi 75055812	70	>74	>53
				MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	62	>69	>41
				myoglobin	HPSDFGADAQAAMSK	gi 27806939	51	>55	>40
13. Frankfurters poultry	cooked, smoked	chicken & turkey MRM 65%	chicken	myosin	GQTVSQVHNSVGALAK	gi 13432175	60	>56	>46
				myosin	TLALLFATYGGEAEGGGGK	gi 13432175	15	>59	>31
				myosin	VAEQELLDATER	gi 13432175	79	>56	>52
				MLC1/3f	DQGTFFDFVEGLR	gi 212330	63	>58	>54
				MLC2f	GADPEDVIMGAFK	gi 223047	62	>60	-
				beta-enolase	AAIAQAGYTDK	gi 46048765	51	>59	>36
14. Frankfurters classics	smoked	pork 71%, milk proteins	pig	myosin-1	SALAHAVQSSR	gi 157279731	61	>63	-
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	71	>57	-
			cattle	casein alpha-S1	HQGLPQEVLENLLR	gi 225632	91	>59	>48
				casein alpha-S1	EPMIGVNQELAYFYPELFR	gi 225632	81	>59	>48
15. Frankfurters	cooked, smoked	veal 50%, pork 28%	cattle	MLC2f	EASGPINFTVFLNMFGEK	gi 115497166	73	>37	>18
			pig	myosin-1	SALAHAVQSSR	gi 157279731	31	>45	>34
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	65	>40	>22
16. Hotdogs	cooked, smoked	pork 40%, chicken 18%	pig	myosin-1	SALAHAVQSSR	gi 157279731	47	>62	>41
				chicken	myosin	VAEQELLDATER	gi 13432175	57	>58
				MLC1/3f	DQGTFFDFVEGLR	gi 212347	81	>68	-
				MLC2f	GADPEDVIMGAFK	gi 223047	90	>60	-
17. Roast turkey	sliced, cooked, roasted	turkey meat 84%	turkey	MLC1f	ALGQNPTNAEMNK	gi 326922419	79	>59	>42
				troponin C	PSMTDQQAEAR	gi 136044	42	>63	>41
18. In-house sausages	smoked, cooked	pork	pig	myosin-1	SALAHAVQSSR	gi 157279731	41	>37	-
				GAPDH	WGDAGATYVVESTGVFTTMEK	gi 329744642	63	>54	-
				beta-enolase	NYPVVSIEDPFDQDDWK	gi 113205498	53	>57	>53

^aMASCOT score at FDR of 1%. ^{b,c}Individual ion scores to indicate identity or extensive homology.

Table 3. Peptides unique to fast and slow type MHC isoforms

Species	Protein	Peptide marker
MHC type II fast		
cattle	myosin-1(2X)	ALEDQLSELK
	myosin-2(2A)	MEIDDLASNVETISK
	myosin-2(2A)	TLAFLFSGTPTGDSEASGGTK
horse	myosin-2(2A)	VVETMQTMLDAEIR
pig	myosin-4(2B)	SALAHAVQSSR
	myosin-1(2X)	
	myosin-2(2A)	TLAFLFSGAQTGEAEAGGTK
MHC type I slow/cardiac		
cattle	myosin-7	SAETEKEIALMK
	myosin-7	GQNVQQVVYAK
horse	myosin-7	GTLEDQIIEANPALEAFGNAK
	myosin-7	MLSNLFANYLGADAPIEK
pig	myosin-7	LLSNLFANYAGADTPVEK




Figure 1. Analytical workflow for fast detection of peptide markers in processed meat products using LESA-MS/MS.




Figure 2. Example of sequenced spectrum of the pork myosin-1 and myosin-4 peptide SALAHAVQSSR (563.67²⁺), fast type isoforms, obtained from frankfurters made from veal and pork (sample 15) using LESA-MS/MS.




Figure 3. Sequenced LESA-MS/MS spectra of slow type 1 myosin isoform; (A) pig myosin-7 peptide LLSNLFANYAGADTPVEK (962.10^{2+}) obtained from cooked ham (sample 1); (B) horse myosin-7 peptide MLSNLFANYLGADAPIEK (984.29^{2+}) obtained from horse sausage (sample 8) cooked in oven.